

THE NEW YORK ACADEMY OF SCIENCES

(Founded in 1817)

COUNCIL 1958

President

BORIS PRIGEL

President Elect

HILARY KOPROWSKI

Vice Presidents

M J KOPAC

C P RHOADS

Recording Secretary

EMERSON DAY

Corresponding Secretary

FREDRICK C NACHOD

Treasurer

ROBERT F LIGHT

Elected Councilors

1956-1958

DONALD B KEYES
WARREN O NITLSON

CHARLES D MARPLE
FREDERICK Y WISELOGIE

1957-1959

GEORGE H MANGUN
MINA REES

HAYDEN C NICHOLSON
WILLIAM W WALCOTT

1958-1960

DAVID A KARNOFSKY
WAYNE W UMBREIT

GUSTAV J MARTIN
JOHN F VANCE

Finance Committee

HAYDEN F TAYLOR *Chairman*

GORDON Y BILLARD

JOHN TEE VAN

Executive Director

EUNICE THOMAS MINER

SECTION OF GEOLOGY AND MINERALOGY

KURT E LOWE *Chairman*

IRFDRICH P YOUNG Jr *Secretary*

SECTION OF BIOLOGY

ALBERT S GORDON *Chairman*

LOUIS G NICKFILL *Secretary*

DIVISION OF MYCOLOGY

M L LITTMAN *Chairman*

KARL MARAMOROSCH *Secretary*

SECTION OF PSYCHOLOGY

ELAINE R GRIMM *Chairman*

WILLIAM W CUMMING *Secretary*

SECTION OF ANTHROPOLOGY

MARGARET MEAD *Chairman*

DOROTHY C JENSEN *Secretary*

SECTION OF PHYSICS AND CHEMISTRY

JOSEPH GRFENSPAN *Chairman*

LIEBE F CAVALIERI *Secretary*

SECTION OF OCEANOGRAPHY AND METEOROLOGY

ABRAHAM S KUSSMAN *Chairman*

CHARLES KNUDSEN *Secretary*

SECTION OF MATHEMATICS AND ENGINEERING

LYLE BORST *Chairman*

NICHOLAS V FEODOROFF *Secretary*

Past Presidents

WILLIAM K GREGORY

HORACE W STUNKARD

HARDEN F TAYLOR

VICTOR K LAMER

M L CROSSLEY

WALTER S ROOT

ROSS F NIGRELLI

The Sections and the Division hold meetings regularly one evening each month during the academic year October to May inclusive. All meetings are held at the building of The New York Academy of Sciences, 2 East Sixty-third Street, New York 21, New York.

Conferences are also held at irregular intervals at times announced by special programs.

ANNALS OF THE NEW YORK ACADEMY OF SCIENCES

VOLUME 70 ART. 3 PAGES 211-262

June 3, 1958

Editor in Chief

OTTO V. ST. WHITELOCK

Managing Editor

FRANKLIN V. FURNESS

Associate Editor

PETER A. STUPPSON

ANIMAL DISEASE AND HUMAN IMMUNITY*

Conference Chairman and Consulting Editor

JAMES LIEBERMAN

CONTENTS

Foreword	By JAMES LIEBERMAN	211
Introduction		
Animal Diseases: Their Relationship to the Health of Man	By J. D. MARTIN	240
Part I The Viral Encephalitides		
Introduction	By WILLIAM McDONALD	247
The Natural History of the Arthropod Borne Encephalitides in the United States	By A. D. HEAL AND JERSTO H. LIDEN	251
Vector Relationships of the Arthropod Borne Encephalitides in North America	By K. A. W. CHAMBERLAIN	31
Host Relationships of the Arthropod Borne Encephalitides	By HUBERT L. HANSON	370
Japanese Encephalitis Vaccine Studies in Pheasants Under Experimental and Field Conditions	By OSCAR S. SHAW, DANIEL COHEN, JAMES L. CRENSHAW AND ROBERT F. F. SLING	325
Part II Other Problems in Virology		
Organ Viruses of Man and Animals	By G. D. HUNG AND JOSEPH I. MELNICK	342
The Significance of Human Antiviral Neutralizing Substances in Animal Sera	By MICHAEL KLEIN	367
Counterparts of Human Viral Disease in Animals	By HILARY KOPROWSKI	369
Comparative Studies on Viral Hepatitis in Animals and in Man	By MERRILL L. LARD	383

This series of papers is the result of a conference on *Animal Disease and Human Health* held by The New York Academy of Sciences in collaboration with the Communicable Disease Center, Public Health Service, Atlanta, Ga., September 11, 12, and 13, 1957.

THE NEW YORK ACADEMY OF SCIENCES

(Founded in 1817)

COUNCIL 1958

President

BORIS PREGEL

President Elect

HILARY KOPROWSKI

Vice Presidents

M J KOPAC

C P RHODES

Recording Secretary

EMERSON DAY

Corresponding Secretary

FREDERICK C NACHOD

Treasurer

ROBERT F LIGHT

Elected Councilors

1956-1958

DONALD B KEYES
WARREN O NELSON

CHARLES D MARPLE
FREDERICK Y WISELOGLE

1957-1959

GEORGE H MANGUN
MINA PEES

HARPEN C NICHOLSON
WILLIAM W WALCOTT

1958-1960

DAVID A KARNOFSKI
WAYNE W UMBRETT

GUSTAV J MARTIN
JOHN E VANCE

Finance Committee

HARDEN F TAYLOR Chairman

GORDON V BILLARD

JOHN TEEVAN

Executive Director

EUNICE THOMAS MINER

SECTION OF GEOLOGY AND MINERALOGY

KURT E LOWE Chairman

FREDERICK P YOUNG Jr Secretary

ALBERT S GORDON Chairman

SECTION OF BIOLOGY

LOUIS G NICHOLL Secretary

M L LITTMAN Chairman

DIVISION OF MYCOLOGY

KARL MARAMOROSCH Secretary

ELAINE R CRISM Chairman

SECTION OF PSYCHOLOGY

WILLIAM W CUMMING Secretary

MARGARET MEAD Chairman

SECTION OF ANTHROPOLOGY

DOPOTHY C JENSEN Secretary

JOSEPH GREENSPAN Chairman

SECTION OF PHYSICS AND CHEMISTRY

LIEBE F CAVALIERI Secretary

ABRAHAM S KUSSMAN Chairman

SECTION OF ORNITHOLOGY AND METEOROLOGY

CHARLES KNUDSEN Secretary

LYLE BORST Chairman

SECTION OF MATHEMATICS AND ENGINEERING

NICHOLAS V FEODOROFF Secretary

Past Presidents

WILLIAM K GREGORY

HORACE W STURKARD

HARDEN F TAYLOR

VICTOR K TAMER

M L CROSSLEY

WALTER S FOOT

ROSS F NIGRELLI

The Sections and the Division hold meetings regularly, one evening, each month, during the academic year, October to May, inclusive. All meetings are held at the building of The New York Academy of Sciences, 2 East Sixty-third Street, New York 21, New York.
Conferences are also held at irregular intervals at times announced by special programs.

ANNALS OF THE NEW YORK ACADEMY OF SCIENCES

VOLUME 70 ART. 3 PAGES 211-762

June 3, 1958

Editor in Chief

OTTO V. ST. WHITLOCK

Managing Editor

FRANKLIN N. FURNESS

Associate Editor

EDITH A. STURGEON

ANIMAL DISEASE AND HUMAN HEALTH*

Conference Chairman and Consulting Editor

JAMES LIEBERMAN

CONTENTS

Foreword	By JAMES LIEBERMAN	219
Introduction		
Animal Diseases	Their Relationship to the Health of Man	By J. D. MARTIN 289
Part I The Viral Encephalitides		
Introduction	By WILLIAM McD. HAMMON	292
The Natural History of the Arthropod Borne Encephalitides in the United States	By A. D. HESS AND IRENE H. HOLDEN	294
Veterinarian Relationships of the Arthropod Borne Encephalitides in North America	By ROY W. CHAMBERLAIN	312
Host Relationship of the Arthropod Borne Encephalitides	By JOSEPH F. KISSLING	320
Equine Encephalitis Vaccine Studies in Pheasants Under Epizootic and Pre-Epizootic Conditions	By OSCAR S. MANN, DANIEL COHEN, JEAN L. GERENDE AND JOSEPH F. KISSLING	378
Part II Other Problems in Virology		
Orphan Viruses of Man and Animals	By G. D. HILL AND JOSEPH L. MELNICK	342
The Significance of Human Antiviral Neutralizing Substances in Animal Sera	By MORITZ KLEIN	362
Counterparts of Human Viral Disease in Animals	By HILARY KOWROWSKI	369
Comparative Studies on Viral Hepatitis in Animals and in Man	By MORRIS P. L. LARD	383

This series of papers is the result of a conference on *Animal Disease and Human Health* held by The New York Academy of Sciences in collaboration with the Communicable Disease Center, Public Health Service, Atlanta, Georgia, September 11, 12, and 13, 1957.

Part III Leptospirosis A Contemporary Problem

Introduction	By DONALD S. MARTIN	391
Animal Reservoirs of Leptospirae	By B. BABUDIERI	393
Leptospirosis Infections in Man	By JOE L. STOCKARD AND THEODORE F. WOODWARD	414
Laboratory Diagnosis of Leptospirosis	By WILLIAM S. GOCHENOUR JR. CHESTER A. GLEISER AND MARTHA K. WARD	421
Epidemiological Patterns of Leptospirosis	By MILDRED M. GALTON ROBERT W. MENGES AND JAMES H. STEELE	427

Part IV Recent Developments in the Epidemiology of Rabies

Introduction	By ERNEST S. TIERKEL	445
Bat Rabies Experimental Host Transmission Studies	KENNETH F. BURNS DOROTHY FRENCH SHELTON AND EARL W. GROGAN	452
Newer Tools for the Prevention of Rabies in Domestic Animals	By MERLIN L. KAEBERLE	467
Prophylaxis Against Rabies in Humans	By JOHN P. FOX	480

Part V Ornithosis

Ornithosis in Domestic Fowl Newer Findings in Turkeys	By JOHN P. DELAPLANE	495
Newer Method for Detection of Avian Ornithosis	By ALBERT A. BENEDICT AND CLARENCE MCCLARLAND	501

Part VI Other Contemporary Disease Problems

The Changing Picture of Murine Typhus in the United States	By HARRY D. LATT	516
Air Borne Transmission of Q Fever The Role of Parturition in the Generation of Infective Aerosols	By HARTWELL H. WELSH EDWIN H. TENNETTE FRANCIS R. ABINANTI AND JOHN F. WINN	528
Human Brucellosis with Special Reference to the Disease in the United States	By NORMAN B. McCULLOUGH	541
Brucellosis in Livestock Control and Eradication	By C. K. MINGLE	557
Parasites of Animals and Human Disease	By THOMAS W. M. CAMERON	564
Industrial Anthrax	By PHILIP SIGMUND BRACHMAN AND FRANCIS ROBERT FEAFY	574
Newcastle Disease	By R. P. HANSON AND C. A. BRANDLY	585
Salmonellosis Observations on Incidence and Control	By P. R. EDWARDS	598
The Exotic Zoonoses	By M. S. SHAHAN AND J. TRAUM	614
Listeriosis A Potential Public Health Problem	By JOHN I. WINN WILLIAM B. CHERRY AND ELIZABETH O. KING	624
The Public Health Importance of Animal Tuberculosis	By I. J. ANDERSON	632
Recent Developments in Animal Ringworm and Their Public Health Implications	By WILLIAM KAILAN ILLICIE K. CFORG AND LIBERO VJELLO	636
Cat Scratch Fever	By J. F. PRIER	650
New Knowledge on the Ecology of Sylvatic Plague	By IFO KARTMAN FRANK M. PRINCE STUART F. QUAN AND HAROLD I. STARR	668
Animal Disease and Biological Warfare	By JACK H. HEMPY	712

Part VII Selected Chronic Diseases

A Comparison of Rheumatoid Like Arthritis in Swine and Rheumatoid Arthritis in Man	By DENNIS SIKES	717
The Pulmonary Mononitosis Complex in Sheep	By FRANCISCO DURAN FERNALD ERWIN JUNGHER ALBERTO CUBA CAPARÓ KEEN A. RAFFERTY JR. AND CHARLES HELMBOLDT	726

Part VIII International Programs

Global Priorities in Animal Disease Control	By MARTIN M. KAILAN	743
The Pan American Zoonoses Center	By BENJAMIN D. BLOD	747

Summary

Summary	By LEONARD M. SCHULMAN	752
---------	------------------------	-----

Copyright 1958 by The New York Academy of Sciences

FORWORD

By James Lieberman

Communicable Disease Center / Public Health Service / Atlanta, Ga.

This monograph comprises a publication on comparative medicine that should prove useful to the professions of the medical sciences.

An editorial that appeared in *The New York Times* (September 12, 1957) said in part of the conference on which this monograph is based: "The relationship of animal disease to human welfare, although recognized through the ages, has not been explored as thoroughly as it might be. There is a tendency to forget that the dawn of bacteriology emerged through the pioneering efforts of such men as Louis Pasteur with his work on anthrax, a disease affecting lower animals as well as man."

The conference is a ten year progress report on the 'zoonoses,' the category of diseases of lower animals transmissible to man. Diseases such as psittacosis, brucellosis, cat scratch fever, toxoplasmosis and a host of others will be reviewed with accompanying progress reports on methods for their diagnosis and control. Other diseases not included in this category will be discussed from the standpoint of their possible relationship to similar conditions in man. For example, pulmonary adenomatosis, which has been decimating sheep flocks in many areas of the world, has several factors in common with lung cancer in humans.

Although the subject of this monograph can hardly be dealt with adequately in a publication of this size, perhaps it will be agreed that the subject matter selected illustrates the inseparability of those diseases that affect both man and the lower animals. The contributors to these pages have not been concerned solely with the acute communicable diseases, but have ventured also into the comparative pathology of several of the chronic diseases and the chronic sequelae to infectious diseases. There are special papers dealing with global activities in this field and with biological warfare. In each instance topics are discussed by persons well versed in their respective specialties. Limitations of space alone have prevented amplifying the scope of this publication.

In a very broad sense, comparative medicine and its subspecialties have been commented upon since the Biblical era. Medical and veterinary texts have devoted much time to this subject, but one of the first sessions devoted to *The Relation of Diseases in the Lower Animals to Human Welfare* was sponsored by The New York Academy of Sciences in 1946; these proceedings were published by the Academy in April 1947. It is expected that this volume will supplement rather than completely supplant the earlier publication and that it will clarify and bring up to date many concepts understood imperfectly or not at all ten years ago.

Introduction

ANIMAL DISEASES THEIR RELATIONSHIP TO THE HEALTH OF MAN

By J. D. Martin*

Section of Epidemiology, Louisiana State Department of Health, New Orleans, La.

Significance of the Relationship of Animal Health to Human Health

The health of humans is inextricably related to the health of animals. Recognition of this fact dates to antiquity, but the closeness and full significance of the relationship is just beginning to be seen in its proper perspective.

Diseases, both human and animal, gravely influence man's affairs. National and international and also personal health and welfare are dependent to a large extent upon the prevention of disease. The need of the United States to be prepared for any eventuality poses a problem for today and tomorrow, and such preparedness requires healthy citizens. This is secured by a healthful economy and, in addition to other public health measures, an adequate supply of safe, wholesome food to meet the world's peacetime and wartime needs, as well as an adequate system for its distribution. The achievement of this state of preparedness possible only when men and animals are healthy, will be attained only when people become aware of the problems confronting them and when they apply all available knowledge and skills to resolve these problems.

More than 200 diseases of animals are communicable; almost half are known to be transmissible from animal to man. There are also more than 100 different parasites that affect both man and animals.

As investigative methods improve, the number of disease entities due to hitherto unidentified infectious and noninfectious processes will increase steadily and will continue to increase as man applies himself to a study of his nature and his environment.

In addition to the direct effect of animal diseases on human health through the impact of infectious diseases transmissible to man, human health and well-being are also significantly influenced by animals through the food and raw materials derived from them, through recreation and sports associated with them, and through the opportunities they offer for research into basic problems affecting man. Animals serve man as food, pets, beasts of burden, and means of transportation. At times they also affect him as enemies intent upon his injury or destruction.

The interrelationship between animals and man is intimate, significant, and far-reaching. If man is to continue to improve his health, he must recognize this relationship, study it carefully, and apply the knowledge gained.

Historical Recognition of the Relationship

Man's knowledge of this relationship of animal disease to human health dates from antiquity. Control measures were developed even before the rec-

ognition of the infectious nature of some of the diseases of animals transmissible to man

The Bible tells us that Abel the son of the first man and woman offered of his flocks to the Lord This account, with that of Adam and Eve in the Garden of Paradise indicates that the relationship of animals to man was recognized from the earliest days of human existence

Study of the history of diseases and their manifestations strongly suggests that indubitable evidences of human disease¹ appeared almost synchronously with the first manifestations of human life on the earth

Mettler² expresses doubt that particular etiologic agents of specific diseases have ever been completely absent He quotes Moodie³ as saying that parasitism is a most antiquated mode of life and has been proved to extend as far back as the paleozoic period

In Biblical times the capture of the Ark of the Covenant by the Philistines was associated with an epidemic probably bubonic plague that killed 50 000 Bethshemites alone That true bubonic plague was recognized from very ancient times appears probable as the association of rats with disease of bubonic nature is also indicated in the cuneiform writings of Babylonia⁴

A celebrated passage in Susruta⁵ is interpreted as indicating that early Indian medicine had knowledge of the relationship between mosquitoes and malarial fever A warning to abandon a dwelling when its rats act queerly and die also suggests to Castiglioni⁶ that the ancient Persians and Indians suspected the relationship between plague and rats

Bubonic plague and louse borne typhus probably were known to the Romans also History suggests frequent epidemics of bubonic plague with thousands of persons dying and Castiglioni tells of the pestilence called the Plague of Antoninus or of Galen that lasted from 164 to 180 A D and was probably louse borne typhus although it could have been bubonic plague Historically the diagnosis is still uncertain

There is evidence too that throughout the classical Greek period numerous endemic foci of plague and malaria existed⁷

We also find in the Bible an account of a disease that in the time of Moses destroyed all the beasts horses asses oxen camels and sheep This disease could have been anthrax

Rabies was one of the first recognized zoonoses Kelser⁸ quotes Aristotle as writing in the Fourth Century B C that dogs suffer from madness which puts them in a state of fury and all animals which they bite when in this condition become also attacked with madness

Malaria too has been known from ancient times Frightful epidemics of malaria wreaked havoc in Rome in the later periods of the Roman Empire and the people associated malaria with swamps and drained them in an effort to control it

Tsutsugamushi fever or mite typhus is believed by Williams⁹ to have been recognized as an entity since the Sixth Century A D Blake and his co workers⁷ believe that it afflicted the natives of south China in the Sixteenth Century

Epidemics of louse borne typhus have probably occurred everywhere and in

Introduction

ANIMAL DISEASES THEIR RELATIONSHIP TO THE HEALTH OF MAN

By J D Martin*

Section of Epidemiology, Louisiana State Department of Health, New Orleans, La

Significance of the Relationship of Animal Health to Human Health

The health of humans is inextricably related to the health of animals. Recognition of this fact dates to antiquity, but the closeness and full significance of the relationship is just beginning to be seen in its proper perspective.

Diseases both human and animal gravely influence man's affairs. National and international and also personal health and welfare are dependent to a large extent upon the prevention of disease. The need of the United States to be prepared for any eventuality poses a problem for today and tomorrow, and such preparedness requires healthy citizens. This is secured by a healthful economy and in addition to other public health measures, an adequate supply of safe, wholesome food to meet the world's peacetime and wartime needs, as well as an adequate system for its distribution. The achievement of this state of preparedness, possible only when men and animals are healthy, will be attained only when people become aware of the problems confronting them and when they apply all available knowledge and skills to resolve these problems.

More than 200 diseases of animals are communicable; almost half are known to be transmissible from animal to man. There are also more than 100 different parasites that affect both man and animals.

As investigative methods improve, the number of disease entities due to hitherto unidentified infectious and noninfectious processes will increase steadily and will continue to increase as man applies himself to a study of his nature and his environment.

In addition to the direct effect of animal diseases on human health through the impact of infectious diseases transmissible to man, human health and well-being are also significantly influenced by animals through the food and raw materials derived from them, through recreation and sports associated with them, and through the opportunities they offer for research into basic problems affecting man. Animals serve man as food, pets, beasts of burden, and means of transportation. At times they also affect him as enemies intent upon his injury or destruction.

The interrelationship between animals and man is intimate, significant, and far reaching. If man is to continue to improve his health, he must recognize this relationship, study it carefully, and apply the knowledge gained.

Historical Recognition of the Relationship

Man's knowledge of this relationship of animal disease to human health dates from antiquity. Control measures were developed even before the rec-

Present address: State of Louisiana Department of Public Welfare, Baton Rouge, La.

ognition of the infectious nature of some of the diseases of animals transmissible to man

The Bible tells us that Abel the son of the first man and woman offered of his flocks to the Lord This account, with that of Adam and Eve in the Garden of Paradise indicates that the relationship of animals to man was recognized from the earliest days of human existence

Study of the history of diseases and their manifestations strongly suggests that indubitable evidences of human disease¹ appeared almost synchronously with the first manifestations of human life on the earth

Mettler² expresses doubt that particular etiologic agents of specific diseases have ever been completely absent He quotes Moodie³ as saying that parasitism is a most antiquated mode of life and has been proved to extend as far back as the paleozoic period

In Biblical times the capture of the Ark of the Covenant by the Philistines was associated with an epidemic probably bubonic plague that killed 50 000 Bethshemites alone That true bubonic plague was recognized from very ancient times appears probable as the association of rats with disease of bubonic nature is also indicated in the cuneiform writings of Babylonia⁴

A celebrated passage in Susruta⁵ is interpreted as indicating that early Indian medicine had knowledge of the relationship between mosquitoes and malarial fever A warning to abandon a dwelling when its rats act queerly and die also suggests to Castiglioni⁶ that the ancient Persians and Indians suspected the relationship between plague and rats

Bubonic plague and louse borne typhus probably were known to the Romans also History suggests frequent epidemics of bubonic plague with thousands of persons dying and Castiglioni tells of the pestilence called the Plague of Antoninus or of Galen that lasted from 164 to 180 A D and was probably louse borne typhus although it could have been bubonic plague Historically the diagnosis is still uncertain

There is evidence too that throughout the classical Greek period numerous endemic foci of plague and malaria existed⁷

We also find in the Bible an account of a disease that in the time of Moses destroyed all the beasts horses asses oxen camels and sheep This disease could have been anthrax

Rabies was one of the first recognized zoonoses Kelsor⁸ quotes Aristotle as writing in the Fourth Century B C that dogs suffer from madness which puts them in a state of fury and all animals which they bite when in this condition become also attacked with madness

Malaria too has been known from ancient times Frightful epidemics of malaria wreaked havoc in Rome in the later periods of the Roman Empire and the people associated malaria with swamps and drained them in an effort to control it

Tsutsugamushi fever or mite typhus is believed by Williams⁹ to have been recognized as an entity since the Sixth Century A D Blake and his co-workers⁷ believe that it afflicted the natives of south China in the Sixteenth Century

Epidemics of louse borne typhus have probably occurred everywhere and in

Introduction

ANIMAL DISEASES THEIR RELATIONSHIP TO THE HEALTH OF MAN

By J. D. Martin*

Section of Epidemiology Louisiana State Department of Health New Orleans La

Significance of the Relationship of Animal Health to Human Health

The health of humans is inextricably related to the health of animals. Recognition of this fact dates to antiquity, but the closeness and full significance of the relationship is just beginning to be seen in its proper perspective.

Diseases both human and animal gravely influence man's affairs. National and international and also personal health and welfare are dependent to a large extent upon the prevention of disease. The need of the United States to be prepared for any eventuality poses a problem for today and tomorrow, and such preparedness requires healthy citizens. This is secured by a healthful economy and, in addition to other public health measures, an adequate supply of safe, wholesome food to meet the world's peacetime and wartime needs, as well as an adequate system for its distribution. The achievement of this state of preparedness possible only when men and animals are healthy, will be attained only when people become aware of the problems confronting them and when they apply all available knowledge and skills to resolve these problems.

More than 200 diseases of animals are communicable; almost half are known to be transmissible from animal to man. There are also more than 100 different parasites that affect both man and animals.

As investigative methods improve, the number of disease entities due to hitherto unidentified infectious and noninfectious processes will increase steadily and will continue to increase as man applies himself to a study of his nature and his environment.

In addition to the direct effect of animal diseases on human health through the impact of infectious diseases transmissible to man, human health and well-being are also significantly influenced by animals through the food and raw materials derived from them, through recreation and sports associated with them, and through the opportunities they offer for research into basic problems affecting man. Animals serve man as food, pets, beasts of burden, and means of transportation. At times they also affect him as enemies intent upon his injury or destruction.

The interrelationship between animals and man is intimate, significant, and far-reaching. If man is to continue to improve his health, he must recognize this relationship, study it carefully, and apply the knowledge gained.

Historical Recognition of the Relationship

Man's knowledge of this relationship of animal disease to human health dates from antiquity. Control measures were developed even before the recognition of the relationship.

Present address: State of Louisiana Department of Public Welfare, Baton Rouge, La.

ognition of the infectious nature of some of the diseases of animals transmissible to man

The Bible tells us that Abel the son of the first man and woman offered of his flocks to the Lord This account with that of Adam and Eve in the Garden of Paradise indicates that the relationship of animals to man was recognized from the earliest days of human existence

Study of the history of diseases and their manifestations strongly suggests that indubitable evidences of human disease¹ appeared almost synchronously with the first manifestations of human life on the earth

Mettler² expresses doubt that particular etiologic agents of specific diseases have ever been completely absent He quotes Moodie³ as saying that parasitism is a most antiquated mode of life and has been proved to extend as far back as the paleozoic period

In Biblical times the capture of the Ark of the Covenant by the Philistines was associated with an epidemic probably bubonic plague that killed 50 000 Bethshemites alone That true bubonic plague was recognized from very ancient times appears probable as the association of rats with disease of bubonic nature is also indicated in the cuneiform writings of Babylonia⁴

A celebrated passage in Susruta⁵ is interpreted as indicating that early Indian medicine had knowledge of the relationship between mosquitoes and malarial fever A warning to abandon a dwelling when its rats act queerly and die also suggests to Castiglioni¹ that the ancient Persians and Indians suspected the relationship between plague and rats

Bubonic plague and louse borne typhus probably were known to the Romans also History suggests frequent epidemics of bubonic plague with thousands of persons dying and Castiglioni tells of the pestilence called the Plague of Antoninus or of Galen that lasted from 164 to 180 A D and was probably louse borne typhus although it could have been bubonic plague Historically the diagnosis is still uncertain

There is evidence too that throughout the classical Greek period numerous endemic foci of plague and malaria existed²

We also find in the Bible an account of a disease that in the time of Moses destroyed all the beasts horses asses oxen camels and sheep This disease could have been anthrax

Rabies was one of the first recognized zoonoses Kelser³ quotes Aristotle as writing in the Fourth Century B C that dogs suffer from madness which puts them in a state of fury and all animals which they bite when in this condition become also attacked with madness

Malaria too has been known from ancient times Frightful epidemics of malaria wreaked havoc in Rome in the later periods of the Roman Empire and the people associated malaria with swamps and drained them in an effort to control it

Tsutsugamushi fever or mite typhus is believed by Williams⁴ to have been recognized as an entity since the Sixth Century A D Blake and his co-workers⁵ believe that it afflicted the natives of south China in the Sixteenth Century

Epidemics of louse borne typhus have probably occurred everywhere and in

every period of history where people have lived under crowded unsanitary conditions

As urbanization developed in the medieval period the city, which was such an important factor in economic prosperity became an equally important factor in the spread of disease. The crowding of people into cities and the almost constant military conflict between these cities with the resultant movement of armies spread disease everywhere. The most serious of the diseases thus spread was bubonic plague. Zinsser⁸ quotes J. F. C. Hecker's estimate that at least 25 million people died in Europe of this disease during this period.

The development of cities also resulted in the adoption of laws to protect the health of mankind. During the Dark Ages history is silent about sanitary regulations concerning animals and animal products until 1350 A.D. when John II of France established sanitary police as a governmental instrument.⁹ Two of the functions of this regulatory force were to keep hogs out of cities and to prevent butchers and fishmongers from selling spoiled meat and fish. After 1350 A.D. step by step the sanitary regulations that presently protect man were developed.

Impact of Animal Diseases on Man

Animal diseases directly and indirectly affect man's physical, mental, emotional, social, economic and political well being. The direct impact is evidenced by the morbidity and mortality resulting from animal pathogens that infect man. The indirect results are shown by the effect of animal diseases on man's economy, his recreational activities, and his social and political aspirations.

To estimate the number of persons who annually become ill or die as a result of infections transmitted by animals is well nigh impossible. Even in the United States many of these conditions affecting man go unrecognized and unreported. In less fortunate countries where these diseases are more common reporting is usually even less reliable so that attempted estimates of morbidity or mortality would probably be greatly in error on the conservative side.

As one studies the staggering total of deaths and hours of absenteeism due to zoonoses by comparison the deaths and disabilities resulting from wars appear negligible. Accurate statistics are not available but it is estimated that 21 million persons died of influenza in the pandemic of 1918 and 1919 while 100 million more were made ill. Whether or not this pandemic can be traced to animals we shall probably never know but some persons believe that the infectious agent that caused it was the swine influenza virus. During and following the Balkan wars and World War I typhus fever appeared in epidemic form in all the countries of eastern and central Europe. It is estimated that at least 1 million persons died in these countries during and following this world war.¹⁰

In our own country yellow fever made its appearance in the late Seventeenth Century. In 1907 Carroll¹¹ estimated that yellow fever had caused 100,000 deaths in the United States since 1793.

The epidemics that decimated the Indian peoples of Mexico and helped to

destroy their civilization could have been due to a large extent to yellow fever bubonic plague or typhus fever. According to the writings of Padre Sabagun¹ a great epidemic of typhus carried off about 2 million Indians in Mexico in 1577 A.D.

Plague a disease of antiquity that has caused millions of human deaths in the years during which it has been prevalent is also a disease of the present. One of the latest epidemics of great magnitude occurred in Manchuria in 1910-1911 killing more than 50 000 persons¹² but bubonic plague smolders in many endemic foci throughout the world and if the right conditions develop it can again become epidemic or pandemic.

Animal diseases transmissible to man are still with us. Some of these are capable of causing large numbers of human deaths inestimable human suffering dislocation of large numbers of people and upsets in emotional social and economic life and of serving as a contributing factor in changes in political power.

Animals benefit man through various uses. Anything therefore that improves the health status of animals contributes clearly to the health and well being of man.

Losses in animals affect humans in several ways by the increased cost of production of domestic animals by the reduced quantity and quality of animal products available to the consumer and through diminished opportunity for recreation.

In addition to losses due to the death of animals diseases in animals also cause losses in milk in meat and egg production in hides because of poor quality in work days for draft animals in wasted feed and labor for animal care and in loss of offspring in some infectious and reproductive diseases¹³. Moreover the livestock owner undergoes considerable inconvenience and the additional cost necessary to replace the sick or dead animal.

This inconvenience and cost is not borne by the livestock owner alone however it is also borne by every man woman and child everywhere in increased taxes and increased costs of food food products and the necessities of life derived from animals.

The United States Department of Agriculture estimates the annual livestock loss from all causes in this country at more than 2 billion dollars a year¹⁴. About one half of this loss is due to parasites. Animal death losses from all causes in the United States in 1934 included more than 15 million cattle 25 million calves 4 million sheep and lambs 105 million hogs and pigs 235 million chickens 12 million turkeys and about 250 thousand horses and mules together with an unknown number of goats fur bearing animals domestic rabbits and miscellaneous poultry¹⁵. Losses in the United States each year are staggering and in many instances unnecessary. In addition to losses on the farm there are others not counted in the preceding statement such as those resulting during transportation from the farm to the market in processing and packaging plants and from the condemnation of carcasses hides edible offal and other by products in whole or part because of disease or parasites. If we add to the costs of the obvious and hidden losses the cost of controlling or preventing diseases and parasites and the cost of protecting the public from un-

wholesome meat and meat products the total may well approximate 5 billion dollars a year and this in a country such as the United States. What then must be the cost of livestock losses to so many of the nations of the world that are not as fortunate? The impact of animal disease on man's economic status is terrific.

Briefly too it is necessary to note the impact of animal diseases on man's recreational activities. Epizootics kill off many of the animals that man hunts for pleasure and so decrease his opportunity to pursue this sport. Certain of these diseases in game animals also endanger human life.

Zoonoses also prevent the full use of several areas in the world that are otherwise well suited to the production of livestock. To feed the world's growing population it is necessary that every acre of land available for livestock production be utilized. Unfortunately much of the land in question is in poor and technically undeveloped countries. Rinderpest prevents the development of the potentially great grazing areas of Africa. Perhaps the most important of the zoonoses of wild animals at this time is African trypanosomiasis which affects man not only directly in its impact on his health but because of its economic effect. Because of this disease an area of grazing land in central Africa approximately 4 million square miles in extent remains economically poor.¹⁶ This area would be wonderful range country for livestock if trypanosomiasis could be eradicated or controlled. Control of African swine disease in coastal Africa would permit hog production in Africa to become a reality and this immense continent could then contribute great quantities of meat to a world starving for animal protein.¹⁷ The problem is not confined to Africa however if mastitis could be controlled and prevented in the dairy herds of Europe and in the United States milk production could be increased appreciably. This is important as there is a direct relationship between longevity and per capita income and the percentage of animal protein in the diet (TABLE 1).¹⁸

We who live in the United States are singularly blessed. There are few other places where food supplies can be taken for granted. Perhaps one half of the world's people exist on diets that are nutritionally poor.¹⁷ Protein malnutrition is found wherever animal diseases and parasites are uncontrolled. Clarkson¹⁷ says that the average person in the world usually has less meat in a year than most citizens of the United States eat in a week. This is to a significant degree the result of uncontrolled parasitism and animal disease.

Extent of the Problem

The problem is steadily increasing in magnitude. About one half of the people of the world are on deficient diets. These people need animal proteins yet unless a concerted effort is made to bring all of the world's technical knowledge, skills, and financial resources into play to control animal diseases and parasites the present problem will be even greater in the future.

To supply an average family of four in the United States three tons of food a year are required.¹⁷ More than one half of this is meat, eggs, poultry, and dairy products. In 1956 25 per cent more of these animal foods were consumed there than in 1935. The United States has only a small percentage of the world's peoples but its farms produce one third of the world's recorded yearly

total of 80 billion pounds of red meat and one fourth of the 100 billion pounds of milk.¹⁷

The world's population is increasing steadily. These people must be fed adequately. To produce the necessary food resources the world over must be utilized to the fullest for the production of livestock. To accomplish this parasites, pestiferous insects and vectors of disease must be controlled and eradicated. Such an achievement would contribute greatly to better health and improved standards of living for all the people of the world.

All animals, domestic and wild, are infested with parasites that contribute to man's burden of disease. In some areas parasitic infestation is so great that man's struggle against these organisms is part of his struggle to survive. Control of parasites, therefore, should be one of man's goals. Fortunately it is an attainable one.

The fight against animal disease is international in scope. Sometimes the battle against disease must be fought by a nation outside its national boundaries. An example of this is the successful campaign that the United States has just waged in Mexico at a cost of more than 134 million dollars to eradicate foot and mouth disease.¹⁸ The Mexican government spent a similar large sum of money in this fight,¹⁹ but foot and mouth disease is now under control in that country. The vast meat resources of Uruguay and Argentina are also barred from the United States because of the danger of importing the same disease.¹⁸

The problem of animal disease and human health has another very serious implication. Today in this world of stress where nations align themselves against one another, the threat of biological warfare is no longer just a theory but a real possibility, and a number of the agents that cause animal disease are adaptable to biological warfare.¹⁹ It behooves us, therefore, to become increasingly aware of this danger, to become familiar with the clinical picture of these diseases in animals and man, to be ready to apply necessary control measures and, in the meantime, to work to develop faster and more accurate ways of identifying agents of disease and of producing and using immunizing agents.

Research

The advancement of medical knowledge has been and is inseparably linked with animal experimentation. Experiments on animals have led and in the future may be expected to lead to great benefits to man and to animals. Man's contest with external nature is expressed in two different ways: adaptation to or changing of a given environment.¹⁹ Research assists in this struggle.

Through the centuries man has been concerned with the development of knowledge to cure diseases. Today science is more concerned with the development of methods of preventing disease by eliminating its cause.

Great promise appears in the use of genetics as a tool for eliminating pathological abnormalities and disease. Studies with animals have shown that certain characteristics can be bred in or out of animals and that certain animals have a genetic or natural resistance to specific disease.²⁰ Such studies must be continued.

Research into the effect of environmental factors is important also in the

TABLE J
LONGEVITY AND ECONOMIC DEVELOPMENT

	A	F lat lon	P lat lon	W lat lon	C lat lon	A lat lon	P lat lon	T lat lon	I lat lon	D lat lon	B lat lon	P lat lon	L lat lon	I lat lon
India	26 7	1931	34	7021	56	9	16	166 0	23 7	34 9	91(10)	23	3	
Egypt	30 9	1971	55	2199	69	8	12	70	30 0	45 8	85(11)	25	3	
China	34 71(1)	1979	23	2201	68	5	7	12 5	76 1(2)	45 0(1)	57	35	3	
Mexico	31 2	1922-33	61	1909	59	20	34	12 5	76 1(2)	43 5	57	35	3	
Korea	37 5	1931-35	61	1904	60	15	21	12 5	76 1(2)	43 5	57	35	3	
Peru	39 0(7)	1933-35	62	2090	58	8	14	12 5	76 1(2)	43 5	57	35	3	
Chile	41 2	1939	14	2481	0	24	34	12 5	76 1(2)	43 5	57	35	3	
Brazil	43 0	1937-40	46	2552	73	26	37	12 5	76 1(2)	43 5	57	35	3	
Columbia	46 3	1939-41	6	1934	62	39	41	12 5	76 1(2)	43 5	57	35	3	
Bulgaria	46 3	1975-28	109	2831	90	18	0	12 5	76 1(2)	43 5	57	35	3	
Japan	48 3	1935-36	93	2268	6	12	18	12 5	76 1(2)	43 5	57	35	3	
Iceland	49 8	1931-32	9	2 07	2	19	26	12 5	76 1(2)	43 5	57	35	3	
Hungary	49 8	1940-31	125	2515	59	24	21	12 5	76 1(2)	43 5	57	35	3	
Greece	50 0	1974	136	2573	65	14	22	12 5	76 1(2)	43 5	57	35	3	
U.S.S.R.	50 0(8)	1926-2	138	2515	71	14	24	12 5	76 1(2)	43 5	57	35	3	
Portugal	50 7	1939-42	134	2461	74	73	31	12 5	76 1(2)	43 5	57	35	3	
Czechoslovakia	53 6	1923-37	134	2461	74	25	35	12 5	76 1(2)	43 5	57	35	3	
Italy	54 9	1930-32	140	2677	81	19	34	12 5	76 1(2)	43 5	57	35	3	
Austria	56 5	1940-33	167	2933	9	37	46	12 5	76 1(2)	43 5	57	35	3	
Ireland	57 0	1931-40	184	2550	50	37	46	12 5	76 1(2)	43 5	57	35	3	
Ireland	57 1	1975-27	248	3184	92	41	45	12 5	76 1(2)	43 5	57	35	3	
Belgium	57 7	1978-32	261	2885	77	32	42	12 5	76 1(2)	43 5	57	35	3	
France	58 8	1931-38	283	3012	87	48	44	12 5	76 1(2)	43 5	57	35	3	
Germany	61 3	1942-34	530	2987	77	34	44	12 5	76 1(2)	43 5	57	35	3	
United Kingdom	61 5	1941	468	3005	80	43	54	12 5	76 1(2)	43 5	57	35	3	
Norway	62 4	1921-31	212	3122	83	41	49	12 5	76 1(2)	43 5	57	35	3	
Switzerland	67 9	1931-41	445	3011	82	48	54	12 5	76 1(2)	43 5	57	35	3	

many of these pests. Man aggravates the problem by himself becoming the transporter of destructive pests and pathogens from farm to farm, state to state, nation to nation, and continent to continent.

Changing environmental conditions also favor the establishment of new pests and insect vectors of disease.

The movement of livestock to auction barns and agricultural fairs has augmented the spread of insect pests, favored the transmission of communicable diseases, and increased the spread of parasites.

Rodents and other wild animals and birds periodically migrate in large numbers and carry dangerous insect life and parasites with them. Wherever trade moves, rats follow, and with the rats, the diseases commonly associated with them. In addition, many other wild animals (for example, the hamster) and wild birds are moving in commercial channels to be sold as pets. Each of these is a potential source of human disease.

Van Houweling³ tells an interesting story that highlights the increasing problem caused by today's transportation systems in the spread of disease. In 1843 a New York milkman, Peter Dunn, bought a milk cow from the captain of the *Washington*, a British ship. It seemed to Dunn that he was getting the cow at a big bargain. As it turned out, however, this cow was just as expensive as the famous cow of Mrs. O'Leary, which supposedly started the fire that destroyed one third of Chicago in 1871. Dunn's cow introduced contagious pleuropneumonia into the United States and cost our livestock owners a fortune before the disease was brought under control and eradicated. In addition, this program cost the national government 5 years of work and \$1,509,100.72 before it was completed. Pertinent to my point, however, is the fact that pleuropneumonia, in spite of its contagiousness, spread slowly because at that time transportation was slow and the movement of livestock was limited. Similarly, hog cholera required 15 years to spread across the land after its identification in 1833, and tick fever moved across North Carolina at a rate of only 4 miles a year in the 1870s.¹⁷ Contrast these spreads with the speed with which, starting in California in 1952, a vesicular exanthema of hogs swept through 20 states in 3 months,¹⁷ and the problem of infectious disease control today becomes obvious.

This does not mean that livestock should not be moved from place to place, but it does suggest that only healthy livestock should be moved, and then only under sanitary conditions.

Relationship to Human Medicine

The influence that animal health exerts on human health and welfare is real. There is a grave need for physicians and other scientists to become much more appreciative of this influence so that they may be prepared to guide people into an awareness of the problem so that, through their cooperative efforts, necessary control measures may be instituted.

As mentioned earlier, there is also a need for physicians to familiarize themselves with the human manifestations of the zoonoses and the best methods of treating these diseases.

Research people and epidemiologists must assist physicians with studies of

study of health and disease. Because of their relatively short lives and their expendability, animals lend themselves readily to use for research involving generations or serial studies.

Many of the diseases to which man is heir, such as abnormalities of growth and development, metabolic and nutritional disorders, poisonings and infections, also occur in animals. Research into the nature and control of animal diseases can precede and be a pattern for similar studies in related human diseases.

As far back as 40 B. C. Celsus¹ foresaw that clinical work with animals would later find a place in the practice of human medicine. This observation was emphasized by Galen 200 years later, but progress along remedial lines was slow and halting for centuries and at the end of the Fifteenth Century, in spite of early descriptions of animal plagues, knowledge of animal diseases was still in the folklore stages.¹ Louis Pasteur, Robert Koch, Theobald Smith, Sir Alexander Fleming, and others in the late Nineteenth and early Twentieth Centuries opened the door to modern medical science. Mohler²¹ has aptly pointed out that the bacteria, viruses, parasites, and toxins that cause diseases operate in accordance with established laws, and that the knowledge gained of the causes of disease is not dependent on the nature of the academic degrees possessed by the research workers or practitioners who do the studies. Rather, we must agree that we are concerned instead with a worker's competence, integrity, and ability to apply the knowledge he develops. Many valuable contributions to human medicine have been and will continue to be made by allied disciplines, especially that of veterinary medicine.

A study of the effect of the ingestion of vegetation and meat by animals may pave the way to a better understanding of the effect of such foods on man. We must know the effect of vitamin- or mineral-deficient food or foods that have taken up an excess amount of chemicals from the soil when ingested by animals and humans. This last is particularly true in this day when so many chemical poisons are being used. Study of the effects of the ingestion of these foods, and of foods contaminated with insecticides, fungicides, pesticides, and drugs may give us a better understanding of the way in which food control should be developed.

Spread of Disease

There is a need, too, to study the potential of the spread of disease today and in the future, and to establish adequate methods to prevent or to control this threat. Today, when the most distant parts of the earth are only hours apart by air travel, it is possible for a person or animal in the incubation stage of a disease to leave a country and arrive in another country, half the world away, before he manifests clinical symptoms of the disease. This greatly complicates the problem of controlling infectious diseases.

Disease control is further complicated by insects that spread from place to place in various ways. Flies and ticks are carried on commodity shipments. Lice, mites, flies, and cattle grub larvae remain on animals during shipment. The larvae of flies are transported in straw and manure. Lice and ticks are spread by contact. Moving livestock to new ranges or feed lots further spreads

and to share in the prevention of the problems of tomorrow that will result as man either adapts to or changes his environment

Summary

This paper discusses the relationship of animal health to human health and welfare. The ways in which animal diseases affect man, the uses that man can make of animals as he studies human diseases, the reasons underlying difficulties in control and some new problems that man can expect to face are presented. Finally, it is pointed out that only the harmonious efforts of scientists and laymen can cope with the challenges to human health and welfare resulting from diseases in animals.

References

1. CATTIGLIONI A. 1947. *A History of Medicine*. 2nd ed. Knopf, New York, N.Y.
2. METTLER C. C. 1947. *History of Medicine*. Blakiston, Philadelphia, Pa.
3. MOODIE R. L. 1923. *Pathology of Animals*. Ill. Quoted by C. C. Mettler 1947. *History of Medicine*. Blakiston, Philadelphia, Pa.
4. SURTA SAMHITA. Latin translation by Haesler 1844. English translation by A. I. R. Hoernle, Calcutta 1927 and by B. M. Kinyalal, Calcutta 1907-1916. Quoted by A. Castiglioni 1947. *A History of Medicine*. 2nd ed. Knopf, New York, N.Y.
5. KELLER R. A. 1947. *Kalies' Diseases Transmitted from Animals to Man*. T. G. Hull, Filadelfia 1921. Thomas, Springfield, Ill.
6. WILLIAMS K. W. 1944. A check list of the mite vectors and animal reservoirs of tsutsugamushi disease. *Am J Trop Med* 24: 355-357.
7. BLANCHET C. K. F. MACY J. F. SADLER JR. G. M. KOHLS & J. J. BELL. 1945. Studies on tsutsugamushi disease (scrub typhus, mite borne typhus) in New Guinea and adjacent islands: epidemiology, clinical observations and etiology in the Lae area. *Am J Hyg* 41: 243-372.
8. ZINSSER H. 1935. *Rats: Their History*. Little, Brown & Co., Boston, Mass.
9. FREEDMAN B. 1957. *Saltarians Handbook—Theory and Administrative Practice*. I. Cress, New Orleans, La.
10. VALZAH R. VAN. 1938. Diseases due to rickettsia. *In Internal Medicine*. Chap. IV. J. H. Musser, Ed. Lea and Febiger, Philadelphia, Pa.
11. CARROLL J. 1901. Yellow fever. *In Modern Medicine*. Sir W. Osler, Ed. 2: 736-759. Lea Brothers, Philadelphia, Pa.
12. WRITINGS OF PADRE SABATON. Quoted by James Tabery. 1930. *In Raters of Plagues*. Scribner's Sons, New York, N.Y.
13. REIMANN H. A. 1938. The bacillary diseases. *In Internal Medicine*. J. H. Musser, Ed. Lea and Febiger, Philadelphia, Pa.
14. LOSSES IN AGRICULTURE. 1954. A preliminary appraisal for review. *Agr. Research Service U.S. Dept. Agr.* 2: 127.
15. NORDQUIST A. V. & C. H. PAL. 1956. Economic losses from animal diseases and parasites. *Yearbook Agr.* 1956. U.S. Dept. Agr. 11: 14.
16. STEELE J. H. 1952. Animal diseases of public health significance. *Ann Internal Med* 36(2): 511-524.
17. CLARKSON M. R. 1956. Food supplies and animal diseases. *Yearbook Agr.* 1956. U.S. Dept. Agr. 7: 10.
18. FREEDMAN B. 1951. Health and longevity. *New Orleans Med. Surg. J.* 104(3): 115.
19. NARR K. J. 1956. Family food producing populations. *In Man's Role in Changing the Face of the Earth*. Thomas et al., Univ. Chicago Press, Chicago, Ill.
20. WATER M. F. 1956. Genetics and disease. *Yearbook Agr.* 1956. U.S. Dept. Agr. 46: 54.
21. MOHLER J. R. 1941. The relation of diseases in lower animals to human welfare. *U. Ia Bicentennial Conf. Univ. Ia Press*. Philadelphia, Pa.
22. SCHÖENING H. W. B. SCHWARTZ & A. W. LINDQUIST. 1956. How diseases and parasites are spread. *Yearbook Agr.* 1956. U.S. Dept. Agr. 40: 45.
23. VAN HEUWELING C. D. 1956. Our battle against animal diseases. *Yearbook Agr.* 1956. U.S. Dept. Agr. 1: 7.

the epidemiology of disease with experiments to develop better and faster ways of identifying infectious agents and disease processes and with better methods of prevention treatment and rehabilitation

Problems Still Facing Man

Obviously a discussion of all the problems that must be studied is not possible but there are several areas of research and control that should be mentioned

Before going on to new programs we must make certain that the gains that have been made in disease control are maintained and extended

Every effort should be made to free our hemisphere of foot and mouth disease Eradication of rabies and brucellosis is a goal that can be achieved Therefore a program to extirpate rabies should be developed and the program to eradicate brucellosis should be accelerated

The control or eradication of the infectious hepatitides of man and animals of leptospirosis trypanosomiasis swine cholera mastitis and rinderpest the control of occupational diseases of animal origin elucidation of the epidemiology of the encephalitides and the control of arthropod borne diseases are some of the challenges facing scientists

As business and industry continue to decentralize their operations the problem of providing safe wholesome food from animal sources will become increasingly difficult because of the difficulties of supervision

Another problem confronting us that must be studied results from an advance in physical science the development of atomic energy The use of atomic energy poses special problems for biological scientists who must now consider and plan for the impact of this force with its radioactivity on human and animal health and food products particularly those of animal origin and for the control of waste products in proximity to humans and animals

Conclusions

The health of animals and the health and well being of humans cannot be separated The animal kingdom has influenced civilization through the ages and will continue to do so Since food and other products of animal origin are among man's basic needs animal diseases are a constant threat to the health and welfare of mankind therefore they must be controlled or eradicated as quickly as possible Once high standards of animal health are achieved they must be maintained as new efforts are exerted to extend and improve them

The control of disease in animals and man requires the integrated efforts of scientists of all disciplines working in harmony with the public health services the animal health agencies and the public information service

To assure an adequate control of each individual disease procedures should be developed to assure proper identification of the disease and the reporting of its occurrence to a central agency Studies should be developed to identify the weak links in the disease chain so that these may be broken and the disease eradicated All groups interested in animal health and in human health and welfare must join together in sharing attitudes knowledge information skills methods and plans to eradicate the great animal plagues that affect man

and to share in the prevention of the problems of tomorrow that will result as man either adapts to or changes his environment

Summary

This paper discusses the relationship of animal health to human health and welfare. The ways in which animal diseases affect man, the uses that man can make of animals as he studies human diseases, the reasons underlying difficulties in control and some new problems that man can expect to face are presented. Finally, it is pointed out that only the harmonious efforts of scientists and laymen can cope with the challenges to human health and welfare resulting from diseases in animals.

References

- 1 CASTIGLIONI A. 1947. A History of Medicine. 2nd ed. Knopf, New York, N. Y.
- 2 METTLER C. C. 1947. History of Medicine. Blakiston, Philadelphia, Pa.
- 3 MOODIE K. I. 1923. Pathology. Urdana Ill. Quoted by C. C. Mettler 1947. History of Medicine. Blakiston, Philadelphia, Pa.
- 4 SURTA SAMHITA. Latin translation by Haesler 1844. English translation by A. F. R. Hoernle. Calcutta 1927 and by B. M. Kinnaird. Calcutta 1907-1916. Quoted by A. Castiglioni 1947. A History of Medicine. 2nd ed. Knopf, New York, N. Y.
- 5 KILSEY R. A. 1947. Tables. In Diseases Transmitted from Animals to Man. T. G. Hull. Ed. 3rd ed. 149-211. Thomas, Springfield, Ill.
- 6 WILLIAMS R. W. 1944. A check list of the mite vectors and animal reservoirs of tsutsugamush disease. Am. J. Trop. Med. 24: 355-357.
- 7 BLAKE F. C. & I. MACY J. I. SALTUSK JR. C. M. KOHLS & I. J. BELL. 1945. Studies on tsutsugamush disease (scrub typhus mite borne typhus) in New Guinea and adjacent islands: epidemiology, clinical observations and etiology in Dofalura area. Am. J. Hyg. 41: 243-372.
- 8 ZINSSER H. 1935. Rats, Lice and History. Little Brown & Co. Boston, Mass.
- 9 IRFEDMAN B. 1957. Sanitarians Handbook—Theory and Administrative Practice. Peckless, New Orleans, La.
- 10 VALFARI R. VAN. 1938. Diseases due to rickettsia. In Internal Medicine. Chap. IV. J. H. Mueller, F. J. Lea and Feliger. Philadelphia, Pa.
- 11 CARROLL J. 1907. Yellow fever. In Modern Medicine. Sir W. Osler. Ed. 2: 736-759. Lea Brothers, Philadelphia, Pa.
- 12 WRITINGS OF LADRI SABAGUN. Quoted by James Tabery. 1930. In Riders of Plagues. Scribner's Sons, New York, N. Y.
- 13 REIMANN H. A. 1938. The lacillary diseases. In Internal Medicine. J. H. Mueller, F. J. Lea and Feliger. Philadelphia, Pa.
- 14 LOSSES IN AGRICULTURE. 1954. A preliminary appraisal for review. Agr. Research Service. U. S. Dept. Agr. 2: 127.
- 15 NORDQUIST A. V. & C. H. PAIS. 1956. Economic losses from animal diseases and parasites. Yearbook Agr. 1956. U. S. Dept. Agr. 11-14.
- 16 STEELE J. H. 1952. Animal diseases of public health significance. Ann. Internal Med. 36(2): 511-574.
- 17 CLARKSON M. R. 1956. Food supplies and animal diseases. Yearbook Agr. 1956. U. S. Dept. Agr. 7-10.
- 18 FREEDMAN B. 1951. Health and longevity. New Orleans Med. Surg. J. 104(3): 115.
- 19 NARR K. J. 1956. Early food producing populations. In Man's Role in Changing the Face of the Earth. Thomas et al. Univ. Chicago Press, Chicago, Ill.
- 20 WATERS N. I. 1956. Genetics and disease. Yearbook Agr. 1956. U. S. Dept. Agr. 46-54.
- 21 MOHLER J. R. 1941. The relation of disease in livestock to human welfare. Univ. Pa. Biennial Conf. Univ. Pa. Press, Philadelphia, Pa.
- 22 SCHOENING H. W. B. SCHWARTZ & A. W. LINDQUIST. 1956. How diseases and parasites are spread. Yearbook Agr. 1956. U. S. Dept. Agr. 40-45.
- 23 VAN HOLWELING C. D. 1956. Our battle against animal diseases. Yearbook Agr. 1956. U. S. Dept. Agr. 1-7.

Part I The Viral Encephalitides

INTRODUCTION

By William McD Hammon

*Department of Epidemiology and Microbiology Graduate School of Public Health
University of Pittsburgh Pittsburgh Pa*

The topic of this portion of this monograph namely, viral encephalitides requires some qualifying comment that should serve as a brief orienting introduction As one who has worked a long time in this field the provision of such a commentary is my prerogative and possibly the one useful contribution that I can make at this point

Viral encephalitides is altogether too broad a term for what is to follow It could include such varied maladies as rabies mumps herpes lymphocytic choriomeningitis and others Perhaps a more inclusive term would be 'arthropod borne virus encephalitides' This term was introduced by W C Reeves and myself in 1943^{1, 2} to take the place of a miscellany of names that seemed inadequate in view of the knowledge then available

This name now in common use for fourteen years has I regret to say outlived its usefulness Newer knowledge has indicated that dengue yellow fever Colorado tick fever West Nile fever and Sindbis fever are diseases not ordinarily associated with encephalitis in any recognized natural host and are closely related with the former group either antigenically or by other valid criteria The agents in this broader group are now quite commonly called the arthropod borne viruses omitting the term encephalitis and abbreviated as arbor viruses This appears at first to be a reasonable step but many of us have objections to the shorter term and particularly to its abbreviation Use of the term arthropod borne viruses is not restricted to viruses capable of producing human disease and including swamp fever of horses and many other maladies I feel that the abbreviation is most unfortunate and I hope that its use will not spread The objection is due to the fact that the abbreviation happens to form a Latin word and in international usage will certainly be misconstrued by many to have significance as a tree rather than as an abbreviation for arthropod borne Furthermore a subgroup of which I am chairman of the Virus Subcommittee of the International Nomenclature Committee* is now searching for a suitable short pseudogeneric name such as poliovirus and myxovirus to apply to this group and the concurrent use of arbor virus again would be confusing The new officially accepted name may well turn out to be an abbreviation of arthropod and possibly zoonoses One of the common features of this group of viruses is that they are not only arthropod borne but are dependent for survival in nature upon a viremic vertebrate host generally one lower than man with man usually serving only as an accidental host and not as the true reservoir Thus they are viruses of

The recommendations of this subcommittee appear in the *International Bulletin of Bacterial Nomenclature and Taxonomy* published by the Iowa State College Press Ames Iowa

arthropod borne zoonoses not tree viruses or plain arthropod borne viruses. Since they are for the most part if not entirely zoonoses (for example we are not certain regarding the endemic basic cycle of *Phlebotomus* fever and a few others but this may be parallel to the situation of yellow fever before jungle yellow fever was discovered) I am now inclined to suggest the epidemiological group name of arthropod borne zoonoses for as many as are known to qualify and to wait for the nomenclature committee to agree in the near future on a pseudogeneric short name probably with a Greek or Latin derivation.

I feel it timely to point out that this monograph particularly this section of it deals principally with a few of our own domestic viruses that belong in this group of arthropod borne zoonoses. These are only a very few of those known the world over a group in the process of antigenic classification by the hemagglutination technique that Jordi Casals and his colleagues at the Rockefeller Foundation virus laboratories New York N. Y. are now pursuing with results of great promise.³

In the pages that follow are presented reviews of previous work and summaries of certain current work in this field by well selected investigators from the staffs of the Communicable Disease Center of the Public Health Service at several different installations and by the New Jersey State Health Department Trenton N. J.

References

- 1 HAMMON W. McD. 1943 J Am Med Assoc 121 560-564
- 2 HAMMON W. McD. W. C. REEVES & M. CRAY. 1943 Am J Public Health 33 201-207
- 3 CASALS J. & L. V. BROWN. 1954 J Exptl Med 99 439-449

THE NATURAL HISTORY OF THE ARTHROPOD BORNI ENCEPHALITIDES IN THE UNITED STATES

By A. D. Hess and Preston Holden

Communicable Disease Center, Bureau of State Services, Public Health Service, Atlanta, Georgia

Over a decade ago Bates (1946) pointed out the importance of the natural history viewpoint in investigating the mechanisms of maintenance and dispersal of a disease as compared with the sometimes more restricted concepts of epidemiology and ecology. This is particularly true for diseases such as plague, yellow fever and the encephalitides that have arthropod vectors and wild animal hosts as reservoirs. These diseases have enzootic cycles by which they are perpetuated in nature independently of their transmission to man. Determination of the basic factors responsible for their perpetuation requires coordinated field and laboratory investigations by individuals trained in many disciplines including epidemiology, biology, ecology and immunology.

In any field involving such a variety of disciplines it is inevitable that there will be differences of opinion with regard to terminology. In order to avoid confusion the following definitions are given for some of the principal terms used in this paper.

Host—an animal (vertebrate or invertebrate) that is susceptible to infection with the disease organism (virus).

Inapparent host—an individual in which the virus produces no clinical manifestations of disease.

Apparent host—an individual in which the virus produces a clinical syndrome.

Reservoir—a host that serves as a source of infection for other hosts including vectors.

Dead end host—a host that does not serve as a source of infection for other hosts or vectors.

Vector—an arthropod or vertebrate that transfers infection from one host to another.

Sylvan vector—a vector that transfers infection from wild host to wild host (synonymous with enzootic vector).

Endemic or epidemic vector—a vector that transfers infection to man and/or domestic animals.

Infection chain—the course of transmission of the virus from host to host (in temperate climates there are both summer and winter infection chains).

Transmission cycle—a cyclical infection chain in which transmission repeatedly follows the same type of course.

Basic transmission cycle—the infection chain through which the virus is normally perpetuated in nature.

Primary vectors or primary reservoirs—those that are essential elements in the basic transmission cycle or endemic chains of infection.

Secondary vectors or secondary reservoirs—those which are not essential elements in chains of infection but which may sometimes be involved particularly during epidemics or epizootics.

The authors are stationed in Logan, Utah and Greeley, Colorado respectively.

Natural history of encephalitis the interrelationships of virus vectors, and hosts that are responsible for perpetuation of virus in nature and for its spread to man and domestic animals

Encephalitis (plural encephalitides) inflammation of the brain In the present paper the term encephalitis is used to refer specifically to arthropod borne viral encephalitis

In the United States there are three principal arthropod borne viruses that cause encephalitis in humans those of western equine (WEF) eastern equine (EEE) and St. Louis (SLE) encephalitides As indicated by their names WEF and EEE produce clinical disease in horses as well as in man whereas SLE infections in horses are inapparent EEE also produces a clinical syndrome in birds and is a serious problem in pheasant farms in some of the eastern states

In recent years our knowledge of the distribution of these three viruses has been expanded considerably Prior to 1940 delineation of the geographical area of endemicity for each of these diseases was based largely upon the distribution of cases in man or in horses WEF was considered endemic only in the central plains and the Pacific coastal regions EEE was believed confined to the Atlantic coastal states Epidemics of SLE in man had been known only in the states along the upper Mississippi and lower Ohio valleys Since 1940 however our concept of the respective distribution of each of the three agents in nature has been considerably altered because of the isolation of virus from mosquitoes the detection of inapparent infection and the occurrence of clinical cases (FIGURE 1) We now know that although WEF occurs primarily west of the Mississippi River it is also present in some states along the Atlantic and Gulf coasts as evidenced by antibodies in vertebrate hosts and recoveries of virus from wild birds and mosquitoes (Holden 1955a Alexander and Murray 1957 Kissling *et al.* 1955) SLE occurs in conjunction with WEF in the states west of the Mississippi River but it also has an important extension into the central states EEE occurs in the Atlantic and Gulf coast states from New Hampshire to Texas and as far inland as Wisconsin

It is believed highly improbable that these three diseases have spread to new areas during recent years Much more tenable is the belief that the infections have existed in the same general geographical areas since antiquity This does not mean that there have not been local extensions into areas in which the viruses were previously not present as for example in the millions of acres of formerly arid land in the West that have been brought under irrigation with resultant increases in the populations of bird reservoirs and mosquito vectors of encephalitis It is quite probable that further research will reveal an even wider geographical distribution for each of the three encephalitis viruses

There are certain basic similarities in the natural histories of the three types of encephalitis It is generally believed that birds may serve as natural hosts for all three and that various species of mosquitoes are vectors Humans and horses appear to be dead end hosts or at most secondary hosts for all three viruses Beyond these similarities there are important specific differences




FIGURE 1 The geographical distribution of the arthropod-borne encephalites in the United States. Total virus activity in man and in animals.

in the natural histories of the three diseases and the details of these differences are the topic of this presentation.

Western Equine Encephalitis

Summer infection chains. The summer infection chains for WEE are reasonably well established (FIGURE 2). There seems to be little question that




FIGURE 2 Summer infection chains for western equine encephalitis. The chains for rural St. Louis encephalitis are similar except that horses are inapparent rather than apparent hosts.

Culex tarsalis is the primary vector in the sylvan transmission cycle as well as in the endemic chain of infection in the western United States. It is the one species in which population densities, virus infection and vector ability, blood feeding habits and consistent association with epidemics are adequate to account for WEE encephalitis in humans and horses (Hammon and Reeves 1945; Reeves 1953; Eklund 1954; Cockburn, Sooter and Langmuir 1957). Furthermore, the distribution of this species in North America closely coincides with the general area in which outbreaks of WEE have been known to occur (Jenkins 1950). WEE virus has been isolated many times in nature from *C. tarsalis*. Isolations have also been made from *C. pipiens*, *C. quinquefasciatus*, *C. restuans*, *C. stigmatosoma*, *Aedes melanomm* (*dorsalis*), *A. nigromaculis*, *A. infirmatus*, *A. vexans*, *Inopheles freeborni*, *Culiseta melanura* and *C. inornata* as well as several species of bird mites and an assassin bug, *Triatoma sanguisuga* (a general review of published records of isolations was made by Ferguson 1954; some later isolations were reported by Blackmore and Winn 1954 and Burroughs and Burroughs 1954; Reeves 1953; Kissling *et al.* 1955). Species other than *C. tarsalis* are, however, considered to be of secondary importance. Some of them may, however, be essential in the transmission of WEE in the eastern and southeastern states since this is out of the range of *C. tarsalis*. *C. pipiens* and *C. quinquefasciatus* have been shown experimentally to be poor vectors for WEE (Hammon and Reeves 1943; Chamberlain *et al.* 1954). The primary vectors of WEE in the East are therefore yet to be determined.

Although it seems quite clear that birds are the important natural hosts of WEE (Hammon and Reeves 1945; Reeves 1955; Eklund 1954), it has not yet been determined which species are primary reservoirs and which are of secondary importance. Published and unpublished records indicate that WEE virus has been recovered from about 20 species of birds and 6 species of mammals and that WEE antibodies have been found in more than 75




FIGURE 1 The geographical distribution of the arthropod-borne encephalites in the United States. Total virus activity in man and in animals.

in the natural histories of the three diseases and the details of these differences are the topic of this presentation.

Western Equine Encephalitis

Summer infection chains. The summer infection chains for WEE are reasonably well established (FIGURE 2). There seems to be little question that

increasingly to the importance of nestlings in natural chains of infection of the encephalitides

One possible explanation of the usual *tarsalis* bird feeding association is this many birds nest and roost in trees and as foliage gives off carbon dioxide at night when photosynthesis has ceased *tarsalis* which is attracted by carbon dioxide (Reeves 1951 Reeves 1953a) is thus brought into the sphere of attraction of roosting or nesting birds when it is seeking a blood meal. Additional studies are under way at the Logan Field Station of the Communicable Disease Center (CDC) to evaluate the importance of carbon dioxide in the orientation of blood seeking *tarsalis*. In atypical situations it would not be at all surprising to find mammals serving as the natural hosts for *tarsalis*. For example Philip Bell and Larson (1955) found that almost 14 per cent of the black tailed jack rabbits in an arid area of northern Nevada had WEE antibodies thus suggesting a high degree of feeding by *tarsalis*.

The longevity of the mosquito vectors and the extent of repeated blood feeding are important factors with regard to summer transmission cycles of encephalitis. Obviously the mosquito must take an infective blood meal survive through the extrinsic incubation period and then take a second blood meal if it is to transfer infection from one vertebrate host to another. In observations with *Iedes* mosquitoes Chamberlain Corristan and Sikes (1954a) found that optimal transmission of WEE virus occurred in as short a period as 5 days and that high transmissibility was maintained for about a month. It is difficult to obtain longevity data on mosquitoes under natural conditions but from the general information available it appears that a large percentage of summer populations of *C. tarsalis* probably survives long enough to effect at least one transmission and under favorable conditions adults might survive long enough for several transmissions. Current observations at the Bakersfield Cal Logan Utah and Greeley Colo field stations of CDC indicate that higher humidity in the adult mosquito's microhabitat greatly favors increased longevity of both summer and winter populations of *C. tarsalis*. Thus the vast irrigation developments in the western states may not only result in greatly increased production of *C. tarsalis* but may also create conditions favoring increased mosquito longevity and therefore greater likelihood for the build up of epidemics of WEE encephalitis.

As previously indicated the natural vertebrate reservoirs of encephalitis in a particular area must develop sufficient viremia for infection of the mosquitoes that transmit virus from host to host. It has been shown in certain situations that *tarsalis* may feed as much on the larger mammalian dead-end hosts such as horses cattle and sheep as on birds (Reeves and Hammon 1944). Where there is an abundance of these dead-end hosts in relation to avian hosts they may deflect blood seeking vectors from the human population (zoophylaxis) decrease the extent of feeding upon infected avian reservoirs and drain off virus that otherwise would reinforce the sylvan avian cycles of encephalitis transmission. Thus the chances of human infection would be decreased both by the lower infection rates in mosquitoes and by their lower attack rates upon humans. Potentialities for this type of zoophylaxis exist in a number of

species of wild birds and a half dozen species of wild mammals as well as most of the common domestic birds and mammals. The problem then is to determine which of this bewildering array of species are essential reservoirs for the enzootic perpetuation of WEE under natural conditions and whether others may form necessary links in endemic chains of infection.

R. W. Chamberlain, R. E. Kissling, and their co-workers have done some highly significant research on encephalitis virus vector and virus host relationships under controlled laboratory conditions as described elsewhere in this monograph. Such studies are invaluable in establishing potential vector-host combinations that may be important for the perpetuation of encephalitis viruses in nature. However, the mere fact that a particular vector has a sufficiently low infection threshold to become infected by biting a host that develops a certain level of viremia does not mean that this vector-host combination is important in nature. An analogous situation exists with regard to malaria in that most *Anopheles* mosquitoes can develop and transmit malaria plasmodia in laboratory tests; however, relatively few *Anopheles* species actually are vectors under natural conditions. In addition to having the required vectoring and reservoiring potentials, the primary vectors and reservoirs of encephalitis must be appropriately associated in both time and space.

Recent observations indicate that *Culex tarsalis* exhibits a peak of biting activity at dusk, whereas the *Aedes* mosquitoes in the same area reach biting peaks earlier in the evening (Beadle 1955). It may be epidemiologically significant that the fiercely biting *Aedes* cause many people to seek shelter within their screened dwellings before they are exposed to the principal biting attack of *C. tarsalis*, a primary vector of encephalitis. Since *C. tarsalis* has its peak biting activity at dusk, this should be a critical period in the transmission of WEE virus, and the primary reservoirs of WEE are probably among the hosts that are available to *C. tarsalis* at this time of day. It is a common observation of field workers that the evening flights of birds to their nocturnal roosting sites take place a short while prior to this peak of the biting activity of *tarsalis*.

Precipitin and other tests with field-collected mosquitoes usually indicate that *tarsalis* feeds largely on birds, but there is ample evidence that it also feeds on a variety of mammalian and even reptilian hosts (Reeves and Hammon 1944; Dow, Reeves, and Bellamy 1957). Our general observations, including unpublished studies by R. P. Dow, indicate that *tarsalis* does not necessarily prefer to feed upon birds, but that it will feed on mammals as readily as on birds if both are equally available. It also appears that the inherent behavior characteristics of the available hosts, particularly their degree of quiescence, are probably more important than the particular species that are at hand. For example, unpublished observations by J. S. Blackmore and R. P. Dow show that engorgement rates for *C. tarsalis* were much higher on relatively inactive nestlings than on the more active adult birds. The poorer venture of the nestlings may also have been a factor in the higher feeding rates. It is generally recognized that embryos and very young animals are more susceptible to infection with encephalitis viruses. Thus, cumulative evidence points

1955 Reeves *et al*, 1955) and it is generally agreed that mites do not play any significant role in the perpetuation of WLF encephalitis virus in nature

(3) There is much evidence to support the possibility that WFL virus may pass the winter in mosquitoes. W. C. Reeves and his co-workers (in press) have isolated WFL virus from naturally infected *Culex tarsalis* in Kern County Calif. during all months of the year except December and Blackmore and Winn (1956) isolated the virus from *C. tarsalis* collected in Colorado during December. Bellamy, Reeves and Scrivani (in press) recovered WFL virus from experimentally infected *C. tarsalis* after the mosquitoes had been kept for 113 days in an unheated cellar during the winter. *C. tarsalis* completed extrinsic incubation and transmitted WEL virus to chickens after 97 and 109 days of winter incubation.

Two aspects of the biology of *C. tarsalis* are of particular importance in relation to its potential role as an overwintering reservoir for WEL virus: namely its overwintering habits and its prehibernation feeding habits. *C. tarsalis* may be found during winter months in abandoned mines and cellars (Keener 1952). There is indication, however, that the temperatures in most such shelters are too high for successful hibernation and the normal winter habitat for this mosquito in the more northern latitudes has yet to be discovered (Dow, Mail and Richards 1956).

If *Culex tarsalis* is to serve as an overwintering reservoir for encephalitis viruses it must have an infective blood meal before going into hibernation. There is, however, strong evidence of a prehibernation decline in blood feeding by *tarsalis* with a possible switch to carbohydrate sources of food (Mail and Smith in press). There is even a possibility that a prehibernation blood meal may be detrimental to overwinter survival of the mosquito (Hammon, Reeves and Calindo 1945). These relationships of prehibernation feeding to overwintering of *C. tarsalis* obviously would decrease its potentialities as an overwinter reservoir for virus. Investigations to obtain additional information on these relationships are now under way at the Creeley Logan and Bakersfield field stations of CDC.

(4) There is also strong experimental evidence to support the possibility that WEE virus may pass the winter in resident avian hosts. W. C. Reeves, R. E. Bellamy and G. A. Hutson have conducted extensive investigations in Kern County Calif. that show that WEE virus can be recovered from experimentally infected birds after extended periods that are considerably greater than the normal hibernation period for the mosquito vectors. The results of these studies are now being summarized for publication.

Some virus diseases such as those of psittacosis and herpes simplex have extended periods of latency. Observations on some of the encephalitis viruses also support the possibility that there may be long periods during which the agent remains in the host in a latent or avirulent condition. For example, Corrigan, LaMotte and Smith (1956) showed that bats infected with Venezuelan equine encephalitis and stored at hibernating temperatures (10°C) maintained a low order viremia for at least 90 days and that the viremia level rose rapidly when the animals were returned to room temperatures. This suggests that latent virus of other encephalitides might be reactivated in verte

irrigated areas in the West, where dairying and the raising of cows and other domestic mammals constitute the predominant agricultural industries. To our knowledge however there has never been any attempt to correlate mosquito and bird infection rates in these areas with population densities of cattle or other dead end hosts of encephalitis.

The primary reservoirs in the sylvan cycle of WFL encephalitis may not necessarily be the same as those involved in the chain of transmission to man and domestic animals. For example in a particular situation sylvan WEL encephalitis might be perpetuated continuously by an association of *C. tarsalis* with blackbirds without any involvement of human infections. The addition to this complex of a more domestic avian host such as the English sparrow however, might result in a build up of infection rates in populations of *tarsalis* associated with man and thus lead to human infection.

With a given mosquito infection rate the chances for human infection will be correlated directly with the vector's biting or attack rates on man. These attack rates are obviously related in turn to mosquito population densities. Recent studies with caged birds indicate that the percentage of *tarsalis* feeding on avian hosts decreases as the number of mosquitoes increases (Dow, Reeves and Bellamy 1957). This would suggest that at times of unusually high mosquito prevalence there may be a much greater deflection of *tarsalis* from avian hosts to mammals including man and domestic animals. Similarly, increased human attack rates might result from a decrease in population of avian hosts despite the fact that the mosquito population remained constant.

From the above discussions it is apparent that whereas the identity of the primary vector of WEL seems to be reasonably well established much more information is needed on the particular avian species that serve as primary summer reservoirs and on the critical ecological relationships between hosts and vectors. Such information is needed if we are to understand the causative mechanisms of WEE epidemics.

Winter infection chains of WFL. In temperate North America where vector populations are inactive during the winter months it is obvious that there must be some reservoiring mechanisms that enable encephalitis viruses in the absence of active transmission to survive this period. Five possibilities have received investigative attention: (1) transovarian passage, (2) overwintering in infected mites, (3) overwintering in infected mosquitoes, (4) overwintering in infected birds, and (5) reintroduction each spring by migrating birds. The potentialities for each of these mechanisms with regard to WEE will be discussed separately.

(1) Transovarian passage in ticks is recognized as a mechanism for the perpetuation of louping ill and of Russian spring summer encephalitis. Since the primary vectors of WEE appear to be *Culex* mosquitoes that overwinter as adult females the possibility of transovarian passage as an overwintering mechanism for this particular virus seems to be eliminated.

(2) WEE virus has been isolated from the mites of wild birds (Keeves *et al* 1947; Miles *et al* 1951) and for a time it was believed that they might constitute the sylvan vector as well as a long term reservoir. Recent laboratory investigations however have given negative results (Chamberlain and Sikes




FIGURE 3 Summer infection chains for urban St. Louis encephalitis. The chains for rural SLE are similar to those for WEE (FIGURE 2).

1954 Reeves 1953) but most isolations have been from *C. tarsalis* or the *C. pipiens* complex which appear to be the primary rural and urban vectors respectively.

As in the case of WEE, present information is insufficient for determination of the particular species that are primary reservoirs of SLE. Until recently SLE virus had not been isolated from any wild or domestic birds (Eklund 1954) but has now been recovered from several species (Kanzenhofner *et al* 1957). SLE neutralizing antibodies have been found in most of the common domestic birds and mammals as well as in about 55 wild birds and a half dozen wild mammals. The problem of determining which of these many species are primary reservoirs is therefore similar to that for WEE. For SLE however the problem is further complicated by the possibility that the primary reservoirs for rural and urban SLE may be different. Birds may serve as natural reservoirs for urban SLE but it is known that they develop relatively low viremia levels with this virus (Hammon Reeves and Sather 1951 Chamberlain *et al* 1957). These levels are sufficient to infect the more susceptible mosquito species such as *C. tarsalis* but the extent to which the less susceptible *C. pipiens* complex may become infected by feeding on birds has not been fully determined. The possibility that mammals may be involved as reservoirs of urban SLE cannot be overlooked. It is known that other arthropod borne encephalitis viruses may have mammalian reservoirs. For example Venezuelan equine encephalitis will produce viremia levels in bats well above the threshold levels for infection of some mosquito species (Corristan LaMotte and Smith 1956). Observations on the recently discovered Kyasanur Forest disease in India suggest that monkeys may serve as the natural reservoir

brates or arthropod hosts at times of physiological change as, for example during the mating and egg laying season of avian hosts or when mosquitoes emerge from hibernation.

(5) Reintroduction of encephalitis virus each year into endemic areas by avian hosts migrating from tropical areas of continuous transmission offers a possible means of overwintering for which there is some circumstantial evidence (Kissling *et al.* 1955). With regard to WEE however it is generally believed that resident avian hosts or arthropod vectors are the more likely winter reservoirs at least in the Far West. On the basis of their observations in Kern County to date Reeves, Bellamy and Scrivani (in press) conclude that "The possibility that a vertebrate host is the immediate source of winter time vector infection is as likely as the alternative that virus survives solely in the vector in winter."

St. Louis Encephalitis (SLE)

Summer infection chains of SLE In the Far West it seems apparent that the summer transmission cycle for SLE in rural situations is similar to that for WEE *C. tarsalis* serving as the primary vector and birds as the natural vertebrate reservoir (FIGURE 2). The basic ecologic interrelationships for rural SLE would thus be similar to those outlined for WEE but this does not imply that there are not important differences between the two. One such difference is that WEE outbreaks usually occur in midsummer whereas SLE outbreaks take place in the late summer or fall even when the two occur in the same geographical area. This might be due to longer intrinsic or extrinsic incubation periods for SLE which in turn would limit its build up in the shorter summer seasons at higher altitudes or more northerly latitudes. Such inherent differences in the two agents may require different specific interrelationships between vectors, hosts and viruses despite the fact that their natural chains of infection are similar.

Although the infection chains for WEE and rural SLE appear to be quite similar there is increasing evidence that the transmission cycle for SLE in urban situations of the central states and possibly other areas is quite different (FIGURE 3). The *Culex pipiens* complex is being increasingly implicated as the primary vector for urban SLE. The importance of these household mosquitoes was first postulated by Lumsden (1934) and was based on his investigation of the 1933 epidemics in the vicinities of St. Louis and of Independence, Mo. Subsequent findings of other workers have further supported Lumsden's original hypothesis. The *C. pipiens* complex in the United States includes *pipiens*, *molestus*, *quinquefasciatus* and possibly others (Mattingly *et al.* 1951). Under laboratory conditions members of this complex have been known for some time to be capable of transmitting SLE virus. High population densities of one or more species have been found in vicinities in which SLE epidemics have occurred and collections of these mosquitoes from localities in which epidemics were in progress have yielded specimens from which SLE viruses were isolated (Kanzenhofer *et al.* 1951; Beadle *et al.* 1951).

In addition to the *C. pipiens* complex and *C. tarsalis* SLE virus has been isolated in nature from *C. stigmatosoma* and *C. melanicon* (*dorsalis*) (Ferguson


PRIMARY ENDEMIC VECTOR

FIGURE 4 Summer infection chains for eastern equine encephalitis

siana. Five recoveries of FEE virus have been reported from *C. melanura* in New Jersey (Holden Miller and Jobbins 1954 Burbutis and Jobbins in press). Isolations of FEE virus in nature have also been made from *Mansonia perturbans* (Howitt *et al.* 1949) *Anopheles crucians* (Kissling *et al.* 1955 Karstad *et al.* 1957) *Culex salinarius* (Burbutis and Jobbins in press) *Iedes mitchellae* and a pool of *Culicoides* that had not been identified as to species (Karstad *et al.* 1954). Laboratory attempts to transmit EEE virus with *C. salinarius* and *A. crucians* have not been successful (Chamberlain *et al.* 1954). The geographical distributions of *A. mitchellae* and salt marsh species of *Culicoides* in the United States coincide in general with the distribution of FEE and these species merit further attention as potential vectors of EEE.

The distribution of salt marsh mosquitoes *Aedes sollicitans* and *I. taeniorhynchus* is also similar to that of ELE and circumstantial evidence would indicate that they might be involved as endemic vectors of EEE as for example in the outbreak which took place in the Dominican Republic in 1948 and 1949 (Eklund Brennan and Bell 1950 Eklund Bell and Brennan 1951). In laboratory tests with *I. sollicitans* Chamberlain *et al.* (1954) have given this species a vector potential rating of excellent.

In assessing the importance of *C. melanura* as a vector of WEE and EFE viruses it is necessary to take into account that this mosquito has rarely been observed to bite man even in situations where this insect was known to be present in considerable abundance. Further information is needed on the feeding habits of *C. melanura*. If it truly does not bite man the endemic vector of FEE must be another species of mosquito or some other arthropod

from which this Group B type virus is transmitted by arthropods to man (Work and Trapido 1957). It is also well established that mammals are natural reservoirs of Russian spring summer encephalitis and other arthropod borne virus diseases such as yellow fever and dengue. Further study may indicate that mammals (including man) may possibly serve as reservoir links in SLE chains of infection.

The *C. pipiens* complex is quite variable biologically and the different variants are quite labile and difficult to distinguish taxonomically (Mattingly *et al.* 1951). For example, true *pipiens* probably does not feed upon mammals, whereas *molestus* even though it is autogenic may feed avidly upon man and other mammals. Further work on the summer transmission cycles for urban SLE must therefore include detailed studies on the feeding habits of the *C. pipiens* complex as well as studies of the vertebrate hosts.

In situations where small urban communities are surrounded by irrigated rural areas it is possible that there may be mixed epidemics of rural and urban SLE in which both *C. tarsalis* and the *C. pipiens* complex are involved as primary vectors. This may have been true for the SLE outbreak in the Texas Panhandle during 1956 (Alexander and Murray 1957). Studies during previous years had shown that both *C. tarsalis* and *C. quinquefasciatus* were abundant in the area of the outbreak (Harmston *et al.*, 1956) and SLE virus has been recovered from both these species in several other areas.

Winter infection chains of SLE. The possible winter reservoirs for *C. tarsalis* transmitted rural SLE appear to be quite similar to those for WEE. The winter reservoir mechanisms for urban SLE may also be similar to those for WEF, although involving different primary reservoirs and vectors. Positive evidence on the possibility that hibernating mosquitoes serve as winter reservoirs for SLE has been obtained by Bellamy, Reeves, and Scrivani (in press) who found that SLE virus survived for 116 days in experimentally infected *C. quinquefasciatus* that were kept in an unheated cellar during the winter months. Hurlbut (1950) has also shown that *C. quinquefasciatus* can carry the virus of Japanese B encephalitis through a 91 day period of simulated hibernation and retransmit it to young albino mice.

Resident birds are also possible winter reservoirs for SLE. Chamberlain and his co-workers (1951) recovered SLE virus from the gizzard of a cowbird 38 days after infection.

The likelihood that migrating birds may reintroduce SLE virus into some endemic areas of the United States each spring requires further investigation.

Eastern Equine Encephalitis (EEE)

Summer infection chains of EEE. The summer infection chains for EEE are quite different from those for WEE and SLE and some of the important links are yet to be discovered (FIGURE 4). The identity of the primary vectors is particularly puzzling. Most of the accumulated evidence suggests that *Culiseta melanura* is the primary sylvan vector responsible for bird to bird transmission in nature. Chamberlain *et al.* (1951) reported one isolation of EEE virus from naturally infected *C. melanura* mosquitoes collected in Loui-

function among wild bird species in their natural habitat. However, unless the mode of transmission once demonstrated is shown to be peculiar to the activities of confined pheasants, it should be further considered as a possible means of survival for the virus in nature (FIGURE 4).

Winter infection chains of EEE. Of the three agents under discussion, EEE virus has been studied least in regard to its mechanism of survival during the winter months. To our knowledge, no detailed studies have been made on the ability of any arthropod to serve as the winter reservoir. *Culiseta melanura*, the mosquito thought to be the principal sylvan vector, is thought to hibernate in the larval stage. This belief is based upon the finding of viable larvae under winter conditions, but there is no specific information available on the fate of the adult at the end of the active mosquito season.

The possible role of birds in the overwintering of EEE virus likewise has received only limited study. There are no published reports of attempts to demonstrate latent virus in birds that previously had been infected with the agent. Kissling *et al.* (1955) reported on an attempt to test the hypothesis that LEE and WEE virus were introduced into the United States annually by migrating birds. These workers found a higher antibody rate against each agent in birds migrating into the tropics during the winter months than in permanent or winter resident birds in Louisiana. Although the results of this study lend support to the hypothesis that the equine viruses are reintroduced each year by migrating birds, the evidence is by no means conclusive.

Conclusion

In looking back over the investigations of encephalitis of the past decade, it appears that perhaps too little attention has been given to the vertebrate reservoirs in nature. In our preoccupation with the arthropod vectors, we have frequently neglected the vertebrate hosts. Probably our thinking in this regard has been too much conditioned by experiences with other mosquito-borne diseases such as malaria and yellow fever. In the western states at least, there has been a recent trend toward giving more research attention to the vertebrate hosts. Eventually we may find that for the arthropod-borne encephalitides, an attack upon the primary reservoirs is more feasible than vector control in breaking the chain of infection. It is important, therefore, that we give full consideration to all of the critical interrelationships in the natural history of these diseases if we are to accomplish our ultimate objective of prevention or control of encephalitis.

Summary

The three principal arthropod-borne viruses in the United States are western equine (WEE), eastern equine (EEE), and St. Louis (SLE). WEE and SLE occur primarily in the twenty-two western states, whereas EEE occurs primarily in the Atlantic and Gulf coast states. Birds serve as natural hosts and mosquitoes as vectors for all three viruses. Man and horses are accidental hosts. Clinical disease in man is produced by all three viruses; in horses by WEE and EEE, and in birds by EEE alone.

that feeds on man and horses as well as on birds. Fortunately for man mosquitoes of the *C. pipiens* complex are poor vectors of IFE (Davis 1940 Chamberlain *et al.* 1954) and are probably not involved as endemic vectors.

Information on the natural hosts of IIF is even more meager than for WIF and SLE. A few recoveries of IIF virus have been made from wild and domestic birds and mammals and antibodies have been found in a wide variety of species including 40 or 50 kinds of birds. Laboratory studies with EEL infections in horses (Kissling *et al.* 1954) indicate that they develop relatively low viremia levels and it seems most likely that birds serve as the primary reservoirs in the basic infection chain. The specific identity of these reservoirs is yet to be established. Some birds such as red winged blackbirds, cardinals and English sparrows may be apparent hosts in which IIF infections are almost always fatal; others such as American egrets, snowy egrets and white ibis appear to be inapparent hosts (Kissling *et al.*, 1954a).

In summary it seems reasonable to assume that the enzootic foci for EEE are fresh water swamps that harbor the wild bird reservoirs and in which the primary sylvan vector *C. melanura* breeds. The endemic chains of infection through which FEF is carried from these enzootic foci to man and domestic mammals are however yet to be determined (Holden 1954).

LIT infections in ring necked pheasants In the process of elucidating the natural histories of many diseases epidemiologists have frequently observed phenomena for which no explanation was readily available on the basis of pre-existing knowledge or belief. For example it was originally thought that the infection chain of yellow fever consisted solely of a man mosquito-man transmission cycle. Furthermore *Aedes aegypti* was believed to be the only vector. It was subsequently observed however that the causal virus survived admirably in areas where man was infected too infrequently to serve as the reservoir and/or where the known vector had been effectively controlled. These observations led to the discovery of transmission cycles in which man and *Aedes aegypti* had no part. From these findings have evolved the now well known concepts of so called urban and jungle yellow fever (Bates 1946).

The natural history of the encephalitides is also complicated by findings that are not satisfactorily explained by the hypotheses that have been discussed. The results of studies on the epizootiology of EEE in confined flocks of ring necked pheasants (*Phasianus colchicus torquatus*) are illustrative of findings that are inconsistent with the hypothesis that the encephalitides are invariably mosquito borne.

Reports on the behavior of the disease in naturally occurring outbreaks in pheasants (Beaudette and his co workers 1939 1941 1945 1948 and 1952) and under controlled experimental conditions (Holden 1955) provide conclusive evidence that the infection may be spread in these hosts by some means of direct contact that does not involve transmission by arthropods. We can only speculate upon the significance of these findings. It is realized that the ring necked pheasant is not native to those areas in which EEF is epizootic. Furthermore studies have been made only on flocks reared in captivity. Thus it is possible that the mechanisms of spread among captive pheasants cannot

- BLACKMORE J S & J I WINN 1954 *Aedes nigromaculis* (Willow) mosquito naturally infected with western equine encephalomyelitis virus. *Proc Soc Exptl Biol Med* 87 378-329
- BLACKMORE J S & J I WINN 1956 A winter isolation of western equine encephalitis virus from hibernating *Culex tarsalis* Coq. *Proc Soc Exptl Biol Med* 91 146-149
- BIRBUTIS J P & D M JOBBINS 1957 *Culiseta melanura* (Coq) and eastern equine encephalomyelitis in New Jersey. 44th Ann Meeting Proc N J Mosq Extermination Assoc. In press
- BURROUGHS A L & R N BURROUGHS 1954 A study of the ecology of western equine encephalomyelitis virus in the upper Mississippi river valley. *Am J Hyg* 60 2-36
- CHAMBERLAIN R W H KUBIN R F KISLING & M F EDSON 1951 Recovery of virus of eastern equine encephalomyelitis from a mosquito *Culiseta melanura* (Coquillett). *Proc Soc Exptl Biol Med* 77 346-39
- CHAMBERLAIN R W & R K SIKES 1955 Laboratory investigations on the role of bird mites in the transmission of eastern and western equine encephalitis. *Am J Trop Med Hyg* 4 106-118
- CHAMBERLAIN R W R K SIKES D B NELSON & W D SCUDIA 1954 Studies on the North American arthropod borne encephalitudes. VI Quantitative determinations of virus-vector relationships. *Am J Hyg* 60 218-285
- CHAMBERLAIN R W F C CORRIAN & R K SIKES 1954a The extrinsic incubation of eastern and western equine encephalitis in mosquitoes. *Am J Hyg* 60 269-277
- CHAMBERLAIN R W K F KISLING D D STAMM & W D SCUDIA 1954 Virus of St. Louis encephalitis in the species of wild birds. *Am J Hyg* 65 110-118
- COCKBURN T A C A SOOTER & A D LANGMUIR 1957 Ecology of western equine and St. Louis encephalitis viruses: a summary of field investigations in Weld County, Colorado 1949 to 1953. *Am J Hyg* 65 130-146
- CORRIAN F C I C LAMOTTE Jr & D C SMITH 1956 Susceptibility of rats to certain encephalitis viruses. *Feederation Proc* 15 1
- DAVIS W A 1940 A study of birds and mosquitoes as hosts for the virus of eastern equine encephalomyelitis. *Am J Hyg* 32 45-59
- DOW R P C A MAIL & C S RICHARDS 1956 Observations on the overwintering of *Culex tarsalis* in northern Utah. *Proc 9th Ann Meeting Utah Mosquito Abatement Assoc* 12 15
- DOW R P W C REEVES & R F BELLAMY 1957 Field tests of avian host preference of *Culex tarsalis* Coq. *Am J Trop Med Hyg* 6 294-303
- FALUND C M 1954 Mosquito-transmitted encephalitis viruses: a review of their insect and vertebrate hosts and the mechanisms for survival and dispersion. *Folia Parasitol* 3 285-305
- FALUND C M J F BELL & J M BRENNAN 1951 Antibody survey following an outbreak of human and equine disease in the Dominican Republic caused by the eastern strain of equine encephalomyelitis virus. *Am J Trop Med* 31 312-329
- FALUND C M J M BRENNAN & J F BELL 1950 Final report to the Pan American Sanitary Bureau regarding the 1948-49 outbreak of eastern equine encephalitis in the Dominican Republic. *Bol Ofic sanit panam* 29 493-508
- FERGUSON F F 1954 Biological factors in the transmission of American arthropod borne virus encephalitudes. *Public Health Monogr* No 23
- HAMMON W McD W C REEVES & G I SATHER 1951 Western equine and St. Louis encephalitis viruses in the blood of experimentally infected wild birds and epidemiological implications of findings. *J Immunol* 67 35-366
- HAMMON W McD & W C REEVES 1943 Laboratory transmission of western equine encephalomyelitis virus by mosquitoes of the genera *Culex* and *Culiseta*. *J Exptl Med* 78 425-434
- HAMMON W McD & W C REEVES 1945 Recent advances in epidemiology of arthropod borne virus encephalitudes including certain exotic types. *Am J Public Health* 35 994-1004
- HAMMON W McD W C REEVES & P GALINDO 1945 Epidemiologic studies of encephalitis in the San Joaquin Valley of California 1943 with the isolation of viruses from mosquitoes. *Am J Hyg* 42 299-306
- HARMSTON F C C R SCHULTZ R B EAD & C C MENZIE 1956 Mosquitoes and encephalitis in the irrigated high plains of Texas. *Public Health Repts U S* 71 759-766
- HOLDEN P B J MILLER & D M JOBBINS 1954 Isolations of eastern equine encephalomyelitis virus from mosquitoes (*Culiseta melanura*) collected in New Jersey 1953. *Proc Soc Exptl Biol Med* 87 457-459

Wild birds serve as the primary reservoir and *C. tarsalis* as the primary sylvan and endemic vector, for WEL and rural SIE. Mosquitoes of the *C. pipiens* complex appear to be the primary sylvan and endemic vectors of urban SIE. Mammals as well as birds are suggested as possible reservoirs. Wild birds also are primary reservoirs for IFF and *C. melanura* seems to be the primary sylvan vector. The endemic vector for IFE is yet unknown. Direct bird to bird transmission of IFF among pheasant flocks has been proved, but its possible role among wild birds has not been determined.

Either the mosquito vectors or the avian hosts appear to be possible overwintering reservoirs for encephalitis viruses. Studies with WFE and SLE have shown that both of these viruses can survive for extended periods in either mosquitoes or birds. The possible reintroduction of virus into endemic areas each season by migrating birds requires further investigation. Overwintering in mites or transovarian passage is not believed to be the mechanism for the winter survival of encephalitis viruses.

Ecological interrelationships in time and space determine the actual role of potential vectors or hosts in the epizootiology and epidemiology of encephalitis. Of primary importance are the population densities, distribution and nesting, resting and roosting habits of the avian hosts and the population densities, summer and winter longevity, blood feeding habits and host preferences of the arthropod vectors. An abundance of available dead end hosts, such as domestic mammals, may provide a considerable degree of zoophylaxis for the human population.

It is believed that the role of specific vertebrate hosts in the natural history of the encephalitides deserves greater investigative attention and that control of vertebrate reservoirs as well as vector control offers promise in preventing or limiting epidemics of encephalitis among man and domestic animals.

References

- ALEXANDER, E. P. & W. A. MURRAY, JR. 1957. Surveillance of arthropod borne encephalitis in the United States—1956. Mimeogr. Rept. Epidemiol. Branch, CDC, U. S. Public Health Service.
- BATES, M. 1946. The natural history of yellow fever in Colombia. *Sci. Monthly* 63: 42–52.
- BEADLE, L. D. 1955. Man biting habits of *Culex tarsalis* and associated mosquitoes in northern Utah. Proc. 8th Ann. Meeting Utah Mosquito Abatement Assoc. 3.
- BEADLE, L. D., G. C. MUEZIES, C. R. HAYES, JR., F. J. VON ZUBEN, JR. & I. B. EADS. 1957. St. Louis encephalitis in Hidalgo County, Texas: vector evaluation and control. *Public Health Repts. U. S.* 72: 531–535.
- BEAUDETTE, F. R. 1939. Equine encephalomyelitis in a feline host. *Proc. 43rd Ann. Meeting U. S. Livestock San. Assoc.* 185–203.
- BEAUDETTE, F. R., J. J. BLACK & C. B. HUDSON. 1941. Another case of equine encephalomyelitis in New Jersey pheasants. *J. Am. Vet. Med. Assoc.* 98: 449–451.
- BEAUDETTE, F. R. & C. B. HUDSON. 1945. Additional outbreaks of equine encephalomyelitis in New Jersey pheasants. *J. Am. Vet. Med. Assoc.* 107: 384–386.
- BEAUDETTE, F. R. & J. J. BLACK. 1948. Equine encephalomyelitis in New Jersey pheasants in 1945 and 1946. *J. Am. Vet. Med. Assoc.* 112: 140–144.
- BEAUDETTE, F. R., J. J. BLACK, C. B. HUDSON & J. A. BIVINS. 1952. Equine encephalomyelitis in pheasants from 1947 to 1951. *J. Am. Vet. Med. Assoc.* 121: 478–483.
- BELLAMY, R. E., W. C. REEVES & R. I. SCRIVANI. 1957. Relationships of mosquito vector to winter survival of encephalitis viruses. II. Under experimental conditions. *Am. J. Hyg.* In press.

- BLACKMORT J S & J F WINN 1954 *Aedes nigromaculis* (Dull & W.) mosquito naturally infected with western equine encephalomyelitis virus. *Proc Soc Exptl Biol Med* 87: 378-379.
- BLACKMORT J S & J F WINN 1956 A winter isolation of western equine encephalitis virus from hibernating *Culex tarsalis* Coq. *Proc Soc Exptl Biol Med* 91: 146-149.
- BURBULTI I P & D M JOBBINS 1957 *Culiseta melanura* (Coq.) and eastern equine encephalomyelitis in New Jersey. 44th Ann Meeting Assoc N J Mosq Extermination Assoc. In press.
- BURROUGHS A L & R N BURROUGHS 1954 A study of the ecology of western equine encephalomyelitis virus in the upper Mississippi river valley. *Am J Hyg* 60: 27-36.
- CHAMBERLAIN R W H RUBIN R F KISLING & M F FETTERSON 1951 Recovery of virus of eastern equine encephalomyelitis from a mosquito *Culiseta melanura* (Coquillett). *Proc Soc Exptl Biol Med* 77: 336-337.
- CHAMBERLAIN R W & R K SIKES 1955 Laboratory investigations on the role of *Aedes* mites in the transmission of eastern and western equine encephalitis. *Am J Trop Med Hyg* 4: 106-118.
- CHAMBERLAIN R W R K SIKES D B NELSON & W D SUDIA 1954 Studies on the North American arthropod borne encephalitides. VI Quantitative determinations of virus vector relationships. *Am J Hyg* 60: 28-285.
- CHAMBERLAIN R W F C CORRISSAN & R K SIKES 1954a The extrinsic incubation of eastern and western equine encephalitis in mosquitoes. *Am J Hyg* 60: 269-277.
- CHAMBERLAIN R W R F KISLING D D STAMM & W D SUDIA 1955 Virus of St. Louis encephalitis in three species of wild birds. *Am J Hyg* 65: 110-119.
- COCARTON T A C A SOOTER & A D LANCHEIR 1957 Ecology of western equine and St. Louis encephalitis viruses: a summary of field investigations in Weld County Colorado 1949 to 1953. *Am J Hyg* 65: 130-146.
- CORRISSAN F C I C LAMOTTE JR & D C SMITH 1956 Susceptibility of bats to certain encephalitis viruses. *Federation Proc* 15: 1.
- DAVIS W A 1940 A study of birds and mosquitoes as hosts for the virus of eastern equine encephalomyelitis. *Am J Hyg* 32: 45-51.
- DOW R P G A MAIL & C S RICHARDS 1946 Observations on the overwintering of *Culex tarsalis* in northern Utah. *Proc 9th Ann Meeting Utah Mosquito Abatement Assoc* 12: 15.
- DOW R P W C REEVE & R E BELLAMY 1957 Field tests of avian host reference of *Culex tarsalis* Coq. *Am J Trop Med Hyg* 6: 294-303.
- FALUND C M 1954 Mosquito-transmitted encephalitis viruses: a review of their insect and vertebrate hosts and the mechanisms for survival and dispersion. *Fertil Parasitol* 3: 285-405.
- FALUND C M J F BELL & J M BRENNAN 1951 Antibody survey following an outbreak of human and equine disease in the Dominican Republic caused by the eastern strain of equine encephalomyelitis virus. *Am J Trop Med* 31: 312-378.
- FALUND C M J M BRENNAN & J F BELL 1950 Final report to the Pan American Sanitary Bureau regarding the 1948-49 outbreak of eastern equine encephalitis in the Dominican Republic. *Bol Ofic sanit panam* 29: 493-508.
- FERGUSON F F 1954 Biological factors in the transmission of American arthropod borne virus encephalitides. *Public Health Monographs* No 23.
- HAMMON W MCD W C REEVES & G E SATHIER 1951 Western equine and St. Louis encephalitis viruses in the blood of experimentally infected wild birds and epidemiological implications of findings. *J Immunol* 67: 357-366.
- HAMMON W MCD W C REEVES 1943 Laboratory transmission of western equine encephalomyelitis virus by mosquitoes of the genera *Culex* and *Culiseta*. *J Exptl Med* 78: 425-434.
- HAMMON W MCD W C REEVE 1945 Recent advances in epidemiology of arthropod borne virus encephalitides including certain exotic types. *Am J Public Health* 35: 994-1004.
- HAMMON W MCD W C REEVES & P CALINDO 1945 Epidemiologic studies of encephalitis in the San Joaquin Valley of California 1943 with the isolation of viruses from mosquitoes. *Am J Hyg* 42: 279-306.
- HARMSTON F C C R SCHULTZ R B EADS & G C MENZIES 1956 Mosquitoes and encephalitis in the irrigated high plains of Texas. *Public Health Reports U S* 71: 759-66.
- WILDEN P B J MILLER & D M JOBBINS 1954 Isolations of eastern equine encephalomyelitis virus from mosquitoes (*Culiseta melanura*) collected in New Jersey 1953. *Proc Soc Exptl Biol Med* 87: 457-459.

- HOLDEN I 1954 Epizootiological studies on eastern equine encephalomyelitis in ring necked pheasants in a localized area New Jersey 1953 44-45 Doctoral thesis Univ Pittsburgh Pittsburgh Pa
- HOLDEN P 1955 Transmission of eastern equine encephalomyelitis in ring necked pheasants Proc Soc Exptl Biol Med 88 601-610
- HOLDEN I 1955a Recovery of western equine encephalomyelitis virus from naturally infected English parrots of New Jersey 1953 Proc Soc Exptl Biol Med 88 490-492
- HOWITT B I H P DODGE I K BISHOP & R H CORRIE 1949 Recovery of the virus of equine encephalomyelitis from mosquitoes (*Wansonia perturbans*) collected in Georgia Science 110 141-142
- HURLBUT H S 1950 The transmission of Japanese B encephalitis by mosquitoes after experimental hibernation Am J Hyg 61 265-268
- JENKINS D W 1950 Bionomics of *Culex tarsalis* in relation to western equine encephalomyelitis Am J Trop Med 30 909-916
- KARSTAD I H O K FLETCHER J SPALATIN R I ROBERTS & R I HANSON 1957 Eastern equine encephalomyelitis virus isolated from three species of Diptera from Georgia Science 125 395-396
- KEENER G G JR 1952 Observations on overwintering of *Culex tarsalis* Coquillett (Diptera: Culicidae) in western Nebraska Mosquito News 12 205-209
- KISHLING R L R W CHAMBERLAIN M F EIDSON R K SIKES & M A BECCA 1954 Studies on the North American arthropod borne encephalitides II Eastern equine encephalitis in horses Am J Hyg 60 237-250
- KISHLING R L R W CHAMBERLAIN R K SIKES & M F EIDSON 1954a Studies on the North American arthropod borne encephalitides III Eastern equine encephalitis in wild birds Am J Hyg 60 251-265
- KISHLING R L P W CHAMBERLAIN D B NELSON & D D STAMM 1955 Studies on the North American arthropod borne encephalitides VIII Equine encephalitis studies in Louisiana Am J Hyg 62 233-254
- LEMSDEN L L 1934 Observations on the epidemiological features of epidemic encephalitis lethargica Off Surgeon General U S Public Health Service Unpublished
- MAIL G A & J V SMITH 1957 Seasonal activity patterns of mosquitoes in Cache Valley Proc 10th Ann Meeting Utah Mosquito Abatement Assoc In press
- MATTINGLY P F L L ROZEBOOM K L KNIGHT H IAVEN F H DRUMMOND S F CHRISTOPHERS & P G SHUTE 1951 The *Culex pipiens* complex Trans Roy Ent Soc London 102 331-382
- MILES V I B F HOWITT R GORRIE & T A COCKBURN 1951 Encephalitis in mid west V Western equine encephalomyelitis virus recovered from mites *Dermanyssus americanus* Fwing Proc Soc Exptl Biol Med 77 395-396
- PHILIP C B J F BELL & C L LARSON 1955 Evidence of infectious diseases and parasites in a peak population of black tailed jack rabbits in Nevada J Wildlife Management 19 226-233
- RANZENHOFER L R E R ALEXANDER L D BEADLE A BERNSTEIN & R C PICKARD 1957 St Louis encephalitis in Calvert City Kentucky 1955 An epidemiologic study Am J Hyg 65 147-161
- REEVES W C 1951 Field studies on carbon dioxide as a possible host stimulant to mosquitoes Proc Soc Exptl Biol Med 77 64-66
- REEVES W C 1953 The knowns and the unknowns in the natural history of encephalitis Proc 21st Ann Conf Calif Mosquito Control Assoc 53-59
- REEVES W C 1953a Quantitative field studies on a carbon dioxide chemoatant of mosquitoes Am J Trop Med Hyg 2 375-381
- REEVES W C 1955 Epidemiological aspect of encephalitis under field conditions Proc Calif Mosquito Control Assoc 23rd Ann Conf & 11th Ann Meeting Am Mosquito Control Assoc 48-50
- REEVES W C W MCD HAMMON W H DOETSCHMAN H F MCCLURE & C SATHRE 1955 Studies on mites as vectors of western equine and St Louis encephalitis in California Am J Trop Med Hyg 4 90-105
- REEVES W C & W MCD HAMMON 1944 Feeding habit of the primary and possible mosquito vectors of western equine and St Louis encephalitis in the Yakima Valley Washington Am J Trop Med 24 131-134
- REEVES W C W MCD HAMMON D P FURMAN H E MCCLURE & B BRADMAN 1947 Recovery of western equine encephalomyelitis virus from wild bird mites (*Ornithonyssus sylvarum*) in Kern County California Science 105 411-412

- RYAN, W. C. & F. BELLAMY & K. I. SCRIVANI. 1957. Relation of mosquito vectors to winter survival of encephalitis viruses. I. Under natural conditions. Am. J. Hyg. In press.
- WORK, T. H. & H. TRAPIDO. 1957. Kyasanur forest disease—a new virus disease in India. Summary of preliminary report of investigations of the Virus Research Centre on an epidemic disease affecting forest villagers and wild monkeys of Shimoga District, Mysore. Indian J. Med. Sci. 11: 1-2.

VECTOR RELATIONSHIPS OF THE ARTHROPOD BORNE ENCEPHALITIDES IN NORTH AMERICA

By Roy W. Chamberlain

Communicable Disease Center, Public Health Service, Montgomery, Ala.

Although much remains to be learned about the ecology of the arthropod borne encephalitides in North America, past and recent studies have provided considerable knowledge of these diseases. Infection in horses and man has been associated with large mosquito populations and infection in wild birds and birds have been shown to be important hosts^{1,2}. Many arthropod species have been tested in the laboratory for their susceptibility to virus infection and for their ability to transmit and certain mosquitoes associated with birds have been indicted as vectors^{1,3,12}. Antibody and virus isolation studies in wild vertebrates have indicated the field host ranges and virus isolations from field caught mosquitoes have revealed those species that actually feed upon infected animals under natural conditions^{1,3,6,9,11,14,1}. Further information on the feeding habits of mosquitoes has been gained through field and laboratory observations and precipitin tests on naturally acquired blood meals. Correlation of mosquito susceptibility with viremias attained in various vertebrates has furnished clues concerning favorable host vector combinations^{10,13}. These are only a few of the important contributions that have been made.

As a result of these contributions certain vector relationships now appear reasonably clear. By making the most of this knowledge a logical selection of the most probable vectors is possible.

Eastern equine encephalitis (EEE). Field investigations suggest strongly that the foci of EEE infection are fresh water swamps. Outbreaks of EEE in horses or man have generally originated in the vicinity of swamps and the presence of the virus in birds and mosquitoes during intraepidemic and intraepizootic periods has been established in swampy areas. A high proportion of swamp inhabiting birds becomes immune during the transmission season and numerous isolations of virus have been made from their blood^{8,15,16,17,18}. Furthermore laboratory studies have shown wild birds to be highly susceptible to infection and to produce levels of virus in their blood sufficiently high to infect certain species of mosquitoes^{15,14}. Complete bird mosquito bird transmission cycles have been accomplished in the laboratory.

Attempts to isolate EEE virus from horse and man biting mosquitoes and other insects during epidemics or epizootics have almost always met with failure although thousands of individuals of such common species as *Aedes sollicitans*, *A. triseriatus* and *Psorophora ferox* have been tested. On the other hand tests on a swamp breeding mosquito *Culiseta melanura* have been highly productive. Eleven isolations of the virus have been made from this species in Louisiana¹⁵, New Jersey^{11,19} and Massachusetts⁵ always in association with active infection in the swamp-frequenting bird population. This species has been shown in the laboratory to be susceptible to infection and to transmit by its bite¹⁰.

A lesser number of isolations has been made from other species of mosquitoes. *Mansonia perturbans* collected in Georgia yielded virus on one occasion²⁴ As it is also a swamp breeder will feed upon birds and can transmit with moderate efficiency in the laboratory¹⁰ it falls under suspicion as a possible vector. *Anopheles crucians* has yielded virus on two occasions but only in trace amounts^{14, 25} The importance of these isolations is negligible since this species has been shown to be incapable of transmitting infection in the laboratory¹⁰ Virus has been isolated twice from blooded *Culex salinarius* taken in a swamp¹¹ This species has also been shown to be refractory to EEE infection in the laboratory¹⁰ and as the infection rate of associated birds was high at the time the isolation was made the virus detected was most likely in recently ingested bird blood One isolation has been made from *A. mitchellae* and another from a pool of *Culicoides* both apparently associated with infection in horses²⁷

The virus isolations from *Culiseta melanura* are made more significant by the fact that relatively few specimens were tested An isolation has been made from about every 200 to 400 specimens checked despite the fact that they were collected near their breeding sites and that young, previously unfed individuals made up a high proportion of the catches

Culiseta melanura is known to breed only in fresh water swamps or in heavily shaded seepage areas swamplike in character Although it is a little known species it is common in its preferred breeding areas While it has been found as far west as Colorado and Oklahoma²⁸ its range is generally restricted to the eastern half of the United States and coincides with the distribution of EEE in this country Precipitin tests on specimens engorged with blood when captured have shown that birds are preferred hosts^{11, 29}

The evidence at hand points therefore to fresh water swamps as enzootic foci to birds as primary hosts and to *C. melanura* as the principal enzootic vector In view of the restricted habitat of *C. melanura* however and its apparent preference for bird hosts epizootics in horses and the occasional cases in man cannot logically be attributed to this species It seems more likely that other vectors are required to carry the infection from birds to horses and to man

The infrequency of infections in man rules him out as a likely source of infection for more mosquitoes^{9, 30} Horses likewise appear to be of limited importance in epizootic spread of the disease since they rarely produce enough virus in their blood to infect mosquitoes efficiently²¹ They apparently receive their infections mostly from the bites of mosquitoes that have fed previously upon birds

In view of these restrictions certain conditions appear necessary to permit an epizootic in horses (and an epidemic in man) to occur Bird infection would need to be at high level furnishing a relatively great source of virus for mosquito infection This condition would depend upon an adequate population of *Culiseta melanura* (and assisting species such as *Mansonia perturbans*) and a low immunity rate in the bird population The latter would depend in turn upon a light infection rate the previous year As some of the infected birds fan out into the adjacent areas they become diluted with noninfected birds

not associated with the swamp foci. This, together with the fact that a relatively small proportion of the common horse feeding mosquitoes find opportunity to feed upon birds, make it necessary that the population of these mosquitoes be high. If, in addition to all this, there is a low immunity rate in the horse population conditions are favorable for the epizootic. Now, with large numbers of the epizootic species of mosquito present, the likelihood of additional horse infections through mosquitoes infected from an occasional horse with higher viremia becomes possible.²⁰ A high horse infection rate also favors occasional mechanical transmissions from horse to horse by the interrupted feedings of biting insects such as *Culicoides*²¹ horseflies, deer flies, stable flies and mosquitoes. That mosquitoes and stable flies are capable of transmitting mechanically between laboratory animals has been shown experimentally.²²

Mechanical transmission between caged pheasants by means of pecking has been demonstrated.²³ There is no proof however that this mode of transmission is of much consequence among wild, unconfined birds. It seems unlikely to influence the natural cycle of the disease appreciably.

Western equine encephalitis (WEE) The ecology of this disease has undergone intensive investigation over many years.^{19, 24} Hundreds of isolations of the virus from *Culex tarsalis* attest to the close association of this mosquito species with natural sources of infection. Serologic and virus isolation studies in wild birds have firmly established their role as the most important hosts. Inoculation of wild birds in the laboratory has confirmed their susceptibility and revealed viremias adequate to infect susceptible mosquitoes.^{24, 25} Transmissions to barnyard fowl²⁶ and wild birds²⁵ by mosquitoes have also been accomplished in the laboratory.

Studies on the feeding habits of *C. tarsalis* have shown that while this species feeds most frequently upon birds it is also attracted to man, horses and numerous other vertebrates. It is closely associated with birds in the West for it breeds in natural temporary water sources in dry land areas and in irrigation water overflows in cultivated areas the same sites in which many bird species concentrate. Because of its numbers, its close association with birds, its avidity for man and horses, and its great susceptibility as proved in laboratory tests *C. tarsalis* fulfills requirements for both an endemic and an epidemic vector. While many other species of mosquitoes are susceptible particularly the *Aedes*, very few isolations of virus have been made from them. It is possible that *A. dorsalis* and *A. nigromaculis* which are abundant during the encephalitis season might assist in the spread of WEE in horses once an outbreak is under way in the same manner that certain *Aedes*, *Psorophora* or other genera may spread EEE infection from an occasional high viremia horse. At most however this mode of spread would appear to be only supplementary to transmission from birds to horses by *C. tarsalis* as WEE viremias in horses are apparently even lower than EEE viremias.²⁷ Occasional transmissions from birds to horses by *Aedes* probably also occur but with considerably less frequency than by *C. tarsalis* because of the differences in their bird feeding habits.

Until a few years ago the distribution of WEE was believed to be limited to *Culex tarsalis*. Now we know that this is not the case. While disease in

horses and man appears generally to be limited to areas where *C. tarsalis* is the vector infection in wild birds and mosquitoes in swamps in the East are known to occur in the absence of *C. tarsalis*. In 1952 several isolations of WEE virus were made from birds collected in and adjacent to a swamp in Louisiana this same survey¹⁶ also yielded an isolation of this virus from *Culiseta melanura* and two others from *Aedes infirmatus* a species breeding on the swamp margin. In 1953 Holden²⁸ isolated WIF virus from a sparrow collected adjacent to a swamp in New Jersey, and Kissling⁴ detected WLI antibody in resident birds and horses in Massachusetts. An isolation of WEL from freshly engorged *C. pipiens quinquefasciatus* in North Carolina was made in 1955.¹¹ This isolation established the presence of WFE in that area. The importance of this mosquito species as a vector is greatly in doubt however as it has been shown to be refractory to WEE infection in the laboratory.¹⁰ The virus isolated was undoubtedly from the blood recently ingested and did not represent true mosquito infection. In 1956 several WIF isolations were made from the blood of birds collected in and near a swamp in New Jersey.²¹ Again a WEF virus isolation was made from *C. melanura*.¹¹

This evidence indicates that the enzootic status of WEE in the East is similar to that of EEE. Its unimportance as a disease producing agent in man and horses is probably due chiefly to the absence of its highly efficient vector of the West. Lower viremias in birds are produced than by EFE making infection of various feds and other horse and man feeding mosquitoes that may occasionally feed upon birds less likely. Also the viremia of WEE is lower in horses than that of EEE²⁷ so that horse to-horse transmission must be an even more rare occurrence. In addition laboratory studies have shown that when animals immune to EEE were infected with WEF they produced significantly lower viremias than nonimmune animals.²⁴ This factor may be important in areas where a high proportion of birds and horses possess FEE antibody. A combination of these deterrent effects probably serves to keep WEE within its enzootic foci in the East.

St. Louis encephalitis (SLE) Strong evidence indicates that SLE like EEE and WEE is primarily a bird disease transmitted by mosquitoes.^{1, 8, 9, 22, 29} Apparently similar factors foster its successful propagation namely adequate populations of susceptible birds and vector mosquito species and close association between the birds and the vectors. Again man seems to be an unfortunate accidental host.

The virus of SLE differs from those of EEE and WEE however in having an almost completely reversed mosquito susceptibility range. EEE and WEE viruses develop readily in all species of *Aedes* and *Psorophora* thus far tested but do so poorly or not at all in most *Culex* except *C. tarsalis*.^{10, 13} SLE virus on the other hand is highly infectious for all species of *Culex* that have been tested but does not develop well in *Aedes* and *Psorophora*.^{11, 13} These vector differences are reflected in the epidemiology of SLE not in the West where *C. tarsalis* abounds but farther east in the central states where *tarsalis* is scarce or absent.

Extensive studies in the Far West particularly in the Central Valley of California have shown with little doubt that *C. tarsalis* is the main SLE

vector in that area and is responsible for most of the transmission to wild birds, domestic fowl and man. Up to 1953, 69 isolations of the virus had been made from wild caught *C. tarsalis* as against only 3 for *C. quinquefasciatus*, 2 for *C. stigmatosoma* and 1 for *C. dorsalis*.³ In 1954 alone, 87 additional isolations from *C. tarsalis* were reported.¹⁷ The greatest outbreaks have coincided with an abundance of *C. tarsalis* and a high proportion of the cases have been rural rather than urban.⁴⁰ As might be expected, the epidemiology is similar in many respects to that of WEF.

A different epidemiological picture has been presented in the central and east central states, namely, in the outbreaks in St. Louis, Mo., 1933, Hidalgo County, Texas, 1954, Calvert City, Ky., 1955, and Louisville, Ky., 1956. In those areas *C. tarsalis* was scarce or absent. There the human cases were mostly urban or suburban and they were associated with heavy breeding of *C. pipiens* or *C. quinquefasciatus*.^{5, 6, 20, 25, 41} These mosquito species rarely find conditions as suitable for breeding in the more natural countryside. This fact, together with the sparser human populations in the country, may account for the relative infrequency of truly rural cases. Sylvan transmission to wild birds and fowl is possibly accomplished by a widespread nondomestic species such as *C. salinarius*, which has been shown to be an excellent transmitter in the laboratory.¹¹

The mechanics of an urban epidemic appear clear. Infected wild birds probably introduce the virus to high populations of suburban and urban *C. pipiens* or *C. quinquefasciatus*, which rapidly spread it to city dwelling birds and chickens. These in turn serve as a ready source of infection for additional numbers of the mosquitoes. A consequence of the resultant epizootic is exposure of a large segment of the human population to infection by mosquito bite.

A perplexing feature of the epidemiology of SILE is the importance of the *C. quinquefasciatus pipiens* complex in the East in the absence of *C. tarsalis*, but its relative unimportance in its presence in the Far West. Obviously, the balance of ecological factors that determine vector efficiency is tipped in its favor in the East but against it in the West. Although many contributing factors may be involved, the most likely appear to be strain differences in either the mosquitoes or the virus in the respective areas.

Studies comparing the transmitting efficiencies of *C. tarsalis* and 7 strains of *C. pipiens* and *C. quinquefasciatus* from various parts of the country showed that all transmitted a California strain of SILE virus from chick to chick with nearly 100 per cent efficiency.^{11, 13} The threshold of infection of *C. tarsalis* appeared to be the lowest, however, with virus in such low concentration as to be undetectable in 1:10 diluted blood adequate for infection.¹ That this finding may be significant is suggested by comparing the results of studies on the viremias produced in birds by the California virus strain¹⁹ and a strain from Kentucky.⁸ The Kentucky strain of SILE virus was found to develop about a tenfold greater concentration in the blood.

In a discussion of this rather flimsy but provocative evidence with W. C. Reeves of the School of Public Health, University of California, Berkeley, Calif., a hypothesis was developed to account for SILE vector differences in

California and the East. If *C. tarsalis* is abundant and more susceptible than the other mosquito species there may be little natural selection for high bird viremia virus strains. As a consequence the virus strains might characteristically produce viremias in birds too low for efficient infection of California *C. quinquefasciatus*. In the East however in the absence of *C. tarsalis* it is possible that the only virus strains propagated would be those that produced viremias in birds adequate to infect *C. quinquefasciatus pipiens* or other natural vectors of similar susceptibility with the resultant establishment of relatively high bird viremia strains in that area.

In spite of a considerable knowledge of vector relationships control measures through vector abatement are often impracticable from an economic point of view. Only urban SIE gives immediate promise of relatively inexpensive control. It is obvious therefore that the reasons for continued investigations on the ecology of encephalitis are not merely academic. There is a definite need to gain additional information that might lead to more effective control. Phases particularly needing study are the overwintering mechanisms, the finer points of vector host relationships and the influence of climatic and immunological factors upon disease spread. If these phases were fully understood outbreaks might be predictable and control more surely and economically effected.

References

1. FERGUSON F. F. 1954. Biological factors in the transmission of American arthropod-borne virus encephalites: a summary. U. S. Public Health Serv. Public Health Monograph No. 23.
2. BEADLE I. D. 1952. Eastern equine encephalitis in the United States. Mosquito News 12: 107-107.
3. REEVES W. C. 1953. The knowns and the unknowns in the natural history of encephalitis. Proc. & Papers. 23rd Ann. Conf. Calif. Mosquito Control Assoc.
4. LOOMIS F. C. 1955. Adult mosquito occurrence and human infection encephalitis cases in California. Proc. & Papers. 23rd Ann. Conf. Calif. Mosquito Control Assoc. 4th 11th Ann. Meeting. Am. Mosquito Control Assoc. 51-52.
5. LUDEN L. L. 1934. Observations on the epidemiological features of epidemic encephalitis lethargica. Office Surgeon General U. S. Public Health Service. Unpublished report.
6. RAZENTHOFER F. K., F. R. ALEXANDER, I. D. BEADLE, A. BERNSTEIN & R. C. HICKARD. 1951. St. Louis encephalitis in Calvert City, Kentucky, 1950. An epidemiologic study. Am. J. Hyg. 65: 14-161.
7. COHEN D. & O. SUGSMAN. 1951. Equine encephalomyelitis in New Jersey. Public Health News. New Jersey State Dept. Health 37: 270-237.
8. KI-LING R. F. & D. D. STAMM. Unpublished data.
9. FAHEND C. M. 1954. Mosquito-transmitted encephalitis viruses. A review of their insect and vertebrate hosts and the mechanisms for survival and dispersion. Exptl. Parasitol. 3: 285-305.
10. CHAMBERLAIN R. W., R. K. SIKES, D. B. NELSON & W. D. SUDIA. 1954. Studies on the North American arthropod-borne encephalites. VI. Quantitative determinations of virus-vector relationships. Am. J. Hyg. 60: 278-285.
11. CHAMBERLAIN R. W. & W. D. SUDIA. Unpublished data.
12. BARNETT H. C. 1956. The transmission of western equine encephalitis virus to the mosquito *Culex tarsalis* Coquillett. Am. J. Trop. Med. Hyg. 5: 86-98.
13. CHAMBERLAIN R. W. & W. D. SUDIA. 1951. The North American arthropod-borne encephalitis viruses in California. Coquillett. Am. J. Hyg. 66: 151-159.
14. CHAMBERLAIN R. W., A. C. SCHOTER & A. D. FAHEND. 1955. Ecology of western equine and St. Louis encephalitis viruses. A summary of field investigations in Weld County, Colorado, 1949 to 1953. Am. J. Hyg. 65: 130-146.
15. KI-LING R. E., R. W. CHAMBERLAIN, R. K. SIKES & M. E. EDSON. 1954. Studies on

vector in that area and is responsible for most of the transmission to wild birds domestic fowl and man. Up to 1953 69 isolations of the virus had been made from wild caught *C tarsalis* as against only 3 for *C quinquefasciatus* 2 for *C stigmatosoma* and 1 for *C dorsalis*.³ In 1954 alone 87 additional isolations from *C tarsalis* were reported.¹⁷ The greatest outbreaks have coincided with an abundance of *C tarsalis* and a high proportion of the cases have been rural rather than urban.⁴⁰ As might be expected the epidemiology is similar in many respects to that of WLF.

A different epidemiological picture has been presented in the central and east central states namely in the outbreaks in St. Louis Mo. 1933 Hidalgo County, Texas 1954 Calvert City Ky. 1955 and Louisville Ky. 1956. In those areas *C tarsalis* was scarce or absent. There the human cases were mostly urban or suburban and they were associated with heavy breeding of *C pipiens* or *C quinquefasciatus*.^{5, 6, 20, 25, 41} These mosquito species rarely find conditions as suitable for breeding in the more natural countryside. This fact, together with the sparser human populations in the country, may account for the relative infrequency of truly rural cases. Sylvan transmission to wild birds and fowl is possibly accomplished by a widespread nondomestic species such as *C salinarius* which has been shown to be an excellent transmitter in the laboratory.¹¹

The mechanics of an urban epidemic appear clear. Infected wild birds probably introduce the virus to high populations of suburban and urban *C pipiens* or *C quinquefasciatus* which rapidly spread it to city dwelling birds and chickens. These in turn serve as a ready source of infection for additional numbers of the mosquitoes. A consequence of the resultant epizootic is exposure of a large segment of the human population to infection by mosquito bite.

A perplexing feature of the epidemiology of SLE is the importance of the *C quinquefasciatus pipiens* complex in the East in the absence of *C tarsalis* but its relative unimportance in its presence in the Far West. Obviously the balance of ecological factors that determine vector efficiency is tipped in its favor in the East but against it in the West. Although many contributing factors may be involved the most likely appear to be strain differences in either the mosquitoes or the virus in the respective areas.

Studies comparing the transmitting efficiencies of *C tarsalis* and 7 strains of *C pipiens* and *C quinquefasciatus* from various parts of the country showed that all transmitted a California strain of SLE virus from chick to chick with nearly 100 per cent efficiency.^{11, 12} The threshold of infection of *C tarsalis* appeared to be the lowest however with virus in such low concentration as to be undetectable in 1/10 diluted blood adequate for infection.²⁴ That this finding may be significant is suggested by comparing the results of studies on the viremias produced in birds by the California virus strain¹⁰ and a strain from Kentucky.⁸ The Kentucky strain of SLE virus was found to develop about a tenfold greater concentration in the blood.

In a discussion of this rather flimsy but provocative evidence with W. C. Reeves of the School of Public Health, University of California, Berkeley, Calif., a hypothesis was developed to account for SLE vector differences in

infected English sparrows of New Jersey 1953 Proc Soc Exptl Biol Med 88 490-492

- 3) CHAMBERLAIN R W K I KISLINC D D STAMM & W D STELLA 1955 Virus of St. Louis encephalitis in three species of wild birds Am J Hyg 65 110-118
- 40 LONGHORN W D JR I M STEVENS A C HOLLIFFER JR A CUTLER JEN & I H FENNETTE 1956 Epidemiologic observations on acute infectious encephalitis in California, with special reference to the 1952 outbreak Am J Hyg 63 69-86
- 41 CHEN T D A C R HERLICH R F WHITE D M MAON & M I FERGUSON 1957 St. Louis encephalitis in Hidalgo County Texas Epidemiological features Public Health Repts U S 72 512-518

- the North American arthropod borne encephalitis. III Eastern equine encephalitis in wild birds. *Am J Hyg* 60 251-265
- 16 KISSLING R E R W CHAMBERLAIN D B NELSON & D D STAMM 1955 Studies on the North American arthropod borne encephalitis. VIII Equine encephalitis studies in Louisiana. *Am J Hyg* 62 233-254
 - 17 TENNETTE I H 1955 Etiologic aspects of the infectious encephalitis. *Proc & Papers 23rd Ann Conf Calif Mosquito Control Assoc Also 11th Ann Meeting Am Mosquito Control Assoc* 43-47
 - 18 CHAMBERLAIN R W H RUBIN R I KISSLING & M F FIDSON 1951 Recovery of the virus of eastern equine encephalomyelitis from a mosquito *Culiseta melanura* (Coquillett). *Proc Soc Exptl Biol Med* 77 396-399
 - 19 HOLDEN P B J MILLER & D M JOHNS 1954 Isolations of eastern equine encephalomyelitis virus from mosquitoes (*Culiseta melanura*) collected in New Jersey 1953. *Proc Soc Exptl Biol Med* 87 457-459
 - 20 BEADLE L D G C MENZIES C R HAYES F J VON ZUBEN & R B GADDS 1957 St. Louis encephalitis in Hidalgo County, Texas. Vector evaluation and control. *Public Health Repts U S* 72 531-535
 - 21 STAMM D D R E KISSLING O SCHUMAN & D COHEN 1957 Equine encephalitis in wild birds. New Jersey 1956. To be published
 - 22 SCHAEFFER M & I H ARNOLD 1954 Studies on the North American arthropod borne encephalitis. I Introduction. Contributions of newer field laboratory approaches. *Am J Hyg* 60 231-236
 - 23 KISSLING R E H RUBIN R W CHAMBERLAIN & M F FIDSON 1951 Recovery of virus of eastern equine encephalomyelitis from blood of a purple grackle. *Proc Soc Exptl Biol Med* 77 398-399
 - 24 STAMM D D & R E KISSLING 1957 The influence of reciprocal immunity on eastern and western equine encephalomyelitis infection in horses and English parrots. *J Immunol* 79 342-347
 - 25 ALEXANDER J R & W A MURRAY 1957 Surveillance of arthropod borne encephalitis in the United States—1956. Mimeographed report. Public Health Service Communicable Disease Center. Atlanta Ga.
 - 26 HOWITT B I H R DODGE L K BISHOP & R H GORRIE 1949 Recovery of the virus of eastern equine encephalomyelitis from mosquitoes (*Mansonia perturbans*) collected in Georgia. *Science* 110 141-142
 - 27 KARSTAD L H O K FLETCHER J SPALATIN K ROBERTS & R P HANSON 1957 Eastern equine encephalitis virus from three species of Diptera from Georgia. *Science* 125 395-396
 - 28 CARPENTER S J W W MIDDLEKAUFF & R W CHAMBERLAIN 1946 The mosquitoes of the Southern United States east of Oklahoma and Texas. *Am Midland Naturalist Monogr No* 3
 - 29 BLUBBIS P 1957 New Jersey Agr Exptl Sta. Personal communication
 - 30 SCHAEFFER M R F KISSLING R W CHAMBERLAIN & J M VANELLA 1954 Studies on the North American arthropod borne encephalitis. IV Antibody in human beings to the North American arthropod borne encephalitis. *Am J Hyg* 60 266-269
 - 31 KISSLING R E R W CHAMBERLAIN M F FIDSON R K SIKES & M A BLCCA 1954 Studies on the North American arthropod borne encephalitis. II Eastern equine encephalitis in horses. *Am J Hyg* 60 237-250
 - 32 SUDIA W D D D STAMM R W CHAMBERLAIN & R F KISSLING 1956 Transmission of eastern equine encephalitis to horses by *Aedes triseriatus* mosquitoes. *Am J Trop Med Hyg* 5 802-803
 - 33 HOLDEN P 1955 Transmission of eastern equine encephalomyelitis in ring-necked pheasants. *Proc Soc Exptl Biol Med* 88 607-610
 - 34 HAMMON W MCD W C REEVES & G E SATHER 1951 Western equine and St. Louis encephalitis viruses in the blood of experimentally infected wild birds and epidemiological implications of findings. *J Immunol* 67 357-366
 - 35 KISSLING R E R W CHAMBERLAIN W D SUDIA & D D STAMM 1957 Western equine encephalitis in wild birds. *Am J Hyg* 66 48-55
 - 36 HAMMON W MCD & W C REEVES 1943 Laboratory transmission of western equine encephalomyelitis virus by mosquitoes of the genera *Culex* and *Culiseta*. *J Exptl Med* 78 425-434
 - 37 STAMM D D & R F KISSLING 1957 Experimental western equine encephalitis in horses. To be published
 - 38 HOLDEN P 1955 Recovery of western equine encephalomyelitis virus from naturally

Stamm has demonstrated that a 4 week old CIW strain of laboratory mice has a demonstrable viremia for 3 days following subcutaneous inoculation with titers ranging from 10^{-3} to 10^{-7} I D₅₀. However these latter rodents represent an artificial situation having been bred and selected for special susceptibility to infection with the neurotropic viruses.

Birds have been shown to circulate fairly small amounts of SLL virus after experimental infection. Hammon, Keesee and Suther² showed that the virus could be detected in the blood of house finches, English sparrows, tricolored red winged blackbirds and white crowned sparrows. In all except the white crowned sparrows virus was demonstrated only in the undiluted blood or rarely in the 10^{-1} dilution. The white crowned sparrows however showed viremia titers as high as $10^{1.5}$ and $10^{1.8}$ LD₅₀. The antibody response was negligible during the first 2 months after inoculation. Chamberlain *et al.*⁴ followed the response of individual birds infected by both subcutaneous inoculation and by the bite of infected mosquitoes. Cowbirds, red winged blackbirds, English sparrows and the domestic pigeon were the species included in the study. The pigeons and a portion of the cowbirds and red wings were tested by inoculating a 10^{-1} dilution of the blood into mice. Of 15 birds so tested, none of the subcutaneously inoculated cowbirds or red wings showed a demonstrable viremia in this diluted blood. One pigeon showed traces of virus in the 10^{-1} dilution of blood on the fourth and fifth days following infection and the other on the seventh day. One of 3 red wings infected by mosquito bite showed a trace of virus in diluted blood on the first day, as did a second individual on the seventh day. One of 2 cowbirds similarly infected showed a titer of $10^{1.5}$ I D₅₀ on the fourth day. A second series of birds was inoculated subcutaneously and the blood tested as undiluted material in mice for detection of virus. Three red wings circulated virus intermittently for 5 to 8 days following inoculation; usually insufficient virus was present to kill all mice inoculated. The highest titer observed in the red wings was $10^{1.5}$ LD₅₀ found in one bird on the sixth day. Of 4 cowbirds similarly tested, 2 failed to have any demonstrable virus in the undiluted blood for 14 days subsequent to inoculation. One cowbird had a trace of virus in the undiluted blood through 7 days after inoculation while the fourth bird had a maximum titer of $10^{1.1}$ LD₅₀ on the sixth day. One of 8 English sparrows also tested by inoculation of undiluted blood into mice failed to show a demonstrable viremia during the first 7 days after inoculation. One sparrow had a trace of virus in its blood on the seventh day, 1 a trace on the fifth day, and another had traces of blood virus on the first 2 days after inoculation. The remaining 4 sparrows had maximum virus titers of $10^{1.3}$ LD₅₀ to $10^{-0.5}$ LD₅₀ occurring on the fourth and fifth days after inoculation. These 4 birds had viremic periods lasting for from 4 to 6 days. Antibody response in all except the pigeons was either absent or very weak when tested at 25 to 30 days after inoculation.

All of the previously mentioned experimental infections of birds with SLE virus were done using viruses isolated in California. A strain of SLE virus isolated from a naturally infected bird from Kentucky produced somewhat higher viremia levels in cowbirds.⁵ Three birds inoculated subcutaneously had viremia for 7 days following inoculation. Two of these had titers above

HOST RELATIONSHIP OF THE ARTHROPOD BORNE ENCEPHALITIS VIRUS

By Robert F. Kissling

Communicable Disease Center Public Health Service Montgomery Ala

One of the distinguishing characteristics of the arthropod borne viruses is that of a fairly wide host range. By definition these viruses are capable of infecting both invertebrates and vertebrates. Among the vertebrates many species of mammals and birds may be infected with these viruses. However each virus tends to have one class that will furnish the more important hosts.

The importance of a vertebrate as a host for the encephalitis viruses depends upon several factors. First the individual must circulate virus in its blood in sufficient concentration and for a reasonable length of time to allow the invertebrate vector to become infected by ingesting the virus containing blood meal. Moreover the vertebrate host to be of consequence in the continuing life cycle of the virus must choose a habitat also suitable for the invertebrate vector and by its habits allow sufficient opportunity for the arthropod to obtain blood meals.

When the principal vectors are determined many otherwise potential vertebrate hosts can be dismissed because their preferred habitat does not coincide with that of the vector. In the case of eastern equine encephalitis (EEE) in which the principal endemic vector appears to be *Culiseta melanura* (a swamp-breeding mosquito) certain upland game birds, burrowing rodents or any other vertebrate preferring a relatively dry habitat could not enter into the endemic cycle of the virus. Also since *C. melanura* is preferentially a bird feeder mammals such as otters and swamp rabbits even when they choose a swamp habitat are not likely to become involved in the cycle of EEE infection. In fact serologic studies on small mammals resident in swamps endemic for EEE have failed to show any evidence of experience with the virus.

Although serologic surveys indicate which individuals are exposed to the vector and which are capable of being infected by the virus this information does not tell us which species can serve as hosts capable of passing the virus to additional vectors.

St. Louis encephalitis (SLE) virus is found at very low levels in the blood of any species during the viremic phase of infection. This is compensated however by the fact that the arthropod vectors are able to become infected on blood meals containing very small amounts of virus.

Cox, Philip and Kilpatrick¹ found no virus circulating in the blood of 3 horses inoculated intracerebrally with SLE virus. Stamm found that only 1 of 4 cottontail rabbits circulated SLE virus after subcutaneous inoculation. This 1 animal had a minimal amount of virus present in its blood during a 3 day period after inoculation. Only 1 of 5 mice died when inoculated with the 1:2 dilution of blood. Three of the 4 rabbits developed neutralizing antibody. Six roof rats (*Rattus rattus*) were also inoculated subcutaneously with SLE virus. None showed demonstrable virus in a 10^{-4} dilution of blood during the 4 days after inoculation nor did antibody develop in any of the rats.

In some areas of the country chickens have a high rate of natural antibody to WEE while in other areas where the virus is also present negligible numbers possess antibody.

In the case of WEE virus it again appears that birds are the primary hosts of the virus wild birds being especially incriminated. Domestic poultry may play a secondary role depending upon the vectors' accessibility to them.

Eastern equine encephalitis (LEE) virus has a somewhat wider host range among mammals than either SLE or WEE viruses. Birds appear to be uniformly susceptible.

Horses have been found to circulate EEE¹¹⁻¹⁴ virus in their blood after experimental infection. The virus appears with a fair degree of regularity but its presence is of short duration usually for 1 to 3 days only. The maximum virus titers observed have ranged from $10^{0.4}$ to $10^{6.2}$ LD₅₀. Titers in the range of 10^3 LD₅₀ are the maximum usually observed. Horse-to-horse transmission through the bite of mosquitoes has been shown experimentally¹¹⁻¹⁴ but in at least one case the infecting horse displayed an unusually high titer of blood virus at the time of the mosquito feeding.

Davis¹⁵ has shown that mourning doves, red-winged blackbirds, grackles, cowbirds, pigeons, and English sparrows have a viremia period following inoculation with EEE virus. Kissling and his co-workers¹⁶⁻¹⁸ added the white ibis, American egret, snowy egret, cardinal, and pheasant to the list of birds susceptible by subcutaneous inoculation. Stamm¹⁹ has found that the yellow-crown night heron, white-throated sparrow, blue jay, tufted titmouse, meadow lark, song sparrow, towhee, hermit thrush, and brown thrasher are also susceptible. In short, no bird so far tested that is free of antibody to the virus of EEE has failed to respond to inoculation of virus with a period of viremia.

The usual period of viremia in birds infected with EEE virus is 4 days. The maximum blood virus titers obtained vary with the species of bird. Large birds such as the herons and pheasants will have titers reaching from 10^7 to $10^{8.9}$ LD₅₀. The smaller birds such as grackles, cardinals, and sparrows will have maximum viremia titers from $10^{6.0}$ to $10^{8.0}$ LD₅₀.

Domestic poultry have fairly low viremia titers, and serologic surveys of chickens in endemic and epidemic areas have shown only an occasional individual with antibody to EEE, indicating very little association with the vectors of the virus.

Serologic surveys have limited the natural endemic hosts of EEE to birds, more especially wild birds. Of the wild birds, the smaller species are more efficient hosts than are the larger birds, as shown by the height of viremia that develops during infection. Also, small birds are more numerous and occur in greater concentration. Horses may occasionally enter into the epidemic maintenance of EEE and since they are closely associated with humans, they could assume some importance as an indirect source of human infection.

Venezuelan equine encephalitis (VEE) virus is one of the arthropod-borne viruses that seem to multiply with greater efficiency in mammals than in birds.

Horses infected experimentally with VEE virus circulate virus for 4 to 6 days.²⁰ Maximum blood virus titers were found to range from $10^{4.0}$ to $10^{6.0}$.

$10^{1.0}$ LD₅₀ Fourteen English sparrows also were inoculated with the Kentucky virus, using a small dosage. These birds were tested for viremia on the third day after inoculation and then at 6- to 7-day intervals through the twenty third day. At 3 days 8 of the birds were viremic. At 9 days 4 of these were still viremic and an additional previously negative bird had viremia at this time. At 16 and 23 days 1 bird still had viremia when this bird died on the twenty seventh day, SLE virus was isolated from the spleen and kidneys but not from the brain or liver.

Chickens have been observed to develop a high rate of antibody to SLE virus during the course of an epidemic. Experimental studies on chickens show that the viremia levels of adults are comparable to those found in wild birds; exceptionally high titers are observed in young birds.

Thus even with the limited information available we are able to narrow the list of important vertebrate hosts of SLE to birds, with special emphasis on the smaller perching birds such as sparrows and finches. Domestic poultry in areas where the principal vector of SLE is *Culex pipiens quinquefasciatus* are also severely suspect as important epidemic hosts since they are a preferred host for the vector.

Western equine encephalitis (WEE) virus appears to multiply to higher levels in the blood of birds than does SLE virus and, in addition, it causes clinical disease in more species of mammals than does SLE virus.

It has been found difficult to produce clinical disease of WEE in horses by subcutaneous inoculation of virus.⁶ The viremia levels in horses so inoculated are very low and fleeting. Usually virus is not present in sufficient quantities to infect mice by intracerebral inoculation and it has been necessary to resort to the inoculation of one half day old chicks⁷ to demonstrate the traces of virus which are present in the blood of horses during the viremic phase.

Other mammals have not been investigated as far as WEE viremia levels are concerned.

Using a California strain of WEE virus Hammon *et al.*⁸ found that viremia was present in English sparrows, house finches, tricolored red winged black birds, and white crowned sparrows following subcutaneous inoculation of virus. Although individual birds were not followed daily, virus was recovered from some individuals as late as 9 and 10 days after inoculation. Most recoveries of virus were made during the first 4 days after inoculation. The maximum titers of $10^{4.0}$ to $10^{6.0}$ LD₅₀ were found during the first 2 days after inoculation. Kissling *et al.*⁸ followed the daily viremia patterns of individual birds including towhees, cardinals, English sparrows, white throated sparrows, blue jays, and purple finches after subcutaneous inoculation with a WEE virus isolated from Louisiana. Viremia was detected in each species with maximum virus titers comparable to those observed by Hammon and his co-workers.⁸ The periods of demonstrable viremia ranged from 2 to 6 days, the majority of the birds being viremic for 4 days. Kaplan, Winn, and Palmer⁹ found that the common pigeon has blood virus levels and a viremic period similar to those of the other species of birds tested. Domestic fowl have been shown to circulate sufficient virus in their blood to allow infection of *Culex tarsalis*.¹⁰ The blood virus levels in WEE infected chickens are generally lower than those found in wild birds.

In some areas of the country chickens have a high rate of natural antibody to WEF while in other areas where the virus is also present negligible numbers possess antibody.

In the case of WEE virus it again appears that birds are the primary hosts of the virus wild birds being especially incriminated. Domestic poultry may play a secondary role depending upon the vectors accessibility to them.

Eastern equine encephalitis (EEF) virus has a somewhat wider host range among mammals than either SLE or WEF viruses. Birds appear to be uniformly susceptible.

Horses have been found to circulate EEE^{11, 12} virus in their blood after experimental infection. The virus appears with a fair degree of regularity but its presence is of short duration usually for 1 to 3 days only. The maximum virus titers observed have ranged from $10^{2.5}$ to $10^{3.5}$ LD₅₀. Titers in the range of $10^{2.0}$ LD₅₀ are the maximum usually observed. Horse to-horse transmission through the bite of mosquitoes has been shown experimentally^{11, 14} but in at least one case the infecting horse displayed an unusually high titer of blood virus at the time of the mosquito feeding.

Davis¹⁵ has shown that mourning doves, red-winged blackbirds, grackles, cowbirds, pigeons, and English sparrows have a viremia period following inoculation with EEE virus. Kissling and his co-workers^{16, 17} added the white ibis, American egret, snowy egret, cardinal, and pheasant to the list of birds susceptible by subcutaneous inoculation. Stamm¹⁸ has found that the yellow-crown night heron, white-throated sparrow, blue jay, tufted titmouse, meadow lark, song sparrow, towhee, hermit thrush, and brown thrasher are also susceptible. In short, no bird so far tested that is free of antibody to the virus of EEE has failed to respond to inoculation of virus with a period of viremia.

The usual period of viremia in birds infected with EEE virus is 4 days. The maximum blood virus titers obtained vary with the species of bird. Large birds such as the herons and pheasants will have titers reaching from $10^{2.0}$ to $10^{3.0}$ LD₅₀. The smaller birds such as grackles, cardinals, and sparrows will have maximum viremia titers from $10^{1.0}$ to $10^{2.0}$ LD₅₀.

Domestic poultry have fairly low viremia titers, and serologic surveys of chickens in endemic and epidemic areas have shown only an occasional individual with antibody to EEE indicating very little association with the vectors of the virus.

Serologic surveys have limited the natural endemic hosts of EEE to birds, more especially wild birds. Of the wild birds the smaller species are more efficient hosts than are the larger birds, as shown by the height of viremia that develops during infection. Also, small birds are more numerous and occur in greater concentration. Horses may occasionally enter into the epidemic maintenance of EEE and since they are closely associated with humans they could assume some importance as an indirect source of human infection.

Venezuelan equine encephalitis (VEE) virus is one of the arthropod-borne viruses that seem to multiply with greater efficiency in mammals than in birds.

Horses infected experimentally with VEE virus circulate virus for 4 to 6 days.⁹ Maximum blood virus titers were found to range from $10^{4.0}$ to $10^{7.5}$.

LD₅₀ These high titers were maintained for longer periods of time than were the maximum blood virus titers in horses infected with EEE virus. No difficulty was experienced in infecting certain species of mosquitoes by allowing them to feed upon horses in the viremic stage.

Very little has been done with other species of mammals but it has been shown that the usual laboratory mammals are quite susceptible to peripheral inoculation of virus and at least in guinea pigs viremia titers approximate those found in horses.

Wild birds especially those native to North America when infected with VEE virus do not have viremia levels as high or of as long duration as horses.²¹ In fact occasional individuals especially doves and pigeons seem to be refractory to infection. Birds in the viremic phase of the disease seldom had sufficient blood virus present to allow mosquito infection. VEE viremia periods in these birds rarely exceeded 3 days.

The United States probably has little to fear concerning VEE since it is apparently dependent upon a mammal reservoir. Such a disease would have difficulty in becoming permanently established in areas where the mammalian wildlife has been reduced to the low point now prevailing in the United States. Also due to the restricted movement of mammals as compared with that of birds such a disease would probably remain very limited geographically should it be introduced.

In summary the information now available seems to indicate that mammals do not circulate enough virus in their blood during the viremic phase of the disease to be of any importance as a source of virus for epidemic arthropod vectors of SLE or WEE viruses. Horses may occasionally provide an infective blood meal for mosquitoes during the viremic phase of EEE virus infection. However the blood of horses infected with VEE virus is an excellent source of virus for mosquitoes. Birds are excellent sources of virus for mosquitoes when infected with SLE, WEE, and EEE viruses but are rather poor hosts in this respect during the viremic phase of VEE infection.

Although many or perhaps all species of birds are capable of circulating the encephalitis viruses in quantities sufficient for the infection of mosquitoes other factors must be considered in assaying the relative importance of any particular species. These factors are those characteristics or habits that allow exposure to the virus. Thus it has been observed that gregarious species such as white ibis and yellow crowned night herons show a high rate of antibody to EEE virus.^{6, 15} In addition to other factors such as longevity which could account for this observation the heavy concentration of young susceptible birds in a small area during the nesting season no doubt offers an optimum opportunity for the virus to become disseminated. The gregariousness of a species must however be associated with the selection of a suitable habitat since it has been observed that gulls or even herons nesting on coastal islands show very little or no antibody to EEE. Although these latter birds are exposed to many mosquitoes the species are those that may serve as epidemic vectors rather than those species found in endemic situations. If the virus were introduced under these conditions such birds could add an important

impetus to the spread of an epidemic but they are of little or no importance in the long term survival of the virus

The selection of a particular type of habitat is also of importance in smaller species. During endemic times isolations of EEE virus have been confined to such birds as thrushes, catbirds, and grackles, species preferring moist swampy woodland. Under epidemic conditions isolations of EEE have been made from English sparrows, pigeons, wild pheasants, and other such birds, species that are normally found in more open areas. Surveillance for infection in these latter species could probably furnish warning for an incipient epidemic in humans or horses.

Whether or not these viruses may acquire properties that favor epidemic spread or whether the occurrence of an epidemic is dependent solely upon the favorable coincidence of virus, heavy mosquito populations, and a dense susceptible bird population is not definitely known. It is interesting, however, that a strain of EEE virus isolated under an endemic condition produced in apparent infections and maximum viremia titers of $10^{2.5}$ to $10^{3.5}$ I.D.₅₀ in pheasants, while a strain isolated from pheasants during an epidemic period produced maximum viremia titers in pheasants of $10^{4.2}$ to 10^5 I.D.₅₀, accompanied by clinical disease. Likewise, a strain of SLE virus isolated from a bird during an epidemic in Kentucky produces higher viremia titers in birds than does a California strain isolated during an endemic period.

In the experimental infections of birds observed at the Montgomery laboratory, virus has never been recovered from birds later than two weeks after the initial infection, except in two instances, both of which were SLE infections. One would expect that a second viremia would be of a fleeting nature, cut short by a rapid antibody recall, and that, as such, it would offer very limited opportunities for the infection of mosquitoes. However, mosquito transmission may not be the only method of transfer of virus from bird to bird, since transmission of EEE occurs with ease among captive pheasants even in the absence of arthropod vectors. The poor antibody response in birds following SLE infection should encourage the exploration of the latency problem in this particular disease. In the cases of EEE and WEE, however, it has been our experience that antibody response in birds is rapid following the termination of viremia and persists for long periods.

Working on the hypothesis that the latent virus may be masked in some form, a limited experiment was carried out in which the tissues of sparrows which had recovered from SLE infections were grown in tissue cultures. In this manner it would be possible to free the tissues of antibody, and the rapid proliferation of tissue might encourage a build up of virus. These experiments met with negative results.

However, the question of latency in the vertebrate host cannot be dismissed since many possible mechanisms have yet to be explored.

Thus far the vertebrate hosts have been neglected in the application of control measures for the encephalitides, yet in the absence of a suitable host population the virus could not exist.

Due to the type of habitat in which EEE virus is found, it is unlikely that

mosquito abatement could ever be practicable for the control of this disease. It is only under unusual circumstances that a human infection or even a horse infection would be expected to result from an exposure to mosquitoes feeding on swamp inhabiting birds. The birds that live in proximity to human habitations are the more likely sources for virus that infects man and his domestic animals. Therefore an effort to reduce the populations of certain pest birds such as English sparrows, starlings, and pigeons would be in order whenever this disease threatens.

Much more must be known concerning the identity of the endemic hosts for SLF virus. However, the severity of epidemics of SLF in urban areas of the central United States might be lessened by population control of the less desirable birds that serve as epidemic hosts of the virus.

For any of the arthropod borne encephalitides, a combined attack on both the vectors and epidemic hosts should reduce the activity of the virus to a level below that rendered potentially possible by efforts directed toward only one member of the infection cycle. More information is needed concerning the bird population concentration that is needed to support epizootic activity of these viruses.

References

- 1 COX H R C B PHILIP & J W KILPATRICK 1941 Susceptibility of horses to St Louis encephalitis virus. *Public Health Repts U S* 56 27 1391-1397
- 2 STAMM D D Susceptibility of rodents to St Louis encephalitis virus. To be published
- 3 HAMMON W McD W C REEVES & G L SATHER 1951 Western equine and St Louis encephalitis viruses in the blood of experimentally infected wild birds and epidemiological implications of findings. *J Immunol* 67 357-367
- 4 CHAMBERLAIN I W R E KISSLING D D STAMM & W D SLIDIA 1957 Virus of St Louis encephalitis in three species of wild birds. *Am J Hyg* 65 110-118
- 5 STAMM D D Unpublished data
- 6 STAMM D D W D SLIDIA R W CHAMBERLAIN & R E KISSLING Experimental western equine encephalitis in horses. To be published
- 7 CHAMBERLAIN R W R K SIKES & R E KISSLING 1954 Use of chicks in eastern and western equine encephalitis studies. *J Immunol* 73 106-114
- 8 KISSLING I E I W CHAMBERLAIN W D SLIDIA & D D STAMM 1957 Western equine encephalitis in wild birds. *Am J Hyg* 66 48-55
- 9 KAPLAN W J F WINN & D F PALMER 1955 Susceptibility of the pigeon (*Columba livia*) to infection with western equine encephalomyelitis virus. I Blood virus levels following subcutaneous inoculation. *J Immunol* 76 225-226
- 10 HAMMON W McD & W C REEVES 1943 Laboratory transmission of western equine encephalomyelitis by mosquitoes of the genera *Culex* and *Culiseta*. *J Exptl Med* 78 425-434
- 11 TEN BROECK C & M H MERRILL 1935 Transmission of equine encephalomyelitis by mosquitoes. *Am J Pathol* 11 847
- 12 KISSLING I E R W CHAMBERLAIN M E EIDON R K SIKES & M A BUCCA 1954 Studies on the North American arthropod borne encephalitides. II Eastern equine encephalitis in horses. *Am J Hyg* 60 237-250
- 13 GILTNER L F & M S SHAHAN 1936 The present status of infection, equine encephalomyelitis in the United States. *J Am Vet Med Assoc* 88 363-374
- 14 SLIDIA W D D D STAMM R W CHAMBERLAIN & R E KISSLING 1956 Transmission of eastern equine encephalitis to horses by *Aedes triseriatus* mosquito. *Am J Trop Med* 6 807-808
- 15 DAVIS W A 1940 A study of bird and mosquitoes as hosts for the western equine encephalomyelitis. *Am J Hyg* 32 45-59
- 16 KISSLING R E R W CHAMBERLAIN R K SIKES & M E EIDON 1954 Studies on the North American arthropod borne encephalitides. III Eastern equine encephalitis in wild birds. *Am J Hyg* 60 251-265

- 17 KISSLING K F M F TIPSON & D D STAMM 1954 Transfer of maternal neutralizing antibodies against eastern equine encephalomyelitis virus in birds *J Infectious Diseases* **85** 177-181
- 18 KISSLING K F 1957 Eastern equine encephalomyelitis in pheasants 94th Ann Meeting Am Vet Med Assoc To be published
- 19 STAMM D D Unpublished data
- 20 KISSLING K F K W CHAMBERLAIN D B NELSON & D D STAMM 1956 Venezuelan equine encephalomyelitis in horses *Am J Hyg* **63** 274-287
- 21 CHAMBERLAIN R W K F KISSLING D D STAMM D B NELSON & R A SIKES 1956 Venezuelan equine encephalomyelitis in wild birds *Am J Hyg* **63** 61-273

EQUINE ENCEPHALITIS VACCINE STUDIES IN PHEASANTS UNDER EPIZOOTIC AND PRE EPIZOOTIC CONDITIONS

By Oscar Sussman

Bureau of Veterinary Public Health New Jersey State Department of Health Trenton N J

Daniel Cohen

Communicable Disease Center Public Health Service Atlanta Ga

Jean E C erende

*Bureau of Vital Statistics and Administration New Jersey State Department
of Health Trenton N J*

and

Robert E Kissling

*Virus and Rickettsia Section Communicable Disease Center Public Health
Service Montgomery Ala*

Eastern equine encephalitis (EEE) was first reported in ring necked pheasants by Tyzzer Sellards and Bennett in Connecticut in 1938¹ Beaudette *et al* described 27 outbreaks from 1939 to 1953 occurring in pheasant flocks raised commercially in New Jersey² Beaudette concluded EEE occurs with sufficient frequency to pose a serious economic problem to breeders of ring necked pheasants in New Jersey and perhaps in other highly endemic areas For this reason and because of its public health importance further effort should be made to elucidate the epidemiology of EEE and to devise practical methods for its control³

Prior field trials with respect to the efficacy of an EEE vaccine in pheasants were made by Holden⁴ and by Beaudette⁵ Holden stated with respect to a planned vaccine study in 1953 Results of the immunization procedure and the experimental manipulations of the birds were not conclusive A November storm destroyed the outside rearing pens releasing most of the birds therein Therefore, it was impossible to determine relative attack rates in immunized and control birds Beaudette stated The outcome in nonvaccinated and vaccinated at Forked River in 1946 was rather spectacular in that the disease did not appear in any vaccinated pen In previous studies however Beaudette *et al*⁵ had noted that Proportions in excess of ninety per cent of the population in an individual pen may show signs of disease Other birds of the same age in contiguous pens may escape disease for weeks or perhaps altogether It is therefore possible that birds in the pen in which all were vaccinated were never under viral challenge In the same year Beaudette⁶ reported an outbreak among vaccinated birds on the New Gretna N J farm investigated No actual mortality figures were available

Under controlled laboratory conditions Kissling¹⁰ observed a survival rate of 67 per cent (6/9) in a group of pheasants that were vaccinated with a single

dose of formalized LEE vaccine after challenge with a strain of virus that caused a 10 per cent (7/10) mortality in nonvaccinated pheasants. This apparent protection occurred in spite of the fact that no demonstrable neutralizing antibody appeared following use of a single dose of formalized vaccine.

The work discussed in this paper is an attempt under controlled field conditions to evaluate the usefulness of a formalized chick embryo vaccine in the control of naturally occurring IFE within individual pens of commercially raised pheasants. The study was made during the 1956 epizootic in New Jersey.

Materials and Methods

In August 1956 a commercial pheasant breeding farm at Deans, N. J. with approximately 5000 pheasants had lost more than 1500 birds. The virus of FEE had been isolated by I. R. Beaudette of Rutgers University, New Brunswick, N. J. from some of these birds. FIGURE 1 illustrates the distribution of pens at the farm.

The disease first appeared August 16 in Pen 3 which contained 1800 birds 11 weeks of age (hatched May 28, 1956). By September 4 the epidemic in this pen had subsided after a mortality of 18.9 per cent. The disease next appeared August 22 in Pen 1 (A and B) which contained 1800 birds 13 weeks of age (hatched May 21, 1956). In FIGURE 1 it can be seen that Section A of Pen 1 and Pen 3 are contiguous for a distance of 150 feet. All pens were constructed of chicken wire, were 7 feet high and had supporting posts 12 feet apart. A ceiling of chicken wire or twine netting supported by posts 12 feet apart.




FIGURE 1. Diagram of the layouts of the pheasant farms at Deans and Jamesburg, N. J.

TABLE 1

COMPARISON OF MORTALITY RATES OF PHEASANTS DURING THE FIRST SEVEN DAYS OF AN EPIDEMIC OF EEE IN PEN 1 AND PEN 7 SUMMER OF 1956

L to	First 7 days of outbreak of EEE				
	Living	Dead	Total	Percentage mortality	Standard error of t
Pen 1	1563	237	1800	13.17	± 0.79
Pen 7	524	74	598	12.37	± 1.37
Total	2087	311	2398	12.97	± 0.69

Critical ratio = 0.4995 $\chi^2 = 0.2495$ Interpretation: no difference

feet apart prevented the birds from escaping. At the time of our initial visit on August 27, 1956 238 birds in Pen 1 (A and B) had died.

Adjacent to Pen 1 (A and B) was Pen 2A which contained 860 pheasants (hatched June 26 1956) showing no signs of encephalitis on August 27, 1956. The pheasants in Pens 1, 2, and 3 were raised on the ground.

Pens 4 and 5 were on platforms 4 feet from the ground. The birds kept in them were thus said to be raised on wire and were never in contact with the ground. Pen 4 contained 400 birds (hatched July 4 1956) and Pen 5 contained 100 birds (hatched May 7 1956). Pen 6 (on the ground) contained 600 pheasants retained as breeders from the previous year (1955). Pen 4 was 250 feet from both Pen 1A and Pen 3. Pen 6 was 33 feet from Pen 1A.

On September 19 the disease broke out at a farm near Jamesburg four miles distant from the farm at Deans and separated from it by a swamp (Igeon Swamp). Both the Deans and Jamesburg establishments were located on sandy soil typical of the coastal plain of New Jersey. The Jamesburg farm obtained its pheasants (hatched May 28 1956) as day old chicks from the Deans farm. The Jamesburg pen will be referred to as Pen 7 and is included in the vaccination studies. Statistical analysis of the Jamesburg outbreak (TABLE 1) establishes the comparability of the mortality rate of Pen 7 and the mortality rate in Pen 1 (A and B) at Deans.

The vaccine was utilized under two conditions: (1) in affected Pens 1 (A and B) and 7 seven days after outbreak, and (2) in pre-epidemic Pen 2A and nonaffected Pen 4. Each bird received 0.2 ml of undiluted formolized chick embryo bivalent vaccine* injected intramuscularly into the pectoral region. A repeating type vaccinating syringe was used†. One needle (1½ in. 20 gauge) was used for 50 to 75 birds. In Pens 1A and 2A birds were handled in alternating groups of 10 to 15. The vaccinates were stained with an alcoholic methyl green solution. The nonvaccinated controls were stained with an alcoholic carbol fuchsin solution in both instances on the medial surface of the left wing. The nonvaccinated pheasants were allowed to run with the vaccinated birds in the same area. Birds failing to respond to drives in Pen

This vaccine was generously provided by Samuel Bozeman of Pitman Moor, Fairport, Indianapolis, Ind.

† Pol. Intramuscular Vaccinator produced by Vineland Poultry Laboratory, Vineland, N. J.

1A were classified as 'nonvaccinated—not handled'. In Pen 4 vaccine was injected into all the birds. Pen 5 was maintained as a nonvaccinated control for the birds raised on wire.

Dead birds were picked up daily and the pens in which they died their vaccination status and the date of death were recorded. All heads were frozen for virus isolation. Birds were opened and observed for macroscopic (post mortem) lesions of diseases other than encephalitis. Blood samples were taken from representative birds in all pens at varying intervals.

The procedure employed was to vaccinate the pheasants in the unaffected pens before entering those in which the epidemic was present. The birds first vaccinated were the 400 raised on wire in Pen 4; those in Pen 5 were left unvaccinated as its control. The next group handled comprised the birds in Pen 2A. 530 (61 per cent) of the 860 total number of birds were vaccinated by August '8. On September 19 (21 days later) the breeder moved 402 of the best feathered, least picked birds to Pen 2B. Of this group 243 (60.4 per cent) had been vaccinated. Pen 2B had not been used the previous season and the birds in it were debeaked prior to placing them in the new pen. On September 21 (23 days after vaccination) the disease broke out in Pen 2A. Of the 458 birds that had been left in this pen 280 (61.1 per cent) had been vaccinated.

Before vaccinating in Pen 1 where an epidemic prevailed the pen was divided into Sections 1A and 1B. Prior to this division there was feed and water in both sections of the pen and the birds passed freely from one part to the other. No attempt was made to count or to select birds for either section. The separation was made by erecting a chicken wire fence from ground to roof netting. Four hundred fifty (450) birds were left in Pen 1B to serve as nonvaccinated controls for Pen 1A. In Pen 1A there were 1113 pheasants. Birds were driven in successive drives into a catching enclosure. Of this handled group 590 were vaccinated and marked with green dye. 288 were not vaccinated and were marked with red dye. The remaining 785 birds that did not respond to the drives were noted to be in poorer condition than the handled birds. These nonvaccinated birds were subsequently referred to as 'not handled'—no color in the mortality records. In Pen 1A as a whole 48.5 per cent of the 1113 birds were vaccinated on August 29 and 30.

Pen 7 at Jamesburg containing 598 pheasants experienced an outbreak of EEE on September 19. The diagnosis was confirmed by virus isolation at Rutgers University by Beaudette. On September 26 the remaining 324 birds in Pen 7 were vaccinated.

There were thus 4 groups of pens involved.

Group 1 representing 3 proportions of vaccinated to nonvaccinated pheasants 7 days after outbreaks of the disease in the pen: all raised on the ground (A) Pen 1B 0.0 per cent vaccinated (B) Pen 1A 48.5 per cent vaccinated and (C) Pen 7 100.0 per cent vaccinated.

Group 2 consisting of 2 groups of birds vaccinated when no EEE disease was present (A) Pen 2A 61.1 per cent vaccinated (birds raised on the ground) (B) Pen 2B 60.4 per cent vaccinated (birds raised on the ground) and (C) Pen 4 100.0 per cent vaccinated (birds raised on the wire).

Group 3 included 2 pens of nonvaccinated and healthy birds (A) Pen 5 birds on wire and (B) Pen 6 breeders (on the ground)

Group 4 the first pen to be affected in which no attempt had been made to curb the disease (A) Pen 3

Results

Pens vaccinated 7 days after outbreak The daily mortality in Pens 1A 1B and 7 following vaccination is given in TABLE 2. The results are graphically illustrated in FIGURE 2 which shows the survival curves of pheasants plotted according to the actuarial method employed in life tables as described by Berkson and Gage.¹¹ The survival curves of the vaccinated groups level off about the ninth day after vaccination whereas that of the nonvaccinated birds in Pen 1B continued its downward descent to the thirteenth day. The differences in the three are statistically significant (TABLE 3).

FIGURE 3 shows the survival curves within Pen 1A where 48.5 per cent of the 1113 birds available were vaccinated. As previously indicated the non

TABLE 2
TABULATION OF DEATHS AMONG BIRDS IN PENS VACCINATED SEVEN DAYS
AFTER OUTBREAK

N d y afte nati	P 1A					Pe 1B 100% All d d b d	P n 7 100% cc All dead b d
	51.5% n c nated			V 48.5% ted %	T t 11A All d d b d		
	Ha d d d	N t dled l	T tal				
1	8	24	32	10	42	26	11
2	12	14	26	12	38	12	9
3	20	30	50	29	79	30	45
4	23	19	42	26	68	23	18
5	33	37	70	32	102	30	15
6	11	39	50	11	61	26	25
7	6	22	28	7	35	24	16
8	12	27	34	11	45	36	5
9	2	4	6	—	6	38	2
10	—	3	3	2	5	41	3
11	—	5	5	—	5	12	3
12	—	1	1	—	1	10	—
13	—	1	1	—	1	3	2
14	—	1	1	—	1	3	—
15	—	1	1	—	1	2	—
16	—	—	—	—	—	—	1
17	—	—	—	—	—	—	1
20	—	1	1	1	2	—	—
22	2	—	2	—	2	1	—
24	—	—	—	—	—	3	—
Number of dead birds	129	224	353	141	494	320	156
Number of birds at time of vaccination	288	285	573	540	1113	450	524




FIGURE 2 Survival curves of pens of pheasants vaccinated seven days after the outbreak of the epizootic

TABLE 3

NUMBER AND PER CENT OF PHEASANTS LIVING OR DEAD VACCINATED SEVEN DAYS AFTER OUTBREAK OF FEF SUMMER 1956

L cat	Number			Percentage	
	Living	Dead	Total	Percentage	Standard Error
Pen 7 100% vaccinated	368	156	524	29.8	± 2.00
Pen 1A 48.5% vaccinated	619	494	1113	44.4	± 1.49
Pen 1B 0% vaccinated	130	320	450	71.1	± 2.14

$N = 170$ Interpretation: reliable difference

vaccinates that were not handled were in poorer physical condition and their survival rate was low (21.4 per cent). Of the handled birds the vaccinated portion had the highest survival rate (73.9 per cent) whereas the nonvaccinated handled portion had a lower survival rate (55.2 per cent) a statistically significant difference (TABLE 4). The nonvaccinated total group represents both the handled and nonhandled birds.

FIGURE 4 presents survival curves of vaccinated and nonvaccinated portions of the birds in Pen 1A as compared to those of the totally vaccinated birds in Pen 7 and the totally nonvaccinated occupants of Pen 1B. The survival rates of those comprising the vaccinated portion of Pen 1A and the 100.0 per cent vaccinated Pen 7 were similar and TABLE 5 indicates that no statistical difference exists between these two rates. The survival rates for the pheasants in the nonvaccinated portion of Pen 1A and the entire population of Pen 1B in which no birds were vaccinated were statistically different (TABLE 6). It appears that the benefits of the vaccine do not confine themselves merely to birds vaccinated; a more favorable survival rate is found among susceptible birds remaining in the pen. In a disease spread by bird to bird contact the




FIGURE 3 Survival curves of pheasants in Pen 1A vaccinated seven days after the outbreak of the epizootic

TABLE 4
COMPARISON OF MORTALITY RATES OF VACCINATED AND NONVACCINATED HANDLED BIRDS
PEN 1A

Pheasant in P 1A	Living	Dead	Total	Percentage	Standard Error
Vaccinated green	399	141	540	26.1	± 1.9
Nonvaccinated red	159	129	288	44.8	± 2.9
Total	558	270	828	32.6	± 1.6

Critical ratio = 5.2 $\chi^2 = 27.5$ Interpretation: reliable difference

greater the number of protected birds the more difficult it becomes for the virus to reach the decreased number of susceptibles. In birds vaccinated during this epidemic, results indicate that regardless of the proportion of individuals vaccinated, the survival rates in the vaccinated birds remained fairly constant.

FIGURE 5 indicates the per cent occurrence of deaths of pheasants in Pens 1B, 1A, and 7. It will be noted that the peak occurrence of mortality occurred on the third day following vaccination in the 100 per cent vaccinated group (Pen 7), on the fifth day following vaccination in the 48.5 per cent vaccinated group (Pen 1A), and on the tenth day following vaccination in the 0.0 per cent vaccinated group (Pen 1B). Vaccination in the midst of this epidemic shortened the time necessary to reach the peak of epidemic loss and decreased the number of individuals that succumbed.




FIGURE 4—Survival curves of vaccinated and nonvaccinated birds of Pen 1A as compared to those of totally vaccinated birds of Pen 7 and of the totally non-vaccinated birds of Pen 1B

TABLE 5

COMPARISON OF MORTALITY RATES OF VACCINATED PHEASANTS IN PENS 1A AND 7

	L i k	D e d	T o t a l	P e r c e n t m o r t a l i t y	S t a n d a r d d e v i a t i o n
Vaccinated Pen 1A	399	141	540	26.1	± 1.9
Vaccinated Pen 7	368	156	524	29.8	± 2.0
Total	767	297	1064	27.9	± 1.4

Critical ratio = 1.33 $\chi^2 = 1.77$ Interpretation: no difference

TABLE 6

COMPARISON OF MORTALITY RATES OF NONVACCINATED PHEASANTS IN PENS 1A AND 1B

	L i k	D e d	T o t a l	P e r c e n t m o r t a l i t y	S t a n d a r d d e v i a t i o n
Nonvaccinated Pen 1A	220	353	573	61.6	± 2.0
Nonvaccinated Pen 1B	130	320	450	71.1	± 2.1
Total	350	673	1023	65.8	± 1.5

Critical ratio = 3.2 $\chi^2 = 10.1$ Interpretation: different

TABLE 7 and FIGURE 6 summarize the results in this group of birds vaccinated seven days after an outbreak occurred.

Pen vaccinated 23 days prior to outbreak TABLE 8 presents the daily mortality in Pens 2A and 2B. The outbreak first appeared in Pen 2A on September 21—23 days after 61.1 per cent of the birds had been vaccinated. The birds in Pen 2B had been moved from 2A on September 19—2 days prior to epidemic onset in Pen 2A.

FIGURE 7 indicates the survival curves for Pen 2A. The significant dif




FIGURE 3 Survival curves of pheasants in Pen 1A vaccinated seven days after the outbreak of the epizootic

TABLE 4
COMPARISON OF MORTALITY RATES OF VACCINATED AND NONVACCINATED HANDLED BIRDS
PEN 1A

Pheasant Pen 1A	Living	Died	Total	Percentage	Standard Error
Vaccinated green	399	141	540	26.1	±1.9
Nonvaccinated red	159	129	288	44.8	±2.0
Total	558	270	828	32.6	±1.6

Critical ratio = 5.2 $\chi^2 = 27.5$ Interpretation reliable difference

greater the number of protected birds the more difficult it becomes for the virus to reach the decreased number of susceptibles. In birds vaccinated during this epidemic results indicate that regardless of the proportion of individuals vaccinated the survival rates in the vaccinated birds remained fairly constant.

FIGURE 5 indicates the per cent occurrence of deaths of pheasants in Pens 1B, 1A and 7. It will be noted that the peak occurrence of mortality occurred on the third day following vaccination in the 100 per cent vaccinated group (Pen 7) on the fifth day following vaccination in the 48.5 per cent vaccinated group (Pen 1A) and on the tenth day following vaccination in the 0.0 per cent vaccinated group (Pen 1B). Vaccination in the midst of this epidemic shortened the time necessary to reach the peak of epidemic loss and decreased the number of individuals that succumbed.

TABLE 7
NUMBER AND PER CENT OF PHEASANTS LIVING AND DEAD VACCINATED SEVEN DAYS AFTER
OUTBREAK OF EE SUMMER 1966

Locat	Number			Percent			
	Living	Dead	Total	Survival	Mortality	Standard error of rate	Vaccinated
Pen 7	368	156	524	60.2	39.8	± 2.0	100.0
Pen 1B	130	320	450	28.9	71.1	± 2.1	0.0
Pen 1A	619	494	1113	55.6	44.4	± 1.5	48.5
Vaccinated green	399	141	540	73.9	26.1	± 1.9	100.0
Nonvaccinated	220	353	573	38.4	61.6	± 2.0	0.0
Handled red	157	129	286	55.2	44.8	± 2.9	0.0
Not handled no color	61	224	285	21.4	78.6	± 2.4	0.0

TABLE 8
TABULATION OF DEATH AMONG BIRDS VACCINATED 23 DAYS PRIOR TO OUTBREAK

Day following outbreak	Pen 2A			Pen 2B
	Nonvaccinated	Vaccinated	All dead birds	All dead birds
1	4	1	5	1 (nonvaccinate) 2 (nonvaccinates)
2	4	—	4	
3	3	—	3	
4	4	1	5	
5	11	—	11	
6	10	2	12	
7	10	—	10	
8	17	4	21	
9	11	1	12	
10	11	4	15	
11	11	5	16	3
12	8	4	12	
13	4	6	10	
14	5	3	8	
15	3	—	3	
16	1	3	4	
17	2	—	2	
18	3	1	4	
19	1	—	1	
20	—	—	—	
21	1	—	1	
22	—	—	—	
Number of dead birds	124	35	159	3
Number of birds	178	280	458	402

FIGURE 9 summarizes the findings in Pen 2A. Of the 178 nonvaccinated birds, 124 or 69.7 per cent succumbed, whereas of the 280 vaccinated only 36 or 12.5 per cent died.

In Pen 2B three birds, all nonvaccinated, died from the disease. Since




FIGURE 5 Per cent occurrence of deaths in pens in which the birds were vaccinated seven days after the outbreak of the epizootic




FIGURE 6 Survival and mortality rates of birds vaccinated seven days after the outbreak of the epizootic

ference in survival rates between the vaccinated (87.5 per cent) and the non vaccinated portions (30.3 per cent) are readily observable. FIGURE 8 illustrates the percentage of deaths occurring in Pen 2A. The epidemic ran its normal 21 day course and the peak mortality rate occurred on the eighth day of the epidemic for the nonvaccinated group and on the thirteenth day for the vaccinates. Of the vaccinated group the birds succumbing resisted the challenge for a longer interval of time. This would suggest that a vaccine break occurred only after a sufficient virus challenge had been built up within the pen.

TABLE 7

NUMBER AND PER CENT OF PHEASANTS LIVING AND DEAD VACCINATED SEVEN DAYS AFTER OUTBREAK OF EEE SUMMER 1956

Locality	Number			Percent			
	Living	Dead	Total	Survival	Mortality	Standard error of t	Vaccinated
Pen 7	368	156	524	70.2	29.8	± 2.0	100.0
Pen 1B	130	370	500	26.0	74.0	± 2.1	0.0
Pen 1A	619	494	1113	55.6	44.4	± 1.5	48.5
Vaccinated green	399	141	540	73.9	26.1	± 1.9	100.0
Nonvaccinated	220	353	573	38.4	61.6	± 2.0	0.0
Handled red	159	129	288	55.2	44.8	± 2.9	0.0
Not handled no collar	61	224	285	21.4	78.6	± 2.4	0.0

TABLE 8

TABULATION OF DEATHS AMONG BIRDS VACCINATED 23 DAYS PRIOR TO OUTBREAK

Day of outbreak	Pen 2A			Pen 2B
	Nonvaccinated	Vaccinated	All dead birds	All dead birds
1	4	1	5	
2	4	—	4	
3	3	—	3	1 (nonvaccinated)
4	4	1	5	
5	11	—	11	2 (nonvaccinated)
6	10	2	12	
7	10	—	10	
8	17	4	21	
9	11	1	12	
10	11	4	15	
11	11	5	16	
12	8	4	12	
13	4	6	10	
14	5	3	8	
15	3	—	3	
16	1	3	4	
17	2	—	2	
18	3	1	4	
19	1	—	1	
20	—	—	—	
21	1	—	1	
22	—	—	—	
Number of dead birds	124	35	159	3
Number of birds	178	280	458	402

FIGURE 9 summarizes the findings in Pen 2A. Of the 178 nonvaccinated birds 124 or 69.7 per cent succumbed whereas of the 280 vaccinated only 36 or 12.5 per cent died.

In Pen 2B three birds all nonvaccinated died from the disease. Since




FIGURE 7 Survival curves of birds in Pen 2A vaccinated twenty three days prior to the outbreak of the epizootic




FIGURE 8 Per cent occurrence of deaths of birds in Pen 2A vaccinated twenty three days prior to the outbreak of the epizootic

these birds were moved two days before the onset of the epidemic in Pen 2A it is possible that they succumbed to an infection contracted in Pen 2A prior to their transfer. It is difficult to ascertain whether the failure of the disease to reach epidemic proportions in the new Pen 2B should be attributed to (1) the debeaking performed as the birds were transferred (2) the fact that they were placed in a new clean pen or (3) the better condition of the birds at the time they were selected.

No losses due to the disease were noted in Pens 4, 5 or 6. Sera obtained from Pens 5 and 6 indicate negative results in these disease free pens of non vaccinated birds. TABLE 9 summarizes the results for these pens and for Pens 2A and 2B.




FIGURE 9. Survival and mortality rates of birds in Pen 2A vaccinated twenty three days prior to the outbreak of the epizootic

TABLE 9
NUMBER AND PER CENT OF PHEASANTS LIVING AND DEAD IN PENS UNAFFECTED AT TIME OF VACCINATION SUMMER 1956

Locality	Number			Percent			
	Living	Dead	Total	Survival	Mortality	Standard Error	Vitality
Pen 2A	299	159	458	65.3	34.7	2.22	61.1
Vaccinated	245	35	280	87.5	12.5	1.98	100.0
Nonvaccinated	54	124	178	30.3	69.7	3.45	0.0
Pen 2B	399	3	402	99.3	0.7	0.42	60.4
Vaccinated	243	0	243	100.0	0.0	—	100.0
Nonvaccinated	156	3	159	98.1	1.9	1.08	0.0
Pen 4	400	0	400	100.0	0.0	—	100.0
Pen 5	100	0	100	100.0	0.0	—	0.0
Pen 6	600	0	600	100.0	0.0	—	0.0

It is apparent from the results in Pen 2A that vaccination prior to outbreak did in fact offer substantial protection (87.5 per cent survival) to those birds vaccinated. Only under the severe challenge produced by allowing susceptible controls to remain within the pen did some single vaccinated birds break (12.5 per cent). The authors feel that if 100 per cent of the birds in the pen had been vaccinated prior to exposure the viral build up would have been limited and the vaccine breaks would have occurred less frequently if at all. The question of debeaking versus vaccination must be settled by wildlife management personnel and commercial breeders on the basis of ability of the birds to survive in nature upon release and by the economics of the industry.

Summary

The efficacy of vaccine in an epidemic and pre epidemic situation of EEE was evaluated in New Jersey during the 1956 epizootic. The vaccine was proved effective under the conditions of the experiment.

Under controlled conditions pens of pheasants that had undergone the first week of the 3 week disease cycle were vaccinated. The resulting mortality




FIGURE 7 Survival curves of birds in Pen 2A vaccinated twenty three days prior to the outbreak of the epizootic




FIGURE 8 Per cent occurrence of deaths of birds in Pen 2A vaccinated twenty three days prior to the outbreak of the epizootic

these birds were moved two days before the onset of the epidemic in Pen 2A it is possible that they succumbed to an infection contracted in Pen 2A prior to their transfer. It is difficult to ascertain whether the failure of the disease to reach epidemic proportions in the new Pen 2B should be attributed to (1) the detoxing performed as the birds were transferred (2) the fact that they were placed in a new clean pen or (3) the better condition of the birds at the time they were selected.

No losses due to the disease were noted in Pens 4, 5 or 6. Sera obtained from Pens 5 and 6 indicate negative results in these disease free pens of non vaccinated birds. TABLE 9 summarizes the results for these pens and for Pens 2A and 2B.




FIGURE 9 Survival and mortality rates of birds in Pen 2A vaccinated twenty three days prior to the outbreak of the epidemic

TABLE 9

NUMBER AND PER CENT OF PHEASANTS LIVING AND DEAD IN PENS UNAFFECTED AT TIME OF VACCINATION SUMMER 1956

Location	Number			Percent			
	Living	Dead	Total	Survival	Mortality	Standard Error	Vaccinated
Pen 2A	299	159	458	65.3	34.7	2.27	61.1
Vaccinated	245	35	280	87.5	12.5	1.98	100.0
Nonvaccinated	54	124	178	30.3	69.7	3.45	0.0
Pen 2B	329	3	402	92.3	0.7	0.42	60.4
Vaccinated	243	0	243	100.0	0.0	—	100.0
Nonvaccinated	156	3	159	98.1	1.9	1.08	0.0
Pen 4	400	0	400	100.0	0.0	—	100.0
Pen 5	100	0	100	100.0	0.0	—	0.0
Pen 6	600	0	600	100.0	0.0	—	0.0

It is apparent from the results in Pen 2A that vaccination prior to outbreak did in fact offer substantial protection (87.5 per cent survival) to those birds vaccinated. Only under the severe challenge produced by allowing susceptible controls to remain within the pen did some single vaccinated birds break (12.5 per cent). The authors feel that if 100 per cent of the birds in the pen had been vaccinated prior to exposure the viral build up would have been limited and the vaccine 'breaks' would have occurred less frequently if at all. The question of debeaking versus vaccination must be settled by wildlife management personnel and commercial breeders on the basis of ability of the birds to survive in nature upon release and by the economics of the industry.

Summary

The efficacy of vaccine in an epidemic and pre epidemic situation of EEE was evaluated in New Jersey during the 1956 epizootic. The vaccine was proved effective under the conditions of the experiment.

Under controlled conditions pens of pheasants that had undergone the first week of the 3 week disease cycle were vaccinated. The resulting mortality




FIGURE 7 Survival curves of birds in Pen 2A vaccinated twenty three days prior to the outbreak of the epizootic




FIGURE 8 Per cent occurrence of deaths of birds in Pen 2A vaccinated twenty three days prior to the outbreak of the epizootic

these birds were moved two days before the onset of the epidemic in Pen 2A it is possible that they succumbed to an infection contracted in Pen 2A prior to their transfer. It is difficult to ascertain whether the failure of the disease to reach epidemic proportions in the new Pen 2B should be attributed to (1) the debeaking performed as the birds were transferred (2) the fact that they were placed in a new clean pen or (3) the better condition of the birds at the time they were selected.

No losses due to the disease were noted in Pens 4, 5 or 6. Sera obtained from Pens 5 and 6 indicate negative results in these disease free pens of non-vaccinated birds. TABLE 9 summarizes the results for these pens and for Pens 2A and 2B.

Discussion of the Paper

S. A. GOLDBERG (*Jewish Hospital, Newark, N. J.*) While examining the late D. H. Udall's collection of slides from an enzootic disease of horses in Kansas thought to be caused by a certain toxin from a plant in the pastures it was noted that the principal lesion in the brain was perivascular lymphocytic infiltration mantling which is similar to the lesion in encephalitis.

Because lymphocytes are extremely radiosensitive it was thought that radiotherapy should be useful in the treatment of acute encephalitis.

To verify this hypothesis mice were inoculated intranasally with the St. Louis strain of encephalitis virus. Treatment consisted of one fourth erythema dose started one day after inoculation.

Of 38 mice treated the incubation period varied from 6 to 11 days. Nine survived. Of the others 1 survived 6 days, 4 survived 8 days and 14 survived from 9 to 11 days.

Of 53 mice similarly inoculated but not treated the period of incubation varied between 5 and 8 days. All but 1 lived only 1 day after symptoms appeared. Only 1 survived.

Of 7 human cases treated all recovered and none developed postencephalitic Parkinson's syndrome after 10 years of observation.

Portman and Lough tested this treatment on 49 cases of acute encephalitis. Of these 59 per cent were considered recovered and 30 per cent improved.

References

- GOLDBERG, S. A., M. BRODIE & I. STANLEY. 1935. Effect of X-ray on experimental encephalitis in mice inoculated with the St. Louis strain. *Proc. Soc. Exptl. Biol. Med.* **32**: 537-500.
- GOLDBERG, S. A., C. F. BAKER & J. W. HERFF. 1934. The rationale of X-ray treatment in encephalitis lethargica. *Radiology* **22**(6): 663-667.
- PORTMAN, U. A. & R. LOUGH. 1945. Roentgen therapy for acute encephalitis. *Am. J. Roentgenol. Radium Therapy* **53**(6): 597.

rates were as follows: 29.8 per cent (100 per cent vaccinated), 44.4 per cent (48.5 per cent vaccinated), 71.1 per cent (0 per cent vaccinated). Analysis of the results within the 48.5 per cent pens of vaccinated birds indicates that the vaccine protected the vaccinated birds and, to a degree, the nonvaccinated birds as well. The use of vaccine influenced the epidemic curve so that the peak of mortality occurred earlier in pens of vaccinated birds as opposed to the pens in which the birds were nonvaccinated.

The utilization of a vaccine on 61.1 per cent of the birds in a pen 23 days prior to an outbreak indicated that the vaccine produced sufficient immunity in that period to protect 87.5 per cent of the birds vaccinated as opposed to a survival rate of 30.3 per cent of the nonvaccinated controls. A study of daily mortality indicates that the peak mortality in the vaccinates occurred in the second week of the 3 week epidemic as opposed to a first week peak noted in the susceptible controls. This suggests that a "vaccine break" occurred only after a sufficient virus challenge had been built up within the pen.

Acknowledgments

The authors gratefully acknowledge the assistance rendered this project by William C. Carter and Robert F. Goldsboro of the New Jersey State Department of Health; Milton C. Clark and Olive and Evelyn Oberman, owners of the Deans farm; Foster Abele, owner of the Jamesburg farm; A. Heaton Underhill and Lester C. MacNamara of the New Jersey Division of Fish and Game; Fred R. Beaudette (deceased) and David C. Tudor, Viral Laboratory of Rutgers University; Raymond Fagan of the Wyeth Research Laboratory; Richard E. Shope of The Rockefeller Institute for Medical Research, New York, N. Y.; and Marguerite F. Hall, Carl E. Weigle and Elmer L. Shaffer, all of the New Jersey State Department of Health.

References

1. TYZZER, F. E., A. W. SELLARDS & B. L. BENNETT. 1938. The occurrence in nature of equine encephalomyelitis in the ring-necked pheasant. *Science* 86(2291): 505-506.
2. BEAUDETTE, F. R. 1940. Neutralization tests with the sera of pheasants surviving an outbreak of equine encephalomyelitis. *Proc. 3rd Intern. Congr. Microbiol.* 93.
3. BEAUDETTE, F. R. 1940. Equine encephalomyelitis in avian hosts. *Proc. 43rd Ann. Meeting U. S. Livestock San. Assoc.* Dec. 6-8, 1939: 185-201.
4. BEAUDETTE, F. R., J. J. BLACK & C. B. HUDSON. 1941. Another case of equine encephalomyelitis in New Jersey pheasants. *J. Am. Vet. Med. Assoc.* 98: 447-451.
5. BEAUDETTE, F. R. & C. B. HUDSON. 1945. Additional outbreaks of equine encephalomyelitis in New Jersey pheasants. *J. Am. Vet. Med. Assoc.* 107: 384-386.
6. BEAUDETTE, F. R. & J. J. BLACK. 1948. Equine encephalomyelitis in New Jersey pheasants in 1945 and 1946. *J. Am. Vet. Med. Assoc.* 112: 140-147.
7. BEAUDETTE, F. R., J. J. BLACK, C. B. HUDSON & J. A. BIVINS. 1952. Equine encephalomyelitis in pheasants from 1941 to 1951. *J. Am. Vet. Med. Assoc.* 121: 478-483.
8. BEAUDETTE, F. R., P. HOLDEN, J. J. BLACK, J. A. BIVINS, C. B. HUDSON & D. C. TUDOR. 1954. Equine encephalomyelitis in pheasants in 1952-53. *Proc. 58th Ann. Meeting U. S. Livestock San. Assoc.* 309-321.
9. HOLDEN, P. 1954. Epidemiological studies on EEE in ring-necked pheasants in a localized area, New Jersey, 1953. Doctoral Thesis, Univ. Pittsburgh, Pittsburgh, Pa.
10. KISSLING, R. F. Eastern equine encephalomyelitis in pheasants. 94th Ann. Meeting Am. Vet. Med. Assoc. To be published.
11. BRAXSON, J. & R. P. GAGG. 1950. Calculation of survival rates for cancer. *Proc. Staff Meetings Mayo Clinic* 25: 270-286.

Discussion of the Paper

S. A. GOLDBERG (*Presbyterian Hospital Newark N. J.*) While examining the late D. H. Udall's collection of slides from an enzootic disease of horses in Kansas thought to be caused by a certain toxin from a plant in the pastures it was noted that the principal lesion in the brain was perivascular lymphocytic infiltration mantling which is similar to the lesion in encephalitis.

Because lymphocytes are extremely radiosensitive it was thought that radiotherapy should be useful in the treatment of acute encephalitis.

To verify this hypothesis mice were inoculated intranasally with the St. Louis strain of encephalitis virus. Treatment consisted of one fourth erythema dose started one day after inoculation.

Of 38 mice treated the incubation period varied from 6 to 11 days. Nine teen survived. Of the others 1 survived 6 days, 4 survived 8 days and 14 survived from 9 to 11 days.

Of 53 mice similarly inoculated but not treated the period of incubation varied between 5 and 8 days. All but 1 lived only 1 day after symptoms appeared. Only 1 survived.

Of 7 human cases treated all recovered and none developed postencephalitic Parkinson's syndrome after 10 years of observation.

Portman and Lough tested this treatment on 49 cases of acute encephalitis. Of these 59 per cent were considered recovered and 30 per cent improved.

References

- GOLDBERG S. A., M. BRODIE & I. STANLEY. 1935. Effect of X-ray on experimental encephalitis in mice inoculated with the St. Louis strain. *Proc. Soc. Exptl. Biol. Med.* **32**: 587-590.
- GOLDBERG S. A., C. F. BAKER & J. W. HERTF. 1934. The rationale of X-ray treatment in encephalitis lethargica. *Radiology* **22**(6): 663-66.
- PORTMAN L. V. & R. LOUGH. 1945. Roentgen therapy for acute encephalitis. *Am. J. Roentgenol. Radium Therapy* **53**(6): 517.

Part II Other Problems in Virology

ORPHAN VIRUSES OF MAN AND ANIMALS*

By G. D. Hsiung and Joseph L. Melnick†

Yale University School of Medicine, New Haven, Conn.

INTRODUCTION

The increased use of tissue culture methods in the field of virology has resulted in the discovery of many cytopathogenic agents. These viral agents may be recovered from the tissues and excreta of man and the lower animals; many of them defy detection by methods other than cell cultivation. Although we have excellent methods for the detection of these newly recognized viruses, we are only beginning to learn about the diseases, if any, that they may produce. These agents have been called 'orphan' viruses because we know so little about the diseases with which they may be associated. As so many of them have been isolated from the intestinal tract, they have been considered with the enteroviruses. It is well known that families of viruses such as those of herpes and the pox group exist in different species, and now it appears that we also have enteroviruses for a number of different animal species, such as enteric cytopathogenic human orphan virus (ECHO), enteric cytopathogenic monkey orphan virus (ECMO), enteric cytopathogenic bovine orphan virus (ECBO), and enteric cytopathogenic swine orphan virus (ECSO).

ECHO Viruses¹

The human enteric viruses to which the orphan viruses belong are not like those enteric bacteria that inhabit the intestinal tract for the life of the individual. The proportion of persons carrying enteric viruses decreases as age increases. Observations to date indicate that the enteric viruses produce only transitory infections, the seriousness of which depends upon the type and strain of virus.

Although some ECHO viruses have been recovered from cases of infant diarrhea and from patients with aseptic meningitis, the epidemiological criteria regarding all of the ECHO viruses as etiological agents of these syndromes have not yet been met. However, certain members of this group have been associated with outbreaks of aseptic meningitis^{2,4} and certain types, notably 5, 6, and 9, and perhaps others, have been isolated from the cerebrospinal fluid. In Europe in 1956, ECHO Type 9 accounted for thousands of cases of aseptic meningitis, often with rash^{5,7} and in 1957 the epidemic spread into the United States. Other ECHO viruses, Types 4 and 16, have also been associated with cases of rash, as well as with aseptic meningitis. The epidemic strains of ECHO Type 9 prevalent in Europe in 1955 and 1956 seem to be antigenic variants of the prototype strain.⁴

In addition to the association of the currently prevalent ECHO Type 9

* The work reported in this paper was supported in part by a grant from The National Foundation for Infantile Paralysis, Inc., New York, N. Y.

† Address as of June 1958: Baylor University College of Medicine, Houston, Texas.

strains with outbreaks of aseptic meningitis one other important fact has come out of these studies namely that certain of these virulent strains after passage in monkey kidney tissue produce paralysis and myositis in infant mice. Mice become paralyzed when inoculated by subcutaneous or other routes and histologically exhibit a myositis similar to that found in animals inoculated with Group A Coxsackie viruses. In view of these mouse pathogenic variants the question must be raised at this time (although it cannot be answered from the available evidence) whether ECHO Type 9 should be classified with the ECHO viruses or with the Coxsackie viruses.

If a scheme of classification rests on the host range of the virus then the host range of the natural virus before laboratory manipulation might well be the criterion of choice. This procedure certainly would make the test for host range easier to carry out than one that required as a preliminary to the test an indefinite number of tissue culture passages. This leads to another problem in that we have isolated a number of Coxsackie A9 and Coxsackie B strains that fail to infect infant mice unless these agents are first passaged in tissue culture. Obviously we are only at the beginning in the study of these agents which ultimately may well be grouped together for purposes of classification as enteroviruses.

NEW DEVELOPMENTS IN TISSUE CULTURE METHODS FOR STUDYING VIRUSES OF MAN AND ANIMALS

Morphologic Characteristics of Plaques

The growth of animal viruses in cells under agar produces areas of necrosis resulting in virus colonies or plaques. Differences in plaque patterns produced by various animal viruses have been noted by many investigators for instance Dulbecco with Newcastle virus and western equine encephalitis virus⁸ Sellers with vesicular stomatitis and foot and mouth disease⁹ and Dubes with different strains of poliovirus¹⁰. Our own previous studies on the enteroviruses had shown^{11, 12} that poliovirus Types I, II, and III produce large circular plaques with clear centers and sharp boundaries within 4 or 5 days after seeding on rhesus monkey (*Macaca mulatta*) kidney cultures (FIGURE 1). Coxsackie virus plaques (FIGURE 2) were also round resembling those of the polioviruses except for their delayed appearance which usually occurs 6 or 7 days after seeding. Apparently healthy cells stained with neutral red could be seen within the area of degeneration; this was especially marked at the periphery of the plaques of Coxsackie A9. In order to demonstrate the slow growing plaques of ECHO viruses it was necessary to adapt Dulbecco's technique⁸ to stoppered bottle cultures¹¹ in which monkey kidney cells are stable for 14 days or longer under agar. As shown in FIGURE 3a, ECHO virus plaques are more irregular in shape than poliovirus plaques and their boundaries are diffuse; exceptions are Types 7 and 12 (FIGURES 3b and 5a) which form plaques that are almost impossible to distinguish from those of the polioviruses. Measles is another virus that was found to produce typical plaques in bottle cultures as shown in FIGURE 4. Minute plaques with sharp boundaries appeared as early as 4 days after inoculation into bottle cultures of *patas* monkey (*Erythrocebus patas*) kidney cells and enlarged although slowly thereafter.



FIGURE 1 Plaque morphology of polioviruses 8 days after seeding of rhesus cultures poliovirus Type I (Mahoney) poliovirus Type II (MEF) and poliovirus Type III (Saukett)



FIGURE 2 Plaque morphology of Coxsackie viruses 8 days after seeding of rhesus cultures Coxsackie viruses A9 (Crig) B1 (Connecticut-5) B3 (Nancy) B4 (Texas-130) and B5 (Faulkner)

Host Cell Susceptibility

Important in the characterization of a virus is the determination of its host range. Thus it has been the general experience that adenoviruses may be more readily isolated in HeLa cells than in monkey kidney cultures while




FIGURE 3 (a) Plaque morphology of representatives of Group A ECHO viruses on rhesus cultures 11 days after seeding: Type 3 (Berar), Type 4 (Pesack), Type 6 (D Meo) and Type 9 (Quigley). (b) Plaque morphology of representatives of the 2 groups of ECHO viruses on rhesus cultures 7 days after seeding: right Group A ECHO Type 13 (Hampill), left Group B ECHO Type 12 (Traas).

the opposite is true for the ECHO viruses. A comparative study on the susceptibility of kidney cells from different monkey species to poliovirus, Coxsackie and ECHO viruses using the plaque method has been reported in detail¹². Of the species investigated, only the kidney cells of rhesus and




FIGURE 4. Plaques of measles virus (17th *patas* passage) in *patas* bottle cultures 11 days after seeding. Dilutions: undiluted tissue culture fluid, 10^{-1} , 10^{-2} , and 10^{-3} .

TABLE 1

COMPARATIVE SUSCEPTIBILITY OF RHESUS AND *PATAS* MONKEY KIDNEY CULTURES TO DIFFERENT VIRUSES

Virus	Concentration of fluid	Average number of plaques per bottle—0.1 ml inoculum	
		Rh	<i>pat</i>
Poliovirus	10^{-8}	8.9	23.5
Coxsackie A9	10^{-6}	18.5	0
Coxsackie B1	10^{-6}	17.0	33.5
ECHO Type 1	$10^{-4.0}$	24.2	0
ECHO Type 7	$10^{-6.0}$	25.3	23.0
Measles	$10^{-6.0}$	0	8.5

At the virus dilutions used in this experiment no plaques were formed; however, in cases where large virus inocula were used, tiny delayed plaques occasionally appeared.

patas the African red grass monkey will be considered here. As shown in TABLE 1, *patas* cells were found to be twice as susceptible as rhesus cells to poliovirus and to Coxsackie B1 virus, and about equal to rhesus cells in susceptibility to ECHO Type 7. In contrast, Coxsackie A9 and ECHO Type 1 failed to form plaques in *patas* cultures. Measles produced distinct plaques on *patas* monolayer, but on rhesus culture it gave an irregular response at best.

Isolation and Identification of ECHO Viruses

If we take advantage of differences in plaque morphology and cell susceptibility of rhesus and *patas* cultures, ECHO viruses can be placed into two

TABLE 2
GROUPING OF ECHO VIRUSES

Group A (<i>patas</i> negative)	
1 Small irregular plaques with hazy centers on rhesus only	ECHO 3 (Berardi strain) 4 6 6 9 11 14 15 16 17 18 19
2 Small irregular plaques with clear centers on rhesus (occasionally tiny delayed plaques on <i>patas</i>)	FCHO 1 13
3 No plaques on either rhesus or <i>patas</i>	FCHO 2 3 5 6 10
Group B (<i>patas</i> positive)	
1 Large circular plaques on rhesus and <i>patas</i>	ECHO 7
2 Small or medium sized plaques on rhesus and tiny delayed plaques on <i>patas</i>	FCHO 8
3 Large circular plaques on rhesus and tiny delayed plaques on <i>patas</i>	ECHO 12

Except for the 6 prime strains and the Berardi strain of Type 3 all viruses listed are prototype strains

groups (TABLE 2) Group A contains ECHO virus Types 1 3 (Berardi strain) 4 6 9 11 13 14 15 16 17 18 and 19 these produce small irregularly shaped hazy plaques in rhesus cultures but are negative in *patas* cultures. Plaques of FCHO Type 1 and ECHO Type 13 are clearer in the centers occasionally with large inocula of these types tiny delayed plaques appear on *patas* cultures. Group B contains ECHO Types 7 8 and 12 these ECHO types are positive in *patas* cultures as well as in rhesus cultures. ECHO Type 7 (FIGURE 5a) produces large circular plaques resembling those of poliovirus in both rhesus and *patas* cultures. ECHO Type 8 (FIGURE 5b) produces smaller and more irregular plaques on rhesus cultures and tiny delayed plaques on *patas* cultures. ECHO Type 12 produces large circular plaques on rhesus but small irregular plaques on *patas* cultures. The prototype strains of ECHO Types 2 3 5 6 and 10 have not produced plaques in tube cultures they were found to be cytopathogenic for rhesus but not for *patas* cells and are considered to be members of Group A. The reasons for the failure of these ECHO viruses to produce plaques are obscure particularly in the case of the Type 6 prototype strain (D Amori) which produces complete cytopathic changes in 48 hours when cells are kept in a fluid medium. Although the cell sheet degenerated under agar when the inoculum of this strain was large no clear plaques have been observed.

Difficulties have been encountered in identifying different strains of ECHO Type 8 isolated from various stool samples. As shown in FIGURE 6a stool No. 269 which was typed as ECHO Type 8 produced small irregular plaques on rhesus cells at 10^{-2} concentration and gave no plaques when an inoculum one thousandfold greater was placed on *patas* cultures. Thus this strain of ECHO Type 8 is like those belonging to Group A. On the other hand stool No. 271 (FIGURE 6b) which was also typed serologically as ECHO Type 8 produced circular plaques on rhesus as well as on *patas* cells thus resembling FCHO Group B viruses. A rhesus tissue culture passage of stool No. 279 (FIGURE 6c) that produced plaques of medium size on rhesus cells but tiny delayed plaques on *patas* cells was grouped as ECHO Type 8 by its biological characteristics as well as by serologic typing.

Presumptive grouping of stool isolates on the basis of plaque morphology




FIGURE 4 Plaques of measles virus (17th *patas* passage) in *patas* bottle cultures 11 days after seeding. Dilutions: undiluted tissue culture fluid, 10^{-1} , 10^{-2} , and 10^{-3} .

TABLE 1
COMPARATIVE SUSCEPTIBILITY OF RHESUS AND *PATAS* MONKEY KIDNEY CULTURES TO DIFFERENT VIRUSES

Virus	Cult. fl. d.	A. sg. numb. f. pl. q. per bottle—	
		Rh	<i>p. t. s.</i>
Poliovirus	$10^{-5.7}$	89	23.5
Coxsackie A9	$10^{-6.7}$	18.5	0*
Coxsackie B1	10^{-4}	17.0	33.5
ECHO Type 1	10^{-5}	24.2	0
ECHO Type 7	10^{-5}	25.3	23.0
Measles	$10^{-3.0}$	0	8.5

At the virus dilutions used in this experiment no plaques were formed; however, in cases where large virus inocula were used, tiny delayed plaques occasionally appeared.

patas the African red grass monkey will be considered here. As shown in TABLE 1, *patas* cells were found to be twice as susceptible as rhesus cells to poliovirus and to Coxsackie B1 virus, and about equal to rhesus cells in susceptibility to ECHO Type 7. In contrast, Coxsackie A9 and ECHO Type 1 failed to form plaques in *patas* cultures. Measles produced distinct plaques on *patas* monolayer, but on rhesus culture it gave an irregular response at best.

Isolation and Identification of ECHO Viruses

If we take advantage of differences in plaque morphology and cell susceptibility of rhesus and *patas* cultures, ECHO viruses can be placed into two

TABLE 2
GROUPING OF ECHO VIRUSES

Group A (<i>patas</i> negative)	
1 Small irregular plaques with hazy centers on rhesus only	ECHO 3 (Berardi strain)
4 6 9 11 14 15 16 17 18 19	
2 Small irregular plaques with clear centers on rhesus (occasionally tiny delayed plaques on <i>patas</i>)	ECHO 1 13
3 No plaques on either rhesus or <i>patas</i>	ECHO 2 3 5 6 10
Group B (<i>patas</i> positive)	
1 Large circular plaques on rhesus and <i>patas</i>	ECHO 7
2 Small or medium sized plaques on rhesus and tiny delayed plaques on <i>patas</i>	ECHO 8
3 Large circular plaques on rhesus and tiny delayed plaques on <i>patas</i>	ECHO 12

Except for the 6 prime strains and the Berardi strain of Type 3 all viruses listed are prototype strains

groups (TABLE 2). Group A contains ECHO virus Types 1 3 (Berardi strain) 4 6 9 11 13 14 15 16 17 18 and 19 these produce small irregularly shaped hazy plaques in rhesus cultures but are negative in *patas* cultures. Plaques of ECHO Type 1 and ECHO Type 13 are clearer in the centers occasionally with large inocula of these types tiny delayed plaques appear on *patas* cultures. Group B contains ECHO Types 7 8 and 12 these ECHO types are positive in *patas* cultures as well as in rhesus cultures. ECHO Type 7 (FIGURE 5a) produces large circular plaques resembling those of poliovirus in both rhesus and *patas* cultures. ECHO Type 8 (FIGURE 5b) produces smaller and more irregular plaques on rhesus cultures and tiny delayed plaques on *patas* cultures. ECHO Type 12 produces large circular plaques on rhesus but small irregular plaques on *patas* cultures. The prototype strains of ECHO Types 2 3 5 6 and 10 have not produced plaques in tube cultures they were found to be cytopathogenic for rhesus but not for *patas* cells and are considered to be members of Group A. The reasons for the failure of these ECHO viruses to produce plaques are obscure particularly in the case of the Type 6 prototype strain (D'Amori) which produces complete cytopathic changes in 48 hours when cells are kept in a fluid medium. Although the cell sheet degenerated under agar when the inoculum of this strain was large no clear plaques have been observed.

Difficulties have been encountered in identifying different strains of ECHO Type 8 isolated from various stool samples. As shown in FIGURE 6a stool No. 269 which was typed as ECHO Type 8 produced small irregular plaques on rhesus cells at 10^{-4} concentration and gave no plaques when an inoculum one thousandfold greater was placed on *patas* cultures. Thus this strain of ECHO Type 8 is like those belonging to Group A. On the other hand stool No. 271 (FIGURE 6b) which was also typed serologically as ECHO Type 8 produced circular plaques on rhesus as well as on *patas* cells thus resembling ECHO Group B viruses. A rhesus tissue culture passage of stool No. 249 (FIGURE 6c) that produced plaques of medium size on rhesus cells but tiny delayed plaques on *patas* cells was grouped as ECHO Type 8 by its biological characteristics as well as by serologic typing.

Presumptive grouping of stool isolates on the basis of plaque morphology




FIGURE 5 (a) Plaques of LCHO Type 7 (Wallace) on rhesus (left) and on palus (right) 7 days after seeing (b) Plaques of LCHO Type 8 (Bryson) on rhesus (left) and on palus (right) 7 days after seeing

and cell susceptibility followed by typing with specific antisera was carried out with a total of 70 enteric cytopathogenic agents that had been found to be negative for poliovirus and negative for Coxsackie virus in suckling mice. The results are shown in TABLE 3. Of these 70 strains 6 were identified as Coxsackie viruses despite the fact that they had been negative in suckling mice




FIGURE 6 (a) Plaques of stool No. 269 on rhesus (3 bottles at right dilutions 10^{-1} , 10^{-2} and 10^{-3}) and on *patas* (bottle at left undiluted) 9 days after seeding. (b) plaques of stool No. 271 on rhesus (left) and on *patas* (right) 9 days after seeding. (c) plaques of rhesus kidney passage from stool No. 269 on rhesus (2 bottles at left dilutions 10^{-1} and 10^{-2}) and on *patas* (2 bottles at right dilutions undiluted and 10^{-1}) 7 days after seeding.

TABLE 3
IDENTIFICATION OF ECHO VIRUSES

Source of sample	Number of stool isolates	Number typed as ECHO virus	Number typed as Coxsackie virus	Number unidentified
Phoenix Ariz	34	26	0	8
Charleston W Va	25	20	2	3
Chicago Ill	11	7†	4‡	1
Total	70	53‡	6‡	12

Negative for poliovirus and negative in suckling mice

† Classified among Types 1-19

‡ One stool sample contained a mixture of Coxsackie Type B4 and ECHO Type 4

Fifty three strains fell into the 19 known ECHO types and only 12 have remained as unidentified agents. Selection of the proper typing sera was made easier by the preliminary grouping based upon the above mentioned biological properties.

Mixed Infections

Mixed infections were easily and quickly recognized by the use of the plaque technique. FIGURE 7 shows 2 distinct kinds of plaques in a rhesus culture bottle inoculated with a stool suspension. In *patas* cultures only large circular plaques were observed. Because of the greater numbers of the large plaques on *patas* cultures it was assumed that the large circular plaques were those of



FIGURE 7. Plaques of mixed infection (stool No. D11) on rhesus (left) and on *patas* (right) 7 days after seeding. On the rhesus culture smaller ECHO plaques are visible in the lower portion of the bottle as well as the large circular poliovirus plaque in the upper portion. Only poliovirus plaques appear on *patas*.




FIGURE 8. Technique for picking plaques: center, a bottle culture from which 2 kinds of plaques have been picked; right, capillary pipette that is used for picking; left, the tube culture in which material from the plaque is subcultured.

polioviruses and because it failed to grow in *patas* cultures the virus producing the smaller irregular plaques on rhesus culture was tentatively classed as ECHO Group A. Single plaques were picked as shown in FIGURE 8 using the capillary pipette and were passed into rhesus tube cultures in order to obtain a larger yield of virus for further identification. In this case it was determined by use of specific antisera that poliovirus Type I and ECHO virus Type 1 were present in this specimen. Thus we see that by the plaque method 2 viruses were readily distinguished and a pure line of each virus could be established by passage and identified by typing with specific antisera.

The problem of mixed infections is complicated when the ratio of poliovirus



FIGURE 9 Mixed infection. In the absence of poliovirus Type I antiserum poliovirus plaques of stool No. 50(2) appeared on rhesus (left) and on *palas* (second from left). In the presence of this antiserum FCHO plaques appeared only on rhesus and not on *palas* (right).

to ECHO virus is excessive. FIGURE 9 indicates such an example. Stool No. 50 grew on both rhesus and *palas* cultures. However, in the presence of polio Type I antiserum, a certain amount of the virus failed to be neutralized. These non-neutralizable virus particles grew on rhesus but not on *palas* cultures. Passage of plaques from the rhesus bottle containing poliovirus antiserum yielded a virus that was typed as ECHO Type 18.

TABLE 4 illustrates some of the problems that a diagnostic laboratory encounters in regard to mixed enteric virus infections. Six fecal specimens were tested by the conventional tube method in our laboratory (listed as Laboratory A) and three of these were also tested in a laboratory in another city (listed as Laboratory B). The first 3 specimens were typed in tube cultures as poliovirus Type I with the ECHO component, which was present at low concentration being missed. Poliovirus was missed from stool I C 6, presumably because in this instance the relative amount was very low. The last 2 specimens listed in the table were identified in the tube test as mixtures of poliovirus and ECHO virus, and here it is seen that the relative amounts of the viruses were more nearly equal. Laboratory B missed the poliovirus component in the first 2 specimens listed, presumably because the virus that it isolated could not be neutralized by polio antisera. Stool II A 21(2) gave an interesting result. A cytopathic agent was obtained after this sample was treated with poliovirus Type I antiserum. Although the nonpolio agent thus obtained reacted with ECHO Type 4 antiserum, it was also noted that it produced plaques on *palas* cultures. Further study was therefore made, and it was found that this specimen contained Coxsackie B4 and ECHO Type 4, in addition to poliovirus Type I.

TABLE 4
COMPARISON OF METHODS FOR THE ISOLATION AND IDENTIFICATION OF ENTERIC VIRUSES
IN MIXED INFECTIONS

Stool source	Titer method (CFP + neutralization)		Plaque morphology + host cell susceptibility + neutralization		
	Laboratory A	Laboratory B	Mix	Plaque	
				PI	ECHO
50(2) II A 21(2)	Polio I Polio I	I CHO I CHO	Polio I + FCHO 18 (Group A) Polio I + Croup B 4 + FCHO 4 (Group A)	87 81 (polio + Croup B) (ackie)	13 16
5414	Polio I	Polio I + I CHO	Polio I + FCHO 7 (Group A)†	99	1
IC 6 ID 11	I CHO Polio I + I CHO	N D N D	Polio I + FCHO 17 (Group A) Polio I + FCHO 17 (Group A)	5 67	95 38
III B-40	Polio I + I CHO	N D	Polio I + FCHO 11 (Group A)	75	25

ND = not determined

† Not typed

It can be seen that by use of the plaque method and host-cell susceptibility mixed infections could be recognized more readily and the different agents could be separated into pure clones. However if this method is used for initial virus isolation rhesus tube cultures should be inoculated in parallel with bottle cultures in order to detect FCHO viruses that do not produce plaques.

ECMO Viruses*

Enteric cytopathogenic monkey orphan (ECMO) viruses are included among the simian viruses isolated from monkey kidney cultures by Hull and his colleagues^{14, 15}. These viruses have also been recovered from rectal swabs and fecal materials by Cheever¹⁶ and by Riordan¹⁷. In one series of 21 monkeys studied in New Haven Conn. 25 ECMO viruses were recovered. Some of these viruses were excreted for a period of weeks from the time of the first isolation. The high prevalence of these viruses in monkey stools has led to the question of whether viruses apparently recovered from the kidneys or from other tissues of the monkey are really viruses of these tissues or whether they are ECMO viruses that had contaminated the tissue in the course of their removal from the body. If these viruses are as widespread as the present evidence suggests then the skin and hair of monkeys must be heavily contaminated and contamination of the tissue must be considered as a possible source of these so-called kidney agents. A cooperative program with the laboratories of Hull and of Cheever is now under way in an attempt to classify

Although foamy agents as reported by Rustian and Brown are also simian viruses they are not included in this discussion. It may be said however that in our studies we found that kidney cells of a number of African monkeys in contrast to those from Asia failed to contain foamy agents.




FIGURE 10 (a) Bottles inoculated with an FCMO virus of Group C (No. 8006) on rhesus culture (left dilution 10^{-5}) and palas culture (right undiluted) 7 days after seeding. There are typical plaques visible only on rhesus but no plaques on palas. (b) Bottles inoculated with FCMO Group B virus (No. SV-16) on rhesus culture (left) and palas culture (right) 7 days after seeding. Large circular plaques with islets are visible in cultures of both species.

the numerous agents that have been isolated either from kidney cultures or from the stools of monkeys

Biologic and Serologic Properties of Simian Orphan Viruses

A total of 51 strains of simian orphan viruses (isolated either from kidney cultures or from stool samples) were tested for their plaque morphology in rhesus bottle cultures and for their *patas* cell susceptibility

Some simian viruses were found to produce plaques resembling those of the ECHO viruses described earlier others have produced distinctive plaques characterized by the presence of islets of apparently healthy cells within the area of degeneration as shown in FIGURE 10 Ten strains were found to produce distinct plaques with islets on rhesus cells and were negative in *patas* bottle cultures these were placed in Group C (TABLE 5 and FIGURE 10a)

TABLE 5

PRESUMPTIVE GROUPING OF SIMIAN VIRUSES ACCORDING TO PLAQUE MORPHOLOGY AND *PATAS* CELL SUSCEPTIBILITY

Group A small delayed plaques on rhesus cultures and on <i>patas</i> cultures			
7852	SV ₁		P 4
7853	SV ₄		P 5
7854	SV		P 6
7855A	SV		P 7
7999A	SV ₁		P 8
	SV ₇	SV ₁	P 9
	SV ₁₁	SV ₇	P 10
	SV ₁₂	SV ₁₁	
Group B large circular plaques with islets on rhesus cultures and on <i>patas</i> cultures			
7848	SV ₁		P 1
7849	SV ₄		P 3
7850	SV		P 11
7998			
8000			
8226†			
8278			
8229†			
8422			
8423			
8427			
8429			
Group C large circular plaques with islets on rhesus cultures but no plaques on <i>patas</i> cultures			
855B	SV ₄		P 2
7999B			
8006			
8227			
8425			
8426			
8431			
8432			

The strains listed in the column at the left were isolated in our laboratory the SV series in the center column were made available by R. N. Hull, Lilly Research Laboratories, Indianapolis, Ind., and the P series in the column at the right were supplied by F. S. Cheever, University of Pittsburgh, Pittsburgh, Pa.

† Mixed strains



FIGURE 11 Bottles inoculated with a Group A ECMO virus (No SV₁) Rhesus culture (2 at left) and *patas* culture (2 at right) 10 days after seeding. With the large inoculum (10^{-1}) generalized virus degeneration of the cell sheet occurred as soon as any plaques appeared. With the smaller inoculum on rhesus cultures as well as in the *patas* cultures plaques were tiny and delayed.

Group B consisted of strains that produced distinct plaques with islets on both rhesus and *patas* cultures (FIGURE 10b). Group A included those that failed to produce plaques rapidly, although with large inocula the cell sheet degenerated under agar. However, tiny delayed plaques like those of measles virus usually occurred on *patas* as well as on rhesus cultures 10 days or more after inoculation (FIGURE 11). Some strains in the latter group gave positive complement fixation (CF) responses when allowed to react with human serum that previously had been shown to contain adenovirus antibodies.

Viruses 8226 (FIGURE 12) and 8229 were found to be mixtures. Purification of these virus mixtures was undertaken by picking single plaques and further studies are under way.

Neutralization tests among the ECMO viruses may be carried out by the plaque method as shown in FIGURE 13. Type SV₈ of Hull and our strain No 8227 were placed in Group C because of their plaque morphology and their failure to produce plaques on *patas* cultures. Neutralization of the No 8227 virus by SV₈ antiserum indicated antigenic relationship. Similar correlations were found for SV₁₆ and our strain No 8000 which produced large clear plaques on both rhesus and *patas* (Group B) cultures and for Type SV₃ and our strain 7834 which were placed in Group A because they were *patas* positive but produced only tiny delayed plaques. Thus with some of these viruses the plaque characteristics and host cell range could be correlated with antigenicity.

On the other hand, although some of the viruses were placed in different




FIGURE 12 FCSO viruses (No. 8226) on rhesus (left) and on palas (right) 7 days after seeing. Two kinds of plaques were noted in rhesus tissue cultures, some with and some without islets. Only plaques without islets were observed on palas culture.

groups according to their palas cell susceptibility (such as No. 1855B which was palas negative and SV₂ which was palas positive) they were nevertheless found to be serologically related. Cheever's 13 agent was a palas positive strain but it was neutralized by antiserum to SV₆ (a palas negative strain).

ECHO Viruses *

These viruses are agents isolated from the enteric tract of the bovine species. From the fecal samples of healthy dairy cattle Kunin and Minuse⁹ isolated 8 viral agents in tissue cultures of bovine kidney. None of these agents was neutralized by polio, Coxsackie or ECHO antiserum. Moll and Finlayson¹ reported recently on a cytopathogenic viral agent isolated from feces of cattle with fever. M. Klein (personal communication) and Flein and Earley²⁰ have isolated about 10 viruses, all but one from the feces of apparently healthy cows. One virus was latent in calf kidney culture. None of the 35 agents tested were neutralized to any significant degree by polio antisera, although neutralizing antibodies to all 3 types of poliovirus were noted in bovine sera.²¹ All of these agents were found to grow readily in bovine kidney cultures and in addition some have been found to propagate in monkey kidney but not in HeLa cultures. It is not unlikely that they may have a biological pattern in the bovine species similar to that of the ECHO viruses in man.

Enterocytopathogenic swine orphan (FCSO) viruses were isolated from the stools of newborn swine by Moscovice²² in Italy. These agents grew rapidly in monkey and swine kidney cells but HeLa cells failed to respond.




FIGURE 13 (a) Neutralization test in bottles with ECMO virus 8227 against SV antiserum in rhesus cultures 8 days after seeding. Virus control bottle at left contained 20 plaques. When SV antiserum was present at dilutions of 1/10 and 1/100 no plaques appeared. With an antiserum dilution of 1/1000 14 plaques were observed. (b) Neutralization test in bottles with ECMO virus No. 8000 against SV antiserum in rhesus cultures 6 days after seeding. Virus control bottle at left contained 42 plaques. No plaques appeared when SV antiserum was present at dilutions of 1/10 and 1/100 with an antiserum dilution of 1/1000 1 plaques appeared.

That viral agents exist in the bovine species was already inferred from the fact that bovine sera neutralize certain viruses. For example as mentioned earlier naturally occurring substances that neutralize the three types of poliovirus have been found in normal adult bovine sera.⁵ This phenomenon is discussed more fully elsewhere in these pages by Morton Klein. Sabin

and Fieldsteel³ felt that these inhibitory substances are true antibodies which has led to the suggestion that some of the bovine orphan viruses may be related to polioviruses. Recently Takemori and his colleagues¹⁴ from Tokyo, Japan, have shown that one can distinguish between virus inhibitors in bovine sera and the true antibodies of poliovirus.

By several passages of poliovirus Types I and II in HeLa cell cultures containing 20 to 40 per cent inhibitory normal bovine serum, mutants were obtained that were able to produce large plaques in the presence of the inhibitory bovine serum, in contrast to the small plaques of the parent virus.¹⁵ These mutants, although resistant to the inhibitory normal bovine serum, were no more resistant to true polio antibodies (as present in monkey immune serum) than were the parent viruses. Therefore, it may be concluded that the properties of the parent virus that were changed by passage in the inhibitory serum were not their specific polio antigenic characters. Thus, the inhibitory substances in normal bovine serum cannot necessarily be attributed to a response of the animal to infection with a poliovirus. Only additional work can clarify the meaning of these viral inhibitory substances in bovine sera.

CONCLUDING REMARKS

We have attempted to review some trends in research on orphan viruses. New tissue culture methods continue to yield new viruses at an almost alarming rate from virtually every animal species studied.

The orphan virus category was originally conceived as a place for the temporary classification of new agents until their diseases were found. It may well be that with so many orphan viruses already recognized, a number of them may turn out to infect man or lower animals without ever causing disease, as is the case with saprophytic bacteria. Other orphan viruses, such as ECHO Types 6 and 9, cause aseptic meningitis. As this syndrome is caused by a variety of viruses (including poliomyelitis, Coxsackie, and lymphocytic choriomeningitis), no single agent can be called *the* virus of aseptic meningitis. This poses a difficult problem with reference to removing a virus from the orphan classification, even when it is known to produce a disease, if the same syndrome can be caused by other agents.

For the next decade, at least, virologists will necessarily be concerned with working out methods for the recognition and classification of these agents. We believe that colony or plaque morphology and cell susceptibility, the traditional criteria for classification of bacteria, are of similar value for classification of viruses. Our recent studies with classifying 10 strains of ECHO and 51 strains of ECHO viruses may serve as an example. We were able to group them provisionally by plaque morphology and host cell susceptibility without resorting to the tedious and time-consuming serologic procedures required if serology alone had been used in relating them to the many antigenic types already known among the enteroviruses.

References

1. COMMITTEE ON THE ECHO VIRUSES OF THE NATIONAL FOUNDATION FOR INFANTILE PARALYSIS, INC. 1955. Enteric cytopathogenic human orphan (ECHO) viruses. *Science* **122**: 1187-1188.

- 2 DAVIS D C & J L MELNICK 1956 Association of echo virus type 6 with aseptic meningitis. *Proc Soc Exptl Biol Med* 92 839-843
- 3 KARZON D T, A I BARRON, W WINKELSTEIN & S COHEN 1956 Isolation of echo virus type 6 during outbreak of seasonal aseptic meningitis. *J Am Med Assoc* 162 1298-1303
- 4 McLEAN D M & J L MELNICK 1957 Association of mouse pathogenic strain of echo virus type 9 with aseptic meningitis. *Proc Soc Exptl Biol Med* 94 656-660
- 5 QUERIN THIRY I, F NIHALL & F DEKKING 1957 Echo virus type 9 as a cause of epidemic meningitis. *Science* 125 744-745
- 6 COTTFREDSEN A & H VON MAGNUS Isolation of echo virus type 9 from cerebrospinal fluid. In press
- 7 BOISSARD G P, B L J STOKES, A D MACRAE & F O MACCALLUM 1957 Isolation of viruses related to echo virus type 9 from outbreaks of aseptic meningitis. *Lancet* 272 500
- 8 DELLBECO I 1952 Production of plaques in monolayer tissue cultures by small particles of animal virus. *Proc Natl Acad Sci* 38 447-452
- 9 SELLERS R I 1955 Growth and titration of the virus of foot and mouth disease and vesicular stomatitis in kidney monolayer tissue culture. *Nature* 176 547-549
- 10 DUBES G R 1956 Differences among strains of poliovirus in plaque size on monkey kidney cells. *Virology* 2 284-289
- 11 HSIUNG G D & J L MELNICK 1955 Plaque formation with poliomyelitis Coxsackie and orphan (ECHO) viruses in bottle cultures of monkey epithelial cells. *Virology* 1 533-535
- 12 HSIUNG G D & J L MELNICK 1957 Morphologic characteristics of plaques produced on monkey kidney monolayer cultures by enteric viruses (poliomyelitis Coxsackie and echo groups). *J Immunol* 78 128-136
- 13 HSIUNG G D & J L MELNICK 1957 Comparative susceptibility of kidney cell from different monkey species to enteric viruses (poliomyelitis Coxsackie and echo groups). *J Immunol* 78 137-146
- 14 HULL R N, J R MINNER & J W SMITH 1956 New viral agents recovered from tissue cultures of monkey kidney cell. *Am J Hyg* 63 204-215
- 15 HULL P N & J R MINNER 1957 New viral agents recovered from tissue cultures of monkey kidney cells. II. Problems of isolation and identification. *Ann N Y Acad Sci* 67(8): 413-423
- 16 CHEEVER F S 1957 Discussion Part IV. *Ann N Y Acad Sci* 67(8) 427-429
- 17 RIORDAN J Unpublished data
- 18 RUSTIGIAN R P, JOHNSTON & H PESHART 1956 Infection of monkey kidney tissue cultures with virus like agents. *Proc Soc Exptl Biol Med* 88 8-16
- 19 BROWN L V 1957 Pathogenicity for rabbit kidney cell cultures of certain agents derived from normal monkey kidney cultures. I. Isolation and propagation. *Am J Hyg* 65 189-209
- 20 KACIN C M & E MINLSE 1956 The isolation in tissue culture and chick embryo of filtrable agents from healthy dairy cattle. *Conf of Public Health Vet*
- 21 MOLL T & A V LINLAYSON 1957 Isolation of cytopathogenic viral agent from feces of cattle. *Science* 126 401-402
- 22 KLEIN M & F EARLEY 1957 The isolation of enteric cytopathogenic bovine phan (ECBO) virus from calves. *Bacteriol Proc* 31 73
- 23 BARTELL P & M KLEIN 1955 Neutralizing antibody to virus of poliomyelitis in sera of domestic animals. *Proc Soc Exptl Biol Med* 90 571-601
- 24 MOSCOVICI C, A CINEVRI & V MAZZARACCHIO 1956 Isolamento un coltura di virus utili un virus da enteritis suina. *Zooproflassi* July
- 25 SABIN A B & A H FIELDSTEPL 1953 6th Intern Congr Microbiol 2 343
- 26 TAKEMORI S, S NOMURA, M NAKANA, Y MORIOKA, M HEMMI & M KITAKOJI 1957 Mutations of polioviruses to resistance to the neutralizing substances in normal human serum. *Science* 125 1196-1198

Discussion of the Paper

HENRY M CELFAND (*School of Medicine Tulane University New Orleans La*) William Flynn and I at the Tulane University School of Medicine probably shall be able soon to add another group to the list of viruses presented by Hsiung and Melnick. We believe that we may have isolated several members

of an ECDO group enteric cytopathogenic dog orphan viruses. At least two strains have been obtained from a group of fecal specimens collected from apparently healthy stray dogs. They were isolated in tissue cultures of dog kidney and seem to be adaptable to monkey kidney cells but not to HeLa cells. Work has just begun with these agents but preliminary results indicate that they are not neutralized by rabbit hyperimmune sera against the 3 types of poliovirus against Coxsackie viruses A9 and B1 to B5 or against ECHO viruses Types 1 to 13. These ECDO viruses are neutralized by a majority of the dog sera against which they have been tested and also by a number of human sera. They have not yet been tested for neutralization by specific sera against distemper virus or infectious canine hepatitis (ICH) virus but this will be done soon. However it seems unlikely that these agents will prove to be either distemper or ICH virus. Rabbits are being immunized for a more detailed antigenic study of relationships to human enteric agents.

- 2 DAVIS D C & J L MELNICK 1956 Association of echo virus type 6 with aseptic meningitis. *Proc Soc Exptl Biol Med* 92 839-843
- 3 KARZON D T, A I BARRON, W WINKELSTEIN & S COHEN 1956 Isolation of echo virus type 6 during outbreak of seasonal aseptic meningitis. *J Am Med Assoc* 162 1298-1303
- 4 McLFAN D M & J L MELNICK 1957 Association of mouse pathogenic strain of echo virus type 9 with aseptic meningitis. *Proc Soc Exptl Biol Med* 94 656-660
- 5 QUERFIN THIRY L, F NIHAUL & F DEKKING 1957 Echo virus type 9 as a cause of epidemic meningitis. *Science* 125 744-745
- 6 GOTTFREDSEN A & H VON MAGNUS Isolation of echo virus type 9 from cerebrospinal fluid. In press
- 7 BOISSARD G I, B L J STOKES, A D MACRAE & F O MACCALLUM 1957 Isolation of viruses related to echo virus type 9 from outbreaks of aseptic meningitis. *Lancet* 272 500
- 8 DULBECCO R 1952 Production of plaques in monolayer tissue cultures by single particles of animal virus. *Proc Natl Acad Sci* 38 747-752
- 9 SELLERS I F 1955 Growth and titration of the virus of foot and mouth disease and vesicular stomatitis in kidney monolayer tissue culture. *Nature* 176 547-549
- 10 DUBES G R 1956 Differences among strains of poliovirus in plaque size on monkey kidney cells. *Virology* 2 284-288
- 11 HSIUNG G D & J L MELNICK 1955 Plaque formation with poliomyelitis Coxsackie and orphan (ECHO) viruses in bottle cultures of monkey epithelial cell. *Virology* 1 533-535
- 12 HSIUNG G D & J L MELNICK 1957 Morphologic characteristics of plaques produced on monkey kidney monolayer cultures by enteric viruses (poliomyelitis Coxsackie and echo groups). *J Immunol* 78 128-136
- 13 HSIUNG G D & J L MELNICK 1957 Comparative susceptibility of kidney cells from different monkey species to enteric viruses (poliomyelitis Coxsackie and echo groups). *J Immunol* 78 131-146
- 14 HULL R N, J R MINNER & J W SMITH 1956 New viral agents recovered from tissue cultures of monkey kidney cells. *Am J Hyg* 63 204-215
- 15 HULL R N & J R MINNER 1957 New viral agents recovered from tissue cultures of monkey kidney cells. II Problems of isolation and identification. *Ann N Y Acad Sci* 67(8) 413-473
- 16 CHEEVER F S 1957 Discussion Part IV. *Ann N Y Acad Sci* 67(8) 427-429
- 17 RIORDAN J Unpublished data
- 18 MUSTIGIAN I, P JOHNSTON & H REINHART 1955 Infection of monkey kidney tissue cultures with virus like agents. *Proc Soc Exptl Biol Med* 88 8-16
- 19 BROWN L V 1957 Pathogenicity for rabbit kidney cell cultures of certain agents derived from normal monkey kidney cultures. I Isolation and propagation. *Am J Hyg* 65 189-209
- 20 KUNIN C M & E MINLSE 1956 The isolation in tissue culture and chick embryo of filtrable agents from healthy dairy cattle. *Conf of Public Health Vet*
- 21 MOLL T & A V FINLAYSON 1957 Isolation of cytopathogenic viral agent from feces of cattle. *Science* 126 401-402
- 22 KLEIN M & E FARLEY 1957 The isolation of enteric cytopathogenic bovine orphan (TCBO) virus from calves. *Bacteriol Proc* 31 73
- 23 BARTELL P & M KLEIN 1955 Neutralizing antibody to viruses of poliomyelitis in sera of domestic animals. *Proc Soc Exptl Biol Med* 90 591-601
- 24 MOSCOVICI C, A GINEVRI & V MAZZARACCHIO 1956 Isolamento un cultura di virus da enteritis suina. *Zooprofassi* July
- 25 SABIN A B & A H FIELDSTEEL 1953 6th Intern Congr Microbiol 2 313
- 26 TAKEMORI N, S NOMURA, M NAKANA, Y MORIOKA, M HEMMI & M KITAOKI 1957 Mutation of polioviruses to resistance to the neutralizing substances in normal bovine serum. *Science* 125 1196-1198

Discussion of the Paper

HENRY M GELFAND (*School of Medicine Tulane University New Orleans Ia*) William Flynn and I at the Tulane University School of Medicine probably shall be able soon to add another group to the list of viruses presented by Hsiung and Melnick. We believe that we may have isolated several members

TABLE 1
NEUTRALIZING ANTIBODIES IN COWS TO THREE TYPES OF POLIOVIRUS

Form	Number tested	Percent positive
A	91	43
B	32	70
C	18	72
D	25	69

Sera were tested at a dilution of 1:4 against 50 tissue culture doses of virus

TABLE 2
INCIDENCE OF THREE TYPES OF NEUTRALIZING ANTIBODIES TO POLIOVIRUS IN COWS

Type I	Type II	Type III
35	63	11

for our discussion the question of how one interprets neutralizing substances to certain human viruses in the sera of cows.^{8,10} In our own studies we observed that there were present in the sera of several domestic animals, horses, hogs and particularly cows neutralizing substances to Types I, II and III polioviruses. We have been following about 150 cows on 4 farms in the Philadelphia area as shown in TABLE 1; the neutralizing substance to polioviruses is a widely distributed agent. As shown in TABLE 2, all 3 types of virus are neutralized. In addition we have found neutralizing substances to Types 4 and 5 Coxsackie B viruses and there have been recent reports of neutralizing substances to the adenoviruses in the sera of cows.¹¹ We may then ask this question: Are these neutralizing substances true antibodies that indicate an unusually wide distribution in animal sera of human viruses or their bovine antigenic relatives? The problem of interpreting these serologic reactions is rather awkward because we have no clinical picture to guide us, no epidemiological pattern and no acute and convalescent sera to detect a rise in antibody titer. We are confronted with individual sera from apparently healthy cows that neutralize these human viruses. When studied further, these sera yielded the following information:

- (1) The neutralizing substance is heat stable, resisting 60°C for 30 min.
- (2) The inhibitor is in the globulin fraction obtained by precipitation with ammonium sulphate one third saturation.
- (3) The inhibitor is type specific; some sera inhibit Type I, others Type II or Type III, still others combinations of all three or none at all.
- (4) There is evidence of passive transfer; newborn calves have neutralizing substances that disappear in a month or so and gradually reappear as the animal matures, which is the traditional antibody pattern.
- (5) Neutralizing titers vary from 0 to greater than 1:250 against 50 tissue culture doses of virus, which is again a pattern consistent with an antibody interpretation.

Do the above data permit us to conclude that cows have antibodies to Types I, II and III polioviruses or do other substances exist in serum that mimic this antibody pattern?

THE SIGNIFICANCE OF HUMAN ANTIVIRAL NEUTRALIZING SUBSTANCES IN ANIMAL SERA*

By Morton Klein

Department of Microbiology Temple University School of Medicine Philadelphia Pa

In considering infections that are common to man and animals probably the greatest unexplored area is that of the viruses. Tissue culture has given virologists the golden touch and there are now pouring forth an enormous number of newly recognized human and animal viruses. If one considers that about eighty infections¹ of animals are already known to be shared in common with man one can predict that a study of these newly isolated human and animal viruses will reveal many examples of identity or at least, of antigenic relationships.

As we all know one of the most sensitive and specific tools for determining relationships between viruses is the antigen antibody reaction. In virology we use three basic procedures the complement fixation test the inhibition of hemagglutination test and the neutralization test. An essential requirement of all these tests is that they be specific for if one explores the sera of an animal species and finds something that neutralizes a test virus one would very much like to say the virus is or has been here. Unfortunately serologic tests are never carried out as pure antigen antibody reactions they take place in the rather unpredictable menstruum of serum and there are many things in sera that can give grossly misleading serologic results. It is the intent of this discussion to consider those substances that may disturb the specificity of these serologic reactions.

The literature on unexplained inhibitors in animal sera is a collection of many miscellaneous scattered observations with no central theme discernible other than the fact that there is a variety of inhibitors. One can consider these inhibitors from different points of view their role in nonspecific resistance to infection and the problems they raise when used as media in virus tissue culture work. For the present discussion I have arbitrarily organized the material to try to answer this specific question. How can one recognize a true antibody among a collection of unknown inhibitors?

I shall exclude from any detailed consideration a large number of nonantibody inhibitors that may be extracted from body cells. One can obtain from phagocytic cells such microbial inhibitors as leukins² and phagocytin³ one can extract from red blood cells an agent that inhibits viral hemagglutination⁴ certain polypeptides obtained from tissues are known to be inhibitory⁵ a mucopolysaccharide from the intestinal tract inhibits Theiler's virus⁶ aqueous extracts from brain and spleen⁷ are known inhibitors of microbial activity. None of these cellular inhibitors need concern us here since as far as we know they do not appear in the sera of normal animals.

We shall consider the problem of nonspecific inhibitors by using as the basis

The work reported in this paper is supported in part by a grant from the National Foundation for Infantile Paralysis Inc New York N Y

feres with the hemagglutination test of arthropod viruses²¹ the particle is so large it can be removed by high speed centrifugation or slow filtration (3) in dengue there is a serum inhibitor of the hemagglutination test that must be removed with acetone²² In the encephalomyocarditis group of viruses a nonspecific inhibitor of hemagglutination is frequently observed²³ These nonspecific inhibitors of hemagglutination need not be confused with true antibodies both for the reasons stated above and because they do not in general neutralize infectivity

As for the neutralization test I know of only one heat stable nonglobulin inhibitor of viral neutralization that is the serum lipids reported to give very low levels of neutralization against certain viruses²⁴ It is well to note that highly abnormal sera might give nonspecific neutralization reactions For example sera obtained from jaundiced patients with yellow fever or infectious hepatitis may contain enough bile products to neutralize a virus in fact it does so in the case of yellow fever However our concern is with reasonably normal sera Further a good general procedure to eliminate many of these agents is to determine if the inhibitory activity is associated with the gamma globulin fraction as obtained by ammonium sulfate by alcohol or most elegantly by electrophoresis If the inhibitor is not in the globulin fraction one does not have an antibody If it is in the globulin fraction one may or may not have a specific antibody which brings us to our third category of inhibitors and here matters become considerably more complex

Inhibitors that are globulins The fact that we have a neutralizing substance against the polioviruses that is heat stable and in the gamma globulin fraction does not permit us to conclude that we have an antibody stimulated by the virus of poliomyelitis There are examples of gamma globulin fractions active in the inhibition of hemagglutination test that are not even true antibodies but I am not aware of any gamma globulin fractions having neutralization activity that are not true antibodies I feel we can conclude that in the case of the neutralizing substances to poliovirus all the data are compatible with the interpretation that we are dealing with a true antibody I believe the burden of proof is upon those who would suggest an alternative interpretation If the neutralizing substance is not an antibody what is it?

However the decision that one has an antibody merely raises another question What was the antigenic stimulus? The problem that confronts us is that of serologic cross reactions for different viruses may share common antigens and give rise to different but closely related antibodies Cows have antibodies to the virus of smallpox because they had cowpox mice because they had mousepox Cows may have antibodies to the virus of psittacosis because they have been infected with the agent of sporadic bovine encephalomyelitis²⁵ The viruses of canine distemper and measles cross react²⁶ One may find antibodies to the virus of Japanese encephalitis B because an animal has been infected with the West Nile virus for we now know that there are two large families of arthropod borne viruses each of whose members are antigenically related It is quite easy to see that if we were not aware of these cross reacting viruses a variety of neutralizing substances would simply be called nonspecific inhibitors One can cite many other examples The

There are substances other than antibody that neutralize viruses. There are many of them, most of them poorly defined, many with unusual properties. These substances defy anything but a loose classification, and I propose to consider them under three general headings: (1) heat labile inhibitors of virus that are obviously not antibody; (2) heat stable inhibitors of virus that are obviously not antibody; and (3) heat stable inhibitors of viruses that may or may not be antibody.

Heat labile inhibitors. A variety of heat labile activities have been observed, the most thoroughly studied and perhaps the most common being properdin. This inhibitor has been found active against the viruses of St. Louis encephalitis, herpes, Murray Valley fever, mumps, influenza, and Newcastle disease^{12, 13} which strongly indicates that unheated sera can be a source of considerable confusion. However, the nonspecific activity of unheated sera can be removed simply by heating sera to 56° C for 30 min. I cannot too strongly emphasize that in any serologic survey sera should first be heated for this immediately eliminates a large number of nonspecific inhibitors. There are many studies in the literature that cannot be interpreted because the authors failed to heat their sera or failed to indicate whether or not the sera were heated. There is one recorded error subsequently corrected by the author¹⁴ in which antibodies to Newcastle virus were found regularly in human sera until the sera were heated when the antibodies promptly disappeared. There is some uncertainty about always heating sera for in the case of certain viruses higher neutralizing titers have been reported with unheated sera. We may cite as examples neutralization tests done with the viruses of western equine encephalitis, dengue, and mumps^{15, 16}. However, there is reason to believe that such unheated sera do not offer a more sensitive indicator of antibody by virtue of the additional presence of complement but rather a summation of two distinct inhibitory effects, one being antibody and the other a heat labile inhibitor, perhaps properdin. Heating may thus lower the observed activity but it probably defines the antibody more precisely. Although the heat labile inhibitors may be important in resistance to infection, little more need be said about these agents from our particular point of view for heating to 56° C for 30 min immediately removes them from consideration as antibodies. Certainly then, our inhibitor in bovine sera is not properdin or any other heat labile substance.

Heat stable inhibitors that are not globulins. We use the term heat stable to refer to any inhibitory activity that remains after heating sera to 60° C for 30 min. A variety of serologically reactive heat stable nonantibody inhibitors have been described in the sera from many animal species and observed under several serologic conditions: in the complement fixation test, in agglutination, flocculation, neutralization, and in the inhibition of hemagglutination test. In virology it is the specificity of the inhibition of hemagglutination test that is most commonly obscured. In nearly all cases where the inhibition of hemagglutination test is used, nonspecific inhibitors occur. To cite a few examples: (1) there is an inhibitor in normal sera¹⁷ active against the viruses of influenza, and it must be removed with RDE (receptor destroying enzyme), sodium periodate, or trypsin;¹⁸ (2) there is a large particle in sera that inter-

related to human viruses, for they are neutralized by human gamma globulin and there is clearly an area of animal and human relationships to be worked out with the bovine viruses

Discussion and summary There are a large number of unknown inhibitors in animal sera, and there is no simple method that allows us readily to identify these agents. We have discussed procedures that will help more precisely to define the antibody status of these inhibitors and they are summarized below.

To determine whether an inhibitor is an antibody for our test organism the following observations are helpful:

(1) Sera should retain their inhibitory activity after heating at 60° C for 30 min and the inhibitory activity should be in the globulin fraction. It may also be helpful to heat the sera to 80° C for 20 min, since antibodies would be inactivated at this temperature while heat stable nonantibody inhibitors such as lysozyme would remain active.

(2) Inhibitors should show serologic specificity and titers should vary among animals with a pattern of passive transfer and an increase in incidence with the age of the animal.

(3) If the inhibitor is a true antibody the titer should not be affected when sera are treated with RDE, sodium periodate, trypsin or fat solvents. Centrifugation at high speed should not remove the inhibitor.

(4) If sera are positive by one serologic procedure they should be tested by other serologic procedures. For example if sera positive by the inhibition of hemagglutination reaction also give a positive complement fixation or neutralization reaction one has good evidence of an antibody.

(5) A booster response of the inhibitor when the animal is injected with the test organism is also evidence for a true antibody.

(6) If available clinical and epidemiological data with a rise in inhibitor titer are almost conclusive evidence of a true antibody.

It should be noted that some of these suggested procedures are not always feasible and one may have a true antibody without being able to demonstrate all of the above reactions, for example an antibody may be demonstrable by the neutralization test but not by the complement fixation test. Moreover one may have a nonspecific inhibitor that possesses some of the above properties for example resistance to the agents listed in the third observation cited above. However I think the guide is useful for a nonspecific inhibitor would have very few of the above properties while a true antibody may possess most if not all of the above reactions.

Finally though one may have an antibody that neutralizes a test virus proof that the test organism was also the original antigenic stimulus requires that one isolate the suspected virus from the animal species. Only thus can the problem of cross reactions be resolved.

References

1. WORLD HEALTH ORGANIZATION 1951 Technical Rept. Ser. No. 40
2. SHARNES R. C. and D. W. WATSON 1956 Characterization of leukin an artificial factor from leukocytes active against Gram positive pathogens. J. Exptl. Med. 104: 829

point to be emphasized is that cross reactions confuse our serologic data perhaps more than we realize. We have only a very limited concept of the total number of cross reacting species, for a systematic study of cross reacting patterns among the thousands of bacterial and viral species of man and animals would involve an astronomical number of tests. The fact that even in germ free animals antibodies can apparently develop against orally ingested dead bacilli suggests the enormous range of antigenic stimuli that daily bombard our immune mechanism.⁸ There is no way of knowing how many of our unknown inhibitors represent cross reacting antibodies though I feel that they should loom very large in our consideration of the meaning of unknown inhibitors.

On the basis of the serologic evidence alone one then can merely say that this is an antibody that reacts with the test organism and two interpretations remain (1) the antibody is due to a stimulus received from the polioviruses or (2) from an antigenic relative of the polioviruses. A precise answer can be obtained only by the isolation of the responsible agent.

We have been carrying out extensive isolation studies in an attempt to discover the origin of these antibodies to the polioviruses. We have not yet discovered it but I might summarize our study briefly. As in the case of other investigators we have found feces a fertile source of viruses in calves overwhelmingly so. We have isolated over 75 viruses from the feces of apparently normal calves and cows. These agents are not yet classified. We do not know how many distinct species are in the group though there are certainly several on the basis of their cytopathogenic effects in calf kidney tissue culture. All but two of the viruses were isolated from young animals in the age group 6 months to 2 years and it is quite clear that in humans as in calves there is a period of intense viral infection early in life.

None of these 75 viruses are agents of human poliomyelitis quite simply because they all grow in calf kidney tissue culture and we have never succeeded in getting any of the 3 strains of poliovirus to grow in this medium. There remains the question as to whether any of these viruses are antigenic relatives of the agents of poliomyelitis. What we have done is to screen these agents against sera containing antibodies to Types I, II and III polioviruses. In an as yet incomplete study we found 2 bovine agents that were neutralized by pooled heated sera. However normal heated monkey sera were also found to neutralize our bovine viruses to approximately the same titer (1:16) and so obviously we did not have any bovine relatives of the polioviruses.

In order to avoid the problem of unknown inhibitors in monkey sera we tried to immunize rabbits with our bovine strains and polioviruses for rabbits do not commonly have normal inhibitors to polio, Coxsackie or ECHO viruses.⁹

However we found that rabbits also have a heat stable inhibitor that neutralized several of our bovine viruses in titers up to 1:32. These inhibitors apparently prevented any significant antibody response in spite of rather intense antigenic stimulation. We are currently working on the problem of finding a suitable animal for immunization purposes.

In an incomplete survey we do know that two of our bovine viruses are

COUNTERPARTS OF HUMAN VIRAL DISEASE IN ANIMALS

By Hilary Koprowski
The Wistar Institute, Philadelphia, Pa.

Introduction

*Feyfoble sulat h s t a c t e r s t e l l h u m a n d c a l l s h e r d i v s c i e n c e o m m u n i t y a s c o m m u n i t y a s f a i t h c a l l s d i d i n r e v e l a t i o n*¹

Animal counterparts of diseases occurring in humans offer a fruitful approach to experimental studies of such entities in the animals leading to a better understanding of the human counterparts. For obvious reasons large scale experimentation with humans as subjects is not feasible although a great deal of knowledge has been obtained in the past by studies in volunteers or by heroic experiments performed by scientists on themselves. One of the most recent examples that can be cited is the study of the Coxsackie and adenovirus groups of viruses by the deliberate exposure of volunteers to the agent or agents isolated from subjects in the course of illness. Infectious hepatitis is another entity that has baffled all attempts at thorough study because of the lack of suitable animal hosts and the knowledge gained thus far has been based almost entirely on studies on voluntary participants.

While experiments can be conducted in volunteers with diseases that run a mild course great caution must be exercised in the case of diseases that might endanger the lives of the volunteers. For the most part this limitation does not apply to animals. Here the field is open to pursue study of the clinical syndrome encountered in nature and to attempt to induce its counterpart by exposure of genetically homologous species to the agent or agents isolated in the field. The time factor is an additional advantage in the use of animals as the period required for study of a new disease entity is considerably shorter.

The purpose of this paper is to present in a schematic form aspects of animal diseases for comparison with the human viral entities that can be considered as their counterparts. For apparent reasons such comparison of necessity must be based more on the clinical symptomatology and pathology than on etiology. This approach no doubt presents many pitfalls, the knowledge gained from naturally acquired or experimentally induced diseases in animals can serve only as an adjunct to our understanding of the possibly similar entities in humans.

There is a wealth of wisdom in the terse admonition addressed by Koch to his students and which they have passed down to us: advice which is pertinent even today and should be kept in mind by all students of virology. *Meine Herren vergessen Sie nie dass die Mause keine Menschen sind* *

Infections of the Central Nervous System

There are numerous human counterparts of animal encephalitides. This group of illnesses has probably the least to offer from the standpoint of con-

Gentlemen never forget that mice are not human beings

- 3 HIRSCH J G 1956 Phagocytin a bactericidal substance from polymorphonuclear leucocytes J Exptl Med 103 589
- 4 DE BURGH I M P C YU C HOWE & M BOVARNICK 1948 Preparation from human red cells of a substance inhibiting virus hemagglutination J Exptl Med 87 1
- 5 MILLER B F R ABRAMS A DORFMAN & M KLEIN 1942 Antibacterial properties of protamine and histone Science 96 428
- 6 MANDEL B & F RACKER 1953 Inhibition of Theiler's encephalomyelitis virus (GDVII strain) of mice by an intestinal mucopolysaccharide J Exptl Med 98 399
- 7 NUTINI I G & F M LYNCH 1946 Further studies on effects of tissue extracts on *Staphylococcus aureus* J Exptl Med 84 247
- 8 HAMMON W McD W A MACK & W C REEVES 1947 The significance of protection tests with the serum of man and other animals against the Lansing strain of poliomyelitis virus J Immunol 57 285
- 9 SABIN A B & A H FIELDSTFEL 1953 6th Intern Congr Microbiol Rome Italy Sections 6-7 373
- 10 BARTELL I & M KLEIN 1955 Neutralizing antibody to viruses of poliomyelitis in sera of domestic animals Proc Soc Exptl Biol Med 90 597
- 11 GOLD L & H S GINSBERG 1957 Inhibition of adenoviruses by normal animal sera Federation Proc 16 414
- 12 SULLIN F 1957 Personal communication
- 13 WEDGEWOOD R H GINSBERG & L PILLENER 1956 The properdin system and immunity J Exptl Med 104 107
- 14 HOWITT B 1950 A nonspecific heat labile factor in the serum neutralization test for Newcastle disease virus J Immunol 64 73
- 15 MORGAN I 1945 Quantitative study of the neutralization of western equine encephalomyelitis virus by its antiserum and the effect of complement J Immunol 50 359
- 16 WHITMAN L 1947 The neutralization of western equine encephalomyelitis virus by human convalescent serum The influence of heat labile substances in serum on the neutralization index J Immunol 56 97
- 17 SABIN A B 1950 The dengue group of viruses and its family relationships Bacteriol Rev 14 225
- 18 LEYMASTER G & T WARD 1949 The effect of complement in the neutralization of mumps virus J Immunol 61 95
- 19 FRANCIS T 1947 Dissociation of hemagglutinating and antibody measuring capacities of influenza virus J Exptl Med 85 1
- 20 JENSEN K E 1956 Diagnostic procedures for virus and rickettsial diseases 250 Am Public Health Assoc New York N Y
- 21 CASALS J & L BROWN 1954 Hemagglutination with arthropod borne viruses J Exptl Med 99 429
- 22 SABIN A B 1956 Diagnostic procedures for virus and rickettsial diseases 384 Am Public Health Assoc New York N Y
- 23 GARD S & L HELLER 1951 Hemagglutination by Col MM virus Proc Soc Exptl Biol Med 76 68
- 24 UTZ J 1948 Effect *in vitro* of specific lipid fractions of animal sera on poliovirus Proc Soc Exptl Biol Med 69 186
- 25 UTZ J 1949 Studies on the inactivation of influenza and Newcastle disease viruses by a specific lipid fraction of normal animal sera J Immunol 63 273
- 26 WENNER H R MANGES & G HARSHFIELD 1954 Sporadic bovine encephalomyelitis J Infectious Diseases 94 284
- 27 IMAGAWA D T & J M ADAMS 1957 Immunological relationship between measles and distemper viruses Federation Proc 16
- 28 LOBUND REPORTS No 2 1949 Germ free life studies the chicken 143 Univ Press Notre Dame Ind
- 29 GRAY A Personal communication

TABLE 2
EXAMPLES OF INFECTIONS OF MAN THAT OCCUR IN ANIMALS USUALLY IN LATENT FORM
AND THAT ARE INTERTRANSMISSIBLE

V	Vect	Species affected	Symptomatology
St. Louis	Mosquito Mite	Chicken Man	None Encephalitis
Yellow fever	Mosquito	South American pri- mates Man	Occasional liver involve- ment Liver involvement
LCM	?	Mice Man	None Meningitis pneumonia

Lymphocytic choriomeningitis

In Brazilian jungles moribund marmosets were found to be infected with yellow fever virus but whether the South American primates constitute a natural reservoir of the virus has yet to be proved. The ubiquitous nature of the lymphocytic choriomeningitis virus is well known to students of disease. Mice have been chosen as an example of an animal host simply because we know thanks to Traub⁷ and to Armstrong, Wallace and Ross⁸ more about their role than those of the other hosts. The symptomatology in man is variable and the sporadic nature of the infection is quite apparent. The disease may occur in man in virulent form⁹ but these cases are probably exceptions to the usual latent infection.

Poliovirus like Diseases

The poliovirus like diseases constitute a strange group and originally they were grouped together (TABLE 3)¹⁰. New studies of the Teschen disease seem to indicate that it does not resemble human poliovirus as closely as it was originally thought. Although it can be induced by the ingestion of infectious material according to Horstmann¹ the virus is strangely absent from feces. Because of its ubiquity in outbreaks among animals it is impossible to determine the ratio of symptomatic cases to latent forms and also to calculate accurately the mortality ratio which seem to be higher than in the case of human poliovirus. Furthermore the disease of the animal takes on a more acute form than does the pathologically similar condition in man.

TABLE 3
THE STRANGE CASE OF POLIOVIRUS LIKE DISEASES OF ANIMALS AND MAN

V	Species	Host	Transmission	Excretion in feces	Symptomatology
Teschen	Swine	?	Feeding	—	Encephalomyelitis
Theiler	Mice	?	Feeding	++	Anterior horn
Polio	Man	Latent?	Feeding	++	Anterior horn and brain stem

TABLE 1

EXAMPLES OF INFECTIONS OF CENTRAL NERVOUS SYSTEM FOR WHICH COUNTERPARTS ARE FOUND IN HUMAN MEDICINE AND WHICH ARE INTERTRANSMISSIBLE

Virus	Natural Hosts	Vector	Species affected	Symptomatology	Site of Infection
Arbor A	Birds	Mosquito	Horses Mules Man	Encephalitis Encephalitis Encephalitis	Brain Brain Brain
Japanese B	Birds ²²	Mosquito	Horses Swine Man	Encephalitis Encephalitis Encephalitis	Brain Brain Brain
Louping ill	??	Tick	Sheep Man	Encephalitis Encephalitis	Brain Brain

Arthropod borne encephalitis serologic group A data refer chiefly to the North American equine encephalitis

troverſy The pathways of the infectious process are well known the etiologiſal agents are recognized and intertransmiſſibility between man and animals is well defined However, the natural reſervoir of infection in ſome caſes ſtill preſents a problem

Among members of the Group A arthropod borne (Arbor) encephalitis³ the eaſtern and weſtern equine encephalitis viruses have been moſt thor- oughly inſtiſgated (TABLE 1) Man the accidental hoſt is infected by the ſame vector as are the animals The incidence courſe and outcome of the human illneſs are ſimilar to thoſe of horſes and mules The third member of the Arbor A group Venezuelan equine encephalitis is highly conta- gious for man and can be tranſmitted either by a vector³ or by direct contact with infected fomites⁴ Strangely this virus which is highly virulent for horſes and mules ſeems to provoke a rather mild type of infection in humans who acquire it probably by inhaling the infected material⁵ The only known fatal caſe occurred in a man who became infected in nature during an out- break of the diſeaſe among horſes² It may be queſtionable to ſpeculate on the ſlight ſtatistical evidence of one caſe as to whether the courſe of human illneſs is conditioned by the manner in which this infection is acquired Rela- tively little is known about the other members of the Arbor A group

The pathway of infection with the Japanese B virus courſe of illneſs and outcome are ſimilar in man and animals The relationship between louping ill and Ruſſian ſpring ſummer encephalitis viruses is very cloſe and human beings are known to have become infected with theſe agents Theſe viruses are tranſmiſſible between different ſpecies and cauſe diſeaſe of ſimilar ſympto- matology

The diſeaſes thus far diſcuſſed are known to be the cauſe of frequent epizootics of quite ſevere character but on TABLE 2 are ſhown examples of infections that uſually remain latent in their animal hoſts When tranſmitted to man they may cauſe either ſevere illneſs or inapparent infection The tremendous variation in the clinical picture of human St. Louis encephalitis may be cited as an example⁶

TABLE 2

EXAMPLES OF INFECTIONS OF MAN THAT OCCUR IN ANIMALS USUALLY IN LATENT FORM AND THAT ARE INTERTRANSMISSIBLE

Virus	Vector	Species affected	Symptomatology
St. Louis	Mosquito	Chicken Man	None Inapparent
Yellow fever	Mosquito	South American primates Man	Occasional liver involvement Liver involvement
LCM	"	Mice Man	None Meningitis pneumonia

Lymphocytic choriomeningitis

In Brazilian jungles moribund marmosets were found to be infected with yellow fever virus but whether the South American primates constitute a natural reservoir of the virus has yet to be proved. The ubiquitous nature of the lymphocytic choriomeningitis virus is well known to students of disease. Mice have been chosen as an example of an animal host simply because we know thanks to Traub⁷ and to Armstrong, Wallace and Ross⁸ more about their role than those of the other hosts. The symptomatology in man is variable and the sporadic nature of the infection is quite apparent. The disease may occur in man in virulent form⁹ but these cases are probably exceptions to the usual latent infection.

Poliomyelitis like Diseases

The poliomyelitis like diseases constitute a strange group and originally they were grouped together (TABLE 3)¹⁰. New studies of the Teschen disease seem to indicate that it does not resemble human poliomyelitis as closely as it was originally thought. Although it can be induced by the ingestion of infectious material according to Horstmann¹¹ the virus is strangely absent from feces. Because of its ubiquity in outbreaks among animals it is impossible to determine the ratio of symptomatic cases to latent forms and also to calculate accurately the mortality ratio which seem to be higher than in the case of human poliomyelitis. Furthermore the disease of the animal takes on a more acute form than does the pathologically similar condition in man.

TABLE 3

THE STRANGE CASE OF POLIOMYELITIS LIKE DISEASES OF ANIMALS AND MAN

Disease	Host	Agent	Transmission	Incubation	Pathology
Teschen	Swine	?	Feeding		Lucey haemorrhagic
Theiler	Mice		Feeding	++	Anterior horn
Eliot	Man	Cow	Feeding	++	Anterior horn and brain stem

Theiler's virus infection of mice runs a course similar to that of human poliomyelitis. Infectivity of fecal material and lesions in the anterior horn of the spinal cord are factors that suggested the classification of this disease as mouse poliomyelitis. In this connection it is worthwhile to mention that the mouse adapted human polioviruses do not behave at all similarly to the true mouse poliomyelitis virus. The latter causes intestinal infection and is excreted during a prolonged period whereas the mouse adapted strains can infect their murine hosts only when introduced directly into the central nervous system.

There is no immunological relationship between the agents causing poliomyelitis like disease of swine, mice and man, whether some animal acts as a reservoir for human polioviruses has not yet been determined. Bartell and Klein¹² and Morton Klein elsewhere in these pages have called attention to the possible role of poliovirus antibodies in the sera of cattle. In this connection I should like to relate a curious experience which if confirmed may throw more light on the problem. Five calves 4 weeks old were placed under observation in separate stalls. Once a week stools were collected from each animal and a blood sample was obtained. Suspensions of the fecal material were inoculated into tissue cultures of monkey renal epithelium, from one sample obtained after 4 weeks of observation there was isolated a virus which was cytopathogenic for tissue cells (FIGURE 1). This virus was subsequently identified as Type I human poliovirus. Results of the neutralization test performed against Type I standard poliovirus serum of the same animal indicated the presence of Type I antibodies at the time of virus isolation. The concentration of antibodies was low but the difference in titers seemed to be insignificant. In order to exclude the possibility of laboratory contamination during the process of isolating the virus the same experiment was repeated using the same stool suspension as source of virus. Again Type I poliovirus was isolated. It is quite obvious that many more factors must be elucidated before absolute confirmation of this curious incident may be obtained. I have cited this experience here in order to encourage other investigators to search




FIGURE 1 The isolation of polio virus Type I from the feces of a calf

for evidence which if confirmed may well point to cattle as hosts for another human pathogen in moderate form

Hepatitis of Man and of Animals

Today many infectious diseases are suppressed both epidemiologically and financially. However infectious hepatitis of man stands out as an important disease entity the etiological agent of which thus far has defied the efforts of the most zealous investigators. Although it has been isolated by passage in human volunteers failure to find for it a suitable animal host and to develop specific serologic reactions seriously impedes progress in the study of this disease.

There are several hepatotropic agents that cause hepatitis syndrome in animals and 4 of these will be discussed briefly (TABLE 4). Three viruses causing hepatitis of mice were detected almost simultaneously in 3 laboratories in 3 different strains of mice¹³⁻¹⁵. One strain was isolated from IRI mice suffering from leukemia¹⁵. There was a tremendous variation in the susceptibility of mice of different strains to induced infection with this agent. The murine hepatitis virus (MHV) of Gledhill and Andrewes¹⁴ can produce fatal hepatitis in mice only in association with *L. peritrichus*¹⁶. The analogy with the role of *Hemophilus influenzae* in swine influenza infection is quite obvious. An association of a normally harmless parasite with a moderate virus resulting in a virulent form of hepatitis has not been observed in other hosts.

Although Rift Valley fever infection of man runs a dengue like course its virus is hepatotropic for sheep and cattle. Focal necrosis of the liver follows a fulminant type of infection in young lambs. In man the morphologic changes following infectious hepatitis are variable but postnecrotic cirrhosis seems to be the most prevalent.

TABLE 4
EXAMPLES OF INFECTIOUS HEPATITIS OF ANIMALS AND OF MAN

Species	Transmission	Latency—day	Incubation—day	Fatality rat	Artificial transmission	Significance
Mouse	Contact	Probably life long	5 to 8		None	None
Sheep and cattle	Mosquito	?	1 to 3	>20/10	Mice hamsters monkeys etc	Rift Valley fever infectious for man†
Dog	Contact	161	6 to 9	10	Foxes tissue culture	Fox encephalitis
Duck	Contact	Unknown	1 to 3	35 to 85	None	None
Man	Contact	>480	20 to 40	0.5	None	None

None if murine hepatitis virus (MHV) alone is injected approaching 100 per cent if together with *Eperitrichus* *coccoides*.

† Apparently not hepatotropic in man dengue-like symptomatology

Comparisons between canine and human infectious hepatitis have been reported by qualified observers and need not be repeated here. If only the etiological agents of these rather similar syndromes were intertransmissible methods of prevention seemingly successful in canine hepatitis could then be applied to the human disease.

Hepatitis of the duck is the newest member to be added to the hepatotropic family.¹⁷⁻¹⁹ Although different in character from the human disease one common feature was observed: the protective effect of antibody containing serum which appears to be similar to that of gamma globulin administered to man.²⁰

Respiratory Viruses

The sense of frustration hounding the author while playing the 'counter parts' of human viral diseases attacking the respiratory tract, was in no small measure occasioned by the prevailing ignorance of the characteristic features of many clinical entities compounded by our lack of knowledge of the numerous agents isolated in the course of respiratory illnesses of either man or animal which are still in search of a disease. The naming (or rather misnaming) of newly discovered entities only adds to the confusion.

For example there is nothing characteristic about pneumonia of man caused by filtrable agents. The roentgenographic findings vary widely in character and distribution and the pulmonary lesions reveal an interstitial type of bronchopneumonia sometimes limited to one lobe sometimes involving two. This condition may be encountered in psittacosis, Q fever, smallpox, chicken pox, measles and lymphocytic choriomeningitis. In addition there are 4 other diseases including the so-called atypical pneumonia in which this ubiquitous bronchopneumonia is seen—diseases caused by an agent or agents unknown. Finding a counterpart of this human disease in animals is like equating an unknown with an unknown. There are certainly many conditions which clinically and pathologically resemble human viral pneumonias; one of these distemper may cause pneumonia in dogs. The problem of cattle pneumonia is even more complicated since it appears that almost every disease of cattle particularly of young calves is characterized by diarrhea and ends in pneumonia (TABLE 5). Calf pneumonia refers to a syndrome first described by Baker¹ in 1942. It is characterized by bilateral pneumonia of the anterior and

TABLE 5
INFECTIONS OF THE RESPIRATORY TRACT OF ANIMALS WITH THEIR HUMAN COUNTERPARTS
(STRETCHING THE POINT OF COMPARISON)

Virus	Animal	Counterpart
Distemper	Dog	Atypical viral or Q fever pneumonia
Calf pneumonia	Cattle	Atypical viral or Q fever pneumonia
Pink eye	Horses	Adenovirus infections
Bullnose	Swine	Common cold
Rhinotracheitis	Cattle	If mild resembles common cold
Snoutsiekte	Cattle	If very mild resembles common cold

ventral lobes and occurs mostly during the winter months. It is apparently caused by an agent that can be adapted to the mouse lung.

In their severe forms infectious rhinotracheitis of Colorado²² and the upper respiratory disease of California²³ will also cause bronchopneumonia. These two conditions apparently caused by immunologically identical agents in their mild form may resemble a very severe type of common cold. However can a disease with a fever of 104.1° accompanied by excessive salivation, accelerated and shallow breathing and anorexia still be classified as a common cold?

There is no dissolution of the turbinate bones in the human common cold and therefore rhinitis of swine is not a counterpart for this condition. Bullnose is believed to be a mild form of the disease characterized by swelling of the mucosa without subsequent distortion of the snout. Whether this resembles the condition of the snout in the common cold is a question that must be left to the imagination rather than to science.

It was difficult to find even a single counterpart for the numerous conditions caused by the adenoviruses in man. The illness of horses referred to in TABLE 5 is really equine influenza. However if we call it pink-eye and stress that the conjunctivitis is accompanied by a catarrhal syndrome we may achieve some success in comparing the two entities. However it should be mentioned that in pink-eye occasionally the function of one or both eyes may be lost and sometimes pneumonia causes death. This is not the case in adenovirus infections.

Gastrointestinal Infections

In a complicated field such as that comprising the so-called virus infections of the gastrointestinal tract our ignorance is truly encyclopedic. Although the alternate problems of diarrhea and constipation are probably as important to man as to animals my feeling is that we are wallowing in a morass of unknowns. The diarrhea of suckling mice⁴ is a bane to every laboratory engaged in mouse breeding (TABLE 6). The explosive outbreaks of this illness often ruin carefully planned experiments in which young mice are to be used as experimental hosts.

TABLE 6
INFECTIONS OF ANIMALS PRIMARILY AFFECTING THE GASTROINTESTINAL TRACT

Species	Age Interval	Fatality rate	Factors affecting	Characteristics	Etiology
Mouse	10 to 15 days	0 to 100	Age season strain	Inclusion bodies	Virus?
Cattle	1 to 2 days	Nearly 100	Age season colostrum feeding	—	CPE virus
	Up to 9 months All ages	0 4 to 8	Age season colostrum feeding Not defined	Elementary bodies —	Pastorosis group Virus?
Swine	<10	100	Age	—	TGE† virus
	All ages	0		—	

Calf pneumonia enteritis

† Transmissible gastroenteritis

Comparisons between canine and human infectious hepatitis have been reported by qualified observers and need not be repeated here. If only the etiological agents of these rather similar syndromes were intertransmissible methods of prevention seemingly successful in canine hepatitis could then be applied to the human disease.

Hepatitis of the duck is the newest member to be added to the hepatotropic family.¹⁻¹⁹ Although different in character from the human disease one common feature was observed: the protective effect of antibody containing serum which appears to be similar to that of gamma globulin administered to man.⁹

Respiratory Viruses

The sense of frustration hounding the author while playing the counter parts' of human viral diseases attacking the respiratory tract, was in no small measure occasioned by the prevailing ignorance of the characteristic features of many clinical entities compounded by our lack of knowledge of the numerous agents isolated in the course of respiratory illnesses of either man or animal which are still in search of a disease. The naming (or rather misnaming) of newly discovered entities only adds to the confusion.

For example there is nothing characteristic about pneumonia of man caused by filtrable agents. The roentgenographic findings vary widely in character and distribution and the pulmonary lesions reveal an interstitial type of bronchopneumonia sometimes limited to one lobe sometimes involving two. This condition may be encountered in psittacosis, Q fever, smallpox, chicken pox, measles and lymphocytic choriomeningitis. In addition there are 4 other diseases including the so called atypical pneumonia in which this ubiquitous bronchopneumonia is seen—diseases caused by an agent or agents unknown. Finding a counterpart of this human disease in animals is like equating an unknown with an unknown. There are certainly many conditions which clinically and pathologically resemble human viral pneumonias; one of these distemper may cause pneumonia in dogs. The problem of cattle pneumonia is even more complicated since it appears that almost every disease of cattle particularly of young calves is characterized by diarrhea and ends in pneumonia (TABLE 5). Calf pneumonia refers to a syndrome first described by Baker¹ in 1942. It is characterized by bilateral pneumonia of the anterior and

TABLE 5

INFECTIONS OF THE RESPIRATORY TRACT OF ANIMALS WITH THEIR HUMAN COUNTERPARTS
(STRETCHING THE POINT OF COMPARISON)

Virus	Animal	Counterpart in man
Distemper	Dog	Atypical viral or Q fever pneumonia
Calf pneumonia	Cattle	Atypical viral or Q fever pneumonia
Pink-eye	Horses	Adenovirus infections
Bullnose	Swine	Common cold
Rhinotracheitis	Cattle	If mild resembles common cold
Snoutsiekte	Cattle	If very mild resembles common cold

ventral lobes and occurs mostly during the winter months. It is apparently caused by an agent that can be adapted to the mouse lung.

In their severe forms infectious rhinotracheitis of Colorado²² and the upper respiratory disease of California²³ will also cause bronchopneumonia. These two conditions, apparently caused by immunologically identical agents in their mild form, may resemble a very severe type of common cold. However, can a disease with a fever of 104° F. accompanied by excessive salivation, accelerated and shallow breathing, and anorexia still be classified as a common cold?

There is no dissolution of the turbinate bones in the human common cold, and therefore rhinitis of swine is not a counterpart for this condition. Bullnose is believed to be a mild form of the disease characterized by swelling of the mucosa without subsequent distortion of the snout. Whether this resembles the condition of the snout in the common cold is a question that must be left to the imagination rather than to science.

It was difficult to find even a single counterpart for the numerous conditions caused by the adenoviruses in man. The illness of horses referred to in TABLE 5 is really equine influenza. However, if we call it pink-eye and stress that the conjunctivitis is accompanied by a catarrhal syndrome, we may achieve some success in comparing the two entities. However, it should be mentioned that in pink-eye, occasionally the function of one or both eyes may be lost, and some times pneumonia causes death. This is not the case in adenovirus infections.

Gastrointestinal Infections

In a complicated field such as that comprising the so called virus infections of the gastrointestinal tract, our ignorance is truly encyclopedic. Although the alternate problems of diarrhea and constipation are probably as important to man as to animals, my feeling is that we are wallowing in a morass of unknowns. The diarrhea of suckling mice⁴ is a bane to every laboratory engaged in mouse breeding (TABLE 6). The explosive outbreaks of this illness often ruin carefully planned experiments in which young mice are to be used as experimental hosts.

TABLE 6
INFECTIONS OF ANIMALS PRIMARILY AFFECTING THE GASTROINTESTINAL TRACT

Sp	Age at onset	Fatality rate, per cent	Factors	Characteristics	Etiology
Mouse	10 to 15 days	0 to 100	Age season strain	Inclusion bodies	Virus?
Cattle	1 to 2 days	Nearly 100	Age season colostrum feeding	—	CPE virus
	Up to 9 months All ages	0 4 to 8	Age season colostrum feeding Not defined	Elementary bodies —	Psittacosis group virus?
Swine	<10 All ages	100 0	Age	— —	TGE† virus

Calf pneumonia enteritis

† Transmissible gastroenteritis

Comparisons between canine and human infectious hepatitis have been reported by qualified observers and need not be repeated here. If only the etiological agents of these rather similar syndromes were intertransmissible methods of prevention seemingly successful in canine hepatitis could then be applied to the human disease.

Hepatitis of the duck is the newest member to be added to the hepatotropic family.¹⁷⁻¹⁹ Although different in character from the human disease, one common feature was observed: the protective effect of antibody containing serum which appears to be similar to that of gamma globulin administered to man.²⁰

Respiratory Viruses

The sense of frustration hounding the author, while playing the counter parts of human viral diseases attacking the respiratory tract, was in no small measure occasioned by the prevailing ignorance of the characteristic features of many clinical entities compounded by our lack of knowledge of the numerous agents isolated in the course of respiratory illnesses of either man or animal which are still "in search of a disease." The naming (or rather misnaming) of newly discovered entities only adds to the confusion.

For example, there is nothing characteristic about pneumonia of man caused by filtrable agents. The roentgenographic findings vary widely in character and distribution and the pulmonary lesions reveal an interstitial type of bronchopneumonia sometimes limited to one lobe, sometimes involving two. This condition may be encountered in psittacosis, Q fever, smallpox, chicken pox, measles, and lymphocytic choriomeningitis. In addition, there are 4 other diseases including the so-called atypical pneumonia in which this ubiquitous bronchopneumonia is seen—diseases caused by an agent or agents unknown. Finding a counterpart of this human disease in animals is like equating an unknown with an unknown. There are certainly many conditions which clinically and pathologically resemble human viral pneumonias; one of these, distemper, may cause pneumonia in dogs. The problem of cattle pneumonia is even more complicated since it appears that almost every disease of cattle, particularly of young calves, is characterized by diarrhea and ends in pneumonia (TABLE 5). Calf pneumonia refers to a syndrome first described by Baker²¹ in 1942. It is characterized by bilateral pneumonia of the anterior and

TABLE 5
INFECTIONS OF THE RESPIRATORY TRACT OF ANIMALS WITH THEIR HUMAN COUNTERPARTS
(STRETCHING THE POINT OF COMPARISON)

Virus	Animal	Human
Distemper	Dog	Atypical viral or Q fever pneumonia
Calf pneumonia	Cattle	Atypical viral or Q fever pneumonia
Pink-eye	Horses	Adenovirus infections
Bullnose	Swine	Common cold
Rhinotracheitis	Cattle	If mild resembles common cold
Snoutsick	Cattle	If very mild resembles common cold

TABLE
ATTEMPTS TO IDENTIFY IN THE LEAST CONFUSING WAY THE CURRENT STATUS OF VIRAL
DIARRHEAL DISEASE OF MAN

Age	Etiology	Transmission with tool filth		
		Species	Symptoms	Serial passages
Newborn	FDN	Calves?	Diarrhea?	29
Children	SDI†	Rat†	Corneal inflammation	Several
All ages	FNCE	Man	Diarrhea	8
		Cat	Diarrhea	Several

Epileptic diarrhea of newborn

† Stomatitis and diarrhea of infants

Epileptic and bacterial gastroenteritis

by nausea, anorexia, vomiting, abdominal spasms and watery stools.³⁷ Filtrates of stools fed to human volunteers caused a similar condition in the hands of several groups of workers³⁷⁻³⁹ and a claim was made that the agent was transmitted serially when administered orally to a cat.³⁹

It is obvious that the comparative pathology of gastrointestinal viral infections is still in its teens and that many years will pass before we may benefit from studies of these conditions in animals in order to improve our knowledge of their human counterparts.

Infections Inducing Fetal Damage

Infections that result in fetal damage constitute perhaps the most fruitful field of research in comparative pathology. Stringent biological and ethical considerations that limit research studies in man are more than amply compensated by the occurrence of similar syndromes in animals. These may occur in the course of a naturally acquired disease or they may be induced by agents biologically active against the fetus *in utero* but not after parturition.

Enzootic abortion of ewes is such an example of a latent viral infection. The dormant agent resembles psittacosis virus⁴⁰ (TABLE 8). It would seem that pregnancy may help to make the infection manifest, resulting in abortion late in gestation. The apparently normal lambs born of infected ewes harbor the latent virus for eight months or more.⁴¹ Development of a vaccine protecting the ewes against abortion has been claimed.⁴²

Some of the ewes injected during the first thirty days of gestation with the so-called attenuated bluetongue virus will give birth to lambs that show arrested development of the central nervous system and the attendant characteristic malformations.⁴³ Apparently this agent is effective only on a fetal host, as no infection can be induced after parturition. A similar condition was observed by Young⁴⁴ in sows when they were injected with an attenuated leporine strain of hog cholera virus during the first month of gestation. The German measles

Strangely the disease does not affect newborns but only mice 10 to 15 days old. In some outbreaks all mice may die within 48 hours; in some cases the animals recover but their growth is retarded, and there are also instances of complete recovery. Pathological lesions are limited to the ileum and colon and intranuclear inclusion bodies were found in the epithelial cells of the villi which resembled those observed in infectious panleukopenia of cats.²⁴ The peak of the disease occurs in winter and its incidence declines in warm weather a situation quite unlike that presented by the human counterpart of a similar ailment. The genetic constitution of the host plays an important role in susceptibility to infection and it is conceivable that genetically controlled breeding could lead to the establishment of a resistant strain of mice.

In contrast to mouse diarrhea the agent of calf pneumonia enteritis (CPE) has been isolated⁶ and transmitted to susceptible newborn animals in an aerosol form which is apparently the only satisfactory way to transmit the disease.⁷⁻⁹ The sharp change in the resistance to the infection with increase in age makes the condition somewhat similar to the epidemic diarrhea of newborn babies (EDN) which apparently can be transmitted to calves. The omnipresent psittacosis virus seems to be one of the agents causing diarrhea of cattle;³ the agent was always present in feces of experimentally infected animals and could be isolated from different parts of the intestinal tract. As in the case of its predecessor (TABLE 6) colostrum feeding had a protective effect.

The next diarrheal entity of cattle is referred to as all ages disease. In 1936 it consisted of three immunologically distinct subentities: two named for reasons of local patriotism as virus diarrhea of New York and virus diarrhea of Indiana while a third was topographically called simply mucosa disease.²⁰ Although diarrhea is the characteristic feature of the disease oral erosions and lamination complete the syndrome. The fatality rate of 4 to 8 per cent is quoted for the New York entity. The mortality rate of the

Indiana diarrhea is apparently much higher; in the mucosa disease it appears that all of the affected animals die.²⁰ Although I have listed this condition I am not familiar with a human counterpart in which man would become lame because he had ten stools per day.

The transmissible gastroenteritis (TGE) of swine is as well defined as it possibly can be in this confused field. The explosive start of the disease in baby pigs characterized by vomiting, profuse diarrhea with yellowish stools resembling curdled milk,^{21, 22} followed by extreme dehydration resembles greatly the neonatal syndrome in babies first described by Rice.²³

Unfortunately whereas the etiological agent of TGE has been isolated immunologically defined²⁴ grown in tissue culture and possibly attenuated²⁵ the identity of the EDN agent of man still remains uncertain (TABLE 7). Although passaged serially in calves and immunologically identified²⁶ it was apparently not tested for its disease producing faculty in newborns and it may be considered as perhaps only one component of a complex etiological entity. Even less is known about stomatitis and diarrhea in infants (SDI).²⁸ The syndrome has been described in various outbreaks but the identity of the agent passaged on rabbit cornea has not been firmly established.

Epidemic nonbacterial gastroenteritis (ENGE) is a syndrome characterized

TABLE 9
VESICULAR DISEASES OF ANIMALS AND MAN

Virus	Incubation day	Mucosal site	Site affected	Virus type	Convalescence—months
Foot and mouth	2 to 14	Vesicle with out areola	Lips Mouth	7	6 to 24
Vesicular stomatitis	2 to 14	Vesicle with out areola	Tongue Pharynx	2	6 to 24
Vesicular exanthema	2 to 14	Vesicle with out areola	Stomach Coronary band of hoofs	7	6 to 24
Herpangina (Coxsackie A)	2 to 9	Isolated vesicles with red areola	Soft palate Tonsils Pharynx	7	0 (Permanent)

the formation of scars. The site of lesions in the oral cavity is similar for all four of these conditions but of course lesions of the coronary band of hoofs remain characteristic for the animal pathogen. It is interesting to note the multiplicity of the virus types that cause the same illness. Seven types of foot and mouth virus are matched by the same number of vesicular exanthema agents⁴⁸ and Coxsackie A strains causing herpangina.⁴⁹ Degenerative heart muscle changes observed in foot and mouth disease have not been described for Coxsackie A viruses but they were observed in newborn infants after Coxsackie B infection.^{40, 41} Another difference lies in the observation of the length of immunity produced: the fleeting type of convalescing immunity is typical for animals whereas a more lasting type of immunity follows Coxsackie A infection.

Papillomas

No self-respecting writer on problems in pathology would dare to close without mentioning some aspect of tumor investigations. There are animal viruses causing cancers or leukemia that cannot be considered as counterparts of nonexistent forms in humans. However, papilloma will save the day (TABLE 10).

The local epithelial hyperplasia which occurs in crops in different locations of the human body⁵⁰ is due to a virus that causes formation of cytoplasmic and intranuclear inclusions and may be seen apparently under an electron microscope.⁵¹ Attempts to infect other species with the human virus have failed. Conversely, the filtrable agent of bovine papillomatosis was supposedly transmitted to man by Schultz in 1908.⁵² I was unable to find any confirmation of this observation but the bovine material was reported to cause tumor formation in horses.⁵³

Lquine and canine papillomatoses greatly resemble their bovine counterpart and in both instances species specificity of the causative agent was maintained. The Shope papilloma of the rabbit has been so well described that there is no need for repetition. However, one may mention that this virus when passed

TABLE 8

EXAMPLES OF INFECTIONS OF ANIMALS CAUSING DAMAGE TO THE FETUS AND THEIR COUNTERPARTS IN HUMAN MEDICINE

Virus	Species	Results of effects during pregnancy	Time of attack
Ovine enzootic abortion	Sheep	Abortion or normal birth	Late in gestation
Attenuated bluetongue	Sheep	Malformation (crazy lambs)	Early in gestation
Equine rhinopneumonitis	Horse	Abortion stillbirth or normal	Late in gestation
Leporine hog cholera	Swine	Abortion stillbirth malformation	Early in gestation
German measles	Man	Abortion stillbirth malformation†	Early in pregnancy

Arrested development of central nervous system

† Blindness deafness cardiovascular abnormalities

syndrome in man presents a similar picture with abortions, stillbirths and malformations

The identities of the viruses of equine abortion and of equine influenza are still uncertain⁴⁵⁻⁴⁶ but their effect on the fetus is similar to that induced by the agent of ovine enzootic abortion occasional stillbirths but also frequent normal births, and/or abortion late in gestation

Concerning these conditions one might ask this provocative question Are the animals born from virus infected females immunologically normal? Will exposure to the same agent encounter increased resistance or increased susceptibility because of the inability to develop antibodies? In other words does the mechanism of immunological tolerance operate in these fetal virus infections? In the case of ovine enzootic abortion it apparently does not

German measles is of ill omen for a woman during the first trimester of pregnancy⁴⁷ American statistics indicate that abortions stillbirths and malformations consequent to this disease are probably as frequent as normal births The direct toxic role of the agent has not yet been proved and studies in animals would doubtless help to clarify the entire problem

I may add that obviously there are other viral infections of pregnant animals which may cause abortions stillbirths and malformations However the four examples cited above are so characteristic and bear so much more upon the problem of the human syndrome that the others were omitted from consideration

Vesicular Diseases

It was not easy to find counterparts in human medicine to the three vesicular diseases of animals As is well known the host range and type of lesions do not permit any easy differential clinical diagnosis of the three agents foot and mouth disease vesicular stomatitis and vesicular exanthema (TABLE 9) The mucosal lesions are similar to those observed in herpangina of man minute vesicles filled with clear fluid which in the case of foot and mouth virus rupture and leave denuded areas In the case of vesicular stomatitis vesicular exanthema and herpangina the vesicles heal usually promptly and without

TABLE 11
EXAMPLES OF LACK OF COUNTERPARTS

Animal disease with which human counterpart	Human disease with which animal counterpart
Pseudorabies Scrapie	Exanthematous diseases roseola infectious mononucleosis etc

It is an intriguing thought that perhaps one day children will be vaccinated with distemper virus against measles dogs with rinderpest virus against distemper and cattle with measles virus against rinderpest *Se non è vero è ben trovato* *

No Counterparts

As a *gran finale* to this disquisition the following table is presented to show viral diseases for which I was unable to find any counterparts (TABLE 11). The two animal diseases are characterized by an itch. The only human affliction conceivable as a counterpart is the seven year itch and that is a much more pleasant condition although it is perhaps exasperating at times than the animal syndrome.

Conversely animal skin seems to be devoid of ingredients that would make it susceptible to the exanthema with which the human species is blessed during every viral disease of childhood. Thus if chicken pox is the price we must pay for enjoying the seven year itch we should accept this bargain with all humility for the poor sheep while undergoing a three year incubation period of scrapie⁴³ are not recompensed with a delightful itch.

References

- 1 SHAW G B 1934 Preface to The I evolutionist's Handbook and Pocket Companion. In Prefaces Constable & Co London England
- 2 CASALS J 1957 Trans N Y Acad Sci Ser II 19 219
- 3 GILYARD R T 1944 Bull U S Army Med Dept 75 96
- 4 LENNETTE F H & H KOPROWSKI 1943 J Am Med Assoc 123 1094
- 5 KOPROWSKI H & H R COX 1947 New Engl J Med 236 647
- 6 HORSFALL F L 1955 In Cecil's Textbook of Medicine 75-77 Saunders Philadelphia Pa
- 7 TRACB E 1939 J Exptl Med 69 801
- 8 ARMSTRONG C J J WALLACE & L ROSS 1940 U S Public Health Rept 55 1222
- 9 SMADEL J E R H GREEN R M IALITALY & T A GONZALES 1942 Proc Soc Exptl Biol Med 49 683
- 10 DOBERSTEIN J 1940 Z Infektionskrankh parasit Krankh u Hyg Haustiere 48 213
- 11 HORSTMANN D M 1952 J Immunol 69 379
- 12 BARTLETT P & M KLEIN 1955 Proc Soc Exptl Biol Med 90 597
- 13 JORDAN J & G S MIRICK 1951 Bull Johns Hopkins Hosp 89 326
- 14 GLEDHILL A W & C H ANDREWS 1951 Brit J Exptl Pathol 32 559
- 15 NELSON J B 1957 J Exptl Med 96 293
- 16 GLEDHILL A W & J S I NIVEN 1955 Vet Rev Anat 1 82
- 17 LEVINE P P & J FABRICANT 1950 C ell Vet 40 71
- 18 ASPLIN F D & J D McLAUCHLAN 1954 Vet Record 66 456
- 19 HANSON L F & J O ALBERTS 1956 J Am Vet Med Assoc 128 37

If it is not true it is well invented (GIORDANO BRUNO)

TABLE 10
PAPILLOMAS OF ANIMALS AND MAN

Ser	Location	Demonstration
Cow	Head neck	Filtrates induce papilloma in cattle man(?) and horse(?)
Horse	Nose lips	Transmission to horses
Dog	Mouth lips	Filtrates induce papillomas in dogs
Rabbit (cottontail)	Neck shoulders abdomen	Shope papilloma
Man	Palms and soles	Transmission by filtrates to man electron microscopy

into domestic rabbits but not to cottontails gives rise to papillomas that subsequently may undergo malignant transformation⁵⁶ This has not been observed to my knowledge in case of human warts but these can be treated by psychotherapy a method not as yet applicable to cottontails

New Relations Measles Distemper and Rinderpest

For the past several years suspicions had been harbored that the distemper virus may possibly be pathogenic for man Appearance of distemper antibodies in sera of children recovering from unidentified infections of the upper respiratory tract inclined scientists⁵⁷⁻⁵⁸ to believe that an immunological kinship of the two conditions might exist⁵⁹ in recent years the presence of distemper antibodies in the blood of man has been confirmed by Karzon⁶⁰ and by Carlstrom⁶¹ The presence of antibodies in relation to age of the donors revealed a pattern commonly encountered in other viral infections of man newborn infants showed high incidence of antibodies which dropped to insignificant levels by the age of 6 months but from the second year incidence rose rapidly paralleling the age curve and antibodies were present in all of the sera of adults studied⁵⁹⁻⁶¹

The relationship between the presence of distemper antibodies and measles infection of man was established by Carlstrom⁶¹ who found at least a fourfold rise in distemper antibody titer in sera of 11 of 14 patients convalescing from measles In addition there was an evident correlation between the history of measles and the appearance of distemper antibodies in 37 out of 38 patients It is quite obvious that these observations require further investigation to determine the true nature of this most interesting relationship

To complicate the picture further a preliminary report from workers in Kenya Africa would seem to indicate a possible protective effect of induced asymptomatic rinderpest infection against distemper challenge⁶² Four test puppies pretreated with rinderpest virus showed no signs of distemper when challenged with distemper virus whereas all four control puppies developed signs of distemper and one of them died Here too confirmation of this relationship should be sought

COMPARATIVE STUDIES ON VIRAL HEPATITIS IN ANIMALS AND IN MAN*

By Morris Pollard

University of Texas Medical Branch Galveston Texas

Studies with human viral hepatitis have been handicapped by the lack of a susceptible experimental animal and of a specific diagnostic test. This was amply documented at a recent symposium on the laboratory propagation and detection of the agent of hepatitis (National Academy of Sciences-National Research Council Washington D. C. 1954). A subsequent report on the propagation of human viral hepatitis agents in the Detroit 6 strain of human epithelium (Right et al. 1956) has not yet been confirmed. The significance of the cytopathogenic effect noted in the latter publication must yet be supported by human infectivity trials with the agents being propagated.

The embryonated chicken egg has been used as a medium for the propagation of hepatitis agents; however, thus far adaptation of hepatitis viruses to the chick embryo has not been proved (Henle 1954). While hepatitis agents have induced no demonstrable lesions in the embryonated egg, recent observations indicate that influenza and mumps viruses induce an increased number of mononuclear cells in the chorioallantoic fluid (CAF) of inoculated but asymptomatic embryos (Pollard and Diserens 1955, Buchner et al. 1956). Serum specimens from acute cases of human viral hepatitis appeared to induce a similar response (Pollard and Diserens 1956) however, this has not yet been confirmed (Buchner et al. 1956).

Since the virus or viruses of hepatitis in man are not transmissible to experimental animals (MacCallum 1954, Morris, O'Conner and Coburn 1954) since they induce questionable changes in cultures of human tissues and since they bring about controversial changes in the embryonated egg and indistinct serologic reactions, it was deemed advisable to study the characteristics of a laboratory prototype of hepatitis virus. Some useful information from this might be applicable to the laboratory study of viral hepatitis of man.

Murine viral hepatitis shows some resemblance to the human disease (Friend, Wroblewski and LaDue 1955, DeRitis, Coltorti and Giusti 1956). It shows a narrow species susceptibility. It has not been propagated in fertile eggs (Gledhill, Dick and Niven 1953) and its effect in tissue cultures is uncertain. We are indebted to A. W. Gledhill, National Institute for Medical Research, Mill Hill, London, England, for kindly providing us with a strain of mouse hepatitis virus (MHV) that was free of *L. peritrichus* (Niven, Gledhill and Andrewes 1952). This virus was lethal for newborn mice, usually killing them at 6 to 8 days following intra abdominal inoculation and causing extensive degenerative lesions in the liver. We are grateful to J. B. Nelson, The Rockefeller Institute for Medical Research, New York, N. Y. for

*The work reported in this paper was supported by funds provided by Contract AF 41 (657) 123 with the School of Aviation Medicine, USAF, Randolph Air Force Base, Texas.

- 20 HAGAN W A & D W BRUNER 1957 *The Infectious Diseases of Domestic Animals*
Comstock Ithaca N Y
- 21 BAKER J A 1942 *Cornell Vet* 32 202
- 22 JENSEN I I A CRISER F I CHOW & W W BROWN 1955 *Proc Livestock San*
Assoc 69 189
- 23 MCKIRCHER D C J I MOULTON J W KENDRICK & J SAITO 1955 *Proc Live*
stock San Assoc 69 151
- 24 CHIFFAR I S 1956 *Ann N Y Acad Sci* 66(1) 196
- 25 LAFFENHIMER A M & I S CHIFFAR 1948 *J Exptl Med* 88 317
- 26 MOLT I & C A BRANDIV 1954 *Vet Med* 49 512
- 27 MOLT I & C A BRANDIV 1955 *Vet Med* 50 101
- 28 BRANDIV C A & A W MCCURKIN 1956 *Ann N Y Acad Sci* 66(1) 181
- 29 YORK C J & J A BAKER 1951 *J Exptl Med* 93 587
- 30 BRITCHARD W I R C CARLSON H I MOFFS & D I B TAYLOR 1955 *Proc*
Livestock San Assoc 69 173
- 31 BAY W W I I DOYLE & I M HUTCHINGS 1951 *Am J Vet Research* 12 215
- 32 HAUPTMANN I O & I M HUTCHINGS 1956 *Ann N Y Acad Sci* 66(1) 186
- 33 FICK J I W H BEST S IRANI & H ANRAM ON 1947 *J Am Med Assoc* 109
475
- 34 LEE K M 1956 *Ann N Y Acad Sci* 66(1) 191
- 35 HODDS H I 1956 *Ann N Y Acad Sci* 66(1) 215
- 36 BUDDINCH C J & K DODD 1944 *J Infect* 26 105
- 37 CORDON I & I WHITNEY 1956 *Ann N Y Acad Sci* 66(1) 220
- 38 KOFIMA S H IUKUMI & I ISHIMARU 1953 *Japan J Med Sci Biol* 6 69
- 39 YAMAMOTO A H M ZENNYOJI K YANAGITA & S KATO 1948 *Japan Med J*
1 379
- 40 STAM J I 1951 *J Comp Pathol Therap* 61 215
- 41 McI WEN A D A J LITTLEJOHN & A LOCCIE 1951 *Vet Record* 63 489
- 42 McI WEN A D & A LOCCIE 1954 *Vet Record* 68 393
- 43 SCHULTZ C I D DELAY 1955 *J Am Vet Med Assoc* 127 224
- 44 YOUNG C A L I KITCHEN A J LUDAK & J H SAUTTER 1955 *J Am Vet*
Med Assoc 126 165
- 45 MANNINGER R 1949 *Acta Vet Hung* 1 62
- 46 DOTT I R M C RICHARDS & M I WALLACE 1954 *Cornell Vet* 44 133
- 47 SWAN C 1944 *J Pathol Bacteriol* 66 282
- 48 BANKOWSKI I A A C FERRIS I I STUART & M KUMMER 1955 *Proc Live*
stock San Assoc 69 356 *No personal communication*
- 49 FARROTT I H 1957 *Ann N Y Acad Sci* 67(8) 230
- 50 MONTGOMERY J J CEAR M IRIN LOO M KAHN & Z C KIRSCH 1956 *S African*
Med J 29 608
- 51 KIBRICK S & K BENIE CHKE 1956 *New Engl J Med* 255 883
- 52 BLANK H & C LAKE 1955 *Viral and Bacterial Diseases of the Skin Eye and*
Mucous Membranes of Man 156 Little Brown and Co Boston Mass
- 53 MPELNICK J I H BENTING W C BANFIELD M J STRAUSS & W H CAYLORD 1952
Ann N Y Acad Sci 64(1) 1214
- 54 SCHULTZ I 1908 *Deut med W hscrh* 34 423
- 55 OLSON C I H COOK 1951 *Proc Soc Exptl Biol Med* 77 281
- 56 KIDD J C & I LOUS 1940 *J Exptl Med* 71 469
- 57 ADAMS J M 1953 *Pediatrics* 11 15
- 58 ADAM J M & D I IMACAWA 1957 *Pediat Clin N Am* 193
- 59 CARLSTRÖM C 1956 *Vet Pathol* 45 180
- 60 KARZON D T 1955 *Pediatr* 10 809
- 61 CARLSTRÖM C 1957 *Pediatr* 2 344
- 62 FOLDING J B & I M SIMON 1957 *Vet Record* 69 582
- 63 KUMMER I J C 1946 *Cat J Comp Med* 10 49

COMPARATIVE STUDIES ON VIRAL HEPATITIS IN ANIMALS AND IN MAN*

By Morris Pollard

University of Texas Medical Branch Galveston Texas

Studies with human viral hepatitis have been handicapped by the lack of a susceptible experimental animal and of a specific diagnostic test. This was amply documented at a recent symposium on the laboratory propagation and detection of the agent of hepatitis (National Academy of Sciences-National Research Council Washington D C 1954). A subsequent report on the propagation of human viral hepatitis agents in the Detroit 6 strain of human epithelium (Rightsel *et al* 1956) has not yet been confirmed. The significance of the cytopathogenic effect noted in the latter publication must yet be supported by human infectivity trials with the agents being propagated.

The embryonated chicken egg has been used as a medium for the propagation of hepatitis agents; however, thus far adaptation of hepatitis viruses to the chick embryo has not been proved (Henle 1954). While hepatitis agents have induced no demonstrable lesions in the embryonated egg, recent observations indicate that influenza and mumps viruses induce an increased number of mononuclear cells in the chorioallantoic fluid (CAF) of inoculated but asymptomatic embryos (Pollard and Diserens 1955; Buchner *et al* 1956). Serum specimens from acute cases of human viral hepatitis appeared to induce a similar response (Pollard and Diserens 1956) however, this has not yet been confirmed (Buchner *et al* 1956).

Since the virus or viruses of hepatitis in man are not transmissible to experimental animals (MacCallum 1954; Morris O'Connor and Coburn 1954) since they induce questionable changes in cultures of human tissues and since they bring about controversial changes in the embryonated egg and indistinct serologic reactions, it was deemed advisable to study the characteristics of a laboratory prototype of hepatitis virus. Some useful information from this might be applicable to the laboratory study of viral hepatitis of man.

Murine viral hepatitis shows some resemblance to the human disease (Friend, Wroblewski and LaDue 1955; DeRitis, Coltorti and Giusti 1956). It shows a narrow species susceptibility. It has not been propagated in fertile eggs (Gledhill, Dick and Niven 1953) and its effect in tissue cultures is uncertain. We are indebted to A W Gledhill, National Institute for Medical Research, Mill Hill, London, England, for kindly providing us with a strain of mouse hepatitis virus (MHV) that was free of *I perythron coccoides* (Niven, Gledhill and Andrewes 1952). This virus was lethal for newborn mice, usually killing them at 6 to 8 days following intra-abdominal inoculation, and causing extensive degenerative lesions in the liver. We are grateful to J B Nelson, The Rockefeller Institute for Medical Research, New York, N Y, for

*The work reported in this paper was supported by funds provided by Contract AF 41 (657) 123 with the School of Aviation Medicine, USAF Randolph Air Force Base, Texas.

- 20 HAGAN W A & D W BRUNER 1957 The Infectious Diseases of Domestic Animal
Cornstock Ithaca N Y
- 21 BAKER J A 1942 Cornell Vet 32 202
- 22 JENSEN R L A GRINER T L CHOW & W W BROWN 1955 Proc Livestock San
Assoc 59 189
- 23 MCKERCHER D G J E MOULTON J W KENDRICK & J SAITO 1955 Proc Live
stock San Assoc 59 151
- 24 CHEEVER I S 1956 Ann N Y Acad Sci 66(1) 196
- 25 PAPPENHEIMER A M & F S CHEEVER 1948 J Exptl Med 88 317
- 26 MOLL T & C A BRANDLY 1954 Vet Med 49 512
- 27 MOLL T & C A BRANDLY 1955 Vet Med 60 101
- 28 BRANDLY C A & I W MCCLURKIN 1956 Ann N Y Acad Sci 66(1) 181
- 29 YORK C J & J A BAKER 1951 J Exptl Med 93 587
- 30 PRITCHARD W R R C CARLSON H E MOSES & D E B TAYLOR 1955 Proc
Livestock San Assoc 59 173
- 31 BAY W W L P DOYLE & L M HUTCHINGS 1951 Am J Vet Research 12 215
- 32 HAELTERMANN I O & L M HUTCHINGS 1956 Ann N Y Acad Sci 66(1) 186
- 33 RICE J L W H REST S FRANT & H ABRAMSON 1937 J Am Med Assoc 109
475
- 34 LEE K M 1956 Ann N Y Acad Sci 66(1) 191
- 35 HODES H I 1956 Ann N Y Acad Sci 66(1) 215
- 36 BUDDINGH C J & K DODD 1944 J Pediat 25 105
- 37 GORDON I & F WHITNEY 1956 Ann N Y Acad Sci 66(1) 270
- 38 KOJIMA S H FUKUMI & T ISHIMARU 1953 Japan J Med Sci Biol 6 69
- 39 YAMAMOTO A H M ZENNYOJI K YANAGITA & S KATO 1948 Japan Med J
1 349
- 40 STAMP J T 1951 J Comp Pathol Therap 61 215
- 41 McEWEN A D A J LITTLEJOHN & A FOGGIE 1951 Vet Record 63 489
- 42 McEWEN A D & A FOGGIE 1954 Vet Record 68 393
- 43 SCHULTZ G & P D DELAY 1955 J Am Vet Med Assoc 127 224
- 44 YOUNG G A I L KITCHELL A J LUEDKE & J H SAUTTER 1955 J Am Vet
Med Assoc 126 165
- 45 MANNINGER R 1949 Acta Vet Hung 1 67
- 46 DOLL F R M G RICHARDS & M I WALLACE 1954 Cornell Vet 44 133
- 47 SWAN C 1944 J Pathol Bacteriol 56 289
- 48 BANKOWSKI R A A G PERKINS E F STUART & M KUMMER 1955 Proc Live
stock San Assoc 59 356 Also personal communication
- 49 FARROTT R H 1957 Ann N Y Acad Sci 67(8) 230
- 50 MONTGOMERY J J GEAR M PRINSLOO M KAHN & Z G KIRSCH 1956 S African
Med J 29 609
- 51 KIBRICK S & K BENZESCHKE 1956 New Engl J Med 255 883
- 52 BLANA H & G RAKE 1955 Viral and Rickettsial Diseases of the Skin Eye and
Mucous Membranes of Man 156 Little Brown and Co Boston Mass
- 53 MELNICK J I H BLUNTING W G BANFIELD M J STRAUSS & W H GAYLORD 1952
Ann N Y Acad Sci 54(6) 1214
- 54 SCHULTZ F 1908 Deut med Wochschr 34 423
- 55 OLSON C & R H COOK 1951 Proc Soc Exptl Biol Med 77 281
- 56 KIDD J G & P ROLS 1940 J Exptl Med 71 469
- 57 ADAMS J M 1953 Pediatrics 11 15
- 58 ADAMS J M & D T IMAGAWA 1957 Pediat Clin N Am 113
- 59 CARLSTRÖM G 1956 Acta Pediat 45 180
- 60 KARZON D T 1955 Pediatrics 16 809
- 61 CARLSTRÖM G 1957 Lancet 2 344
- 62 FOLDING J B & R M SIMPSON 1957 Vet Record 69 582
- 63 PILUMMER P J G 1946 Can J Comp Med 10 49

COMPARATIVE STUDIES ON VIRAL HEPATITIS IN ANIMALS AND IN MAN*

By Morris Iollard

University of Texas Medical Branch Galveston Texas

Studies with human viral hepatitis have been handicapped by the lack of a susceptible experimental animal and of a specific diagnostic test. This was amply documented at a recent symposium on the laboratory propagation and detection of the agent of hepatitis (National Academy of Sciences-National Research Council Washington D. C. 1954). A subsequent report on the propagation of human viral hepatitis agents in the Detroit 6 strain of human epithelium (Richtsels *et al.* 1956) has not yet been confirmed. The significance of the cytopathogenic effect noted in the latter publication must yet be supported by human infectivity trials with the agents being propagated.

The embryonated chicken egg has been used as a medium for the propagation of hepatitis agents; however, thus far adaptation of hepatitis viruses to the chick embryo has not been proved (Henle 1954). While hepatitis agents have induced no demonstrable lesions in the embryonated egg, recent observations indicate that influenza and mumps viruses induce an increased number of mononuclear cells in the chorioallantoic fluid (CAF) of inoculated but asymptomatic embryos (Iollard and Diserens 1955, Buchner *et al.* 1956). Serum specimens from acute cases of human viral hepatitis appeared to induce a similar response (Pollard and Diserens 1956) however, this has not yet been confirmed (Buchner *et al.* 1956).

Since the virus or viruses of hepatitis in man are not transmissible to experimental animals (MacCallum 1954, Morris, O'Conner and Coburn 1954) since they induce questionable changes in cultures of human tissues and since they bring about controversial changes in the embryonated egg and indistinct serologic reactions, it was deemed advisable to study the characteristics of a laboratory prototype of hepatitis virus. Some useful information from this might be applicable to the laboratory study of viral hepatitis of man.

Murine viral hepatitis shows some resemblance to the human disease (Friend, Wroblewski and LaDue 1955, DeRitis, Coltorti and Giusti 1956). It shows a narrow species susceptibility. It has not been propagated in fertile eggs (Gledhill, Dick and Niven 1953) and its effect in tissue cultures is uncertain. We are indebted to A. W. Gledhill, National Institute for Medical Research, Mill Hill, London, England, for kindly providing us with a strain of mouse hepatitis virus (MHV) that was free of *I. perythronum coecoides* (Niven, Gledhill and Andrewes 1952). This virus was lethal for newborn mice, usually killing them at 6 to 8 days following intra abdominal inoculation and causing extensive degenerative lesions in the liver. We are grateful to J. B. Nelson, The Rockefeller Institute for Medical Research, New York, N. Y., for

*The work reported in this paper was supported by funds provided by Contract AF 41 (657)-123 with the School of Aviation Medicine, USAF Randolph Air Force Base, Texas.

- 20 HAGAN W A & D W BRUNER 1957 *The Infectious Diseases of Domestic Animals*
Comstock Ithaca N Y
- 21 BAKER J A 1942 *Cornell Vet* 32 207
- 22 JENSEN R L A GRINER T L CHOW & W W BROWN 1955 *Proc Livestock Sci Assoc* 69 189
- 23 MCKENZIE D G J E MOLLITON J W KENDRICK & J SAITO 1955 *Proc Livestock Sci Assoc* 69 151
- 24 CHEEVER F S 1956 *Ann N Y Acad Sci* 66(1) 196
- 25 PAPPENHIMER A M & F S CHEEVER 1948 *J Exptl Med* 88 317
- 26 MOLL T & C A BRANDLY 1954 *Vet Med* 49 512
- 27 MOLL T & C A BRANDLY 1955 *Vet Med* 50 101
- 28 BRANDLY C A & A W MCCLELLAN 1956 *Ann N Y Acad Sci* 66(1) 181
- 29 YOKA C J & J A BAKER 1951 *J Exptl Med* 93 587
- 30 PRITCHARD W R R G CARLSON H E MOSES & D E B TAYLOR 1955 *Proc Livestock Sci Assoc* 69 173
- 31 BAY W W L P DOYLE & L M HUTCHINGS 1951 *Am J Vet Research* 12 215
- 32 HAEUTERMANN F O & L M HUTCHINGS 1956 *Ann N Y Acad Sci* 66(1) 186
- 33 RICE J L W H BEST S FRANT & H ABRAHAMSON 1937 *J Am Med Assoc* 109 475
- 34 LEE K M 1956 *Ann N Y Acad Sci* 66(1) 191
- 35 HODES H I 1956 *Ann N Y Acad Sci* 66(1) 215
- 36 BLDDING C J & K DODD 1944 *J Pediat* 25 105
- 37 GORDON I & E WHITNEY 1956 *Ann N Y Acad Sci* 66(1) 270
- 38 KOJIMA S H FUKUMI & T ISHIMARU 1953 *Japan J Med Sci Biol* 6 69
- 39 YAMAMOTO A H M ZENNYOJI K YANAGITA & S KATO 1948 *Japan Med J* 1 39
- 40 STAMP J T 1951 *J Comp Pathol Therap* 61 215
- 41 McEWEN A D A J LITTLEJOHN & A FOGGIE 1951 *Vet Record* 63 489
- 42 McEWEN A D & A FOGGIE 1954 *Vet Record* 68 393
- 43 SCHULTZ G & P D DELAY 1955 *J Am Vet Med Assoc* 127 274
- 44 YOUNG G A R L KITCHELL A J LUEDKE & J H SAUTTER 1955 *J Am Vet Med Assoc* 126 165
- 45 MANNINGER R 1949 *Acta Vet Hung* 1 67
- 46 DOLL F R M G RICHARDS & M E WALLACE 1954 *Cornell Vet* 44 133
- 47 SWAN C 1944 *J Pathol Bacteriol* 66 289
- 48 BANKOW NI R A A G FERRIS I E STUART & M KUMMER 1955 *Proc Livestock Sci Assoc* 69 356 Also personal communication
- 49 FARROTT R H 1957 *Ann N Y Acad Sci* 67(8) 230
- 50 MONTGOMERY J J CHAR M PRINSLOO M KAHN & Z G KIRSCH 1956 *S Afr J Med* 29 606
- 51 KIBRICK S & K BENIRSCHKE 1956 *New Engl J Med* 255 883
- 52 BLANK H & G RADE 1955 *Viral and Rickettsial Diseases of the Skin Eye and Mucous Membranes of Man* 156 Little Brown and Co Boston Mass
- 53 MELNICK J I H BLINTING W C HANFIELD M J STRAUSS & W H GAYLORD 1952 *Ann N Y Acad Sci* 54(6) 1214
- 54 SCHULTZ F 1908 *Dent med Wochschr* 34 423
- 55 OLSON C & R H COOK 1951 *Proc Soc Exptl Biol Med* 77 281
- 56 KIDD J G & P ROUS 1940 *J Exptl Med* 71 469
- 57 ADAMS J M 1953 *Pediatrics* 11 15
- 58 ADAMS J M & D T IMAGAWA 1957 *Pediat Clin N Am* 193
- 59 CARLSTRÖM G 1956 *Acta Pediat* 45 180
- 60 KARZON D T 1955 *Pediatrics* 16 809
- 61 CARLSTRÖM G 1957 *Lancet* 2 344
- 62 FOLDING J B & R M SIMPSON 1957 *Vet Record* 69 582
- 63 FLOOMER P J G 1946 *Can J Comp Med* 10 49

COMPARATIVE STUDIES ON VIRAL HEPATITIS IN ANIMALS AND IN MAN*

By Morris Pollard

University of Texas Medical Branch Galveston Texas

Studies with human viral hepatitis have been handicapped by the lack of a susceptible experimental animal and of a specific diagnostic test. This was amply documented at a recent symposium on the laboratory propagation and detection of the agent of hepatitis (National Academy of Sciences-National Research Council Washington D. C. 1954). A subsequent report on the propagation of human viral hepatitis agents in the Detroit 6 strain of human epithelium (Rightsel *et al.* 1956) has not yet been confirmed. The significance of the cytopathogenic effect noted in the latter publication must yet be supported by human infectivity trials with the agents being propagated.

The embryonated chicken egg has been used as a medium for the propagation of hepatitis agents; however, thus far adaptation of hepatitis viruses to the chick embryo has not been proved (Henle 1954). While hepatitis agents have induced no demonstrable lesions in the embryonated egg, recent observations indicate that influenza and mumps viruses induce an increased number of mononuclear cells in the chorioallantoic fluid (CAF) of inoculated but asymptomatic embryos (Pollard and Diserens 1955; Buchner *et al.* 1956). Serum specimens from acute cases of human viral hepatitis appeared to induce a similar response (Pollard and Diserens 1956); however, this has not yet been confirmed (Buchner *et al.* 1956).

Since the virus or viruses of hepatitis in man are not transmissible to experimental animals (MacCallum 1954; Morris, O'Conner and Coburn 1954) since they induce questionable changes in cultures of human tissues, and since they bring about controversial changes in the embryonated egg and indistinct serologic reactions, it was deemed advisable to study the characteristics of a laboratory prototype of hepatitis virus. Some useful information from this might be applicable to the laboratory study of viral hepatitis of man.

Murine viral hepatitis shows some resemblance to the human disease (Friend, Wroblewski and LaDue 1955; DeRitis, Coltorti and Giusti 1956). It shows a narrow species susceptibility. It has not been propagated in fertile eggs (Gledhill, Dick and Niven 1953) and its effect in tissue cultures is uncertain. We are indebted to A. W. Gledhill, National Institute for Medical Research, Mill Hill, London, England, for kindly providing us with a strain of mouse hepatitis virus (MHV) that was free of *Eperythrona coccoides* (Niven, Gledhill and Andrewes 1952). This virus was lethal for newborn mice, usually killing them at 6 to 8 days following intra-abdominal inoculation, and causing extensive degenerative lesions in the liver. We are grateful to J. B. Nelson, The Rockefeller Institute for Medical Research, New York, N. Y., for

*The work reported in this paper was supported by funds provided by Contract AF 41 (657) 123 with the School of Aviation Medicine, USAF Randolph Air Force Base, Texas.

- 20 HAGAN W A & D W BRUNER 1957 *The Infectious Diseases of Domestic Animals*
Comstock Ithaca N Y
- 21 BAKER J A 1942 *Cornell Vet* 32 202
- 22 JENSEN R J A GRINER T L CHOW & W W BROWN 1955 *Proc Livestock San*
Assoc 59 189
- 23 McHERCHER D G J E MOULTON J W KENDRICK & J SAITO 1955 *Proc Live*
stock San Assoc 59 151
- 24 CHEEVER F S 1956 *Ann N Y Acad Sci* 66(1) 196
- 25 PAPPENHEIMER A M & F S CHEEVER 1948 *J Exptl Med* 88 317
- 26 MOLL T & C A BRANDLY 1954 *Vet Med* 49 512
- 27 MOLL T & C A BRANDLY 1955 *Vet Med* 50 101
- 28 BRANDLY C A & A W McCLURKIN 1956 *Ann N Y Acad Sci* 66(1) 181
- 29 YORA C J & J A BAKER 1951 *J Exptl Med* 93 587
- 30 PRITCHARD W P R G CARLSON H E MOSES & D E B TAYLOR 1955 *Proc*
Livestock San Assoc 59 173
- 31 BAY W W L P DOYLE & L M HUTCHINGS 1951 *Am J Vet Research* 12 215
- 32 HAEUTERMANN E O & L M HUTCHINGS 1956 *Ann N Y Acad Sci* 66(1) 186
- 33 RICE J L W H BEST S FRANT & H ABRAMSON 1937 *J Am Med Assoc* 109
475
- 34 LEE K M 1956 *Ann N Y Acad Sci* 66(1) 191
- 35 HODES H L 1956 *Ann N Y Acad Sci* 66(1) 215
- 36 RUDDINCH G J & R DODD 1944 *J Pediat* 25 105
- 37 GORDON I & E WHITNEY 1956 *Ann N Y Acad Sci* 66(1) 220
- 38 KOJIMA S H FUKUMI & T ISHIMARU 1953 *Japan J Med Sci Biol* 6 69
- 39 YAMAMOTO A H M ZENNYOJI K IANAGITA & S KATO 1948 *Japan Med J*
1 319
- 40 STAMP J T 1951 *J Comp Pathol Therap* 61 215
- 41 McEWEN A D A J LITTLEJOHN & A FOGGIE 1951 *Vet Record* 63 489
- 42 McEWEN A D & A FOGGIE 1954 *Vet Record* 68 393
- 43 SCHULTZ G & P D DELAY 1955 *J Am Vet Med Assoc* 127 224
- 44 YOUNG G A R L MITCHELL A J IVECKE & J H SAUTTER 1955 *J Am Vet*
Med Assoc 126 165
- 45 MANNINGER R 1949 *Acta Vet Hung* 1 67
- 46 DOLL F R M G RICHARDS & M E WALLACE 1954 *Cornell Vet* 44 133
- 47 SWAN C 1944 *J Pathol Bacteriol* 56 289
- 48 BANKOWSKI R A A G PERKINS E F STUART & M KUMMER 1955 *Proc Live*
stock San Assoc 59 356 Also personal communication
- 49 PARROTT R H 1951 *Ann N Y Acad Sci* 67(8) 230
- 50 MONTGOMERY J J GEAR M PRINSLOO M KAHN & Z G KIRSCH 1956 *S African*
Med J 29 603
- 51 KIRRIKA S & K BENIRSCHKE 1956 *New Engl J Med* 255 893
- 52 BLANK H & G RAKE 1955 *Viral and Rickettsial Diseases of the Skin Eye and*
Mucous Membranes of Man 156 Little Brown and Co Boston Mass
- 53 MELNICK J L H BUNTING W G BANFIELD M J STRALES & W H GAYLORD 1952
Ann N Y Acad Sci 54(6) 1214
- 54 SCHULTZ F 1908 *Deut med Wochschr* 34 423
- 55 OLSON C & R H COOK 1951 *Proc Soc Exptl Biol Med* 77 281
- 56 KIDD J G & P ROLS 1940 *J Exptl Med* 71 469
- 57 ADAMS J M 1953 *Pediatrics* 11 15
- 58 ADAMS J M & D T IMAGAWA 1957 *Pediat Clin N Am* 194
- 59 CARLSTRÖM C 1956 *Acta Pediat* 45 180
- 60 KARZON D T 1955 *Pediatrics* 16 809
- 61 CARLSTRÖM C 1957 *Lancet* 2 344
- 62 FOLDING J B & R M SIMPSON 1957 *Vet Record* 69 582
- 63 PLUMMER P J G 1946 *Can J Comp Med* 10 49

COMPARATIVE STUDIES ON VIRAL HEPATITIS IN ANIMALS AND IN MAN*

By Morris Pollard

University of Texas Medical Branch Galveston Texas

Studies with human viral hepatitis have been handicapped by the lack of a susceptible experimental animal and of a specific diagnostic test. This was amply documented at a recent symposium on the laboratory propagation and detection of the agent of hepatitis (National Academy of Sciences-National Research Council Washington D. C. 1954). A subsequent report on the propagation of human viral hepatitis agents in the Detroit 6 strain of human epithelium (Richtel *et al.* 1956) has not yet been confirmed. The significance of the cytopathogenic effect noted in the latter publication must yet be supported by human infectivity trials with the agents being propagated.

The embryonated chicken egg has been used as a medium for the propagation of hepatitis agents; however, thus far adaptation of hepatitis viruses to the chick embryo has not been proved (Henle 1954). While hepatitis agents have induced no demonstrable lesions in the embryonated egg, recent observations indicate that influenza and mumps viruses induce an increased number of mononuclear cells in the chorioallantoic fluid (CAF) of inoculated but asymptomatic embryos (Pollard and Diserens 1955; Buchner *et al.* 1956). Serum specimens from acute cases of human viral hepatitis appeared to induce a similar response (Pollard and Diserens 1956); however, this has not yet been confirmed (Buchner *et al.* 1956).

Since the virus or viruses of hepatitis in man are not transmissible to experimental animals (MacCallum 1954; Morris, O'Connor and Coburn 1954) since they induce questionable changes in cultures of human tissues, and since they bring about controversial changes in the embryonated egg and indistinct serologic reactions, it was deemed advisable to study the characteristics of a laboratory prototype of hepatitis virus. Some useful information from this might be applicable to the laboratory study of viral hepatitis of man.

Murine viral hepatitis shows some resemblance to the human disease (Friend, Wroblewski and LaDue 1955; DeRitis, Coltorti and Giusti 1956). It shows a narrow species susceptibility. It has not been propagated in fertile eggs (Gledhill, Dick and Niven 1953) and its effect in tissue cultures is uncertain. We are indebted to A. W. Gledhill, National Institute for Medical Research, Mill Hill, London, England, for kindly providing us with a strain of mouse hepatitis virus (MHV) that was free of *Eperythrona coccoides* (Niven, Gledhill and Andrewes 1952). This virus was lethal for newborn mice, usually killing them at 6 to 8 days following intra-abdominal inoculation and causing extensive degenerative lesions in the liver. We are grateful to J. B. Nelson, The Rockefeller Institute for Medical Research, New York, N. Y., for

*The work reported in this paper was supported by funds provided by Contract AF 41 (65)-123 with the School of Aviation Medicine, USAF Randolph Air Force Base, Texas.

- 20 HAGAN W A & D W BRUNER 1957 The Infectious Diseases of Domestic Animals
Comstock Ithaca N Y
- 21 BAKER J A 1942 Cornell Vet 32 202
- 22 JENSEN R L & GRINER T L CHOW & W W BROWN 1955 Proc Livestock San
Assoc 59 189
- 23 MCKERCHER D G J E MOLLTON J W KENDRICK & J SAITO 1955 Proc Live
stock San Assoc 59 151
- 24 CHEEVER F S 1956 Ann N Y Acad Sci 66(1) 196
- 25 PAPPENHEIMER A M & F S CHEEVER 1948 J Exptl Med 88 317
- 26 MOLL T & C A BRANDLY 1954 Vet Med 49 512
- 27 MOLL T & C A BRANDLY 1955 Vet Med 50 101
- 28 BRANDLY C A & A W MCCLURAIN 1956 Ann N Y Acad Sci 66(1) 181
- 29 YORK C J & J A BAKER 1951 J Exptl Med 93 587
- 30 PRITCHARD W R R G CARLSON H E MOSES & D E B TAYLOR 1955 Proc
Livestock San Assoc 59 143
- 31 BAY W W L P DOYLE & I M HUTCHINGS 1951 Am J Vet Research 12 215
- 32 HAELTFEMANN F O & L M HUTCHINGS 1956 Ann N Y Acad Sci 66(1) 186
- 33 RICE J L W H BEST S FRANT & H ABRAMSON 1937 J Am Med Assoc 109
475
- 34 LEE K M 1956 Ann N Y Acad Sci 66(1) 191
- 35 HODES H L 1956 Ann N Y Acad Sci 66(1) 215
- 36 BIDDINGH G J & K DODD 1944 J Pediat 25 105
- 37 GORDON I & I WHITNEY 1956 Ann N Y Acad Sci 66(1) 220
- 38 KOJIMA S H FUJIMU & T ISHIMARU 1953 Japan J Med Sci Biol 6 69
- 39 YAMAMOTO A H M ZENNOJI K YANAGITA & S KATO 1948 Japan Med J
1 349
- 40 STAMP J T 1951 J Comp Pathol Therap 61 215
- 41 McEWEN A D A J LITTLEJOHN & A FOGGIE 1951 Vet Record 63 489
- 42 McEWEN A D & A FOGGIE 1954 Vet Record 68 393
- 43 SCHULTZ G & P D DELAY 1955 J Am Vet Med Assoc 127 224
- 44 YOUNG G A P L KITCHELL A J LUEDAE & J H SAUTTER 1955 J Am Vet
Med Assoc 126 165
- 45 MANNING R 1949 Acta Vet Hung 1 67
- 46 DOLL E R M G RICHARDS & M E WALLACE 1954 Cornell Vet 44 133
- 47 SWAN C 1944 J Pathol Bacteriol 56 289
- 48 BANKOWSKI R A A G PERKINS F F STUART & M KUMMER 1955 Proc Live
stock San Assoc 59 356 Also personal communication
- 49 PARROTT R H 1957 Ann N Y Acad Sci 67(8) 230
- 50 MONTGOMERY J J GEAR M PRINSLOO M KAHN & Z G KIRSCH 1956 S African
Med J 29 608
- 51 KIRICK S & K BENTRSCHKE 1956 New Engl J Med 255 883
- 52 BLANK H & G RAKE 1955 Viral and Rickettsial Diseases of the Skin Eye and
Mucous Membranes of Man 156 Little Brown and Co Boston Ma s
- 53 MELNICK J L H BUNTING W G BANFIELD M J STRAUSS & W H GAYLORD 1952
Ann N Y Acad Sci 54(6) 1214
- 54 SCHULTZ F 1908 Deut med Wochschr 34 423
- 55 OLSON C & R H COOK 1951 Proc Soc Exptl Biol Med 77 281
- 56 KIDD J G & P ROUS 1940 J Exptl Med 71 469
- 57 ADAMS J M 1953 Pediatrics 11 15
- 58 ADAMS J M & D T IMAGAWA 1957 Pediat Clin N Am 193
- 59 CARLSTROM G 1956 Acta Pediat 45 180
- 60 KARZON D T 1955 Pediatrics 16 809
- 61 CARLSTROM C 1954 Lancet 2 344
- 62 FOLDING J B & R M SIMPSON 1957 Vet Record 69 587
- 63 PLUMMER P J G 1946 Can J Comp Med 10 49

TABLE 2
COMPARATIVE TITRATION OF MHV IN CHICK EMBRYOS AND IN NEWBORN MICE

	Virus dilution				
	10^{-1}	10^{-2}	10^{-3}	10^{-4}	10^{-5}
MHV in chick embryos	40	15	4	12	2
MHV in chick embryos	29	7	8		
Above 61 °C /40 minutes	6				
MHV in newborn mice	9/9†			6/7	
Above 61 °C /40 minutes	0/11				

* Histocytes per 100 oil immersion fields. Normal level 0 to 15

† Number of mice infected/number inoculated

sheets of cells developed within 5 days. The nutrient fluid was poured off and 0.1 ml of MHV (Princeton PRI) virus was added to each tube of tissue culture. Ten minutes later 0.9 ml of maintenance solution (calf serum 3 per cent in Eagle's BSS) was added to each tube. The maintenance solution (MS) was changed at 3 or 4-day intervals; however, the cultures were observed microscopically every day. Newborn mice were inoculated with base line culture fluid and with homogenized tissue in the MS at 5-day intervals. Cytopathology was not detected during the 15 days of observation and MHV was not recovered from any specimen except for the base line inoculum (TABLE 3). This experiment was repeated with L-strain cells but cultures were tested at earlier intervals (TABLE 4). MHV was recovered from the culture 3 hours after inoculation and at 9 hours 1 of 6 inoculated mice developed the disease.

TABLE 3
MHV (NELSON PRI) 10^{-5} DILUTION IN L-STRAIN CELL CULTURE

Tissue inoculated	Cytopathology	Mice inoculated
Base line	—	5/5
Five days	0	0/5
Ten days	0	0/5
Fifteen days	0	0/5

Number of mice infected/number of mice inoculated

TABLE 4
MHV (NELSON PRI) 10^{-5} IN L-STRAIN CELL CULTURE

Tissue inoculated	Cytopathology	Mice inoculated
Base line	—	6/6
3 Hours	0	5/5
9 Hours	0	1/6
25 Hours	0	0/4
25 Hours—no tissue	—	0/4
48 Hours	0	0/5

Number of mice showing liver lesions/number of mice inoculated

kindly providing us with a strain of MHV (PRI strain) which was also used in this study. The PRI virus (Nelson, 1952) produces a more fulminant infection in weanling mice in which extensive liver degeneration is manifested.

Three aspects of mouse hepatitis virus are reported on here: (1) the effect of MHV in the embryonated chicken egg, (2) the fate of MHV in cultures of mouse tissues, and (3) the extraction of a complement fixing antigen from MHV infected mouse livers.

(1) Pooled livers from mice acutely infected with MHV (Gledhill strain) were homogenized in a Ten Broeck grinder with Hank's balanced salt solution (BSS) (10 per cent by weight). The emulsion was centrifuged at 2000 rpm for 30 minutes, then at 10 000 rpm for 20 minutes. The supernatant fluid of the latter was then placed in ampoules and stored at dry ice temperature for future use. The MHV, thus prepared, was inoculated into the chorioallantoic fluid (CAF) of 7 day old embryonated eggs. After 4 days of further incubation the inoculated eggs were chill killed by overnight storage at 4°C and the cell content of pooled CAF from these eggs was determined microscopically. The technique has been described elsewhere (Pollard and Diserens, 1955). Control inoculations consisted of MHV heated at 56°C for 30 minutes, 'normal' liver emulsions prepared according to the sequence for MHV above, and CAF from uninoculated embryonated eggs. Each egg inoculated with MHV received approximately 2000 newborn mouse LD₅₀. MHV was titrated simultaneously for infectivity in newborn mice and for cytological response in the CAF of embryonated eggs.

Infective inocula of MHV were associated with increased cells in CAF of inoculated embryos (TABLE 1). Cell counts following inoculations with heat inactivated virus and 'normal' liver extracts were within normal limits, as were the counts from the uninoculated eggs. When the MHV inoculum was diluted beyond 10⁻¹ there was no significant cellular response in the CAF; however, virus effect was still evident in mice at 10⁻⁴ dilution (TABLE 2). Heat inactivated MHV was again noninfectious for newborn mice and negative in chick embryos. Passage of egg material from the 10⁻¹ dilution inoculum into additional eggs was negative.

(2) The L strain of mouse fibroblast (Earle *et al.*, 1943) was propagated in stationary tubes with 20 per cent horse serum in Hank's BSS. Continuous

TABLE 1
COMPARATIVE EFFECT OF MOUSE HEPATITIS VIRUS (MHV) BY HISTIOCYTE RESPONSE IN EGGS AND BY INOCULATION INTO NEWBORN MICE

Inoculum	Histocyte per 100 field	Mice cul t
MHV—liver emulsion	85	25/25
MHV 56°C/30 minutes	22	0/8
MHV 12 liver emulsion	120	5/
MHV 56°C/30 minutes	12	0/2
'Normal' liver emulsion	15	0
Uninoculated control	15	0

Mice infected/mice inoculated

TABLE 2

COMPARATIVE TITRATION OF MHV IN CHICK EMBRYOS AND IN NEWBORN MICE

	Virus dilution				
	10	10 ⁻²	10 ⁻³	10 ⁻⁴	10 ⁻⁵
MHV in chick embryos	40	15	4	12	2
MHV in chick embryos	29	7	8		
Above 61 °C/40 minutes	6				
MHV in newborn mice	9/9†			6/7	
Above 61 °C/40 minutes	0/11				

Histiocytes per 100 oil immersion fields Normal level 0 to 15

† Number of mice infected/number inoculated

sheets of cells developed within 5 days. The nutrient fluid was poured off and 0.1 ml of MHV (1 Princeton IRI) virus was added to each tube of tissue culture. Ten minutes later 0.9 ml of maintenance solution (calf serum 3 per cent in Eagle's BSS) was added to each tube. The maintenance solution (MS) was changed at 3 or 4-day intervals; however the cultures were observed microscopically every day. Newborn mice were inoculated with base line culture fluid and with homogenized tissue in the MS at 5-day intervals. Cytopathology was not detected during the 15 days of observation and MHV was not recovered from any specimen except for the base line inoculum (TABLE 3). This experiment was repeated with L-strain cells but cultures were tested at earlier intervals (TABLE 4). MHV was recovered from the culture 3 hours after inoculation and at 9 hours 1 of 6 inoculated mice developed the disease.

TABLE 3

MHV (NELSON PRI) 1:5 DILUTION IN L-STRAIN CELL CULTURE

Time after inoculation	Cytopathology	Mice inoculated
Base line	—	5/5
Five days	0	0/5
Ten days	0	0/5
Fifteen days	0	0/5

Number of mice infected/number of mice inoculated

TABLE 4

MHV (NELSON PRI) 10⁻⁴ IN L-STRAIN CELL CULTURE

Time after inoculation	Cytopathology	Mice inoculated
Base line	—	6/6
3 Hours	0	5/5
9 Hours	0	1/6
25 Hours	0	0/4
25 Hours—no tissue	—	0/4
48 Hours	0	0/5

Number of mice showing liver lesions/number of mice inoculated

kindly providing us with a strain of MHV (PRI strain) which was also used in this study. The PRI virus (Nelson 1952) produces a more fulminant infection in weanling mice in which extensive liver degeneration is manifested.

Three aspects of mouse hepatitis virus are reported on here: (1) the effect of MHV in the embryonated chicken egg, (2) the fate of MHV in cultures of mouse tissues, and (3) the extraction of a complement fixing antigen from MHV infected mouse livers.

(1) Pooled livers from mice acutely infected with MHV (Gledhill strain) were homogenized in a Ten Broeck grinder with Hank's balanced salt solution (BSS) (10 per cent by weight). The emulsion was centrifuged at 2000 rpm for 30 minutes then at 10 000 rpm for 20 minutes. The supernatant fluid of the latter was then placed in ampoules and stored at dry ice temperature for future use. The MHV thus prepared, was inoculated into the chorioallantoic fluid (CAF) of 7 day old embryonated eggs. After 4 days of further incubation the inoculated eggs were chill killed by overnight storage at 4° C, and the cell content of pooled CAF from these eggs was determined microscopically. The technique has been described elsewhere (Pollard and Diserens 1955). Control inoculations consisted of MHV heated at 56° C for 30 minutes, 'normal' liver emulsions prepared according to the sequence for MHV above and CAF from uninoculated embryonated eggs. Each egg inoculated with MHV received approximately 2000 newborn mouse LD₅₀. MHV was titrated simultaneously for infectivity in newborn mice and for cytological response in the CAF of embryonated eggs.

Infective inocula of MHV were associated with increased cells in CAF of inoculated embryos (TABLE 1). Cell counts following inoculations with heat inactivated virus and 'normal' liver extracts were within normal limits as were the counts from the uninoculated eggs. When the MHV inoculum was diluted beyond 10⁻¹ there was no significant cellular response in the CAF; however, virus effect was still evident in mice at 10⁻¹ dilution (TABLE 2). Heat inactivated MHV was again noninfectious for newborn mice and negative in chick embryos. Passage of egg material from the 10⁻¹ dilution inoculum into additional eggs was negative.

(2) The L strain of mouse fibroblast (Earle *et al.* 1943) was propagated in stationary tubes with 20 per cent horse serum in Hank's BSS. Continuous

TABLE 1
COMPARATIVE EFFECT OF MOUSE HEPATITIS VIRUS (MHV) BY HISTIOCYTE RESPONSE IN EGGS AND BY INOCULATION INTO NEWBORN MICE

Inoculum	Histocyte per 100 field	Mice
MHV—liver emulsion	85	25/25
MHV 56° C/30 minutes	22	0/8
MHV 12 liver emulsion	120	5/7
MHV 56° C/30 minutes	12	0/2
Normal liver emulsion	15	0
Uninoculated controls	15	0

* Mice infected/mice inoculated

TABLE 2

COMPARATIVE TITRATION OF MHV IN CHICK EMBRYOS AND IN NEWBORN MICE

	Virus dilution				
	10^{-1}	10^{-2}	10^{-3}	10^{-4}	10^{-5}
MHV in chick embryos	40	15	4	12	2
MHV in chick embryos	29	7	8		
Above 61 C/40 minutes	6				
MHV in new born mice	9/9†			6/7	
Above 61 C/40 minutes	0/11				

Histocytes per 100 oil immersion fields Normal level 0 to 15

† Number of mice infected/number inoculated

sheets of cells developed within 5 days. The nutrient fluid was poured off and 0.1 ml of MHV (Princeton PRI) virus was added to each tube of tissue culture. Ten minutes later 0.9 ml of maintenance solution (calf serum 3 per cent in Eagle's BSS) was added to each tube. The maintenance solution (MS) was changed at 3- or 4-day intervals; however the cultures were observed microscopically every day. Newborn mice were inoculated with base line culture fluid and with homogenized tissue in the MS at 5-day intervals. Cytopathology was not detected during the 15 days of observation and MHV was not recovered from any specimen except for the base line inoculum (TABLE 3). This experiment was repeated with L-strain cells but cultures were tested at earlier intervals (TABLE 4). MHV was recovered from the culture 3 hours after inoculation and at 9 hours 1 of 6 inoculated mice developed the disease.

TABLE 3

MHV (NELSON PRI) $1:5$ DILUTION IN L-STRAIN CELL CULTURE

Time of inoculation	Cytopathology	Mouse inoculations
Base line	—	5/5
Five days	0	0/5
Ten days	0	0/5
Fifteen days	0	0/5

Number of mice infected/number of mice inoculated

TABLE 4

MHV (NELSON PRI) 10^{-4} IN L-STRAIN CELL CULTURE

Time of inoculation	Cytopathology	Mouse inoculation
Base line	—	6/6
3 Hours	0	5/5
9 Hours	0	1/6
25 Hours	0	0/4
25 Hours—no tissue	—	0/4
48 Hours	0	0/5

Number of mice showing lesions/number of mice inoculated

TABLE 5
MHV (GLEDHILL) 10^{-4} IN L STRAIN CELL CULTURE

Time after inoculation	Cytopathology	Mouse infection
Base line 25 Hours	— 0	10/10 0/5

Number of mice showing liver lesions/number of mice inoculated

TABLE 6
MHV (NELSON PRI) IN MINCED TISSUES FROM NEWBORN MICE
(Virus titration in base line $10^{-4.5}$)

Time after inoculation	Mouse inoculation	
	Liver	Kidney
9 Hours	0/6	3/3
25 Hours	0/3	3/3
72 Hours	0/3	1/3
144 Hours	—	0/2
25 Hours—no cells	1/10	

Number of mice showing liver lesions/number of mice inoculated

Tissue free virus cultures were negative when tested after 25 hours at 37°C . A repeated experiment with MHV (Gledhill strain) was negative at 25 hours and cytopathology was not observed (TABLE 5).

MHV (Princeton PRI) was inoculated into flask cultures of minced liver and minced kidney tissue in MS. The initial virus content of the MS showed mouse infectivity out to $\text{ID}_{50}-10^{-4.5}$. MHV was not recovered from the liver tissue cultures when tested at 9, 25, 72 and 144 hours. MHV was recovered from the kidney cultures at 9 and at 25 hours after inoculation and 1 of 3 mice developed hemorrhagic liver lesions from a 72 hour culture inoculum. The tissue free virus control killed 1 of 10 mice inoculated after 25 hours at 37°C (TABLE 6). It might be argued here that the liver tissue may not have survived for 9 hours thus affording little medium for virus survival; however liver tissue imprints at this time showed histologically intact cells. If MHV entered the mouse cells and manifested an eclipse phase in virus reproduction then either this phase is longer than 15 days or it was extinguished.

During the past year 24 acute serum specimens from viral hepatitis patients have been studied in monolayer cultures of HeLa cell and in accord with the findings of others (Bang and Warwick 1954; Henle 1954; Syverton 1954) cytopathology was not detected. The tissues were homogenized in the MS and inoculated into the CAI of embryonated eggs. All such inoculations were negative. The Detroit 6 strain of cell (Stulberg, Berman and Rudille 1955) was kindly transferred to our laboratory by I. W. McLean of Parke Davis & Company, Detroit, Mich. It was propagated carefully according to the details stressed by McLean and his colleagues (personal communication). R. H. Russell of our laboratory worked with this cell strain for over nine months.

While it is difficult to propagate this strain was adapted satisfactorily to propagation in horse serum. Serum specimens from six selected acute cases of human viral hepatitis have thus far failed to induce cytopathology in the cultures. For obvious reasons we were unable to test the inoculated cultures for virus content nor were the inocula tested for virus content. The inocula did induce significant cytological changes in the C.V. of inoculated eggs.

(3) Pools of liver tissue from newborn Swiss mice acutely ill with MHV (Gledhill) infection were homogenized 10 per cent by weight in Hank's BSS and this material was centrifuged at 2000 rpm for 30 minutes then at 10 000 rpm for 20 minutes at 4° C. Aliquots of supernatant fluid from the latter preparation were then centrifuged at 120 000 g* for 1, 2 and 3 hours. The supernatant fluid was then separated and the sediment was resuspended to original volume in Hank's BSS. Portions of the ultracentrifuged preparations were subjected to (1) testing for infectivity by inoculation into newborn mice (0.05 ml intraperitoneally) (2) desiccation and reconstitution in distilled water (3) desiccation and extraction with chemically pure benzene 3 times at 30 minute intervals and then to reconstitution in distilled water to its original volume (4) mixture with one third volume of diethyl ether at 4° C overnight and rendered ether free by centrifugation and vacuum and (5) heating at 61° C for 30 minutes. Control antigens were processed from liver pools from uninoculated mice.

Antiserum was prepared by inoculating clarified MHV (Gledhill) into adult Swiss mice (0.1 ml intraperitoneally) twice at a 2 week interval and the mice were exsanguinated 2 weeks following the last inoculation. When diluted to 10⁻² the immunizing inocula killed newborn mice.

Control sera were processed from mice of the same colony just prior to the study of MHV infection and from mice inoculated with liver extracts processed from uninoculated mice.

The complement fixation (CF) technique employed was essentially that described by Bengtson (1944) in which 0.2 ml quantities each of antigen and serum and 2 units of complement were used. The hemolytic system was added after incubation overnight at 4° C. Complement fixation was determined by visual inspection. Mouse serum was diluted initially 1/10 in buffered physiological saline and heated at 61° C for 20 minutes.

A box titration of MHV (Gledhill) with serum from immunized and from normal mice is outlined in TABLE 1. Extracts from normal mouse liver did not fix complement in the lowest dilutions of serum and antigen. Further tests with antigens and sera were made with 16 units of antigen and with 8 units of antiserum. Complement fixation tests made with the antigen preparations described above were positive with the supernatant fluid of the preparations centrifuged at 10 000 rpm and at 40 000 rpm (120 000 g) at 1, 2 and 3 hours (TABLE 8). A quantitative decline in CF antigen in the supernatant fluid of the latter procedure was detected following heating, ether extraction, desiccation and desiccation with ether extraction; however these procedures did not cause complete extinction of antigenicity. The sediment

* Model L preparative Spinco centrifuge was used.

- EATON M D W D MURPHY & V L HANFORD 1944 Heterogenetic antibodies in acute hepatitis *J Exptl Med* **79** 537-539
- FRIEND C F WROBLEWSKI J S LADUE 1955 Glutamic oxaloacetic transaminase activity of serum in mice with viral hepatitis *J Exptl Med* **102** 699-704
- GLEDHILL A W S W A DICK & J S F NIVEN 1953 Studies on mouse hepatitis virus (MHV) *Atti 6th Congr intern microbiol Roma* **3** 219-221
- HAVENS W P H LLOYD J L MELNICK & J W COLBERT JR 1949 Complement fixation with fecal antigens in viral hepatitis *J Infectious Diseases* **85** 138-145
- HENLE W 1954 Review of attempts at adaptation of hepatitis virus to the chick embryo *Natl Acad Sci Natl Research Council Publ* **322** 29-34
- LARIN N M 1951 Studies on the agent of canine virus hepatitis (Rubarth's disease) I Complement fixation and precipitin tests *J Hyg* **49** 410-426
- MACCALLUM F O 1954 Review of attempts to transmit hepatitis to other animal hosts and review of recent European efforts at propagation of the virus *Natl Acad Sci Natl Research Council Publ* **322** 68-75
- MANI W 1955 The value of the complement fixation test in the study of canine distemper complex and Rubarth's disease *J Comp Pathol Therap* **65** 291-308
- MORRIS J A J R O CONNER & D R COBURN 1954 Report on a series of attempts to adapt the viruses of human hepatitis to a variety of experimental animals *Natl Acad Sci Natl Research Council Publ* **322** 76
- MILES J A R 1946 A serological investigation in hepatitis using the complement fixation reaction *Brit J Exptl Pathol* **27** 25-33
- NELSON J B 1952 Acute hepatitis associated with mouse leukemia II Etiology and host range of the causal agent in mice *J Exptl Med* **96** 303-312
- NIVEN J S A W GLEDHILL S W A DICK & C H ANDREWES 1952 Further light on mouse hepatitis *Lancet* **2** 1061
- OLITZKI L & H BERNKOPF 1945 Precipitation test in infective hepatitis *J Infectious Diseases* **77** 60-67
- POLLARD M & R H BUSSELL 1957 Complement fixation with mouse hepatitis virus *Science* In press
- POLLARD M & L T DISERENS 1955 The detection of asymptomatic viral infections in the embryonated egg I Studies with influenza mumps and viral hepatitis *USAF School Aviation Med Proj Rept* 55-57
- POLLARD M & L T DISERENS 1956 Immunological studies with viral hepatitis in embryonated eggs *Am J Hyg* **63** 81-94
- RIGHTSEL W A R A KELTSCHE F M TERUSHAN & I W McLEAN JR 1956 Tissue culture cultivation of cytopathogenic agents from patients with clinical hepatitis *Science* **124** 226-278
- STULBERG C S L BERMAN & F H RUDDLE 1955 Detroit-6 strain of human epithelial like cells virus susceptibilities *Proc Soc Exptl Biol Med* **89** 438-441
- SYVERTON J T 1954 Serum hepatitis and infectious hepatitis the use of human cell cultures HeLa strain *Natl Acad Sci Natl Research Council Publ* **322** 17-21

Part III Leptospirosis A Contemporary Problem

INTRODUCTION

By Donald S. Martin

Communicable Disease Center Public Health Service Atlanta Ga

It was just fifty years ago that Stimson¹ described the finding of organisms now accepted as leptospirae in the kidneys of a patient who died supposedly of yellow fever in New Orleans La. A few years later the organism was cultivated successfully from cases of Weil's disease by investigators working independently in Japan² and in Germany.³

Then as in many diseases a period of discovery was followed by a lull. *Leptospira icterohaemorrhagiae* was recognized as the etiological agent of Weil's disease the epidemiology was worked out and the textbooks confidently printed the facts that the organism was carried by rats and that man acquired his infection when he became contaminated by rat urine.

During the past fifteen to twenty years the entire picture has changed. Leptospirosis is now recognized as a world wide public health problem of increasing importance. We now know that the term leptospirosis does not connote a single disease entity and that the clinical pictures may be quite different depending upon the particular serotype causing the infection. We know also that leptospirosis of domestic animals is of great concern to farmers milk producers and breeders of cattle and swine.

The various clinical and diagnostic problems of the leptospiral infections were discussed at length in a symposium held five years ago at Walter Reed Medical Center Washington D. C. The information has been accumulating so rapidly that it is considered most appropriate that we be brought up to date once more.

I take this opportunity to compare the development of our knowledge of leptospirosis with that of the systemic mycoses. Until the last few decades coccidioidomycosis and histoplasmosis were considered as rare but almost always fatal infections with quite characteristic clinical features. Today it is known that infections from these fungi are extremely common in certain areas of the country and that the development of a full blown fatal form of either disease is the exception. Infections from these fungi result in such a variety of clinical manifestations that the true epidemiology of the disease became apparent only when diagnostic skin and serologic tests were developed and applied in large scale studies. It was necessary to impress bacteriologists with the fact that they should not throw out the cultures after the first two or three days—it was necessary to alert physicians to suspect a fungus infection in a patient who did not respond to antibiotic therapy.

I feel that the same principles are applicable today in leptospirosis. Both bacteriologists and physicians must be alerted always to consider the possibility of leptospirosis in any patient with an obscure febrile illness. Blood cultures must be taken early in special media and must be held for weeks

Leptospirosis cannot be excluded because the patient shows no jaundice or denies falling in water contaminated by rat urine. New leptospiral serotypes pathogenic to man are being found in other parts of the world and the clinical picture of infections caused by these serotypes may be quite different from that of classic Weil's disease, canicola fever, pomona meningitis, or pretibial fever. We now are learning also that leptospire of various serotypes can be cultivated from an ever widening range of animal species.

The possibilities are just beginning to unfold and the reader will find in the present section of this monograph the latest information on leptospirosis which is becoming of increasing significance to those who are concerned with infectious diseases of both man and animals.

References

- 1 STIMSON, A. M. 1907. A note on an organism found in yellow fever tissue. *Public Health Repts. U. S.* 22: 541.
- 2 INADA, R., Y. IDO, R. HOSI, R. KANEKO & H. ITO. 1916. The etiology, mode of infection and specific therapy of Weil's disease (*Spirochaetosis icterohaemorrhagica*). *J. Exptl. Med.* 23: 377.
- 3 UHLENHUTH, P. & W. FROMME. 1930. Weilsche Krankheit. *In Handbuch der Pathol. Mikroorg.* Von Kolle, Kraus & Uhlenhuth, Eds. 7: 487. Cited by P. H. Van Thiel. 1948. *The Leptospiroses*. Universitaire Pers. Leiden, Netherlands.

ANIMAL RESERVOIRS OF LEPTOSPIRES

By B. Babudieri

Istituto Superiore di Sanità Rome Italy

Man forms a purely incidental and nonessential link in the chain of the epidemiology of leptospirosis. As a rule the human infection is exhausted within itself and it is only in extremely rare instances that it is the cause of fresh cases of the illness.

Furthermore the pathogenic *Leptospira* are not able to adapt themselves to a saprophytic life and are unable to reproduce in the outside world in which their survival is somewhat limited in time.

Consequently the exclusive sources of the infection direct or indirect are the animal carriers and shedders of the *Leptospira*. A knowledge of these is essential for the determination of the epidemiological characteristics of the infection and in order to be able to provide in advance really efficient prophylactic measures.

It is necessary to define clearly what is meant by the term carrier. In the first place one must distinguish between the condition of the occasional and temporary shedder of leptospires and that of the true carrier. The first is a normal condition for every individual who suffers or has recently recovered from an infection by leptospires. During the illness there is constant leptospiruria which continues long after the recovery but invariably, only for a limited period generally for several weeks.

The temporary shedder of leptospires may constitute a source of natural infection but his epidemiological importance is relatively limited not only because the leptospiruria ceases after a short time but also because in handling the sick or convalescent patient it is easy to carry out the appropriate rules of prophylaxis.

The condition of the permanent carriers is quite different. Here one is dealing with animals that have suffered infection by leptospires (an infection that may have shown morbid manifestations or that may have remained symptomatologically blank) and have then remained shedders of leptospires via the urine for a long period or even for the rest of their lives. It is these carriers that permit the survival of individual serotypes of *Leptospira* and that constitute the most dangerous source of the infection.

Our knowledge of the leptospire carriers is not yet complete. The carriers among the more common animals which more easily come into contact with man are sufficiently well known. Knowledge is much less plentiful however with regard to the rarer animals that are more difficult to study experimentally and are less close to man. In fact at some times when one finds leptospires in the urine or kidney of such animals one does not know whether one is dealing with temporary shedders or with true carriers.

From a practical point of view the importance of the animal carriers depends on different factors. One of these is the reaction of the urine. It is well known that the pathogenic leptospires are very sensitive to variations of pH

Research that I have carried out on some serotypes of these has shown that the extreme limits of pH that may permit the survival of leptospire for a period of at least 6 days in an environment otherwise favorable to their existence are between 6.24 and 8.23. In order that the survival be prolonged to one month these limits must be reduced to 6.35 and 7.96. Some animals including man have a distinctly acid urine and in their case the leptospire excreted with it are either dead or at least unable to reproduce the illness. In such cases and for this reason the animal in spite of being a carrier, does not constitute a danger from contagion. The reaction of the urine may vary, however even in the same animal, according to its diet.

This is the case in the dog which if given a diet of meat has an acid urine fatal for the leptospire when however the dog has a predominantly vegetable diet it has a urine of neutral or even slightly alkaline reaction much more favorable to the survival of the leptospire. According to Zuelzer¹ the rat also generally has a urine with a pH of 5.4 to 5.8 but if it subsists primarily on vegetables the pH rises to a level of 7 to 8.

Generally the herbivorous animals have a neutral or slightly alkaline urine while that of the carnivores is usually acid. It is also necessary to remember that an acid diet at least in dogs does not succeed in disinfecting the animal. The leptospire that colonize the small renal canals survive and reappear in virulent form in the urine as soon as its reaction has again become favorable.

It must also be remembered that leptospire although very sensitive to an acid environment are nevertheless able to survive in it at least within certain limits for a very short period. Thus an animal that excretes a moderately acid urine may be dangerous if the urine is immediately mixed with water or mud of favorable reaction. In this case the leptospire surviving the brief transit in the acid urine find themselves in environmental conditions favorable to their existence.

Another factor of great importance is that of the environment in which the animal carrier lives. Rarely does the infection occur directly through contact with the infected urine. Man nearly always infects himself indirectly by way of water, mud or foods contaminated by the urine. Consequently animal carriers that have the capacity for contaminating their surroundings are particularly important. Those that eliminate leptospire but disperse them in an unfavorable environment where they rapidly die out are of little consequence. For example the importance of small rodents that live in the flooded rice fields is very different from that of the jackal which in Israel lives in arid or semi-desert regions.

Another example is given by the conditions which in the plains of the large rivers of central and eastern Europe permit in some years the outbreak of epidemics of the so-called *Schlammfieber* (mud fever) caused by *L. grippotyphosa*. The usual carrier of this leptospire is the field mouse (*Microtus arvalis*) that lives in holes scattered throughout the fields.

If the season is dry the illness caused by *L. grippotyphosa* is slight because the leptospire eliminated with the urine have little chance of surviving in such surroundings. When however the season is rainy and especially when after great river floods the plains remain muddy and dotted with puddles for

a long time the earth becomes saturated so to speak with leptospires. This condition gives rise to the epidemics of *Schlammfieber* which sometimes affect untold thousands of persons who work in such fields.

Finally a third factor of great practical importance is the possibility of contact direct or indirect between the animal carriers and man. It is easy to comprehend how wild animals that live far from human habitations and cultivated land have much less epidemiological importance if they are carriers than those that live in homes in stables or in the cultivated fields and especially those that experience frequent manual attention.

These three factors variously combined among themselves condition the practical importance of the different animal carriers of leptospires.

THE RELATION BETWEEN LEPTOSPIRES AND THE ORGANISM OF THE ANIMAL CARRIER

What is the mechanism through which the quality of the carriers is established? It is not yet completely known and it still presents difficulties similar to those encountered when the same problems are examined in the field of the bacterial carriers.

In the first place it is worth remarking that at least as far as is known at present the condition of the carrier is not established unless the animal has suffered an infection by leptospires. This infection may give evident clinical manifestations or may pursue its course completely undetected. In both cases however during the course of the infection a transitory state of leptospiremia exists. In this stage it is possible to find leptospires in all organs including the kidney. Leptospires are found in the interstitial tissue in this organ as well as in the vessels. However it is only in relatively small numbers and often simultaneously with more or less extensive lesions of nephritic nature that they penetrate into the renal canals and pass into the urine. When on the other hand an immunized state is established in the patient which happens after seven to ten days the leptospires disappear from the blood and the interstitial tissue and begin to accumulate in the secondary convoluted tubules. These masses true colonies of leptospires adhere to the epithelial cells and sometimes are even included in them without exerting any evident pathogenic action on them with the exception in some cases of a mechanical action that produces a flattening of the epithelia themselves.² The masses are sometimes very large being constituted of an inextricable tangle of spirochetes. It does not seem however that they ever succeed in completely obstructing the tubules in which they colonize a fine canal in the center always persists through which the urine may pass.

Numerous leptospires detach themselves from such masses and are then transported by the urine to the exterior. It may be said that at this stage the leptospires are transformed from internal parasites of the organism into external parasites and that in a sense their life approaches that of the saprophytes. At this point the leptospires are released from the action for them harmful of the defense mechanism of the host. The antibodies circulating in the blood and in the tissues do not succeed in reaching them and are not even able to exert an antigenic action on the host. In fact in animal carriers

Research that I have carried out on some serotypes of these has shown that the extreme limits of pH that may permit the survival of leptospirae for a period of at least 6 days in an environment otherwise favorable to their existence are between 6.24 and 8.23. In order that the survival be prolonged to one month these limits must be reduced to 6.35 and 7.96. Some animals including man have a distinctly acid urine and in their case the leptospirae excreted with it are either dead or at least unable to reproduce the illness. In such cases and for this reason the animal in spite of being a carrier does not constitute a danger from contagion. The reaction of the urine may vary, however even in the same animal according to its diet.

This is the case in the dog which if given a diet of meat has an acid urine fatal for the leptospirae, when however the dog has a predominantly vegetable diet it has a urine of neutral or even slightly alkaline reaction much more favorable to the survival of the leptospirae. According to Zuelzer¹ the rat also generally has a urine with a pH of 5.4 to 5.8 but if it subsists primarily on vegetables the pH rises to a level of 7 to 8.

Generally the herbivorous animals have a neutral or slightly alkaline urine while that of the carnivores is usually acid. It is also necessary to remember that an acid diet at least in dogs does not succeed in disinfecting the animal. The leptospirae that colonize the small renal canals survive and reappear in virulent form in the urine as soon as its reaction has again become favorable.

It must also be remembered that leptospirae although very sensitive to an acid environment are nevertheless able to survive in it, at least within certain limits for a very short period. Thus an animal that excretes a moderately acid urine may be dangerous if the urine is immediately mixed with water or mud of favorable reaction. In this case the leptospirae surviving the brief transit in the acid urine find themselves in environmental conditions favorable to their existence.

Another factor of great importance is that of the environment in which the animal carrier lives. Rarely does the infection occur directly through contact with the infected urine. Man nearly always infects himself indirectly by way of water, mud or foods contaminated by the urine. Consequently animal carriers that have the capacity for contaminating their surroundings are particularly important: those that eliminate leptospirae but disperse them in an unfavorable environment where they rapidly die out are of little consequence. For example the importance of small rodents that live in the flooded rice fields is very different from that of the jackal which in Israel lives in arid or semi-desert regions.

Another example is given by the conditions which in the plains of the large rivers of central and eastern Europe permit in some years the outbreak of epidemics of the so-called *Schlammfieber* (mud fever) caused by *I. griffotyphosa*. The usual carrier of this leptospire is the field mouse (*Microtus arvalis*) that lives in holes scattered throughout the fields.

If the season is dry the illness caused by *I. griffotyphosa* is slight because the leptospirae eliminated with the urine have little chance of surviving in such surroundings. When however the season is rainy and especially when after great river floods the plains remain muddy and dotted with puddles for

a long time the earth becomes saturated so to speak with leptospires. This condition gives rise to the epidemics of *Schlammfieber* which sometimes affect untold thousands of persons who work in such fields.

Finally, a third factor of great practical importance is the possibility of contact direct or indirect between the animal carriers and man. It is easy to comprehend how wild animals that live far from human habitations and cultivated land have much less epidemiological importance if they are carriers than those that live in homes, in stables, or in the cultivated fields and especially those that experience frequent manual attention.

These three factors variously combined among themselves condition the practical importance of the different animal carriers of leptospires.

THE RELATION BETWEEN LEPTOSPIRES AND THE ORGANISM OF THE ANIMAL CARRIER

What is the mechanism through which the quality of the carriers is established? It is not yet completely known and it still presents difficulties similar to those encountered when the same problems are examined in the field of the bacterial carriers.

In the first place it is worth remarking that at least as far as is known at present the condition of the carrier is not established unless the animal has suffered an infection by leptospires. This infection may give evident clinical manifestations or may pursue its course completely undetected. In both cases however during the course of the infection a transitory state of leptospiremia exists. In this stage it is possible to find leptospires in all organs including the kidney. Leptospires are found in the interstitial tissue in this organ as well as in the vessels. However it is only in relatively small numbers and often simultaneously with more or less extensive lesions of nephritic nature that they penetrate into the renal canals and pass into the urine. When on the other hand an immunized state is established in the patient which happens after seven to ten days the leptospires disappear from the blood and the interstitial tissue and begin to accumulate in the secondary convoluted tubules. These masses true colonies of leptospires adhere to the epithelial cells and sometimes are even included in them without exerting any evident pathogenic action on them with the exception in some cases of a mechanical action that produces a flattening of the epithelia themselves. The masses are sometimes very large being constituted of an inextricable tangle of spirochetes. It does not seem however that they ever succeed in completely obstructing the tubules in which they colonize a fine canal in the center always persists through which the urine may pass.

Numerous leptospires detach themselves from such masses and are then transported by the urine to the exterior. It may be said that at this stage the leptospires are transformed from internal parasites of the organism into external parasites and that in a sense their life approaches that of the saprophytes. At this point the leptospires are released from the action for them harmful of the defense mechanism of the host. The antibodies circulating in the blood and in the tissues do not succeed in reaching them and are not even able to exert an antigenic action on the host. In fact in animal carriers

the antibody level gradually diminishes at a rate equal to that in noncarriers until it reaches a level no longer detectable by our methods of investigation. During a search for *L. ballum* in numerous specimens of *Ictodermus sylvaticus* that I carried out in Spain four strains of these leptospire were isolated all from mice that did not show any trace of antibodies in the serum. On the other hand antibodies were found to be present in other mice that however were not carriers. The same phenomenon has been observed in cattle by Gillespie *et al*³ and by Kmety *et al*⁶ in dogs by Alexander *et al*⁴ in pigs by Burnstein and Baker⁵ and in rodents by numerous other authors.⁷⁻¹⁰

However as already mentioned some points are still obscure. One of these arises from the fact that the leptospire are able to survive in the renal tubules even though antibodies against them are present in the urine.^{11,12} However it is true that such antibodies have a low titer and never reach the high level that may be found in the serum.

On the other hand it is known that if a specific serum of not very high titer is introduced into a culture of leptospire one nearly always observes a certain number that are able to survive and multiply giving place by the phenomenon of agglutination to masses that in a sense may even call to mind those observed in the renal tubules.

Another obscure point is the reason for the establishment of the state of the permanent carrier in certain cases while in others the urinary elimination of the leptospira is only a transitory event. Some obscure phenomena of biological affinity certainly exist here by these a state of biological equilibrium is easily established between some serotypes of leptospire and some animal species. This equilibrium is either lacking or much more difficult to establish in other cases. Both components of this symbiotic association play a part in this phenomenon. In fact it is observed that while certain animals easily become carriers of some serotypes of *Leptospira*, they do not become so for others. This confirms anew the biological validity discussed by some investigators of the subdivision of *Leptospira* into serotypes, at the same time it demonstrates the fact that the distinction made initially between temporary shedders and persistent carriers although unconventional is one that rests on a solid if poorly explicable biological basis. Any animal susceptible to infection by leptospire may become a temporary shedder but only those animal species that present a particular condition of biological sympathy for a determined serotype of *Leptospira* can become carriers. For example the rat which is a habitual carrier of *L. icterohaemorrhagiae* is unable to become one for the serologically related *L. canicola*.¹⁴

It should be pointed out that even within the ambit of the same species not all the animals that have suffered infection necessarily become carriers this occurs only in some. Why this occurs it is difficult to explain and it is also difficult to understand why some animals remain carriers all their lives while others after a period of months or even years become autodisinfected. However these phenomena are not only peculiar to the leptospire carriers but are common to all the other categories of carriers of pathogenic agents.

Furthermore even the different strains of *Leptospira* belonging to the same serotype behave differently in the same host. Van der Hoeden¹⁵ has pointed

out that the white mouse may become a carrier of *I. grippolyphasa* on condition that it is inoculated with a sufficient quantity of this agent. The animal may then eliminate the leptospores with the urine either continuously or intermittently according to the strain of the agent.

Animals that are carriers of leptospores do not seem to suffer any damage from their condition¹⁸⁻¹⁹. It is true that partial renal lesions have been described in some carriers (occasionally in cattle and more often in dogs) but it is unlikely that these lesions are to be attributed to the leptospores that the carrier harbors; it is more probable that they are the consequence of the previous acute infection by leptospores. In fact these same lesions may also be found in cattle and in dogs that have recovered from leptospirosis but have not become carriers.

On the other hand one has the impression that the circumstance of being harbored by one particular animal species rather than by another does exert a certain influence on the pathogenic property of some leptospores. In this respect clear experimental results are lacking; nevertheless it is probably true that as noted by some authors the transit of *L. icterohaemorrhagiae* in the rat accentuates its virulence. Thus *I. balantiae* disseminated in Italy by the harvest mouse *Micromys minutus sorcinus* almost invariably produces a light anicteric infection whereas in Indonesia where the principal carrier is the rat *Rattus diardi* it causes a serious illness in which jaundice is very often present.

In the same series of phenomena may be included the fact—one I have repeatedly observed—that when man contracts an infection of *L. pomona* by direct contact with pigs he almost always becomes ill with distinct and prevalent meningitic symptoms whereas these are very often absent in cases of infection due to the same leptospire if it is contracted in the country where it may be acquired from a different source.

Although it is generally accepted that the animal carrier of leptospores harbors these spirochetes in the kidney and eliminates them with the urine the hypothesis has nevertheless been advanced that the leptospores may persist in some other organs for example in the brain or the salivary glands. This latter hypothesis has been advanced because of the repeated observation of infections by a leptospire following a bite from a dog or more frequently from a small rodent¹⁹⁻²¹. This hypothesis is probably untenable. Thus far it has never been demonstrated with certainty that a chronic animal carrier of leptospores may present them in the saliva or the salivary glands²¹. The cases recorded above may be explained either by supposing that the biting animal happens to be in the acute stage of the infection in which case the leptospire may indeed be found in the saliva or by assuming that the wound caused by the bite has been contaminated by infective urine excreted by the animal. This contingency is probable in cases in which the bite has been received from a mouse caught with the hands since it is well known that these small rodents easily excrete urine when frightened.

Another question is that of when and why an animal becomes a carrier or to be more precise contracts the infection.

It is noted from the research of Uhlenhuth and Fromme²² that leptospores

are able to pass through the placenta and to reach the fetus. Sixteen years ago Biocca and I²³ were able to show that sometimes this produced neither abortion nor death of the newborn but simply an illness of short duration in the parent. Nevertheless it has never been demonstrated that an animal has become a carrier through an infection contracted during the fetal stage. On the contrary the observations of all workers occupied with this topic agree that in rats and mice the percentage of carriers is much higher in adult animals than in the young. This indicates that at least in these animals the infection by leptospirae usually occurs neither during the fetal period nor during the period of infancy, when closer contact with the parents possible carriers would suggest that contagion might be easier. It may be that newborn animals acquire from the mother through the placenta or with the milk a temporary passive resistance that protects them for a time. In addition it must not be forgotten that in very young animals leptospirosis takes a more severe course so that many of these victims die before becoming carriers.

In any case all the evidence seems to suggest that the animals more often become infected when they are adults and in the same way in which man may become infected. Furthermore, it is probable that the infection may occur through copulation. It has been experimentally proved in fact that leptospirae may pass through the genital mucosa moreover some cases of human infection by sexual contact have also been described.^{4, 5}

Leptospira in the Arthropoda

The possibility that insects might be carriers of the agent of Weil's disease was considered even before the genus *Leptospira* was discovered and its etiological role recognized. Some epidemiological data made the following hypothesis appear probable. The frequency of the illness in persons living in marshy zones its greater diffusion in the summer and the influence of rain and humidity on the oscillation of the illness are all factors that were assumed to cause the epidemiology of Weil's disease to approach that of malaria. This theory was supported particularly by Hecker and Otto²⁴ and the insects mainly suspected were the mosquito and the horsefly.

Later after the discovery of *Leptospira* this theory was abandoned even though numerous research workers attempted to demonstrate its worth experimentally. These researches have been particularly well summarized and discussed by Uhlenhuth and Fromme² who made a notable contribution to them.

The possible role of *Hematophagus* flies has been studied by Kenter and Ramme² who show that *Haematopota pluvialis* may transmit leptospirae to guinea pigs mechanically. The same possibility has been demonstrated by Uhlenhuth and Kuhn² for *Stomoxys calcitrans*. This horsefly may transmit the illness for a period as long as six days after having bitten a sick animal. In flies however multiplication of the ingested leptospire is not demonstrable and according to workers who have carried out these experiments on guinea pigs the results are not immediately relevant to man nor to the natural epidemiological conditions.

Numerous researches have been carried out on the various species of mos-

quito generally with negative results²⁰⁻²⁶. The only exception is Noguchi²⁷ who maintains that *Ixodes aegypti* may be a carrier and transmitter of leptospirae. According to the unconfirmed observations of Cay and Sellards²⁸ leptospirae are able to survive for at least 22 days in the body of the mosquito.

Attempts to transmit leptospirae by means of lice have likewise been unsuccessful²⁹⁻³¹. Positive results however have been reported by Dietrich³⁰ both with the louse and with a leech and by Blanchard, Lefrou and Laigret³¹ with bedbugs. These arthropods have succeeded in infecting the guinea pig even after a period as long as 38 days subsequent to being fed on a sick human. Bonne³² however has not been able to confirm these last results.

The experiments carried out on fleas and ticks by Huebner and Reiter produced negative results³³. Recently however it has been demonstrated conclusively that ticks may become infected with leptospirae and may not only conserve them for a long time (at least 318 days) in their internal organs³⁴⁻³⁶ but may also emit them with their coxal liquid and pass them on to their descendants³⁴.

Of particular interest are the observations of Burgdorfer³⁴ who not only succeeded in transmitting leptospirosis to guinea pigs by feeding infected examples of *O. turicata* on them but has also found a tick infected in nature by *I. ballum*. These findings were recently confirmed by two Russian authors Krepkogorskaia and Rementsova³⁶ who were able to isolate two strains of *L. grippolyphosa* from specimens of *Dermacentor marginalis* captured in the Kazakh Soviet Socialist Republic a region where leptospirosis is endemic in cattle.

These researches should be worth following up because although it is very improbable that ticks may have importance in the human epidemiology of leptospirosis it is not impossible that they may play a role in the transmission of the infection from animal to animal.

With the exception of these last results it may be said that all these researches were carried out for the most part in the first years that followed the discovery of the *Leptospira* when the epidemiology of the infection was still imperfectly known and that they have not produced any positive results of practical interest. Certainly it must be admitted that in particular experimental conditions some arthropods may become infected with leptospirae may maintain them for a reasonably long time in their bodies and may finally emit them in such condition as to render the infection of a mammal possible. This is however a rare occurrence it can happen only under very particular circumstances or in well-organized experiments but it very improbably occurs in nature.

Vertebrate Carriers of Leptospirae

Very little research has been directed to determine the receptivity of amphibians to leptospirae. Chetti³⁷ and Corrales³⁸ have obtained negative results in the frog and I have searched unsuccessfully for pathogenic leptospirae in some species of amphibian caught in the rice fields of Italy. Uhlenhuth and Fromme³⁷ have found leptospirae in the intestine of the frog but these are probably water leptospirae that are not pathogenic.

Nothing is known of the possible receptivity of fish to infections from leptospires. Altava Barrera and Villalonga⁴⁹ have searched without success for pathogenic leptospires in the organs of minnows (*Gambusia*) living in the infected waters of rice fields in Spain. Dohmen⁵⁰ reports the case of a fisherman who became ill with leptospirosis after being bitten by a pike but probably it was not the fish that infected him directly; the wound caused by the bite formed a point of entry for leptospires present in the water.

As far as I am aware, research on reptiles in this connection is nonexistent.

The possibility that birds may be susceptible to infection with leptospires and thus become carriers has already been examined by Uhlenhuth and Fromme⁵⁷ who have established that chickens and sparrows are resistant to the infection. Martin and Pettit⁵¹, Ghetti⁴⁷ and Oba⁵² have also had negative results with chickens and pigeons, and Oba⁵ has proved that there are no formations of antibodies in the inoculated chicken. Corrales⁴⁸ on the contrary has shown that the serum of the normal chicken exerts a lytic action on leptospires.

In contrast to these results, other more recent workers have succeeded in finding in nature chickens that show a high titer of agglutinins in the serum (Gillespie *et al.*³ for *L. pomona*, Terskich⁵³ for *L. icterohaemoglobinuriae botis* and *L. botis palestiniensis*, Kujumgiev⁵⁴ for *L. andaman*, A. Mochmann⁵⁵ for *L. icterohaemorrhagiae*, *L. grippotyphosa* and *L. canicola*, and Bernkopf and Olitzki⁵⁶ for *L. icterohaemorrhagiae* and for *L. botis*; chickens, ducks, geese, and turkeys). Seropositive geese have been reported by Mochmann⁵⁵ (*L. grippotyphosa*) and by Vissotski and Redkina⁵⁷.

The fact that the chicken embryo may be experimentally infected with leptospires has already been demonstrated by Hallauer and Kuhn⁶³ and it has been shown that the chick may also be infected and may maintain leptospires in the circulation for several days (Hoag, Gochenour and Yager⁵⁹). As a result, chicks are sometimes used for the isolation of leptospires from polluted material.

Gillespie *et al.*³ claim to have shown experimentally that the receptivity of the chicken to infection by leptospires is limited only to the first few days of life. However, Bernkopf⁶⁰ has succeeded in infecting chickens older than six weeks with *L. botis* (*L. grippotyphosa*) producing in them either a morbid state or an obvious tendency toward immunity. The same result has been obtained in chickens and ducks by Kenz and Gillespie⁶¹ with *L. pomonae*. Chalquest⁶ has succeeded in infecting experimentally with *L. pomonae* not only chickens but also ducks, pheasants and partridges. However, he has not succeeded in finding leptospires in their intestines.

It has not been possible thus far to isolate a pathogenic leptospire from a bird infected in nature, with the exception of the recent discovery of Kujumgiev⁵⁴ who claims to have isolated a strain of *L. andaman* from the kidney of a goose. This discovery, however, requires confirmation.

The first reliable discovery of this type was made this summer by my colleague E. Bussinello in the rice fields of the Po valley in Italy. Bussinello (unpublished data) has succeeded in isolating two strains of *L. balantiae* from the intestine and liver of two wading birds frequently found in these rice fields. Other *Leptospira* strains, not yet classified, have been isolated from the kidney

liver and intestine of the following species of aquatic birds *Ardeola minuta* *Ardea cinerea* *Herodias gallicus* *Hydrochelidon nigra* and *Gallinula chloropus*

It is to be noted that *L. bataviae* is the type of leptospire most frequently found in the rice fields of Italy where it is usually transmitted by a small mouse *Micromys minutus sorcinus*. It may be considered that the birds became infected from contaminated water of the rice fields or from devouring mice that were carriers of *Leptospira*. Regarding the discovery of leptospires in the intestine it should be remembered that in birds the ureter flows into the intestine (cloaca).

It is not yet possible to determine whether these birds are true carriers of this leptospire or are only temporary shedders.

If it should be demonstrated that they are carriers the discovery would have very great importance not only because it would modify our knowledge of the epidemiology of leptospirosis but also because it might explain the strange phenomenon that the same serotype of *Leptospira* may be found in widely separated localities in spite of having as habitual carriers animals that live in relatively limited areas. For example *L. bataviae* is found as frequently in Italy as in some Scandinavian countries and in Indonesia. It is also to be noted that the *Grallae* which live in the Italian rice fields are migratory birds that travel very great distances.

Although it is possible that leptospires do infect other animals it is nevertheless beyond doubt that they are essentially parasites of mammals. It is among the mammals in particular the rodents that we find their most important carriers.

Undoubtedly the gray rat *Epimys decumanus* is the most important carrier. Uhlenhuth and Fromme²⁷ have already indicated this rodent as the international carrier of *L. icterohaemorrhagiae*. Wherever researches are carried out it is found with rare exceptions that a notable percentage of rats sometimes 90 per cent in adult animals are carriers and shedders of *L. icterohaemorrhagiae*. When it is considered that the gray rat is extremely common that it is found throughout the world that with ships it can easily migrate from one country to another and that it has close contacts with man and his foodstuffs it is easy to comprehend how *L. icterohaemorrhagiae* is so universally diffused and is of prime importance among the other leptospires both for the frequency and the gravity of the disease it provokes.

The frequency of infection by *L. icterohaemorrhagiae* among sewer cleaners among people bathing in sweet water close to sewer outlets among workers in the rice fields of Spain and in all places where rats abound shows the close relationship that exists between the presence of this animal and human infection.

Also interesting in this respect are the observations made in rice fields in Spain by Altava Barrera and Marin²⁸ who have noted that apart from wide variation in the number of rats that infest rice fields considerable annual variations also exist in the percentage of rats that are leptospire carriers. These percentages may already be determined in the spring before work begins in the rice fields and they give a very reasonable indication of what will be the development of leptospirosis among workers in the following summer. If

the percentage of rats that are carriers is low there will be only sporadic cases of illness, if on the other hand it is high an epidemic is to be expected

Another example of the relation existing between the abundance of rats and the frequency of human infection is given by the epidemic of leptospirosis which, in 1951, hit Kodiak and several other localities in Alaska a territory that until then had been almost free from leptospirosis. The epidemic was studied by W. H. Gaub (unpublished data) and also described in a United States Navy report.⁵

It happened that during the course of World War II the United States armed forces had accumulated in several localities in Alaska very abundant convoys of provisions thus favoring multiplication of rodents. At the end of the war these deposits were demobilized and abandoned. In consequence the rats found there undisturbed an abundant source of food that permitted them to multiply at an extraordinary rate. When the provisions suddenly became exhausted, the rats poured en masse into the small towns of the region and principally through contamination of foodstuffs originated a very considerable number of cases of leptospirosis.

Finally it is worth noting that according to a publication by McKinley,⁶⁴ in the Philippine Republic (in which at least until 1928 human cases of leptospirosis had not been reported) the percentage of rats that are carriers is less than 1 per cent, even in cities such as Manila which have a large exchange of traffic with Japan where the percentage of carrier rats is about 50 per cent and where human cases of the infection are numerous.

The rat represents the classic example of an animal carrier of leptospirosis. It becomes infected at early maturity (the percentage of young rats that are carriers or even simply seropositive is very low) and as has also been shown experimentally it remains a shedder for years practically throughout its life.^{65, 66}

The rat may become a carrier of other pathogenic leptospirae as well as *L. icterohaemorrhagiae* for example *L. pomona*^{67, 68, 69, 71} *L. ballum* and *L. bataviae*.^{7, 73} These are only occasional discoveries however of minor practical interest.

The black rat *Epimys rattus* is likewise a carrier of *L. icterohaemorrhagiae* but less frequently. For example researches carried out in Japan⁷⁴ have shown that almost 40 per cent of the specimens of *Epimys decumanus* are carriers of *L. icterohaemorrhagiae* while scarcely 2 per cent are carriers among *Epimys rattus*. This lesser frequency together with the fact that *Epimys rattus* is notably more rare than *Epimys decumanus* serves to place it in lower order of practical importance among the carriers of *L. icterohaemorrhagiae*.

Other rats having importance as carriers of leptospirae include *Epimys diardi* the principal disseminator of *L. bataviae* in Indonesia *Epimys brevicaudatus*, carrier of *L. autumnalis* and of *L. pyrogenes* and *Epimys culmorum* carrier in Australia of *L. australis*.⁴

Mice are also important carriers of leptospirae. However according to some authors⁷⁵ mice unlike rats are able to eliminate leptospirae for a very long time but normally do not do so all their lives. At a certain period and by a still unknown mechanism they become autosterilized.

Among the mice that are carriers of leptospires the most important is *Microtus arvalis* habitual carrier of *L. grippolyphosa* the leptospire most widely diffused in the great river valleys of Central and Eastern Europe. The number of these mice is subject to notable annual oscillations. When a year of great abundance of mice coincides with abundant rainfall and still more with floods there is nearly always a human epidemic. It is necessary for these two conditions to coincide in order to make possible an epidemic. Thus in 1943 Kathe¹⁴ observed in Silesia that the field mice were very numerous but the season was very dry. Consequently, not only was the epidemic diffusion of the illness in man absent but also the possibility of infection from mouse to mouse was limited. In fact scarcely 1 per cent of the mice were carriers. In the year 1942 on the other hand the season was rainy and although the number of field mice was not excessive it was nevertheless noticed that 18 per cent were carriers and the human illness became an epidemic.

It is known however that Kathe⁴ has his own particular conception of the epidemiology of *Schlammfieber*. Kathe is a supporter of the *biodynamie* according to which pathogenic leptospires should be capable not only of surviving in the external surroundings but also when these present favorable conditions of multiplying in them. According to this theory the high percentage of carrier mice would not be the origin but the consequence of a wide diffusion of pathogenic leptospires in the soil.

The observations of Popp¹⁷ made during a small epidemic caused by *L. grippolyphosa* among pea pickers in Germany are also interesting. In the locality where the epidemic developed Popp found that from 47 to 83 per cent of the adult field mice were carriers of leptospires. In the autumn after the epidemic had subsided he no longer found carrier mice in the area and even the percentage of seropositive mice was rather low (20 per cent). Popp holds that this may be explained by presuming that a good number of the carrier mice died from leptospirosis. This contrasts however with what is known of the relations that usually exist between leptospires and the animal carrier.

It must also be remembered that Van der Hoeden¹⁵ has demonstrated that the white mouse may be made a carrier of *L. grippolyphosa* experimentally and that it sheds the leptospires without showing any apparent ill effects for at least eighteen months.

It is true however that urinary elimination is intermittent especially for some strains of *L. grippolyphosa* which may explain at least in part the results of Popp.

Another species of *Microtus* the *montebelli* is the principal carrier of *L. hebdomadis* the agent of nanukayami or seven day fever and of *L. autumnalis* the agent of hasami or autumnal fever both very common in the Japanese countryside.

Finally in Israel *Microtus guentheri* has been considered the carrier of a particular strain of leptospire called *L. gesseni*.⁷ In the years 1949 and 1950 an enormous increase in the numbers of this mouse in the country regions produced in turn an epidemic of leptospirosis in man. Van der Hoeden¹⁵ has comprehensively shown that the supposed *L. gesseni* is no other than *L. grippolyphosa*.

Microtus may also be an occasional carrier of *L. sejroe*, *L. bataviae*, *L. australis* A⁷⁹ 80 and of *L. 'Sorex'* 80

Micromys minutus sorcinus is the habitual carrier of *L. bataviae* in the rice fields of northern Italy. Mino⁸¹ has counted 1200 nests of these little rodents in 140 hectares (346 acres) of rice fields and among the animals examined he found that 21.9 per cent were carriers. Unlike the other mouse species the *Micromys* does not live on the dykes but constructs its nests by hanging them on the stems of the rice plant. As a consequence its urine falls directly into the semistagnant water of the rice field. The *Micromys* is also a carrier of *L. bataviae* in Finland.⁷⁹

Mus spicilegus is a frequent carrier of *L. sejroe* and *L. saxkoebing* and it may also harbor *L. ballum*.

Mus musculus is the principal carrier of *L. ballum*. This leptospire easily infects the white mouse and many strains of this animal are contaminated with this parasite thus constituting a danger for the experimental worker. In fact laboratory cases of infection by *L. ballum* are relatively numerous. It has been shown that the mouse may remain a carrier for more than 100 days after experimental infection.⁶⁵

Mus musculus is also found to be one of the carriers of *L. sejroe* 79 80 and Parnas⁸⁰ have also isolated *L. grippotyphosa* from it.

L. sejroe like the related *L. saxkoebing* is most frequently spread by mice belonging to the genus *Ipodemus* a genus which together with *Epimys brevis caudatus*, is also responsible for the dissemination of *L. autumnalis* in the Far East and in Europe it may harbor *L. pomona*⁸³ and *L. bataviae*.⁸¹

Arvicola terrestris has been recognized as a carrier of *L. grippotyphosa*, *L. saxkoebing* and *L. Sorex*.⁸⁰

In all probability many other species of mice are carriers of leptospires especially in the tropical regions. Since however in their case it is usually a question of single isolations of the parasites from the kidney it is not yet possible to confirm whether they are in fact true carriers or rather temporary shedders of leptospires.

Among the insectivores note may be made of the genus *Sorex* the so called shrew mouse which at least in Central and Eastern Europe is the carrier of a leptospire corresponding to the type *Poi*.

In the Union of Soviet Socialist Republics the hedgehog—both *Irinaceus europaeus* and *E. auritus*—is a carrier of leptospires the systematic position of which is still badly defined.⁸¹ In Czechoslovakia Kmetz⁸⁵ has isolated from the same species a leptospire similar to *L. australis*. 1. Parnas⁸⁰ has isolated *L. bataviae* and *L. Sorex* from *E. reumanicus*.

Finally it may be mentioned that in the Andaman Islands Collier and Essveld⁸⁶ have isolated from the brain of a bat belonging to the genus *Cynopterus* a leptospire later classified by Collier and Mochtar⁸⁷ as *L. schuiffneri*. The same type was also isolated later from the blood of a patient.

In Java Collier and Mochtar⁸⁷ have also isolated from the kidneys of bats belonging to the genus *Cynopterus* two strains of *Leptospira* belonging to a new serologic type *L. cynopteri*.

With regard to the possibility that bats may be carriers of pathogenic leptospire it is worth mentioning that in central Africa in the region between the rivers Sagan and Omo where Archetti and I¹⁰ established the existence among the inhabitants of infections from leptospira the natives believe that the illness is spread by bats which are very abundant around the wells from which the men and animals drink. The same belief is also reported by Rizzotti¹¹ who records the presence of cases of leptospirosis in the south of Ethiopia. Rizzotti states that the Amhara in northwest Ethiopia call the illness *besid elithof* which means bat sickness.

These opinions of natives merit a certain consideration it must not be forgotten that in Africa the Calla who inhabit British East Africa and Southern Ethiopia had noted correctly the relationship between malaria and the mosquito both of which they were already calling by the same name long before this connection was even suspected in Europe. On the other hand I have searched unsuccessfully for the presence of leptospire in numerous bats captured in the rice growing regions of Spain.¹²

Among domestic animals the dog undoubtedly has the major epidemiological importance for man. The diffusion of leptospire among dogs is very high and their infection is obviously facilitated by their habit of sniffing where other dogs have urinated. The leptospire that most often infect dogs are *L. canicola* and *L. icterohaemorrhagiae*. The first is the species predominant in most European countries in America and in Australia. *L. icterohaemorrhagiae* on the other hand is more prevalent in Italy and Finland. Alexander *et al*¹³ hold that *L. canicola* better adapted to the dog prevails in more developed countries where there is greater opportunity for contact between dog and dog while *L. icterohaemorrhagiae* should be more frequent in less developed countries of predominantly agricultural economy where there is greater opportunity for contact between dogs and rodents. This hypothesis is not easy to support. In fact in Italy there is also a distinct predominance of *L. icterohaemorrhagiae* in dogs that live in the large cities where contact with small rodents is less apt to occur. It is possible however that *L. canicola* predominates where the density of the canine population is higher and the contacts between the dogs are more frequent and easier.

In nature the dog may also become infected with other types of leptospire namely *L. grippityphosa*, *L. sejroe*, *L. australis* & *L. pomona*, *L. autumnalis*, *L. batariae* leptospire of the *hebdomadis* group and with a special serotype isolated by Van Riel¹⁴ in the Congo region of Africa. However it is not known whether they may become carriers of such types.

Dogs may fall ill with a serious form of leptospirosis and their mortality especially in young animals is very high. Sometimes the leptospire provoke a chronic and progressive nephritis which causes the death of the animal after one or two years irrespective of whether it is a leptospire carrier.

After infection by *L. canicola* dogs often become carriers less often after infection by *L. icterohaemorrhagiae*. For example Klarenbeek¹⁵ found leptospiuria in 59 per cent of dogs infected with *L. canicola* and in only 26 per cent of those infected with *L. icterohaemorrhagiae*. Yamamoto¹⁶ found respec

Microtus may also be an occasional carrier of *L. sejroe*, *L. bataviae*, *L. australis* A⁹ 80 and of *L. Sorex*⁷⁸ 80

Micromys minutus sorcinus is the habitual carrier of *L. bataviae* in the rice fields of northern Italy. Mino⁸¹ has counted 1200 nests of these little rodents in 140 hectares (346 acres) of rice fields and, among the animals examined he found that 21.9 per cent were carriers. Unlike the other mouse species the *Micromys* does not live on the dykes but constructs its nests by hanging them on the stems of the rice plant. As a consequence its urine falls directly into the semistagnant water of the rice field. The *Micromys* is also a carrier of *L. bataviae* in Finland.⁷⁹

Mus spicilegus is a frequent carrier of *L. sejroe* and *L. saxkoebing* and it may also harbor *L. ballum*.

Mus musculus is the principal carrier of *L. ballum*. This leptospire easily infects the white mouse and many strains of this animal are contaminated with this parasite thus constituting a danger for the experimental worker. In fact, laboratory cases of infection by *L. ballum* are relatively numerous. It has been shown that the mouse may remain a carrier for more than 100 days after experimental infection.⁶⁸

Mus musculus is also found to be one of the carriers of *L. sejroe*⁷⁹ 80 and Parnas⁸⁰ have also isolated *L. grippolyphosa* from it.

L. sejroe like the related *L. saxkoebing* is most frequently spread by mice belonging to the genus *Ipodemus*, a genus which together with *Lipomys brevis caudatus* is also responsible for the dissemination of *L. autumnalis* in the Far East and in Europe it may harbor *L. pomona*⁷³ and *L. bataviae*.⁸¹

Irricola terrestris has been recognized as a carrier of *L. grippolyphosa*, *L. saxkoebing* and *L. Sorex*.⁸⁰

In all probability many other species of mice are carriers of leptospires especially in the tropical regions. Since however in their case it is usually a question of single isolations of the parasites from the kidney it is not yet possible to confirm whether they are in fact true carriers or rather temporary shedders of leptospires.

Among the insectivores note may be made of the genus *Sorex*, the so called shrew mouse, which at least in Central and Eastern Europe is the carrier of a leptospire corresponding to the type *I. 01*.

In the Union of Soviet Socialist Republics the hedgehog—both *Erinaceus europaeus* and *E. auritus*—is a carrier of leptospires, the systematic position of which is still badly defined.⁸¹ In Czechoslovakia Kmety⁸⁵ has isolated from the same species a leptospire similar to *I. australis*. Parnas⁸⁰ has isolated *L. bataviae* and *I. Sorex* from *E. reumanicus*.

Finally it may be mentioned that in the Andaman Islands Collier and Esseveld⁸⁶ have isolated from the brain of a bat belonging to the genus *Cynopterus* a leptospire later classified by Collier and Mochtar⁸⁷ as *I. schuffneri*. The same type was also isolated later from the blood of a patient.

In Java Collier and Mochtar⁸⁷ 89 have also isolated from the kidneys of bats belonging to the genus *Cynopterus* two strains of *Lepiospira* belonging to a new serologic type *L. cynopteri*.

that in young pigs *L. pomona* may produce an evident illness that is sometimes quite serious and that in the pregnant sow may produce abortion.

Pigs remain carriers of these two types of leptospire for a long period (one year according to Schmidt and Ciovanella¹⁶) and sometimes they eliminate enormous quantities of leptospires with the urine. Persons concerned with the feeding of pigs easily incur infections of leptospira. These infections usually evince the symptoms of the so-called aseptic meningitis and they were known for a long time as "young swineherds meningitis".

The infected urine of the pigs via the drains of the sty may also pollute nearby water courses and ponds thus infecting those who by reason of their work or through bathing may come into contact with such waters.^{17, 18}

An interesting example is described by Mitov, Jankov and Savov¹⁰⁰ in Bulgaria where in 1953 there was a serious epidemic due to *L. pomona* among the inhabitants of the Stalin (Varna) region following a flood that had submerged numerous pigsties existing in the locality.

The pig is subject to infection by numerous other types of leptospire. From the kidneys of pigs Kmety, Plesko and Chvilo⁶ have isolated *L. canicola*, *L. sejroe* and *L. ballum*. Coghlan, Norval and Seiler¹⁰¹ have also isolated *L. canicola* from the urine of the pig. It is not known however whether the pig may actually become a carrier for these types or for what length of time.

An interesting report by Kmety *et al.*⁶ states that pig carriers may be seronegative or else may possess antibodies for a type of leptospire different from that eliminated with the urine.

Leptospirosis is also a serious problem in cattle and although it has been widely studied especially in the United States, the Union of Soviet Socialist Republics and Israel, there does not yet appear to be a satisfactory solution of it. The cattle may fall sick with leptospirosis either in a mild form or to a serious or even fatal degree. A high percentage of abortion in cows is attributed to leptospirosis especially in the United States where the Department of Agriculture has calculated that the annual financial loss caused by *Leptospira* infections in cattle has amounted to as much as \$112 282 000.¹⁰²

I do not know nor shall I discuss the accuracy of this figure since it is irrelevant to my subject. I emphasize merely that the serum of cattle very often shows an anti *Leptospira* agglutinin level that is rather difficult to interpret because it generally relates to healthy animals that have not had morbid manifestations even in the past. Clearly finding anti *Leptospira* antibodies in a cow that has aborted does not justify one in unhesitatingly attributing responsibility for the abortion to the leptospire especially when it is pointed out that the presence of leptospires in the fetus or in stillborn calves is hardly ever proved. In addition the experimental infection of pregnant cattle produces abortion much more rarely than many authors believe.

Nevertheless it is undoubtedly true that cattle may become ill with leptospirosis and also may remain carriers and shedders of leptospires for long periods of time. In the United States and in Australia the type of leptospire most often involved is *L. pomona*; in Israel and in North Africa it is *L. grippolyphosa* and in the Soviet Union an *L. bovis* that probably corresponds to

tively 1.9 per cent and 3.3 per cent. In the United States Alexander *et al.*⁴ found 8 carriers among 92 dogs seropositive for *L. canicola* (8.7 per cent) but none among 32 that were seropositive for *L. icterohaemorrhagiae*.

The state of carrier is maintained in dogs for a considerable time (at least two to six months according to Brunner and Meyer⁵) as has been shown experimentally it never persists however as long as it does in rats. It should also be noted that the emission of leptospire with the urine occurs in a discontinuous manner so that a single negative test is not sufficient to exclude the possibility that a dog is a carrier.

Generally the percentage of dogs that become carriers after infection does not seem to be very high. Recently it fell to my colleagues and to me to examine a training school for Alsatian wolf dogs belonging to the police. Shortly before it had been possible to verify serologically cases of illness due to *L. canicola* that had occurred among the dogs. Sera from all the animals of the school were examined when the acute phase of the small epidemic had already passed. Forty seven dogs almost all of the 52 dogs examined were clearly seropositive for *L. canicola*, often with very high titers of antibodies indicating recent infection. Among 43 of these dogs examined 6 or 14 per cent showed the presence of *L. canicola* in the urine. This is a fairly high percentage but not extremely so.

The high frequency of *Leptospira* infections in dogs contrasts with the rather low frequency of infections of canine origin in man. This is obviously due to the fact that man and his foodstuffs have less occasion to become contaminated with the urine of dogs than with that of the small rodents.

Besides dogs the jackal is also a habitual carrier of *L. canicola*. Van der Hoeden²⁵ has attributed considerable importance to this animal in sustaining the epizootic of *L. canicola* which periodically affects cattle in the south of Israel. It is however probable that this importance is not so great. In fact as Van der Hoeden himself admits²⁶ the almost complete extermination of the jackals in the region has not noticeably influenced the progress of the epizootic. Evidently the infection also spreads from bovine to bovine and from dog to dog.

The cat theoretically greatly exposed to the infection because of its contact with mice and rats is rarely identified as a carrier however. In Indonesia cats have been found that are carriers of *L. bataviae* and *I. jaranica*.²⁷ Klarenbeek¹⁷ has observed cats in The Netherlands that are carriers of *L. canicola* and Gaub (unpublished data) in Alaska found cats that carried *I. hebdomadis*.

Silver foxes may also be carriers of pathogenic leptospire whether of *L. icterohaemorrhagiae* or in the Union of Soviet Socialist Republics of two special types called DVA and DVB which the author has found to correspond respectively to *L. hyos* and *I. pomona*. It is not known whether these animals remain carriers for a short or long time. In any case their epidemiological importance for man is very limited.

Another domestic animal that is often a carrier of pathogenic leptospire and a source of human cases of infection is the pig. The two types of leptospire that most often infect the pig are *I. pomona* and *I. hyos*. Little is known of the morbid manifestations due to this second type. However it is known

that in young pigs *I. pomona* may produce an evident illness that is sometimes quite serious and that in the pregnant sow may produce abortion.

Pigs remain carriers of these two types of leptospire for a long period (one year according to Schmidt and Ciovanella¹⁶) and sometimes they eliminate enormous quantities of leptospores with the urine. Persons concerned with the feeding of pigs easily incur infections of leptospira. These infections usually evince the symptoms of the so-called aseptic meningitis and they were known for a long time as young swineherds' meningitis.

The infected urine of the pigs via the drains of the sty may also pollute nearby water courses and ponds thus infecting those who by reason of their work or through bathing may come into contact with such waters.^{17, 18}

An interesting example is described by Mitov, Jankov and Savov¹⁰⁹ in Bulgaria where in 1953 there was a serious epidemic due to *I. pomona* among the inhabitants of the Stalin (Varna) region following a flood that had submerged numerous pigsties existing in the locality.

The pig is subject to infection by numerous other types of leptospire. From the kidneys of pigs Kmetz, Plesko and Chylo⁴ have isolated *L. canicola*, *L. sejevae* and *L. ballum*. Coghlan, Norval and Seiler¹⁰⁸ have also isolated *I. canicola* from the urine of the pig. It is not known however whether the pig may actually become a carrier for these types or for what length of time.

An interesting report by Kmetz *et al.*⁴ states that pig carriers may be seronegative or else may possess antibodies for a type of leptospire different from that eliminated with the urine.

Leptospirosis is also a serious problem in cattle and although it has been widely studied especially in the United States, the Union of Soviet Socialist Republics and Israel there does not yet appear to be a satisfactory solution of it. The cattle may fall sick with leptospirosis either in a mild form or to a serious or even fatal degree. A high percentage of abortion in cows is attributed to leptospirosis especially in the United States where the Department of Agriculture has calculated that the annual financial loss caused by *Leptospira* infections in cattle has amounted to as much as \$112 282 000.¹⁰²

I do not know nor shall I discuss the accuracy of this figure since it is irrelevant to my subject. I emphasize merely that the serum of cattle very often shows an anti *Leptospira* agglutinin level that is rather difficult to interpret because it generally relates to healthy animals that have not had morbid manifestations even in the past. Clearly finding anti *Leptospira* antibodies in a cow that has aborted does not justify one in unhesitatingly attributing responsibility for the abortion to the leptospire especially when it is pointed out that the presence of leptospores in the fetus or in stillborn calves is hardly ever proved. In addition the experimental infection of pregnant cattle produces abortion much more rarely than many authors believe.

Nevertheless it is undoubtedly true that cattle may become ill with leptospirosis and also may remain carriers and shedders of leptospores for long periods of time. In the United States and in Australia the type of leptospire most often involved is *L. pomona*; in Israel and in North Africa it is *L. grippotyphosa* and in the Soviet Union an *L. botis* that probably corresponds to

L. grippotyphosa and an *L. titulina*, the systematic position of which is still uncertain have been described. It may be noted that strains isolated from cattle in Israel appear to possess some serologic characteristics differing slightly from those of *L. grippotyphosa* so that some authors maintain that it should be considered separately with the name *L. bovis*. In Israel, infections by *L. canicola* are also quite frequent.⁹

Cattle are able to shed leptospire in the urine only for limited periods of time. According to Gillespie *et al.*² such periods average about thirty days but in some cases may exceed one hundred days. Bernkopf and Olitzki¹⁰ in Israel have never succeeded in finding leptospire in the urine of seropositive cows. It is to be noted however that these investigators were limited to a simple direct microscopical search for the spirochetes.

The quantity of leptospire contained in the urine is very high even greater than 1 million per cc. in a case reported by Gillespie *et al.*² it reached 100 million per cc. One can well understand how such enormous quantities of leptospire if coming into contact with water or moist soil may easily spread the contagion to both animals and man.

Even so it seems that *L. pomona* of bovine origin so rarely causes cases of infection in man that in the United States leptospirosis is essentially a veterinary problem and not one of human medicine.

The case for *L. grippotyphosa* is different in fact, both in Israel and in the Union of Soviet Socialist Republics numerous cases of human infection contracted by direct or indirect contact with bovine carriers have occurred. The reason for this difference in infection has not yet been explained.

Another still obscure problem is posed by this question why in the United States do cattle often present agglutinins for *L. sejroe* when this leptospire in contrast to *L. pomona* has never been isolated from them? Indeed in one case, Galton *et al.*¹⁰³ isolated a strain of *L. pomona* from the urine of a cow that was seropositive only for *L. sejroe*.

Similar observations have been made not only in cattle but also in man by Wiesmann¹⁰⁴ in Switzerland. In my own researches on the other hand I have not observed that either human or bovine serum from subjects who have recovered from an infection by *L. pomona* appears able to agglutinate *L. sejroe* to any significant extent.

Sheep and goats may also fall sick with leptospirosis sometimes in a very serious form. Eighteen per cent of the mortality among lambs in New Zealand is due to *L. pomona* according to Webster and Reynolds¹⁰⁵ as much as 44 per cent mortality among goats in Israel according to Van der Hoeden¹⁰⁶ is due to *L. grippotyphosa*. In Turkey Hakioglu¹⁰⁷ has found leptospire in the kidneys of sheep.

Naturally infected sheep may void leptospire for at least nine months according to Webster and Reynolds¹⁰⁵ but Van der Hoeden has not found leptospire in the urine of convalescent sheep.¹⁰⁸

It must therefore be admitted that in Israel sheep in addition to transmitting the infection from one to another contract it from cattle with which they come into contact or with mice that are carrying *L. grippotyphosa*.

The horse is another animal that is very receptive to many types of lepto

spire In addition to becoming ill with the normal sort of infection due to leptospires the horse is also often subject to a particular form of iridocyclitis of relapsing development that often leads to blindness and is called in Italian *malattia della luna* (moon sickness)

Horses however insofar as is known to date do not seem to become true carriers They may eliminate leptospires only for a short time after the infection¹⁰⁸

Pathogenic leptospires have occasionally been isolated from other animals However these have been isolated discoveries in animals that have little contact with man To these cases apart from any doubts as to whether they are true carriers one may attribute a negligible epidemiological significance The following is a list of these animals and the types of leptospire that they have been found to harbor *Citellus citellus* (*I pomona* and *I grippolyphosa*) *Cricetus cricetus* (*I grippolyphosa*) *Herpestes javanicus* (*I javanica*) *Fiber zibethicus* (*I sejroe*) *Apocomus syhaticus* (*I sejroe* and *I Sorex*) *Neomys fodiens* (*I grippolyphosa*) *Castor canadensis* (*I icterohaemorrhagiae*) *Mustela kodiensis* (*I hebdomadis*) *Ondatra* sp (*L icterohaemorrhagiae*) *Clethrionomys rutilus* (*L icterohaemorrhagiae*) *Vulpes herimani* (*I hebdomadis*) *Nesokia bandicota* (*I australis* & *L pyrogenes* and *I icterohaemorrhagiae*) *Myocastor coypus* (*L icterohaemorrhagiae*) and *Eutamias glareolus* (*I grippolyphosa*)

Our discussion of man has been left to the last It has already been noted that man becomes a carrier and a source of infection only rarely In humans leptospires generally disappear from the urine within 40 days from the onset of the infection However Schueffner and Bohlander¹⁰⁹ have found leptospires in the urine of 2 convalescents for 15 days Carnier and Bailly¹¹⁰ ¹¹¹ in 2 other cases for 100 and 103 days and Wimaers and Renoux¹¹ for 3 months Frugoni and Capellani¹¹² have described 2 cases in which healthy subjects who had been in contact with sick patients were eliminating leptospires with their urine Finally Manine Cristau and Plazy¹¹⁴ Cristau¹¹⁵ and Lettit¹¹⁶ described 5 cases of interhuman infection in subjects who had assisted the sick during an epidemic that occurred in Lorient France in 1917 Cases of infection by sexual contact that occurred in Alaska⁴ and one described by Doeleman⁴ are also to be noted

The Identification of Carriers

The simplest way in which carriers may be identified is by the examination of a drop of urine or of the sediment obtained from it after high speed centrifugation in the dark field of the microscope When one is dealing with animals that may be sacrificed one may also make microscopic preparations of a pulp of the renal cortex Instead of direct examination in the dark field one may resort to examination of a preparation that is colored by a method of silver staining

Pulp of the kidney or the urine under examination may also be inoculated into the peritoneum of guinea pigs young chicks or hamsters (in searching for *L canicola*) Then the inoculated animal is kept under observation to see whether it may die show signs of sickness or produce anti *Leptospira* antibodies Better still a blood culture may be made in a suitable medium draw

ing blood from the animal six to seven days after the inoculation. The culture may also be prepared with pieces of renal cortex obtained under sterile conditions.

In my opinion the best method if the animal may be sacrificed is to carry out a direct culture from the kidney; if on the other hand the animal must be kept alive then it is best to inoculate 2 to 3 cc. of urine into the peritoneum of a guinea pig and to carry out a blood culture after 6 days. The blood cultures must be examined after 7, 15, and 30 days. They should be discarded if still negative only after such a period.

When a direct culture from the kidney is done care must be taken to seed only a very small fragment of tissue in each test tube so as to avoid introducing into the culture any antibodies contained in such tissue as these might inhibit the development of any leptospirae that may be present.

When carrying out inoculations of urine it must be remembered that quite often the leptospiruria is intermittent, consequently it is necessary to draw further samples of the urine at different times or on different days. These samples must then be inoculated into the test animal immediately, as any delay might endanger the vitality of the leptospire.

If the animal under examination voids acid urine it is advisable before drawing the sample to feed it on an alkaline diet or to give it some sodium bicarbonate or citrate.

Disinfection of the Carriers

I do not intend to discuss here the various measures that it is expedient to adopt to protect subjects exposed to contagion spread by animal carriers; this is a topic related to prophylaxis of the disease. One thing that concerns the fight against animal carriers more closely is the need to distinguish the animals that are noxious or at least useless and may be killed such as small wild rodents from the useful ones which instead one may attempt to disinfect. However disinfection of an animal carrier is not easy to accomplish. Brunner and Meyer¹⁷ have been particularly concerned with this topic. They have established the fact that although penicillin is effective against acute infection by leptospira in the dog it is nevertheless incapable of disinfecting the animal in spite of the fact that it is found in high concentration in the urine.

These investigators have achieved their best results with the use of streptomycin. Doses of 40 mg./kg. body weight administered 3 to 5 days in succession have always succeeded in achieving the desired purpose. Aureomycin also gave some good results (5 to 40 mg./kg.) for 3 days.

In any case the problem of carrier disinfection especially of dogs and the large domestic animals is one that would be worth studying deeply. Thus far no such study has been undertaken.

References

- 1 ZIEGLER M. 1935. Acta Pathol. Microbiol. Scand. 12: 511.
- 2 SCHLEFFNER W. & W. A. KUENEN. 1923. Ned. Tijdschr. Geneesk. 67: 2018.
- 3 GILLESPIE R. W. H. S. G. KENY L. M. RINGEN & I. K. BRACKEN. 1957. Am. J. Vet. Research 18: 76.

- 4 ALEXANDER A D C & GLIFISER P MALNATI & H YODER 1950 Am J Hyg 65 43
- 5 BERNSTEIN T & J A BAKER 1954 J Infectious Diseases 94 53
- 6 KMETZ F J PLF KO & F CHALO 1956 Zentr Bakteriöl Parasitenk Orig 167 243
- 7 PROEHOFMAN S 1930 Studie over ler epidemiologie van de ziekte van Weil oer haren verwerker en de daaraan verwante organ smen I roefschr Amster lam
- 8 RUY A C J F MINAENKOF & J W WOLFF 1938 Acta 4th Intern Congr Trop Med Amster lam 33
- 9 AOKI Y K KANEKO & T MORIMOTO 1935 Z Hyg Infektionskrankh 117 208
- 10 VAN DER HOEDEN J 1936 Ann Inst Pasteur 56 207
- 11 VAN DER HOEDEN J 1943 Ned Tijdschr Geneesk 79 1943
- 12 VAN DER HOEDEN J 1946 Bull Research Council Israel 6E 1
- 13 RIMPAL W & J KALICH 1948 Berlin u Munich tierärztl Wochschr 3 29
- 14 WALCH SORGDORFER W 1939 Bull Health Org I of N 8 151
- 15 VAN DER HOEDEN J 1943 Trans Roy Soc Trop Med Hyg 47 364
- 16 SCHMIDT G & R GIOVANELLA 1947 Schweiz Arch Tierheilk 89 1
- 17 KLARFNBEE A 1938 Acta 4th Intern Congr Trop Med Amster lam 381
- 18 LOW D C F N BERGMAN C W HIAIT & C A CLEISER 1956 J Infectious Diseases 98 260
- 19 UHLENHUTH P & L ZIMMERMANN 1933 Med Klin Munich 30 1393
- 20 UHLENHUTH I 1935 Zentr Bakteriöl Parasitenk Orig 135 154
- 21a SCHLEFFNER W & H BOHLANDER 1942 Ned Tijdschr Geneesk 86 1341
- 21b SCHLEFFNER W & H BOHLANDER 1947 Zentr Bakteriöl Parasitenk Ori 149 193
- 22 UHLENHUTH P & W FROMME 1916 Z. Immunität forsch 25 31
- 23 BABUDIERI B & F BIOCICA 1941 Keni ist super sanità 4 144
- 24 DOOLEMAN F P J 1932 Ned Tijdschr Geneesk 76 5057
- 25 UNITED STATES NAVY 1953 Report of survey for control of leptospirous in Kodiak and the Aleutian Islands.
- 26a HECKER A & R OTTO 1911 Deut med Wochschr 18 870
- 26b HECKER A & R OTTO 1911 Ver öffentl Militär sanitätsa 46
- 27 UHLENHUTH I & W FROMME 1930 Welche krankheit In Hanfluch der Pathol Mikroorganismen 7(1) 48 G Fischer & L Schwarzenberg Jena & Berlin Germany & Vienna Austria
- 28 REITER H & RAMME 1916 Deut med Wochschr 42 1287
- 29 UHLENHUTH P & P KLIN 191 Z Hyg Infektion krankh 84 517
- 30 SCHLEFFNER W & A MOCHTAR 1977 Arch Schiffs- u Tropen Hyg 31 149
- 31 SCHLEFFNER W & A MOCHTAR S PROEHOFMAN & L HONIG 192 Abhanfl Gel Auslanlk 26 500
- 32 GAY D M & A W SELLARDS 192 Ann Tr p Med Parasitol 21 321
- 33 SAWYER W A & J H BAKER 1978 Am J Trop Med 8 1
- 34 UHLENHUTH P & W FROMME 1919 Z Immunitätsforsch 28 1
- 35 KLEIGER I J & M A JNER 1978 Trans Roy Soc Trop Med 22 181
- 36 BLANC G 1920 Compt rend soc biol 83 485
- 37 NOGUCHI H 195 Zentr ges Hyg u Grenz 11 285
- 38 STOKES J H I RYLE & W H TYTLER 191 Lancet 1 142
- 39 HUBNER H & H REITER 1916 Z Hyg Infektionskrankh 81 171
- 40 DIETRICH W 1917 Z Immunitätsforsch 26 563
- 41 BLANCHARD M G LEFROC & J LAIGRET 1923 Bull soc pathol exotique 16 184
- 42 BONNE C 1974 Compt rend soc biol 91 247
- 43 BURGDORFER W & F G PICKENS 1954 J Infectious Diseases 94 84
- 44 BURGDORFER W 1956 Exptl Parasitol 5 571
- 45a SCHLOSSBERGER H H LANGBEIN & G KRELZ 1957 Z Immunitätsforsch 109 366
- 45b SCHLOSSBERGER H H LA CERN & G KRELZ 1954 Acta Trop 11 300
- 46 KREPAOGORSKAJA T A & M M REMENKOVA 1957 Zhur Mikrotol Epidemol Imm oiol 28 251
- 47 CHIETTI G 192 Ann igiene 22 513
- 48 CORRALES M 1919 Compt rend soc biol 82 14
- 49 ALTAYA V M BARRERA & J VILLALBA C published data
- 50 DOHME A 1939 Med Welt 13 1551 1556
- 51 MARTIN L & A PETTIT 1919 Spi ochétose actér hémorragique Monogr Inst Pasteur Masson Paris France
- 52 OBA S 1973 Arch J Exptl Med 1 1
- 53 TERSKICH Cited by Kujumgiev "

- 54 ALJUNGILA I 1957 *Vet ital* 8 490
- 55 MOCHMANN H 1957 *Berlin u Munch Tierarztl Wochschr* 70 181
- 56 BERNKOPF H & I OLITZKI 1948 *Board Scient Ind Research* 20
- 57 VISSOTSJAI B V & V G REDKINA 1954 *Zhur Mikrobiol Epidemiol Immunobiol* 25 8
- 58 HALLALER C & H ALHN 1947 *Z Hyg Infektionskrankh* 124 125
- 59 HOAG W G W S GOCHINGOUR & R H YAGER 1953 *Proc Soc Exptl Biol Med* 83 112
- 60 BERNKOPF, H 1948 *Report on bovine leptospirosis in Palestine Board Scient Ind Research*
- 61 KENY S G & R W H GILLESPIE 1957 *Vet Med* 310
- 62 CHALQUEST R R 1957 *J Poultry Sci* 36 110
- 63 ALTAYA V M BARRERA & C MARIN 1956 *Rev sanidad hig publ Madrid* 30
- 64 MCKINLEY E B 1928 *Proc Soc Exptl Biol Med* 26 26
- 65 NICOLLE C & C LEBAILLY 1918 *Compt rend soc biol* 81 457
- 66 ROBINSON C H 1974 *Am J Hyg* 4 327
- 67 BABUDIERI B & L BIANCHI 1940 *Z Immunitätsforsch* 98 37
- 68 BABUDIERI B & L BIANCHI 1940 *Rend ist sanità publ Rome* 3 11
- 69 FRAGA DE AZEVEDO J J S VALENTE & J S QUEIROZ 1951 *Anais inst med trop* 8 4 621
- 70 COVALEDA J A PUMAROLA & CANTARELL 1953 *Rev ibérica parasitol* 13 289
- 71 FARINA R & B BABUDIERI 1957 *Rend ist super sanità* 20 857
- 72 ESSEVELD H W A COLLIER & A MOCHTAR 1940 *Mededeel Dienst Volksgezondheid Ned Indie* 29 1
- 73 MOCHTAR A & W A COLLIER 1939 *Schweiz Z allgem Pathol u Bakteriol* 2 319
- 74 IDO Y K HOKI H ITO & H WANI 1917 *J Exptl Med* 26 341
- 75 SCHLEFFNER W & H BOHLANDER 1943 *Z Immunitätsforsch* 104 237
- 76 KATHE J 1950 *Zentr Bakteriol Parasitenk Orig* 155 199
- 77 POPP L 1950 *Z Hyg Infektionskrankh* 131 575
- 78 OLEJNICK K E & S SHAFERSON 1952 *Trans Roy Soc Trop Med Hyg* 46 165
- 79 SALMINEN A 1956 *Ann Med Exptl et Biol Fenniae Helsinki* 34 (suppl 5)
- 80 PARNAS J Unpublished data
- 81 MINO P 1941 *Munch med Wochschr* 88 96
- 82 BABUDIERI B F BUSSINELLO F BAJOCCHI A SALVI & L MASSA 1955 *Riv in fortune e mal profess* 3 453
- 83 BORG PETERSEN C & K L FEMMFSTAD 1956 *J Am Vet Med Assoc* 178 204
- 84 ANANIN V V 1951 *Zool Zhur* 30 375
- 85 KMETZ E 1954 *Zentr Bakteriol Parasitenk Orig* 161 382
- 86 COLLIER W A & H ESSEVELD 1938 *Mededeel Dienst Volksgezondheid Ned Indie* 27 267
- 87 COLLIER W A & A MOCHTAR 1939 *Geneesk Tijdschr Ned Indie* 28 356
- 88 COLLIER W A & A MOCHTAR 1938 *Mededeel Dienst Volksgezondheid Ned Indie* 28 356
- 89 COLLIER W A & A MOCHTAR 1939 *Geneesk Tijdschr N Indie* 79 226
- 90 BABUDIERI B & I ARCHETTI 1940 *Rend ist sanità publ Rome* 3 116
- 91 RIZZOTTI G 1939 *Riv biol coloniale Rome* 2 241
- 92 VAN RIFL J 1952 *Ann soc helge mcd trop* 32 683
- 93 YAMAMOTO S 1943 *Japan J Vet Sci* 5 45
- 94 BRUNNER A T & K F MEYER 1947 *Iroc Soc Exptl Biol Med* 70 450
- 95 VAN DER HOEDEN J 1955 *J Trop Med Hyg* 207
- 96 VAN DER HOEDEN J 1955 *J Am Vet Med Assoc* 126 207
- 97 ESSEVELD H W A COLLIER 1938 *Z Immunitätsforsch* 93 512
- 98 SAVINO E & F FENNELLA 1946 *Dia mcd* 18 564
- 99 TOPCIU V & C DUMA 1956 *Timisoara Med* 1 311
- 100 MITOV A N JANKOV & S SAVIN 1955 *Dof rokaciestveni lej toq irosti Nauka izkumstvo Sofia*
- 101 COLHILL J D J NORVAI & H J SPILFER 1957 *Brit Med J* 1 257
- 102 UNITED STATES DEPARTMENT OF AGRICULTURE 1954 *Losses in agriculture Agr Research Service Washington D C*
- 103 GALTON M M J A ACFFE A LEWIS & F C IRATHER 1956 *J Am Vet Med Assoc* 128 87
- 104 WIESMANN F 1977 *Z Tr penmed u Parasitol* 8 305
- 105 WEBSTER W M & B A LEYNOLDS 1955 *New Zealand Vet J* 47

- 106 VAN DER HOEDEN J 1953 J Comp Pathol Therap 63 101
- 107 HAKIOGLU F 1956 Zentr allgem Pathol u pathol Anat 85 575
- 108 BRYAN J T 1955 Canad Vet 45 16
- 109 SCHLEFFNER W & H BOHLANDER 1942 Zentr Bakteri I Parasitenk Orig 149 191
- 110 CARNIER M & J RAILLY 1917 Comité rend soc Biol 80 38
- 111 CARNIER M & J RAILLY 1918 Presse méd 26 505
- 112 WILMAERS I & I KENCKX 1917 Arch Med Belg 70 115 207
- 113 FRUONI C & CAFFLANI 1917 Genaro Med 164
- 114 MANINE I CRISTAU & LAZY 1917 Comité rend soc Biol 80 531
- 115 CRISTAU I 1917 Comité rend soc Biol 80 178
- 116 PETTIT A 1917 Comité rend soc Biol 80 774 780 848
- 11 BRUNNER K T & K R MEYER 1950 Am J Vet Research 11 89

LEPTOSPIROSIS INFECTIONS IN MAN

By Joe L. Stockard

Department of Preventive Medicine and Rehabilitation, University of Maryland School of Medicine, Baltimore, Md.

and

Theodore E. Woodward

Department of Medicine, University of Maryland School of Medicine, Baltimore, Md.

Introduction Weil is credited with having first described cases of leptospirosis in humans as early as 1886¹ although Landouzy reported his own observations in 1883. The classic disease is characterized by fever, hemorrhagic phenomena, hepatic, renal, and vascular decompensation, and a high fatality rate. In 1915 Inada² isolated the etiologic agent, a spirochete, later identified as a leptospire by Noguchi.³

Studies by various European, Australian, Japanese, Indonesian, and American investigators have identified many immunologically distinct leptospire strains as pathogenic for man and a host of other animals. These specific microorganisms are naturally adapted to unique epidemiologic patterns and they produce highly variable clinical features in animals they infect. The human manifestations are variable, and it behooves physicians to recognize both the common and the unusual clinical features, as well as the environmental hazards that lead to infection of man.

The clinical manifestations of classic Weil's disease, the milder forms of leptospirosis observed in a tropical environment, and the North American types will be described.

Classic Weil's disease (*Leptospira icterohaemorrhagiae*) Following an incubation of 4 to 19 days, the disease begins abruptly, often with a rigor as the first symptom. Fever develops; this continues between 102° and 104° F. for about 8 days and falls by lysis. The symptom complex that usually develops includes prostration, severe and unrelenting headache, anorexia, and myalgia. Muscle tenderness is a characteristic clinical feature, and arthralgia may be intense. Nausea, vomiting, and diarrhea are frequently present and occasionally severe. Mental confusion and delirium may occur earlier in leptospirosis than in other infectious diseases.

Injection of bulbar and palpebral conjunctivae, a very common sign, appears early and persists for several days. Subconjunctival hemorrhages may develop, but increased lacrimation is unusual. Other ocular manifestations during the acute stage of illness include hemorrhagic lesions of the retina, temporary and permanent oculomotor paresis, optic neuritis, and papilledema. Optic atrophy is rare. Leptospiral uveitis is a common complication noted from 2 weeks to 12 months following the onset. Prognosis is good for complete recovery of normal motor and visual functions.

Icteric lesions of the gastric and intestinal mucosa may lead to profuse bleeding from the gastrointestinal tract. Cutaneous lesions may be petechiae or may resemble the exanthems observed in rubella or scarletina. Severe

peripheral vascular decompensation comparable to that observed in the rickettsioses may contribute to a fatal outcome. Myocardial involvement may impair the circulatory status. Jaundice usually appears at the end of the first week simultaneously with defervescence. It may intensify rapidly and persist for a month or more. Deaths occur in jaundiced patients but the intensity of icterus is not necessarily of prognostic significance. Hepatomegaly develops in the jaundiced but splenomegaly is uncommon.

Renal decompensation is characteristic of Weil's disease. Moderate proteinuria may be noted when the illness is mild but in seriously ill patients proteinuria is often heavy. After the first week of disease oliguria or anuria may provoke severe azotemia and electrolyte imbalance of the lower nephron type. Fatalities often result from this serious complication. Diuresis is an obviously favorable omen and fortunately permanent renal damage is unusual.

Central nervous system signs may dominate the clinical picture. Nuchal rigidity can be marked. The spinal fluid pressure is sometimes elevated while microscopic examination reveals a lymphocytic pleocytosis. In jaundiced patients the cerebrospinal fluid may be yellow. Neuritis occasionally complicates the convalescence but usually meningeal signs subside completely.

The peripheral blood usually reveals a leukocytosis with a preponderance of neutrophils. A hypochromic anemia is frequently noted.

A death rate of approximately 30 per cent occurs in jaundiced patients who are severely ill. During the early stages of illness vascular collapse or hemorrhage from the gastrointestinal tract may cause death. Usually the patient succumbs with uremic signs during the second week of illness. The clinical manifestations that develop from derangement of the kidney, liver, or vascular system are the most vital.

Mild cases. Milder forms of leptospirosis occur in Malaya where this diagnosis was confirmed in 244 of 852 febrile illnesses investigated in 1954 and 1955.⁴ Leptospire isolates from the blood in 90 cases were subsequently identified. These represented 12 of the 20 serogroups recognized in the classification proposed by Wolff and Broom.¹⁴ Organisms of the *icterohaemorrhagiae* and *bataviae* serogroups are usually responsible for severe Weil's disease and were isolated from 6 and 10 cases respectively in this study. The remaining strains usually produce milder illness⁵ and did so in Malaya as evidenced by the low mortality rate of 0.8 per cent (TABLE 1).

The clinical manifestations presented by these patients consisted of an acute febrile phase lasting about eight days and a convalescent phase of approximately two weeks duration. Onset abrupt in 10 per cent of cases was often marked by rigor. Shortly thereafter fever, malaise, anorexia, headache, myalgia, and retro-orbital pain rapidly developed (FIGURE 1).

Anorexia and nausea occurred in 96 per cent of the patients. Eighty-five per cent of the cases experienced emesis at least once during the febrile period and many had almost continuous vomiting or retching. Mild diarrhea rarely occurred and except for these few instances bowel function was undisturbed. During the febrile phase headache and retro-orbital pain were continuous. Intensity of these symptoms varied in proportion to fever. Subjective myalgia

TABLE 1

SEROGROUPING OF LEPTOSPIRES OF HUMAN ORIGIN IN MALAYA 1954 AND 1955

Serogroup	Number of strains
<i>pyrogenes</i>	18
<i>hebdomadis</i>	10
<i>bataviae</i>	10
<i>canicola</i>	9
<i>grippolyphosa</i>	8
<i>autumnalis</i>	8
<i>celledoni</i>	8
<i>icterohaemorrhagiae</i>	6
<i>schnuifneri</i>	5
<i>australis</i> 1	4
<i>janica</i>	2
<i>pomona</i>	1
<i>djasiman</i>	1
Total	90

CLINICAL MANIFESTATIONS OF LEPTOSPIROSIS

(182 military patients)

SYMPTOMS

GASTROINTESTINAL SYMPTOMS

96%

HEADACHE

95%

MYALGIA

77%

RESPIRATORY SYMPTOMS

71%

ABRUPT ONSET

70%

RETROBULBAR PAIN

55%

PHOTOPHOBIA

31%

ARTHRALGIA

29%

FIGURE 1 Salient subjective findings in military patients with leptospirosis. Reproduced by permission from *Transactions of the Association of American Physicians* 14

was generalized and most severe in the calf abdominal pectoral and cervical muscle groups. Cough present in half of the patients was productive in 50 per cent of those with this complaint. Hemoptysis occurred in 60 per cent of the cases with a productive cough and was severe in two instances. A few patients complained of dyspnea chest pain coryza and sore throat but these did not constitute prominent features of the illness. Visual disturbances were never detected and photophobia was unusual (FIGURE 2).

Physical signs coincided with those usually observed in leptospirosis. Injection of the conjunctivae was present in 85 per cent of the cases appearing within 24 hours after onset. Its intensity as graded by the physician varied from 1+ to 4+ the majority being graded 4+ (generalized lymphadenopathy was detected in one half of the patients but enlargement of the nodes was less than is ordinarily observed in infectious mononucleosis scrub typhus

CLINICAL MANIFESTATIONS OF LEPTOSPIROSIS

(182 military patients)


SIGNS

FIGURE 2 Objective clinical manifestations of leptospirosis in military patients. Reproduced by permission from *Transactions of the Association of American Physicians*.

or dengue. Objective evidence of muscular tenderness was demonstrated in 43 per cent of the cases. This finding created difficulty in evaluating the significance of pain, particularly over the abdomen, costovertebral angles, and neck. Appendectomy was performed unnecessarily in one instance. Examination of spinal fluid was performed in ten patients who had marked stiffness of the neck. Of these, six revealed lymphocytic pleocytosis ranging from 40 to 400 cells per cu mm.

Hemorrhagic signs appearing from the fourth to eighth days of disease included the following in decreasing order of frequency: hemoptysis, hemateme, epistaxis, petechiae, subconjunctival hemorrhages, melena, hematuria, ecchymoses, and submucosal hemorrhages in the pharynx. Abnormal bleeding was observed in patients infected by all serogroups except *janica pomona* and *senloti*. Bleeding was severe in two patients with hemoptysis, one of whom died on the fourth day after onset. *L. australis B* was isolated from the fatal case. Cough and signs of pulmonary congestion prompted roentgen examination of the chest in forty-two cases. Abnormalities varying from slightly increased vascularity to small localized areas of consolidation were revealed on the films in eleven cases. Of these, six presented physical signs and roentgen evidence of pneumonia. In one instance, these findings dominated the clinical course. This patient's blood yielded *L. grippityphosa* in culture. Other leptospires identified with clinical findings of pneumonia were from the *pyrogenes canicola* and *schuiffneri* serogroups.

Skin eruptions observed in one-fourth of the group appeared between the fourth and eighth days of disease and persisted from a few hours to several days. These lesions consisted of either small macules or maculopapules and

TABLE 1
SEROGROUPING OF LEPTOSPIRES OF HUMAN ORIGIN IN MALAYA 1954 AND 1955

Serotype	Number of strains
<i>pyrogenes</i>	18
<i>hebdomadis</i>	10
<i>bataviae</i>	10
<i>canicola</i>	9
<i>grippolyphosa</i>	8
<i>autumnalis</i>	8
<i>celledoni</i>	8
<i>icterohaemorrhagiae</i>	6
<i>schuettneri</i>	5
<i>australis A</i>	4
<i>javanica</i>	2
<i>pomona</i>	1
<i>djasiman</i>	1
Total	90

CLINICAL MANIFESTATIONS OF LEPTOSPIROSIS

(182 military patients)

SYMPTOMS




FIGURE 1 Salient subjective findings in military patients with leptospirosis. Reproduced by permission from *Transactions of the Association of American Physicians* 18

was generalized and most severe in the calf, abdominal, pectoral, and cervical muscle groups. Cough, present in half of the patients, was productive in 50 per cent of those with this complaint. Hemoptysis occurred in 60 per cent of the cases with a productive cough and was severe in two instances. A few patients complained of dyspnea, chest pain, coryza, and sore throat, but these did not constitute prominent features of the illness. Visual disturbances were never detected and photophobia was unusual (FIGURE 2).

Physical signs coincided with those usually observed in leptospirosis. Injection of the conjunctivae was present in 85 per cent of the cases, appearing within 24 hours after onset. Its intensity, as graded by the physician, varied from 1+ to 4+, the majority being graded 4+. Generalized lymphadenopathy was detected in one half of the patients, but enlargement of the nodes was less than is ordinarily observed in infectious mononucleosis or scrub typhus.

tations cardiovascular renal and hepatic decompensation rarely occur. A high incidence of aseptic type meningitis is characteristic in patients with swinechill disease. In Alabama Schaeffer¹² has reported the clinical features observed in 26 of 50 patients who contracted *L. pomona* infection while swimming in a creek. After an incubation period of 2 to 10 days the illness was characterized by chills and fever, headache, nausea and vomiting, myalgia and meningismus. Some had conjunctival injection and photophobia. Diarrhea, abdominal pain and transient maculopapular eruptions were present in a few cases. Lymphocytic pleocytosis and slight elevations of spinal fluid pressures were found in 6 patients who showed signs of meningitis. After an illness of 5 to 14 days all recovered uneventfully except for a few who experienced a brief uncomplicated relapse.

Beeson and Hankey¹³ in the United States and Broom¹⁴ in England have incriminated leptospirosis as a cause of benign serous type meningitis. Beeson demonstrated significant rises in titers of leptospiral antibodies in 4 per cent of 535 paired sera obtained from patients with meningitis of unknown etiology. Therefore leptospirosis should be considered as a diagnostic possibility along with lymphocytic choriomeningitis, mumps, herpes, poliomyelitis and Coxsackie and ECHO viruses, as well as the various neurotropic encephalitides.

L. autumnalis was isolated first in Japan where the clinical illness was recognized as a mild febrile disorder of seven days' duration occurring in harvesters during the autumn months.¹⁵ Icterus, ocular complications and death occurred rarely.

In 1943 an illness resembling dengue appeared in soldiers at Fort Bragg, N. C. and was described by Daniels and Grennan.¹⁶ The cardinal features consisted of a short febrile period, moderate prostration, splenomegaly and a rash on the anterior aspects of the tibia. After extensive studies by numerous investigators Tatlock¹⁷ adapted the agent thought to be a virus to several laboratory animals and to fertile hens' eggs. The filtrable agent reproduced the characteristic manifestations in volunteers. In 1951 Cochenour¹⁸ identified the etiologic agent as a leptospire immunogenically similar to those of the *autumnalis* serogroup.

Stroup and Riley¹⁹ recently reported the occurrence of maculopapular lesions over the pretibial area in two siblings presumably infected by *L. pomona*. Each patient became ill eight days after swimming in a stagnant pond where pigs watered. The onset was marked by fever, anorexia and malaise, headache, lethargy, and vomiting appeared within two days. Examination on the third day of disease revealed mild muscular tenderness and nuchal rigidity. Proteinuria, microscopic hematuria and mononuclear pleocytosis were also demonstrated. On the fifth day of disease conjunctival injection was noted and maculopapular lesions resembling erythema nodosum appeared over the pretibial area. After forty-eight hours the eruption faded completely. Both patients recovered uneventfully after a total illness of about one week.

Cultures for leptospires were negative. Paired sera from both patients were examined at the Walter Reed Army Institute of Research, Washington, D. C. They revealed a marked rise in agglutination lysis antibody titers for *L. pomona*. Elevated titers for *L. icterohaemorrhagiae* and *L. autumnalis* were in

were distributed over the head and neck, trunk, and upper extremities. Lesions resembling the rose spots identified with typhoid fever were seen in a few patients on the upper extremities, chest, and abdomen.

Splenomegaly and hepatomegaly were unusual, occurring in only 15 per cent of the cases. Icterus detected by inspection alone was rare.

The total leukocyte count was usually normal, but leukocytosis and leukopenia did occur. Neutrophilia was noted in 80 per cent of the patients during the febrile period.

Qualitative proteinuria, casts, and increased cellular elements were demonstrated at some time during the febrile phase in 80 per cent of leptospirosis patients. Oliguria and mild azotemia developed in only 27 cases. It never persisted longer than 3 days.

Liver function was evaluated serially by means of the van den Bergh reaction, total and fractional protein determination, and Kunkel's flocculation test. Liver function, as evaluated by these tests, was disturbed in two thirds of the patients. Elevations of total bilirubin values appeared early and rapidly returned to normal with defervescence. Abnormalities in the albumin globulin ratios and Kunkel's test developed at the beginning of convalescence and persisted for at least 2 weeks. Clinically overt jaundice was observed in only 11 patients, 2 of whom died. Leptospire isolates from the 2 fatal cases were identified subsequently as *L. bataviae* and *L. australis*.⁸ Icterus was also observed in patients infected by organisms in the *hebdomadis canicola schiffneri* and *icterohaemorrhagiae* serogroups.

Defervescence, usually by rapid lysis, marked the beginning of convalescence. All symptoms and signs promptly subsided; urinary findings returned to normal, and icterus, if present, gradually cleared. A few patients experienced a relapse characterized by a return of fever, symptoms, physical signs, and proteinuria for two or three days during the second and third weeks after onset of the acute illness. Convalescence was complicated once by thrombophlebitis and another time by unilateral atrophy of the quadriceps muscles. One man suffered two attacks of leptospirosis separated by four months. On these occasions, cultures of blood yielded *L. schiffneri* and *L. bataviae*, respectively.

With these exceptions, recovery was uncomplicated, and patients resumed their normal activities in three to four weeks.

North American leptospiral types. Small localized outbreaks and sporadic cases of human leptospirosis occur in the United States. The serogroups heretofore reported include *icterohaemorrhagiae canicola pomona*,^{6,7} *bataviae*, and *autumnalis*.⁹

In the United States, as elsewhere, *L. icterohaemorrhagiae* produces illness of variable severity. Leptospire isolates of the *bataviae* serogroup provoke clinical manifestations comparable to classic Weil's disease.⁶ *L. canicola* produces slightly different clinical findings, since jaundice seldom occurs and an aseptic type meningitis is common.¹¹ In other respects, *canicola* fever resembles Weil's disease.

The leptospire *L. pomona*, which is responsible for swineherd's disease, was isolated in Pomona, Australia, by Derrick¹¹ in 1942. The disorder is usually mild with a febrile course of from five to seven days. Hemorrhagic manifes-

LABORATORY DIAGNOSIS OF LEPTOSPIROSIS

By William S. Gochenour, Jr., Chester A. Cleiser, and Martha K. Ward

United States Army Medical Unit, Fort Detrick, Md.

The past decade has seen an increasing awareness of the prevalence of leptospirosis in man, in his domestic animals, and in wildlife in the United States. The public health and livestock sanitary problems posed by these infections and their economic significance to the livestock industry are being increasingly appreciated. This has resulted in an ever growing demand for diagnostic laboratory support in clinical diagnosis, epidemiological investigation, and research. It is the purpose of this paper to review and to evaluate the methods currently available to the leptospiral diagnostic laboratory.

It is perhaps belaboring the obvious to state that to meet the clinician's requirements adequately, the laboratory methods employed must be specific for leptospirosis and reproducible from time to time and in any laboratory, and must unequivocally confirm or negate the clinical impression of leptospirosis within a reasonable period of time. Retrospective diagnostic confirmation may well interest the epidemiologist, but it is of limited value to the clinician. From the practical standpoint, methods for widespread adoption must not be unduly expensive, laborious, or hazardous. Furthermore, they must be within the capabilities of all competent medical laboratory technicians.

Before discussing laboratory methods and techniques, it appears appropriate to review certain characteristics of leptospire and of leptospiral infections in man and in animals that are pertinent to the selection of specimens submitted for examination, their collection and handling, and the choice of the laboratory methods to be employed.

The pathogenic leptospire are fastidious organisms in their environmental and nutritional requirements. They are markedly susceptible to low pH, to the common disinfectants and antiseptics, to drying, and to ordinary pasteurization temperatures. *In vitro* they are sensitive to most antibiotics and are readily overgrown and destroyed by bacterial contaminants in culture media. In a neutral or slightly alkaline environment, however, they may survive for many months at ordinary temperatures, and they have withstood freezing for protracted periods.

In man and in animals, leptospiremia is associated with the initial febrile period of disease. At this same time, leptospire may often be demonstrated in the cerebrospinal fluid. Leptospire are rarely demonstrable in the blood after the abatement of fever and the appearance of detectable circulating antibodies, usually in the second week of illness. However, they may be found for varying periods in the urine, in the kidney tubules, and on occasion in the anterior chamber of the eye.

Laboratory confirmation of the diagnosis of leptospirosis may be made by microscopic visualization of the organisms in appropriate body fluids, by histological demonstration of leptospire and their accompanying pathological changes in affected organs, by isolation of the organisms by direct culture, or

terpreted by Gochenour to represent nonspecific cross immunological reactions. Serum from another sibling who was exposed to the same stagnant pond, revealed antibody titers of 1:100 for *L. pomona*, *L. icterohaemorrhagiae* and *L. sentinelae*.

Summary and conclusion The clinical features of human leptospirosis vary considerably, causing illnesses as severe as fulminant hepatitis and as mild as influenza. The classic manifestations that suggest the diagnosis of Weil's disease are fever, severe prostration, icterus, renal decompensation, hemorrhagic phenomena, and vascular collapse. Milder forms of leptospirosis are more difficult to recognize, but a leptospiral etiology should be considered in any febrile patient when severe headache, vomiting, myalgia, conjunctival injection, proteinuria, and neutrophilia are noted. Leptospirosis is to be suspected further whenever there is a history of exposure to those environmental conditions that favor infection.

References

1. WEIL, A. 1886. Über eine eigenthümliche mit Milztumor, Icterus und Nephritis einhergehende acute Infectionskrankheit. *Deut. Arch. klin. Med.* 30: 209-232.
2. INADA, R., Y. IDO, R. HOKI, R. KANEKO & H. ITO. 1916. The etiology, mode of infection, and specific therapy of Weil's disease (*Spirochaetosis icterohaemorrhagica*). *J. Exptl. Med.* 23: 371-402.
3. NOGUCHI, H. 1917. *Spirochaeta icterohaemorrhagiae* in American wild rats and its relation to the Japanese and European strains. *J. Exptl. Med.* 25: 755-763.
4. McCULLUMB, F. R., JR., J. L. STOCKARD, C. R. ROBINSON, L. H. TURNER, D. C. LEVINS, C. W. MAISEY, M. F. KELLEHER, C. A. GLEISER & J. F. SMADEL. 1957. Leptospirosis in Malaya. I. Sporadic cases among military and civilian personnel. *Am. J. Trop. Med.* 6: 238-256.
5. WALCH, SORCDRAGER, B. 1939. Leptospirosis. *Bull. Health Org. L. of N. S.* 143-356.
6. GOCHENOUR, W. S., JR., R. H. YAGER & P. W. WETMORE. 1950. Antigenic similarity of bovine strains of *Leptospira* (United States) and *Leptospira pomona*. *Proc. Soc. Exptl. Biol. Med.* 74: 199-207.
7. ASHIE, W. F., I. K. PRATT THOMAS & C. W. KUMPE. 1941. Weil's disease: a complete review of American literature and an abstract of the world literature. *Medicine* 20: 145-210.
8. GOCHENOUR, W. S., JR., R. H. YAGER, P. W. WETMORE, L. B. EVANS, T. J. BYRNE, A. ALEXANDER & J. A. HIGHTOWER. 1951. Indonesian Weil's disease in Puerto Rico and the United States. (Abstr.) *Federation Proc.* 10: 408-409.
9. GOCHENOUR, W. S., JR., J. E. SMADEL, F. B. JACKSON, L. B. EVANS & R. H. YAGER. 1952. Leptospiral etiology of Fort Bragg fever. *U. S. Public Health Repts.* 67: 811-813.
10. VAN THIEL, P. H. 1948. *The Leptospirae*. Universitaire Pers Leiden, Netherlands.
11. DERRICK, T. H. 1942. *Leptospira pomona*. *Med. J. Australia* 29 (1): 431.
12. SCHAEFFER, M. 1951. Leptospiral meningitis: investigations of a water borne epidemic due to *L. pomona*. (Abstr.) *Proc. Am. Soc. Clin. Invest. In J. Clin. Invest.* 30: 670-671.
13. BEESON, P. B. & D. D. HANKEY. 1952. Leptospiral meningitis. *N. M. J. Arch. Internal Med.* 89: 575-583.
14. BROOM, J. C. 1951. Leptospirosis in England and Wales. *Brit. Med. J.* 2: 689-697.
15. DANIELS, W. B. & H. A. GRENNAN. 1943. Pretilial fever: an obscure disease. *J. Am. Med. Assoc.* 122: 361-365.
16. TATLOCK, H. 1947. Studies on a virus from a patient with Fort Bragg fever (pretilial fever). *Proc. Soc. Exptl. Biol. Med.* 26: 287-297.
17. STROUP, J. G. & H. D. RILEY, JR. 1957. Pretilial fever due to *L. pomona*: report of 2 cases in siblings. *J. Pediatr.* 50: 716-720.
18. McCULLUMB, F. R., JR., J. L. STOCKARD & T. F. WOODWARD. 1956. Leptospirosis as a major cause of short term pyrexia in a tropical environment. *Trans. Assoc. Am. Phy.* 49: 122.
19. WOLFF, J. W. & J. C. BROOM. 1954. The genus *Leptospira* Noguchi, 1917. Principles of classification and a suggested system based on antigenic analysis. *Documenta Med. Geograph. et Trop.* 6: 18-95.

as to be microscopically demonstrable within a few days after inoculation. Chief among the disadvantages are the costs in laboratory space and personnel and the hazards and laboriousness of the methods. One additional caution should be noted. Many of the Swiss white mouse colonies of the United States are infected with *Ieptospira ballum*. This has caused delay and confusion in the investigation of the etiology of other diseases and must be excluded in instances where only animal isolation of leptospires has proved successful.

Direct culture of patient's blood in the early febrile stage of disease has been employed for more than thirty years. A number of media have been developed for the isolation and maintenance *in vitro* of these organisms. In our experience Fletcher's semisolid medium⁸, Schuffner's modification of Verwoort's medium⁹ and Korthoff's medium⁹ have been quite satisfactory. Blood cultures have been made in the field and at the bedside by inoculation of bottles of medium closed with rubber skirted diaphragm stoppers at the time of venipuncture and in the laboratory from glass bead defibrinated blood. Both methods were equally efficacious in yielding positive cultures. One point of technique warrants special consideration. In our experience it was essential to use minimal inocula (approximately 0.03 ml. per culture tube or bottle) and to inoculate replicate tubes if positive cultures were to be obtained. Larger quantities of the same specimens yielded negative results. We attribute this to the presence of inhibitors in the specimen which if not diluted sufficiently prevented leptospiral multiplication. That these inhibitors may be specific antibodies is suggested by the successful isolation of leptospires with this technique in 12 instances^{8, 9} from blood specimens having a serum leptospiral agglutinating titer of 1:100 or greater.

As leptospires grow slowly *in vitro* it was necessary to incubate cultures for a minimum of twenty-eight days before they could safely be discarded as negative. Of those positive, leptospires were demonstrable in one third by the seventh day of incubation, in one half by the fourteenth day and in the remainder at the twenty-eighth day. The use of the semisolid medium greatly facilitated the periodic examinations required as these cultures could be examined macroscopically for the characteristic disk of leptospiral growth approximately 2 cm. below the surface of the medium obviating in many instances tedious dark ground microscopic examination.

In our experience the sole disadvantage of direct culture is the demonstration of leptospires is its unsuitability for the examination of bacterially contaminated specimens. With suitable specimens, however, our much greater success with direct culture than with animal inoculation in the isolation of leptospires from patient's blood leads us to agree with Fletcher¹⁰ that for this purpose the use of laboratory animals offers no advantages over direct culture but merely adds to the burden of the diagnostic laboratory.

Serologic confirmation of the diagnosis of leptospirosis rests upon demonstration of a significant (at least fourfold) rise in specific antibody in the examination of acute and convalescent sera. While the presence of antibody reflects experience of the host with leptospires, no single serologic positive finding regardless of the titer observed is adequate to substantiate unequivocally a diagnosis of acute leptospirosis. This is particularly true of agglutinating

by animal inoculation or by demonstration of a significant rise in specific antibody through the examination of acute and convalescent sera. These methods will be considered in terms of the requirements mentioned above.

Microscopic demonstration of leptospires in blood, cerebrospinal fluid or urine by dark ground or phase contrast illumination affords the theoretical advantages of rapidity, simplicity and economy. In practice the low concentration of leptospires in these fluids affords small chance of their demonstration.

Further, the ease with which artifacts such as protoplasmic extrusions from formed blood elements and fibrin tendrils moving by Brownian motion may be mistaken for leptospires severely limits the usefulness of the method. Exceptions are the urine of swine and of rodent carriers which are often so heavily populated as to resemble well grown cultures. With these exceptions we feel that microscopic examination of blood and other body fluids is worthless as a diagnostic laboratory procedure and we consider suspect any diagnoses resting solely on this method for confirmation.

Histological demonstration of leptospires in the affected organs of individuals succumbing to the disease may be accomplished by the employment of silver impregnation techniques. The technique of choice at the Armed Forces Institute of Pathology, Washington, D. C. is a modified Warthin-Starry staining method.¹ In arriving at a histopathological diagnosis of acute leptospirosis the pathologist must consider that leptospires unrelated to acute disease in an individual may be harbored in the kidney tubules. This coincidental carrier state should not be imputed to have etiological significance in the case of disease under study. Rather diagnosis must rest on both demonstration of leptospires and compatible and significant pathological changes. A case in point was the first recognition of porcine leptospirosis in the United States. While leptospires were recovered and many of the swine involved had evidence of leptospiral infection, the fulminating and fatal disease seen in the herd was properly attributable to hog cholera and salmonellosis. The leptospiral infection was coincidental and most probably pre-existing.

A wide variety of laboratory animals has been employed in the isolation of leptospires. These include, in addition to the guinea pig, the golden hamster,² the Swiss white mouse,⁴ the *Meriones*,⁵ the baby chick,⁶ and the embryonated hen's egg.⁷ With the exception of the latter, these animals were originally advocated because of their susceptibility to lethal infection by one or more leptospiral serotypes. Recognition of increasing numbers of benign leptospiral serotypes that produce little or no detectable clinical disease in these animals has resulted in their present use largely as accelerated living incubators and as filters for contaminated materials. Animals are inoculated intraperitoneally with relatively large quantities of the specimen (1 to 5 ml.) and peritoneal fluid, heart blood or both are examined for the presence of leptospires microscopically and culturally every day for the first 4 to 6 days postinoculation. The principal advantages of the inoculation of laboratory animals are that materials unsuited for direct cultural examination may be used; that large volumes of a specimen may be inoculated into a single animal; and that, frequently, leptospires may multiply so rapidly in the living host

as to be microscopically demonstrable within a few days after inoculation. Chief among the disadvantages are the costs in laboratory space and personnel and the hazards and laboriousness of the methods. One additional caution should be noted. Many of the Swiss white mouse colonies of the United States are infected with *Leptospira ballum*. This has caused delay and confusion in the investigation of the etiology of other diseases and must be excluded in instances where only animal isolation of leptospires has proved successful.

Direct culture of patient's blood in the early febrile stage of disease has been employed for more than thirty years. A number of media have been developed for the isolation and maintenance *in vitro* of these organisms. In our experience Fletcher's semisolid medium⁸, Schuffner's modification of Verwoort's medium⁹ and Korthoff's medium⁹ have been quite satisfactory. Blood cultures have been made in the field and at the bedside by inoculation of bottles of medium closed with rubber skirted diaphragm stoppers at the time of venipuncture and in the laboratory from glass bead defibrinated blood. Both methods were equally efficacious in yielding positive cultures. One point of technique warrants special consideration. In our experience it was essential to use minimal inocula (approximately 0.03 ml per culture tube or bottle) and to inoculate replicate tubes if positive cultures were to be obtained. Larger quantities of the same specimens yielded negative results. We attribute this to the presence of inhibitors in the specimen which if not diluted sufficiently prevented leptospiral multiplication. That these inhibitors may be specific antibodies is suggested by the successful isolation of leptospires with this technique in 12 instances^{8, 9} from blood specimens having a serum leptospiral agglutinating titer of 1/100 or greater.

As leptospires grow slowly *in vitro* it was necessary to incubate cultures for a minimum of twenty-eight days before they could safely be discarded as negative. Of those positive leptospires were demonstrable in one third by the seventh day of incubation, in one half by the fourteenth day, and in the remainder at the twenty-eighth day. The use of the semisolid medium greatly facilitated the periodic examinations required as these cultures could be examined macroscopically for the characteristic disk of leptospiral growth approximately 2 cm below the surface of the medium obviating in many instances tedious dark ground microscopic examination.

In our experience the sole disadvantage of direct culture to the demonstration of leptospires is its unsuitability for the examination of bacterially contaminated specimens. With suitable specimens however our much greater success with direct culture than with animal inoculation in the isolation of leptospires from patient's blood leads us to agree with Fletcher¹⁰ that for this purpose the use of laboratory animals offers no advantages over direct culture but merely adds to the burden of the diagnostic laboratory.

Serologic confirmation of the diagnosis of leptospirosis rests upon demonstration of a significant (at least fourfold) rise in specific antibody in the examination of acute and convalescent sera. While the presence of antibody reflects experience of the host with leptospires no single serologic positive finding regardless of the titer observed is adequate to substantiate unequivocally a diagnosis of acute leptospirosis. This is particularly true of agglutinating

by animal inoculation, or by demonstration of a significant rise in specific antibody through the examination of acute and convalescent sera. These methods will be considered in terms of the requirements mentioned above.

Microscopic demonstration of leptospires in blood, cerebrospinal fluid, or urine by dark ground or phase contrast illumination affords the theoretical advantages of rapidity, simplicity, and economy. In practice, the low concentration of leptospires in these fluids affords small chance of their demonstration.

Further, the ease with which artifacts such as protoplasmic extrusions from formed blood elements and fibrin tendrils moving by Brownian motion may be mistaken for leptospires severely limits the usefulness of the method. Exceptions are the urine of swine and of rodent carriers which are often so heavily populated as to resemble well grown cultures. With these exceptions we feel that microscopic examination of blood and other body fluids is worthless as a diagnostic laboratory procedure and we consider suspect any diagnoses resting solely on this method for confirmation.

Histological demonstration of leptospires in the affected organs of individuals succumbing to the disease may be accomplished by the employment of silver impregnation techniques. The technique of choice at the Armed Forces Institute of Pathology, Washington, D. C., is a modified Warthin-Starry staining method.¹ In arriving at a histopathological diagnosis of acute leptospirosis the pathologist must consider that leptospires unrelated to acute disease in an individual may be harbored in the kidney tubules. This coincidental carrier state should not be imputed to have etiological significance in the case of disease under study. Rather, diagnosis must rest on both demonstration of leptospires and compatible and significant pathological changes. A case in point was the first recognition of porcine leptospirosis in the United States.² While leptospires were recovered and many of the swine involved had evidence of leptospiral infection, the fulminating and fatal disease seen in the herd was properly attributable to hog cholera and salmonellosis. The leptospiral infection was coincidental and most probably pre-existing.

A wide variety of laboratory animals has been employed in the isolation of leptospires. These include, in addition to the guinea pig, the golden hamster,³ the Swiss white mouse,⁴ the *Mervines*,⁵ the baby chick,⁶ and the embryonated hen's egg.⁷ With the exception of the latter, these animals were originally advocated because of their susceptibility to lethal infection by one or more leptospiral serotypes. Recognition of increasing numbers of 'benign' leptospiral serotypes that produce little or no detectable clinical disease in these animals has resulted in their present use largely as accelerated living incubators and as filters for contaminated materials. Animals are inoculated intraperitoneally with relatively large quantities of the specimen (1 to 5 ml.) and peritoneal fluid, heart blood, or both are examined for the presence of leptospires microscopically and culturally every day for the first 4 to 6 days postinoculation. The principal advantages of the inoculation of laboratory animals are that materials unsuited for direct cultural examination may be used; that large volumes of a specimen may be inoculated into a single animal; and that, frequently, leptospires may multiply so rapidly in the living host

lent correlation with this test was obtained with the live antigen microscopic agglutination test performed in the Division of Veterinary Medicine

While the above mentioned tests are highly serogroup specific one very important fact must be noted Paraspecific responses or cross reactions with heterologous serogroups frequently may be observed at levels higher than those of the homologous organism Particularly notorious in this regard are instances of proved *Leptospira icterohaemorrhagiae* infection wherein antibody levels against *L. canicola* often exceed those against the homologous leptospire Thus while valuable presumptive evidence of the identity of a leptospire may be afforded by the host response elicited following infection definitive establishment of its identity can be effected only by its isolation and characterization In this connection the Division of Veterinary Medicine WRAIR has been designated by the World Health Organization of the United Nations with the concurrence of the Surgeon General United States Public Health Service and the Surgeon General United States Army as the Leptospiral Reference Laboratory for North America Definitive identification of leptospiral strains can be made by this laboratory It is strongly recommended that isolates made from man and animals be sent to it for identification and characterization

The complement fixation test of Randall Wetmore and Warner¹⁴ employing sonically disrupted leptospire as antigens is the oldest of the genus specific serologic tests It has been used extensively in the laboratories of the Division of Veterinary Medicine WRAIR the United States Army Tropical Research Medical Laboratory San Juan Puerto Rico and in the field laboratory in Kuala Lumpur Malaya In these laboratories excellent results were obtained¹⁵ in the diagnosis of human leptospirosis Results in the examination of animal sera have been disappointing and further study of the factors responsible is indicated The advantages of this test are its simplicity of performance and the requirement for employment of only a limited number of antigens in the test battery Its major disadvantage in addition to that noted above is that being genus specific it affords no information as to the presumptive identity of the infecting leptospire

The hemagglutination test of Chang¹⁶ employing erythrocyte sensitizing substances (ESS) and the hemolytic test of Cox¹⁷ are both genus specific In the evaluation study previously mentioned¹³ good correlation with the microscopic live antigen agglutination test was obtained in the examination of human sera Both tests however gave disappointing results in the examination of animal sera Results of continued evaluation studies of the hemolytic test¹⁸ would indicate that it is as efficacious in the laboratory diagnosis of human leptospirosis as is the microscopic live antigen agglutination test Its advantages of simplicity requiring but a single antigen and of freedom from hazard recommended it to the laboratory concerned with diagnosis of human leptospirosis *per se* or as a screening procedure to be employed prior to examination of sera with either living or killed microscopic agglutinating antigens

In summary we feel that laboratory diagnosis of leptospirosis is now within the capabilities of any public health livestock sanitary or hospital laboratory

antibodies which may remain at high levels for many months following infection

The serologic tests employed in the laboratory diagnosis of leptospirosis include microscopic and macroscopic agglutination tests with living or dead leptospiral antigens, complement fixation tests employing intact leptospirally sonically disrupted leptospirally or leptospiral extracts or metabolites as antigens hemagglutination tests using erythrocyte sensitizing substance (ESS) and hemolytic tests using an antigen similar to the latter

The agglutination lysis test of Schuffner and Mochtar¹¹ one of the earliest leptospiral serologic tests has been and is today the most widely used and it is the serologic method against which all others have been evaluated This is a microscopic agglutination test in which living leptospirally are employed as antigens As are all of the leptospiral agglutination tests it is highly serogroup specific This requires the employment of an adequate battery of antigens if antibody against an unknown leptospire is to be detected This battery may range from as many as two dozen antigens in a reference laboratory examining sera from the entire world to as few as eight in a diagnostic laboratory concerned only with specimens derived from man in the United States In addition to the above the hazard attendant on using living antigens the laboriousness of the method and the problems of standardizing antigen sensitivity¹² are seriously limiting factors Further grossly contaminated sera and those containing high concentrations of preservative cannot be examined

The microscopic agglutination test employing killed antigens has been used extensively in this country and abroad The use of killed antigen eliminates the hazards attendant on the agglutination lysis test and also permits the examination of sera too heavily contaminated to be examined by this technique When killed antigens are properly standardized and are periodically checked for sensitivity and absence of autoagglutination their use yields results essentially identical with those obtained with living antigens In collaborative studies we have examined aliquots of the same sera in the Special Bacteriology Laboratories Communicable Disease Center Chamblee Ga and in the Division of Veterinary Medicine Army Medical Service Graduate School (now Walter Reed Army Institute of Research) Walter Reed Army Medical Center Washington D C These studies reported in part by Ward *et al*⁹ illustrate the virtual identity of results obtained by use of these two methods It should be noted however that there are minor differences in interserogroup cross reactions observed with living and with killed antigens and that any selection of the test battery of diagnostic antigens to be employed should take this factor into consideration

Macroscopic plate agglutinating antigens thus far developed have not yet been adequately evaluated to determine whether the shortcomings of lack of sensitivity lack of stability and difficulty of interpretation that characterized earlier macroscopic agglutinating antigens have been overcome

The capillary tube agglutination test of Stoenner has recently been evaluated in a collaborative study with the Division of Veterinary Medicine Walter Reed Army Institute of Research (WRAIR)¹³ In Stoenner's laboratory excel

lent correlation with this test was obtained with the live antigen microscopic agglutination test performed in the Division of Veterinary Medicine

While the above mentioned tests are highly serogroup specific one very important fact must be noted. Paraspecific responses or cross reactions with heterologous serogroups frequently may be observed at levels higher than those of the homologous organism. Particularly notorious in this regard are instances of proved *Leptospira icterohaemorrhagiae* infection wherein antibody levels against *L. canicola* often exceed those against the homologous leptospire. Thus while valuable presumptive evidence of the identity of a leptospire may be afforded by the host response elicited following infection definitive establishment of its identity can be effected only by its isolation and characterization. In this connection the Division of Veterinary Medicine WRAIR has been designated by the World Health Organization of the United Nations with the concurrence of the Surgeon General United States Public Health Service and the Surgeon General United States Army as the Leptospiral Reference Laboratory for North America. Definitive identification of leptospiral strains can be made by this laboratory. It is strongly recommended that isolates made from man and animals be sent to it for identification and characterization.

The complement fixation test of Randall, Wetmore and Warner¹⁴ employing sonically disrupted leptospires as antigens is the oldest of the genus-specific serologic tests. It has been used extensively in the laboratories of the Division of Veterinary Medicine WRAIR, the United States Army Tropical Research Medical Laboratory, San Juan, Puerto Rico, and in the field laboratory in Kuala Lumpur, Malaya. In these laboratories excellent results were obtained¹⁵ in the diagnosis of human leptospirosis. Results in the examination of animal sera have been disappointing and further study of the factors responsible is indicated. The advantages of this test are its simplicity of performance and the requirement for employment of only a limited number of antigens in the test battery. Its major disadvantage in addition to that noted above is that being genus specific it affords no information as to the presumptive identity of the infecting leptospire.

The hemagglutination test of Chang¹⁶ employing erythrocyte sensitizing substances (ESS) and the hemolytic test of Cox¹⁷ are both genus specific. In the evaluation study previously mentioned¹² good correlation with the microscopic live antigen agglutination test was obtained in the examination of human sera. Both tests however gave disappointing results in the examination of animal sera. Results of continued evaluation studies of the hemolytic test¹⁸ would indicate that it is as efficacious in the laboratory diagnosis of human leptospirosis as is the microscopic live antigen agglutination test. Its advantages of simplicity, requiring but a single antigen and of freedom from hazard recommended it to the laboratory concerned with diagnosis of human leptospirosis *per se* or as a screening procedure to be employed prior to examination of sera with either living or killed microscopic agglutinating antigens.

In summary we feel that laboratory diagnosis of leptospirosis is now within the capabilities of any public health, livestock, sanitary or hospital laboratory.

We feel that the success attainable by direct cultural examination of blood in the early febrile period of disease warrants inclusion of this procedure in laboratory studies of febrile illness of undetermined origin. At this time serologic confirmation of the diagnosis of leptospirosis may best be made by microscopic agglutination employing living or killed antigens. When antigens are available the hemolytic test of Cox may well supplement the agglutination tests or supplant them in small hospital laboratories.

Existing knowledge of leptospiral antigen-antibody reactions does not encourage hope of reducing significantly the time required for serologic confirmation of the diagnosis of leptospirosis. More promise lies in the development of methods for the earlier demonstration of leptospire.

References

- 1 BRIDGES C H & L LEE 1957 Kerr's improved Warthin Starry technique. Lab Invest 6(4) 357-367
- 2 GOCHENOUR W S JR, R V JOHNSON, R H YAGER & W S GOCHENOUR 1952 Porcine leptospirosis. Am J Vet Research 13(47) 158-160
- 3 RANDALL R & H A COOPER 1944 The golden hamster (*Cricetus auratus*) as test animal for the diagnosis of leptospirosis. Science 100 133-134
- 4 LARSEN C I 1941 Susceptibility of young mice to *Leptospira icterohaemorrhagiae*. Public Health Rept U S 56 1546-1556
- 5 SYMPOSIUM ON THE LEPTOSPIROSES 1952 Med Sci Publ Army Med Serv Grad School Walter Reed Army Med Center No 1 171
- 6 HOAG W G W S GOCHENOUR JR & R H YAGER 1953 Use of baby chick for isolation of leptospire. Proc Soc Exptl Biol Med 83 712-713
- 7 GLEISER C A, W JAHNKE & R J BYRNE 1955 Avian leptospirosis. Studies on chick embryo culture. Cornell Vet 45 297-304
- 8 GOCHENOUR W S JR, R H YAGER, P W WEYMORE & J A HIGHTOWER 1953 Laboratory diagnosis of leptospirosis. Am J Public Health 43 405-410
- 9 WARD M K, M G McDANIEL, H W TATUM, I E STARR & H R WILLIAMS 1956 An epidemic of canicola fever in man with the demonstration of *Leptospira canicola* infection in dogs, swine and cattle. II Laboratory Studies. Am J Hyg 65 59-69
- 10 FLETCHER W 1927-1928 Recent work on leptospirosis. Tsutsugamushi disease and tropical typhus in the Federated Malay States. Trans Roy Soc Trop Med Hyg 21 265-288
- 11 SCHUFFNER W & A MOCHTAR 1927 The differentiation of leptospiral strains. The course of agglutination and lysis. Trop Diseases Bull 24 714-715
- 12 BORG PETERSEN C & A FAGAELS 1949 The influence of the antigen density and other factors on the serum titer in the agglutination-lysis test for leptospirosis. Acta Pathol Microbiol Scand 26 555-567
- 13 GLEISER C A & A D ALEXANDER 1956 Evaluation of methods for the serodiagnosis of leptospirosis. Walter Reed Army Institute of Research Research Rept 171-56
- 14 RANDALL R, P W WEYMORE & A R WARNER 1949 Sonic vibrated leptospirae as antigens in the complement fixation test for the diagnosis of leptospirosis. J Lab Clin Med 34 1411-1415
- 15 McCORMB F R JR, J L STOCKARD, C R ROBINSON, L H TURNER, D I LEVINS, H F KELLEHER, C A GLEISER & J F SMADEL 1957 Leptospirosis in Malaya. I. Sporadic cases among military and civilian personnel. Am J Trop Med Hyg 6(2) 238-256
- 16 CHANG R S & D F MCCOMB 1954 Erythrocyte sensitizing substances from five strains of leptospirae. Am J Trop Med Hyg 3 481-489
- 17 COX C D 1955 Hemolysis of sheep erythrocytes sensitized with leptospiral extracts. Proc Soc Exptl Biol Med 90 610-615
- 18 COX C D & A D ALEXANDER & I C MURPHY 1957 Evaluation of the hemolytic test in the serodiagnosis of human leptospirosis. Walter Reed Army Institute of Research Research Rept 14 57

EPIDEMIOLOGICAL PATTERNS OF LEPTOSPIROSIS

By Mildred M. Calton, Robert W. Menges and James H. Steele
Communicable Disease Center, Public Health Service, Atlanta, Ga.

Introduction

For many years the leptospiroses were considered rare diseases but it is now well established that they occur in humans and animals throughout the world and that they constitute a major problem in certain domestic animals.^{1, 2} More than forty years of extensive research has gradually unraveled a characteristic epidemiological pattern. For clear understanding of this pattern complete epizootiological data are essential. An adequate knowledge of the distribution of leptospires in potential wild and domestic animal carriers and the factors that influence the role of these carriers in the transmission of leptospiral infections to humans is of particular importance. As pointed out by Van der Hoeden³ the lack of sufficient information may result in misinterpretation of facts but further knowledge will clarify the original picture.

The leptospiral infections are zoonoses, that is, diseases transmissible from animal to animal and from animals to humans.⁴ Infection in humans is accidental and with rare exceptions it represents a dead end in the chain of transmission.⁵

Although rodents and certain domestic animals are considered to be the primary animal carriers, recent investigations have revealed a variety of hitherto unrecognized wild mammal hosts¹⁰⁻¹² and current studies are detecting additional sylvatic leptospiral carriers. Some of the reported isolations of different leptospiral serotypes and their hosts are presented in TABLE 1. In the kidneys of these host animals the leptospires display an affinity for the renal cortex and may be found nesting in the lumina of the convoluted tubules. They may be shed in the urine for long periods. Each leptospiral serotype is believed usually to have a primary animal host for which it is particularly adapted; however, these serotypes will infect other animals as well. Furthermore, a single animal species may become infected with a number of different serotypes or even harbor two types at once.¹⁴

Leptospiral Serotypes Known to Occur in Humans or Animals in the United States

As research has progressed in centers located in widely separated areas of the world, the number of antigenically distinct serotypes of the genus *Leptospira* found to cause leptospirosis has increased. With this increase in recognized serotypes and the detection of an even broader host range, the epidemiology of the disease has become more involved. In 1954, Wolff and Broom¹⁵ published a suggested classification system based on serologic analysis that included 34 leptospiral serotypes isolated from man and/or animals. During the relatively short period since this report appeared, at least 25 additional serotypes have been described in Australia,^{16, 17} Africa,¹⁸ Europe,¹⁹ Malaya,¹¹ and the United States.²⁰

Although the primary problem in the United States appears to be with

TABLE 1
REPORTED OCCURRENCE OF CERTAIN IEPOTOSIRAL SEROTYPES IN MAN AND IN ANIMAL HOSTS

Lept 122		Domestic												Wild												Total	
C	P	Type	C	til	D	G	E	She	Sal	Bat	Ba	co	For	G	Ham	Hedge	f	Me	M	Oposu	Ra	Rat	Sh	Slu	Vol	W	
I		<i>L. actinobacilliformis</i>	94	92	85	86			87					86	88			89	91		90						W
II		<i>L. jamaica</i>	9															93	92								at
III		<i>L. p.</i>	94																								
IV		<i>L. sordidus</i>	86																								
V		<i>L. canicula</i>	86																								
VI		<i>L. texanensis</i>	94																								
VII		<i>L. ballum</i>	91																								
VIII		<i>L. pygmaea</i>	86																								
IX		<i>L. australis B</i>	16																								
X		<i>L. cynopteri</i>	94																								
XI		<i>L. senilis</i>	86																								
XII		<i>L. aulicoides</i>	94																								
XIII		<i>L. djassensis</i>	100																								
XIV		<i>L. australis A</i>	102																								
XV		<i>L. pomona</i>	86																								
XVI		<i>L. grippolyphus</i>	105																								
		<i>L. hebdomadis</i>	94																								
		<i>L. medanensis</i>	94																								
		<i>L. volffii</i>	107																								
		<i>L. seyrone</i>	109																								
VII		<i>L. saxatilis</i>	86	97																							
VIII		<i>L. balatensis</i>	110																								
IX		<i>L. semeranga</i>	112																								
X		<i>L. andamanensis</i>																									
		<i>L. malis</i>																									
		<i>L. hirsuta</i>																									

Numbers in table indicate reference

† This culture died before identification could be completed but serum from the goat agglutinated with *L. p.* no antigen in a 1:2560 dilution

Icterohaemorrhagiae, *I. canicola* and *I. pomona* seven other serotypes have been isolated and serologic findings suggest the presence of at least two others. *I. autumnalis* was identified after an eight year period as the cause of Fort Bragg fever.²¹ *I. ballum* has been found frequently in rural house mice and opossums and occasionally in rats and laboratory mice.^{22, 23} Serologic evidence has suggested the presence of *I. bataviae* infection in humans.²⁴ *L. grippolyphosa* in humans⁵ and cattle²⁵ and *I. sejroe* infection in cattle.^{26, 27} Recent cultural studies¹⁰ done at the Field Station of the Communicable Disease Center (CDC) Newton, Ga. on wild mammals trapped in southwest Georgia have yielded a variety of leptospiral serotypes in animals hitherto unknown as hosts including raccoons, skunks and wildcats. In addition isolations have been obtained from opossums and foxes.

During a 6-month period ending in March 1956,¹⁰ 870 mammals including 14 species were trapped in different areas of 6 southwest Georgia counties and suspensions of the kidneys were cultured. Leptospire were isolated from 44 (5.4 per cent). Of these 16 proved to be *I. ballum*, 14 were *I. pomona*, 2 were *I. australis*, 12 were *L. grippolyphosa*, 2 proved to be a new serotype of the *mitis hyos* serogroup tentatively designated as *I. bakeri*²⁸ and 2 belonged to the *hebdomadis* serogroup. Two additional cultures were found to belong to the *mitis hyos* group but are not identical with *I. mitis*, *L. hyos* or *L. bakeri*. Of the 4 remaining 2 were lost prior to serotyping and the identification of 2 has not been completed. In this series opossums, raccoons, striped skunks, foxes and wildcats were found to harbor 1 or more of these leptospiral serotypes. These 5 species represented 79 per cent of the individuals examined. None of the infected animals showed any clinical signs of leptospirosis and no gross kidney lesions were observed. Only 4 of the 44 infected animals were determined to be less than 1 year old.

The highest incidence of infection appeared in striped skunks with isolations from 18 (14 per cent) of 132 animals. It is notable that 11 (61 per cent) of these cultures proved to be *L. pomona*. The areas studied represented 3 groups of predominant vegetative cover: pine, pine and hardwoods and pastures and cultivated fields. There was no great difference among the vegetative types in incidence of infection of the animals cultured. Continuation of these studies at the Newton Field Station has yielded 7 isolations of *L. autumnalis* from raccoons trapped within a 2 mile area.²⁹ The striped skunk, opossum and raccoon continue to show the highest incidence of infection although occasional isolations have been obtained from the spotted skunk, the gray fox and the wildcat.

Although comparatively little attention has been directed toward the study of wild mammals elsewhere in the United States, Roth⁹ recently isolated *L. pomona* from an opossum in Louisiana and Reilly³¹ reported serologic evidence that suggested *L. canicola* infection in raccoons in the state of New York.

Current information regarding the occurrence of leptospiral serotypes in the United States is summarized in TABLE 2.

Recently 1 proved human case and 3 probable cases of leptospirosis caused by *L. ballum* contracted from Swiss albino mice have been found at the Rocky Mountain Laboratory, Hamilton, Mont.³⁰

TABLE 2
LEPTOSPIRAE IN THE UNITED STATES

Se typ	Orig i lat	k w h t	O cu e n e			
			Ma	Dog	Catti	Sw e
<i>L. icterohaemor</i> <i>rhagia</i>	Japan	Rat	Com mon	Occa sional	?	?
<i>L. canicola</i>	Germany	Dog cattle	Occa sional	Com mon	Rare	Rare
<i>L. pomona</i>	The Nether lands	Swine skunk rac coon wildcat opossum	Occa sional	Rare	Com mon	Com mon
<i>L. autumnalis</i>	Australia	Opossum raccoon	Rare	?	?	?
<i>L. ballum</i>	Denmark	Mice gray fox opossum wild cat rat rac coon striped skunk	?	?	?	?
<i>L. grippolyphosa</i>	Russia	Raccoon	Rare	?	Rare	?
<i>L. balantiae</i>	Indonesia	?	Rare	?	?	?
<i>L. sejroe</i>	Denmark	?	?	?	Spo radic	?
<i>Hebdomadis sero</i> group (LT 11*)	United State	Opossum raccoon	?	?	?	?
<i>L. australis A</i>	Australia	Raccoon opossum	?	?	?	?
<i>L. bakeri</i> †	United States	Opossum	?	?	?	?
<i>Mitis hyos sero</i> group (LT 81)	United States	Opossum raccoon	?	?	?	?

* Serologic evidence only

† *Mitis hyos* serogroup

Transmission

Of primary importance in the transmission of leptospiral infection to humans are the animal carriers that become urinary shedders of leptospires. A diagram illustrating the current pattern of the transmission of leptospirosis in the United States is presented in FIGURE 1.

After acute mild or more frequently inapparent infection the organisms become established in the kidney and are eliminated in the urine.¹ Infections of humans and animals result from direct or indirect contact with the infected urine of these shedders. For example direct contact may occur when individuals care for sick animals or handle the tissues of infected animals in abattoirs. Indirect contact occurs when water or moist soil is contaminated and individuals are subsequently exposed while working or swimming or otherwise come in contact with the contaminated environment.²

Arthropod vectors have not been incriminated as yet in the transmission of leptospirosis in nature.¹⁴ However Schlossberger and Langbein¹⁵ were able to infect the tick *Ornithodoros moubata* by feeding it on guinea pigs infected with *L. icterohaemorrhagiae*. Transmission to normal guinea pigs was effected when the ticks were fed 31 days after infection. These observations were confirmed by Burgdorfer and Pickens.¹⁶ Later Burgdorfer^{17, 18} found that *Ornithodoros turicata* and another tick *Dermacentor andersoni*




FIGURE 1. The epidemiology of leptospirosis in the United States. This chart was modified by G. W. Fischer from Byrne.²⁸ Reproduced by permission from the December 1955 issue of *Public Health Reports*, official journal of the Public Health Service, United States Department of Health, Education and Welfare.

became infected after engorgement upon *L. pomona* infected hamsters and transmitted the disease to normal animals. More recently *L. grippolyphosa* was isolated from the European tick *Dermacentor marginatus* S from cattle in the Union of Soviet Socialist Republics.²⁷ Isolated cases of leptospirosis had occurred among farm animals in the area where the ticks were collected. Thus the detection of natural infection increases the possibility that arthropods are vectors.

While leptospires may be shed in milk during the acute phase of the disease in lactating animals,²⁸ Kirschner²⁹ found that the organisms survive only a few hours in whole milk that is leptospirocidal. In milk diluted 1:20 or higher with neutral water they survive a number of days.⁴⁰ There have been no proved cases of leptospirosis that could be traced to drinking infected milk.

The portals of entry of infection are usually the nasal, oral and conjunctival mucous membranes and abraded skin. There is considerable doubt that leptospires penetrate the intact skin and it is unlikely that the digestive tract is an important portal of entry since the pH of the stomach is usually such that the organism may be destroyed quickly.

Environments that favor the survival and spread of leptospires are wet soil, stagnant water or slow moving streams that are neutral or slightly alkaline when temperatures are 22°C or above. Under such conditions leptospires may survive for several weeks.⁴¹⁻⁴⁴

TABLE 3
REPORTED OUTBREAKS OF LEPTOSPIROSIS IN THE UNITED STATE

Date of onset	Locality	No. of cases	Age range	Probable source	Serotype involved
August 1940 July and August 1942 to 1944	Wrens Ga Fort Bragg N C	35 40/yr	Avg 16 Young adults (soldiers)	Cattle ?	<i>L. pomona</i> † <i>L. autumnalis</i>
August 1942 August 1947	Jackson Hole Wyo Calvary Ga	24 10	15 to 19 13 to 24	Dogs Dead mule in creek	<i>L. canicola</i> † ?
August 1949 July 1950	Swainsboro Ga Geneva Ala	12 50	Young adults Adolescent and young adults	Cattle Swine	<i>L. pomona</i> † <i>L. pomona</i> †
July 1952	Columbus Ga	26	5 to 20 and 2 adults	Swine dogs cattle	<i>L. canicola</i>

With the possible exception of the outbreak at Fort Bragg all outbreaks were attributed to swimming in contaminated water

† Serologic evidence only

Outbreaks and Sporadic Cases of Leptospirosis in Humans in the United States

It is not surprising that large outbreaks may occur after group swimming in contaminated water. Several such outbreaks have been reported in the United States as shown in TABLE 3. The first of these occurred in August 1940 near Wrens, Ga.^{45, 46} and involved 35 patients who had been swimming in Brushy Creek. Clinical signs included fever, nausea, vomiting, headache, some diarrhea, and a rash on the pretibial areas of the legs. It was found that approximately 100 yards upstream from the swimming area offal from slaughtered cattle had been dumped by the local butchers. This unusual disease was given the name of Brushy Creek or pretibial fever. A similar outbreak of pretibial fever occurred among troops at Fort Bragg in August 1942 and again in the summers of 1943 and 1944.^{1, 47, 48}

All patients involved in the outbreaks were quartered in the same area of the post. Many, but not all, had been swimming in the nearby ponds, so the source of their infection remains obscure. Not until 1950 was Fort Bragg fever proved to be a leptospiral infection caused by a member of the *L. autumnalis* group.²¹ After this discovery, McCroan reinvestigated the Brushy Creek outbreak and 2 other similar undiagnosed epidemics that had occurred in Georgia: one* in the late summer of 1947 near Calvary⁴⁶ and the other during the same season in 1949 on the Ochopee River near Swainsboro.⁴⁶ Antibodies to *L. pomona* were demonstrated in serum samples collected at the time of the outbreak and during the subsequent investigation in 1953 from several patients involved in the Brushy Creek and Ochopee River outbreaks.

A follow-up investigation in 1953 of another epidemic of an unknown disease that had occurred in August 1942 in two small communities in Wyoming and Idaho revealed serologic evidence of leptospiral etiology.⁵⁰ Here again all

* Investigated by Elmer Hill and John F. McCroan of the Georgia State Health Department, Atlanta, Ga.

persons affected had been swimming in the same pool located at Jackson Hole Wyo. to which dogs and other animals had access.

In late July 1950 an outbreak occurred in Geneva Ala.⁵¹ involving about 50 persons who became ill with an influenzalike disease after swimming in a creek in which dead swine had been seen floating above the swimming hole. Serologic evidence indicated that *L. pomona* was the cause.

When in late July 1952 an outbreak of illness occurred in a small community near Columbus Ga. involving 26 persons who had been swimming in a dammed up creek leptospirosis was suspected immediately.⁵² The subsequent investigation of this outbreak including laboratory studies⁵³ presented an excellent picture of many of the principles involved in leptospiral epidemiology. *L. canicola* infections thought for many years to be limited to dogs and man were demonstrated in swine cattle and dogs that had access to the creek. All persons who became ill had been swimming in the creek. The data showed a significant correlation between attack rates and the frequency and extent of exposure to the water. Attack rates were consistently higher among those giving a history of diving and immersing their heads.

With the possible exception of the Fort Bragg outbreaks each of these water borne leptospiral epidemics followed a similar pattern. They occurred in the late summer during drought periods there was presumed contamination of a stagnant pond or slow moving creek by urine from infected animals and transmission to humans by immersion in the contaminated water. Such water borne outbreaks following swimming or accidental immersion have been reported in many other areas of the world.⁵⁴⁻⁵⁹

Of even greater importance in the epidemiology of leptospiral infections are the opportunities for exposure encountered by workers in certain occupations particularly abattoir workers cane and rice field workers dairy farmers animal husbandrymen veterinarians sewer workers and trench diggers. For example in Italy Spain the Far East and Pacific regions where the rice fields are flooded and worked by barefoot laborers leptospirosis is an important health and economic problem.⁶⁰⁻⁶² The fields usually abound with mice and rats. Babudieri⁶³ found in a survey of 509 apparently healthy workers in Italy that 20 per cent had leptospiral antibodies. The percentage of infection increased in different groups as the length of service increased until it reached 80 per cent in those who had worked in the rice fields for 20 years or more. Approximately 66 per cent of these infections were due to *L. balariae*. A similar situation exists in Australia among cane field workers. Derrick and his co-workers⁶⁴ found that 57 per cent of the known cases of leptospirosis in North Queensland Australia occurred among cane field workers and were attributed primarily to infection with *L. australis* although 10 other leptospiral serotypes were incriminated. No cases of leptospirosis have been encountered among cane and rice field workers in the United States. Swineherd's disease due to *L. pomona* and *L. mitis* infections is common in certain parts of Europe among butchers abattoir workers and others who have had contact with swine.⁶⁵⁻⁶⁷ Kirschner⁶⁸ reported that 67 per cent of 130 cases in New Zealand occurred among dairy farmers and the remainder in pig breeders or

TABLE 3
REPORTED OUTBREAKS OF LEPTOSPIROSIS IN THE UNITED STATES*

Date	Location	No. of cases	Age range	Probable source	Serologic type
August 1940	Wrens Ga	35	Avg 16	Cattle	<i>L. pomona</i> †
July and August 1942 to 1944	Fort Bragg N C	40/yr	Young adults (soldiers)	?	<i>L. autumnalis</i>
August 1947	Jackson Hole Wyo	24	15 to 19	Dogs	<i>L. canicola</i> †
August 1947	Calvary Ga	10	13 to 24	Dead mule in creek	?
August 1949	Swainsboro Ga	12	Young adults	Cattle	<i>L. pomona</i> †
July 1950	Geneva Ala	50	Adolescent and young adults	Swine	<i>L. pomona</i> †
July 1952	Columbus Ga	26	5 to 20 and 2 adults	Swine dogs cattle	<i>L. canicola</i>

With the possible exception of the outbreak at Fort Bragg all outbreaks were attributed to swimming in contaminated water

† Serologic evidence only

Outbreaks and Sporadic Cases of Leptospirosis in Humans in the United States

It is not surprising that large outbreaks may occur after group swimming in contaminated water. Several such outbreaks have been reported in the United States as shown in TABLE 3. The first of these occurred in August 1940 near Wrens Ga.^{45, 46} and involved 35 patients who had been swimming in Brushy Creek. Clinical signs included fever, nausea, vomiting, headache, some diarrhea, and a rash on the pretibial areas of the legs. It was found that approximately 100 yards upstream from the swimming area offal from slaughtered cattle had been dumped by the local butchers. This unusual disease was given the name of Brushy Creek or pretibial fever. A similar outbreak of pretibial fever occurred among troops at Fort Bragg in August 1942 and again in the summers of 1943 and 1944.^{1, 4, 49}

All patients involved in the outbreaks were quartered in the same area of the post. Many but not all had been swimming in the nearby ponds, so the source of their infection remains obscure. Not until 1950 was Fort Bragg fever proved to be a leptospiral infection caused by a member of the *L. autumnalis* group.²¹ After this discovery, McCroan reinvestigated the Brushy Creek outbreak and 2 other similar undiagnosed epidemics that had occurred in Georgia: one* in the late summer of 1947 near Calvary⁴⁶ and the other during the same season in 1949 on the Ohoopsee River near Swainsboro.⁴⁶ Antibodies to *L. pomona* were demonstrated in serum samples collected at the time of the outbreak and during the subsequent investigation in 1953 from several patients involved in the Brushy Creek and Ohoopsee River outbreaks.

A follow-up investigation in 1953 of another epidemic of an unknown disease that had occurred in August 1942 in two small communities in Wyoming and Idaho revealed serologic evidence of leptospiral etiology.⁵⁰ Here again all

Investigated by Elmer Hill and John F. McCroan of the Georgia State Health Department, Atlanta, Ga.

persons affected had been swimming in the same pool located at Jackson Hole, Wyo. to which dogs and other animals had access.

In late July 1950 an outbreak occurred in Geneva, Ala.⁴¹ involving about 50 persons who became ill with an influenzalike disease after swimming in a creek in which dead swine had been seen floating above the swimming hole. Serologic evidence indicated that *L. pomona* was the cause.

When in late July 1952 an outbreak of illness occurred in a small community near Columbus, Ga. involving 26 persons who had been swimming in a dammed up creek, leptospirosis was suspected immediately.⁴² The subsequent investigation of this outbreak, including laboratory studies⁴³ presented an excellent picture of many of the principles involved in leptospiral epidemiology. *L. canicola* infections, thought for many years to be limited to dogs and man, were demonstrated in swine, cattle, and dogs that had access to the creek. All persons who became ill had been swimming in the creek. The data showed a significant correlation between attack rates and the frequency and extent of exposure to the water. Attack rates were consistently higher among those giving a history of diving and immersing their heads.

With the possible exception of the Fort Bragg outbreaks, each of these water-borne leptospiral epidemics followed a similar pattern. They occurred in the late summer during drought periods; there was presumed contamination of a stagnant pond or slow-moving creek by urine from infected animals and transmission to humans by immersion in the contaminated water. Such water-borne outbreaks following swimming or accidental immersion have been reported in many other areas of the world.⁴⁴⁻⁴⁶

Of even greater importance in the epidemiology of leptospiral infections are the opportunities for exposure encountered by workers in certain occupations, particularly abattoir workers, cane and rice field workers, dairy farmers, animal husbandrymen, veterinarians, sewer workers, and trench diggers. For example, in Italy, Spain, the Far East, and Pacific regions where the rice fields are flooded and worked by barefoot laborers, leptospirosis is an important health and economic problem.⁴⁷⁻⁵¹ The fields usually abound with mice and rats. Babudieri⁴⁷ found in a survey of 509 apparently healthy workers in Italy that 20 per cent had leptospiral antibodies. The percentage of infection increased in different groups as the length of service increased, until it reached 80 per cent in those who had worked in the rice fields for 20 years or more. Approximately 66 per cent of these infections were due to *L. bataviae*. A similar situation exists in Australia among cane field workers. Derrick and his co-workers⁴⁸ found that 57 per cent of the known cases of leptospirosis in North Queensland, Australia, occurred among cane field workers and were attributed primarily to infection with *L. australis*.⁴⁹ Although 10 other leptospiral serotypes were incriminated, no cases of leptospirosis have been encountered among cane and rice field workers in the United States. Swineherd's disease, due to *L. pomona* and *L. mitis* infections, is common in certain parts of Europe among butchers, abattoir workers, and others who have had contact with swine.⁵²⁻⁵⁷ Kirschner⁵ reported that 67 per cent of 130 cases in New Zealand occurred among dairy farmers and the remainder in pig breeders or




FIGURE 2 The geographical distribution of cases of Weil's disease in the period from 1905 to 1941⁷⁰

abattoir workers The apparently lower incidence in the United States of human infection due to occupational exposure may be attributed to higher living standards and to mechanization of industrial and agricultural operations However the recent comment of Kalz⁶⁹ that the number of recognized and reported cases particularly in the United States and in Australia appears to be related directly to the location of interested workers and to the availability of laboratory facilities warrants some thought In recent years there has been an increase in reported cases in the United States The geographic distribution of known cases was summarized by Larson in 1941⁷⁰ and again in 1948 by Molner Meyer and Raskin⁷¹ as shown in FIGURES 2 and 3 In the first period summarized which covered 36 years there were 93 cases in 14 states Seven years later the total number of cases had increased to 299 in 24 states In contrast a summary of cases detected through serologic tests in the CDC Laboratories during a 4-year period (1953 to 1956) plus those reported to the CDC by state health departments or to the National Office of Vital Statistics Washington D C during this time revealed a total of 392 in 39 states In addition 96 cases occurred in outbreaks reported between 1951 and 1956 The distribution of these cases is shown in FIGURE 4

Sporadic cases⁷²⁻⁷⁵ are reported in the United States in veterinarians live stock men abattoir workers sewer workers dog owners and others Haunz and Cardy⁷⁶ recently reported 9 cases in a single family that were attributed to

from bovine leptospirosis were over 112 million dollars or 25 million dollars greater than losses from bovine brucellosis.

In large part the prevention and control of leptospirosis are aspects of the prevention and control of other diseases. Certainly the application of strict sanitary practices will play an important role. Protection of drinking water supplies for domestic animals from contamination, adequate drainage of wet muddy farm areas, and rat control are particularly valuable measures. Several




FIGURE 6. The age distribution of 50 cases of human leptospirosis in the United States from August 1954 to August 1957.




FIGURE 7. The registered occurrence of bovine leptospirosis in the United States during 1951. * Reproduced by permission of *Fl. Proceedings of the United States Live Stock Sanitary Association*.




FIGURE 8 The reported occurrence of bovine leptospirosis during 1956

leptospiral vaccines are available for use^{29, 31} in cattle and while results have been encouraging further evaluation of their effectiveness is needed

In nature leptospirosis has a wide host spectrum that yields a variety of serotypes. As yet *L. pomona* is the only serotype used in the preparation of vaccines for cattle. While most infections in cattle and swine are caused by this serotype *I. canicola* was isolated recently from a sick calf in Alabama⁸ and serologic findings suggest the presence of *I. sejiro*⁶ and *I. grippotyphosa* infections²⁶. Thus the use of a polyvalent vaccine should be considered. The same consideration should be given to canine vaccines which at present are prepared only from *I. canicola* and *I. icterohemorrhagiae* strains.

Prevention and control of the disease in humans particularly in those exposed to it in their occupations depends largely upon improvement of working conditions. Results of vaccination studies have been excellent in the Italian and Spanish rice field workers⁸ among whom the incidence of infection is high. The feasibility of similar immunization of abattoir workers, veterinarians and others at risk should be studied.

Discussion

The leptospiroses spread to humans under favorable environmental circumstances following contact with the tissues or urine of infected animals or contact with water or wet soil contaminated by animal carrier. Higher living standards greatly influence the lower incidence in the general population.

in the United States and other countries that report the disease in animals but have few human cases. In the comparatively few occupations in which the problem is encountered, better sanitary practices, rat control, and protective clothing can play an important role in reducing the hazard. Outbreaks of the disease related to recreational accidents may be expected to occur sporadically.

The detection of leptospiral serotypes hitherto unrecognized in the United States emphasizes the need for a more concerted effort by medical and veterinary laboratories to search for them. As laboratory techniques for the isolation of leptospires are improved and applied, more serotypes will be obtained from humans and from domestic animals. Further study of the new wild mammal hosts to determine the incidence of infection in other parts of the country is indicated. In addition, little is known of the duration of the carrier state in these hosts. Application of the recently developed bladder tap technique²⁴ to secure urine for direct culture would provide some of the information needed. Future research in this direction will throw more light upon the role of these wild mammal hosts in the over-all picture of leptospiral epidemiology and epizootiology.

There has been much speculation as to the reasons for the apparent increase in incidence of leptospiral infections. It may be accounted for in part by a greater awareness of the disease. Current knowledge strongly suggests the existence of an epizootic or panzootic of the leptospires that may flow and ebb for some decades. Until we have further knowledge of the conditions that favor the prevalence of leptospirosis, we have no better explanation than Sydenham's theory of the *genius epidemicus* that contagious diseases are influenced by cosmic or atmospheric conditions.

Summary

Leptospiral infections are known to occur in wild and domestic animals throughout the civilized world. More than forty years of extensive research have revealed a characteristic epidemiological pattern. All leptospires known to infect animals are potentially pathogenic for humans. While some serotypes appear to exhibit a certain host preference, all of the more common types have been found to infect more than one animal species. During the past two years, investigations have detected, in a variety of wild mammal hosts, the presence of 5 leptospiral serotypes hitherto unrecognized in the United States.

Of primary importance in the chain of transmission are the animal carriers of leptospires. After acute, mild, or even inapparent infections, the organisms localize in the kidney and are shed in the urine for varying periods. These carriers serve as foci of infection for other animals and man.

In the United States, reported outbreaks of leptospirosis have been associated primarily with swimming in stagnant pools or creeks presumably contaminated by carrier animals. Sporadic cases that occur are more frequently associated with occupational exposures among veterinarians, abattoir workers, sewer laborers, dog owners, and others exposed to a contaminated environment.

Information concerning 94 sporadic human cases in the United States is presented. These outbreaks and sporadic cases are discussed and compared with the epidemiology of leptospiral infections in other countries. These facts emphasize the need for further research to clarify the significance of the new sylvatic hosts in the over all pattern of leptospirosis.

References

- 1 REINHARD K R 1954 Public health aspects of animal leptospirosis 91st Ann Meeting Proc Am Vet Med Assoc 455-460
- 2 REINHARD K R 1955 Leptospirosis Mich State Univ Cent Symp Rept 12-19
- 3 MORSE F V 1955 The economic significance of leptospirosis in domestic animals 92nd Ann Meeting Proc Am Vet Med Assoc 162-167
- 4 VAN DER HOEDEN J 1953 Leptospirosis of cattle and goats in Israel Proc 15th Intern Vet Congr 1 1-3
- 5 BERNSTEIN T & J A BAKER 1954 Leptospirosis in swine caused by *Leptospira pomona* J Infectious Diseases 94 53-64
- 6 UNITED STATES DEPARTMENT OF AGRICULTURE AGRICULTURAL RESEARCH SERVICE 1954 Losses in agriculture A preliminary appraisal for review 129
- 7 VAN DER HOEDEN J 1955 The epidemiology and epizootiology of leptospirosis in Israel J Trop Med Hyg 58 202-204
- 8 GSELL O R 1953 Epidemiology of the leptospires Symposium on the leptospires Med Sci Publ Army Med Serv Grad School No 1 34-55
- 9 BABUDIERI B 1953 Epidemiology of leptospirosis in Italian rice fields Advance in the control of zoonoses World Health Organization Tech Rept Ser Monograph 19 117-126 Geneva Switzerland
- 10 MCKEEVER S G W GORMAN J F CHAPMAN M M GALTON & D K POWERS 1957 Incidence of leptospirosis in wild mammals from southwestern Georgia with a report of new hosts for six serotypes of *Leptospirae* To be published
- 11 ALEXANDER A D P W WETMORE L B EVANS H JEFFERIES & C A GLEISER 1955 Classification of leptospiral isolates from Malaya Thailand and North Borneo Am J Trop Med Hyg 4 492-506
- 12 YAGER R H & W S GOCHENOUR JR 1952 Leptospirosis in North America Am J Trop Med Hyg 1 457-461
- 13 VAN DER HOEDEN J 1955 Epizootiology of leptospirosis (*canicola*) in the bovine and other species in Israel J Am Vet Med Assoc 126 207-210
- 14 YAGER R H 1953 Epidemiology of the leptospirosis Bull N Y Acad Med 29 650-651
- 15 WOLFF J W & J C BROOM 1954 The genus *Leptospira* Noguchi 1917 Problems of classification and a suggested system based on antigenic analysis Documenta Med Geograph et Trop 6 78-95
- 16 SMITH D J W H F BROWN J I TONGE C N SINNAMON V M MACDONALD C J ROSS & R L DOHERTY 1954 The serological classification of 89 strains of *Leptospirae* from North Queensland including five serotypes new to Australia Australasian Ann Med 3 98-105
- 17 SMITH D J W & H E BROWN 1955 Two additional serotypes of *Leptospirae* from North Queensland Australasian Ann Med 4 287-290
- 18 VAN IJSEL P J L SZPAJAHENDLER & M VAN RIEL 1956 Etude clinique bactériologique et épidémiologique d'un nouveau foyer de Leptospirose au Congo Belge Bull soc pathol exotique 49 118-143
- 19 BABUDIERI B 1956 *Leptospira mini* ein neuer Serotyp pathogener Leptospiren Z Hyg 143 121-176
- 20 GALTON M M D K POWERS S MCKEEVER & G W GORMAN 1957 The identification of two leptospiral serotypes new to the United States Public Health Repts U S 72 431-435
- 21 GOCHENOUR W S JR J E SMADEL E B JACKSON I B EVANS & R H YAGER 1957 Leptospiral etiology of Fort Bragg Fever Public Health Repts U S 67 811-814
- 22 YAGER R H W S GOCHENOUR JR A D ALEXANDER & P W WETMORE 1953 Natural occurrence of *Leptospira ballum* in rural mice and in an opossum Proc Soc Exptl Biol Med 84 589-590
- 23 BROWN J Z 1957 Leptospirosis in feral rodents in southwestern Georgia In preparation

- 24 GOCHENOUR W. S., JR., R. H. YAGER P. W. WETMORE, L. B. EVANS R. J. BYRNE A. ALEXANDER & J. A. HIGHTOWER 1951 *Leptospirosis in Puerto Rico and the United States* Federal Proc. 10 408
- 25 SPAIN R. S. & G. T. HOWARD 1952 Leptospirosis due to *Leptospira interrogans* J. Am. Med. Assoc. 150 1010
- 26 GALTON M. M. J. A. ACREE A. LEWIS & E. C. PRATHER 1956 Leptospirosis in domestic animals in Florida with reference to cattle J. Am. Vet. Med. Assoc. 128 8-91
- 27 YAGER R. H. 1953 Leptospirosis in the United States today Symposium on the leptospiroses. Med. Sci. Publ. Army Med. Serv. Grad. School No. 1 2-1
- 28 MCKEEVER S. G. W. GORMAN M. M. GALTON & A. D. HALL The raccoon *Procyon lotor* as a natural host of *Leptospira interrogans* To be published.
- 29 ROTH E. E. & B. KRIEDEM 1952 The natural occurrence of *L. pomona* in an opossum To be published.
- 30 STÖNNER H. G. & D. MACLEAN 1955 Leptospirosis (bub.) contracted from Swiss mice A. M. A. Arch. Internal Med. In press.
- 31 REILLY J. R. 1954 The raccoon as a wildlife reservoir of *Leptospira interrogans* N. Y. Fish and Game J. 1 279
- 32 BYRNE, R. J. 1955 Canine leptospirosis and public health. Public Health Repts. L. S. 70-1229
- 33 SCHLO-BERGER H. & H. LANGBEIN 1955 Übertragung von *Leptospira interrogans* durch *Oryzias latipes* moulting Z. Immunitätsforsch. 109 366-370
- 34 BURGDORFER W. & E. G. PICKENS 1954 A technique employing eggs of chicken eggs for the injection of arthropod ticks with *Coccidia* & *Bacterium tularensis* *Leptospira interrogans* and western equine encephalitis virus J. Infectious Diseases. 94 84-89
- 35 BURGDORFER W. 1956 The possible role of ticks as vectors of *Leptospira* I Transmission of *Leptospira pomona* by the argasid tick *Oryzia* *devoid* *intracutanea* and the persistence of this organism in its tissues. Entom. Parasitol. 5 51-59
- 36 BURGDORFER W. 1955 Artificial feeding of ixodid ticks for studies in the transmission of disease agents. J. Infectious Diseases. 100 21-214
- 37 KREPKOGORSKAYA T. A. & M. M. REMENKOVA 1955 The isolation of strains of leptospires from the tick *Dermacentor marginatus* S. from cattle Zhur. Mikrob. i. Epidemiol. Immunol. 28 10-12
- 38 LITTLE, R. B. & J. A. BAKER 1950 Leptospirosis in cattle J. Am. Vet. Med. Assoc. 116 105-111
- 39 KIESCHNER L. & T. MAGUIRE. 1955 Anti-leptospirosis effect of milk New Zealand J. Med. J. 54 560-564
- 40 KIESCHNER, L. 1953 Symposium on the leptospiroses. Discussion Med. Sci. Publ. Army Med. Serv. Grad. School No. 1 0
- 41 CHANG S. L. M. BUCKINGHAM & M. P. TAYLOR 1948 Studies on *Leptospira interrogans* IV Survival in water and sewage destruction in water by halogen compounds synthetic detergents and heat. J. Infectious Diseases. 82 26-266.
- 42 SMITH D. J. W. & H. R. M. SELF 1955 Observations on the survival of *Leptospira interrogans* in soil and water J. Hyg. 53 436-444
- 43 OKAZAKI W. & L. M. RINGEN 1955 Some effects of various environmental conditions on the survival of *Leptospira pomona* Am. J. Vet. Research. 15 219-223
- 44 GILLIPIE R. W. H. S. G. KENNY L. M. RINGEN & F. K. BRACKEN 1955 Studies on bovine leptospirosis III Isolation of *Leptospira pomona* from surface waters. Am. J. Vet. Research. 15 6-40
- 45 BOWDWIN C. D. 1947 A new disease entity (?) J. Med. Assoc. Georgia 31 43-438
- 46 McCRAIG J. E. 1957 Division of Epidemiology Georgia Dept. Public Health Personal communication
- 47 DANIELS W. B. & H. A. GREENAN 1943 Pretilial fever an obscure disease J. Am. Med. Assoc. 122 361-365
- 48 TOPPING N. H. C. B. PHILIP & J. R. PAUL 1941 Report of the commission for the study of an unidentified disease at Fort Bragg N. C. Preliminary report submitted to the Bureau General, U. S. Army Washington D. C.
- 49 TAYLOR H. 1941 Studies on a virus from a patient with Fort Bragg fever (pre-tilial fever) J. Clin. Invest. 20 3-79
- 50 COCKBURN T. A. J. D. VAVRA, S. S. SPENCER J. R. DANN L. J. PETERSON & K. R. REINHARD 1954 Human leptospirosis associated with a swimming pool, diagnosed after eleven years. Am. J. Hyg. 60 1-

Information concerning 94 sporadic human cases in the United States is presented. These outbreaks and sporadic cases are discussed and compared with the epidemiology of leptospiral infections in other countries. These facts emphasize the need for further research to clarify the significance of the new sylvatic hosts in the over all pattern of leptospirosis.

References

- 1 REINHARD K R 1954 Public health aspects of animal leptospirosis 91st Ann Meeting Proc Am Vet Med Assoc 455-460
- 2 REINHARD K R 1955 Leptospirosis Mich State Univ Cent Symp Rept 12 19
- 3 MORSE E V 1955 The economic significance of leptospirosis in domestic animals 92nd Ann Meeting Proc Am Vet Med Assoc 167-167
- 4 VAN DER HOEDEN J 1953 Leptospirosis of cattle and goats in Israel Proc 15th Intern Vet Congr 1 1-3
- 5 BURNSTEIN T & J A BAKER 1954 Leptospirosis in swine caused by *Leptospira pomona* J Infectious Diseases 94 53-64
- 6 UNITED STATES DEPARTMENT OF AGRICULTURE AGRICULTURAL RESEARCH SERVICE 1954 Losses in agriculture A preliminary appraisal for review 129
- 7 VAN DER HOEDEN J 1955 The epidemiology and epizootiology of leptospirosis in Israel J Trop Med Hyg 58 207-204
- 8 GSELL O R 1953 Epidemiology of the leptospires Symposium on the leptospires Med Sci Publ Army Med Serv Grad School No 1 34-55
- 9 BABUDIERI B 1953 Epidemiology of leptospirosis in Italian rice field Advance in the control of zoonoses World Health Organization Tech Rept Ser Monograph 19 117-126 Geneva Switzerland
- 10 MCKEEVER S G W GORMAN J F CHAPMAN M M GALTON & D K POWERS 1957 Incidence of leptospirosis in wild mammals from southwestern Georgia with a report of new hosts for six serotypes of *Leptospirae* To be published
- 11 ALEXANDER A D P W WETMORE L B EVANS H JEFFERIES & C A GLEISER 1955 Classification of leptospiral isolates from Malaya Thailand and North Borneo Am J Trop Med Hyg 4 492-506
- 12 YAGER R H & W S COCHENOUR JR 1952 Leptospirosis in North America Am J Trop Med Hyg 1 457-461
- 13 VAN DER HOEDEN J 1955 Epizootiology of leptospirosis (*canicola*) in the bovine and other species in Israel J Am Vet Med Assoc 126 207-210
- 14 YAGER R H 1953 Epidemiology of the leptospirosis Bull N Y Acad Med 29 650-651
- 15 WOLFF J W & J C BROOM 1954 The genus *Leptospira* Noguchi 1917 Problems of classification and a suggested system based on antigenic analysis Documenta Med Geograph et Trop 6 78-95
- 16 SMITH D J W H F BROWN J I TONGE C N SINNAMON V M MACDONALD C J ROSS & R L DOHERTY 1954 The serological classification of 89 strains of *Leptospirae* from North Queensland including five serotypes new to Australia Australasian Ann Med 3 98-105
- 17 SMITH D J W & H F BROWN 1955 Two additional serotypes of *Leptospirae* from North Queensland Australasian Ann Med 4 287-290
- 18 VAN RIEL P J L SZPAJARENDLER & M VAN PIEL 1956 Etude clinique bactériologique et épidémiologique d'un nouveau foyer de Leptospirose au Congo Belge Bull soc pathol exotique 49 118-143
- 19 BABUDIERI B 1956 *Leptospira mini* ein neuer Serotyp pathogener Leptospiren Z Hyg 143 121-126
- 20 GALTON M M D K POWERS S MCKEEVER & G W GORMAN 1957 The identification of two leptospiral serotypes new to the United States Public Health Repts U S 72 431-435
- 21 COCHENOUR W S JR J F SMADEL E B JACKSON I B EVANS & R H YAGER 1957 Leptospiral etiology of Fort Bragg Fever Public Health Repts U S 67 811 814
- 22 YAGER R H W S COCHENOUR JR A D ALEXANDER & P W WETMORE 1953 Natural occurrence of *Leptospira ballum* in rural mice and in an opossum Proc Soc Exptl Biol Med 84 589 590
- 23 BROWN R Z 1954 Leptospirosis in feral rodents in southwestern Georgia In preparation

24. GOCHENOUR W S JR, R H YAGER, P W WFTMORE, L B EVANS, I J BYRNE, A ALEXANDER & J A HIGHTOWER. 1951. Indonesian Weil's disease in Puerto Rico and the United States. *Federation Proc* 10: 409.
25. SPAIN R S & G T HOWARD. 1952. Leptospirosis due to *Leptospira gripp typhi* s.s. *J Am Med Assoc* 150: 1010.
26. GALTON M M, J A ACREE, A LEWIS & F C PRATHER. 1956. Leptospirosis in domestic animals in Florida with reference to cattle. *J Am Vet Med Assoc* 128: 87-91.
27. YAGER R H. 1953. Leptospirosis in the United States today. Symposium on the leptospiroses. *Med Sci Publ Army Med Serv Grad School* No 1: 221.
28. MCKEEVER S G W, GORMAN M M, GALTON & A D HALL. The raccoon *Pyrocyon f. l.* as a natural host of *Leptospira autumnalis*. To be published.
29. ROTH F I & B KNIERDEM. 1957. The natural occurrence of *L. p. mona* in an opossum. To be published.
30. STOENNER H G & D MACLEAN. 1957. Leptospirosis (*ballum*) contracted from Swiss mice. *A M J Arch Internal Med*. In press.
31. RYLLY J R. 1954. The raccoon as a wild life reservoir of *Leptospira canicola*. *N Y Fish and Game J* 1: 220.
32. BYRNE R J. 1955. Canine leptospirosis and public health. *Public Health Lepts U S* 70: 1229.
33. SCHLOS BERGER H & H LANGBEIN. 1952. Übertragung von *Leptospira sclero haemorrhagiae* durch *Ornithodoros morbita*. *Z Immunitätsforsch* 109: 366-370.
34. BURGDORFER W & E G PICKENS. 1954. A technique employing embryonated chicken eggs for the infection of argasid ticks with *Cyella burnetii* *Bacterium l. larente* *Leptospira sclero haemorrhagiae* and western equine encephalitis virus. *J Infectious Diseases* 94: 84-89.
35. BURGDORFER W. 1956. The possible role of ticks as vectors of *Leptospirae*. I. Transmission of *Leptospira pomona* by the argasid tick *Ornithodoros turicata* and the persistence of this organism in its tissues. *Exptl Parasitol* 5: 571-579.
36. BURGDORFER W. 1957. Artificial feeding of ixodid ticks for studies in the transmission of disease agents. *J Infectious Diseases* 100: 212-214.
37. KREPKOGORSKAIA T A & M M REMENKOVA. 1957. The isolation of strains of leptospires from the tick *Dermacentor marginatus* S from cattle. *Zhur Mikrobiol Epidemiol Immunob* 28: 251-257.
38. LITTLE R B & J A BAKER. 1950. Leptospirosis in cattle. *J Am Vet Med Assoc* 116: 103-111.
39. KIRSCHNER L & T MAGUIRE. 1955. Antileptospiral effect of milk. *New Zealand Med J* 64: 560-564.
40. KIRSCHNER L. 1953. Symposium on the leptospiroses. Discussion. *Med Sci Publ Army Med Serv Grad School* No 1: 10.
41. CHANG S L, M BUCKINGHAM & M P TAYLOR. 1948. Studies on *Leptospira sclero haemorrhagiae*. IV. Survival in water and sewage, destruction in water by halogen compounds, synthetic detergents and heat. *J Infectious Diseases* 62: 256-266.
42. SMITH D J W & H R M SEIF. 1955. Observations on the survival of *Leptospira australis* A in soil and water. *J Hyg* 53: 436-444.
43. OKAZAKI W & L M RINGEN. 1957. Some effects of various environmental conditions on the survival of *Leptospira pomona*. *Am J Vet Research* 18: 219-223.
44. GILLESPIE R W, H S G KENY, L M RINGEN & F K BRACKEN. 1957. Studies on bovine leptospirosis. III. Isolation of *Leptospira pomona* from surface waters. *Am J Vet Research* 18: 76-80.
45. BOWDOIN C D. 1942. A new disease entity (?). *J Med Assoc Georgia* 31: 437-438.
46. McCROAN J F. 1957. Division of Epidemiology. Georgia Dept. Public Health. Personal communication.
47. DANIELS W B & H A GRENNAN. 1943. Pre-tibial fever, an obscure disease. *J Am Med Assoc* 122: 361-365.
48. TOPPING N H, C B PHILIP & J R PAUL. 1942. Report of the commission for the study of an unidentified disease at Fort Bragg, N C. Preliminary report submitted to the Surgeon General, U S Army. Washington, D C.
49. TAYLOR H. 1944. Studies on a virus from a patient with Fort Bragg fever (pre-tibial fever). *J Clin Invest* 26: 287-297.
50. COCKBURN T A, J D VAVRA, S S SPENCER, J R DANN, L J PETERSON & K R REITHARD. 1954. Human leptospirosis associated with a swimming pool, diagnosed after eleven years. *Am J Hyg* 60: 1-7.

- 51 SCHAEFFER M 1951 Leptospirosis meningitis Investigation of a water borne epidemic due to *L. pomona* J Clin Invest 30 610
- 52 WILLIAMS H R W J MURPHY J F MCCROAN L F STARR & M K WARD 1956 An epidemic of canicola fever in man with the demonstration of *Leptospira canicola* infection in dog swine and cattle I Clinical and epidemiological studies Am J Hyg 64 46-58
- 53 WARD M K M B MCDANIEL H W TATUM L F STARR & H R WILLIAMS 1956 An epidemic of canicola fever in man with the demonstration of *Leptospira canicola* infection in dogs swine and cattle II Laboratory studies Am J Hyg 64 59-69
- 54 MISAO T S HIROYOSHI K KATSUTA Y NISHIHARA Y KOBAYASHI K KAWASHIMA & M ASO 1955 Canicola fever in Japan Am J Hyg 63 294-307
- 55 WOLFF J W & A C RUY 1953 Change in the frequency of Weil's disease in Amsterdam during the last eighteen years Documenta Med Geograph et Tr p 5 247-252
- 56 BROOM J C 1951 Leptospirosis in England and Wales Brit Med J 2 689-713
- 57 GAULD R I W L CROUCH A I KAMINSKY R L HULLINGHORST W S COCHENOUR JR & R H YAGER 1952 Leptospirosis meningitis report of outbreak among American troops on Okinawa J Am Med Assoc 149 228-231
- 58 VARFOLONEVA A A 1957 Epidemiology and aetiology of an outbreak of leptospirosis Zhur Mikrobiol Epidemiol Immunobiol 28 38-43
- 59 BLAGOVESHCHENSKAYA N M 1957 On the epidemiology of anicteric leptospirosis Zhur Mikrobiol Epidemiol Immunobiol 28 240-244
- 60 IANOVICH T D A G BLIZNICHENKO L V ZABURINA S A MISTIBOVSKII A I BERKOVICH & I P DUSHEVY 1957 Leptospirosis of the canicola type in one of the districts of Rostov on the Don Zhur Mikrobiol Epidemiol Immunobiol 28 259-264
- 61 DURICH J & A PUMAROLA 1955 La endemia de leptospirosis en la provincia de Valencia Med españ 34 84-90
- 62 ALTAVA V M BARRERA I VILLALONGA I GIL C MARIN & B BABUDIERI 1955 Vacunacion como profilaxi en la leptospirosis de los arrozales de Castellon Rev sanidad hig pub 29 1-4
- 63 MISAO T S HIROYOSHI K KATSUTA Y NISHIHARA Y KOBAYASHI K KAWASHIMA & M ASO 1954 Epidemic of canicola fever in Kaizan and Ito villages Fukuoka Prefecture Japan J Med Sci Biol 7 405-415
- 64 DERRICK F H D CORDON C J ROSS R L DOHERTY C N SINNAMON V M MACDONALD & J M KENNEDY 1954 Epidemiological observations on leptospirosis in North Queensland Australasian Ann Med 3 85-91
- 65 GSELL O R 1944 Etiologie der Schweinehuterkrankheit Bull Schweiz Akad med Wiss 1 67
- 66 GSELL O R & E WEISMANN 1948 Leptospirosis Typ *mitis* Schweiz med Wochschr 78 503
- 67 KMETZ F 1955 Leptospirosenherde in der Slowakei Zentr Bakteriol Parasitenk 163 464-476
- 68 KIRSCHNER I 1954 Recent studies on leptospirosis in New Zealand infection with a new type (*Leptospira mitis*) Johnson (Syn *Lept hyos*) in man and animals New Zealand Med J 53 119-128
- 69 KALZ G 1957 The human leptospiroses Am J Med Sci 233 320-333
- 70 FARSON C L 1941 Weil's Disease A report of 51 cases occurring in Puerto Rico and the United States Public Health Repts U S 56 1650-1656
- 71 MOLNER J G K F MEYER & H A RASKIN 1948 Leptospiral infections A survey J Am Med Assoc 136 814-819
- 72 CROSSMAN M M LEVIN & R O'NEILL 1954 *Leptospira pomona* meningitis A report of two cases Calif Med 82 192-193
- 73 KROUSE T B & M M SIGEL 1952 Leptospirosis meningitis Report of a case resembling swineherd's disease due to *Leptospira pomona* in the United States Ind Med Surg 21 121-123
- 74 DYOSKIN S & F W HOOK 1956 Canicola fever with meningitis A report of three cases and an epidemiological study at Fort McClellan Ala A M A Arch Internal Med 97 793-797
- 75 BEESON I B & D D HANLEY 1952 Leptospirosis meningitis Arch Internal Med 89 575-583
- 76 HAUZ I A & J D CARDY 1952 Canicola fever Report of nine cases in one family with abstract of the world literature A M A Arch Internal Med 89 98-993

- 77 HIR CHIBERG N 1956 State Laboratory of Hygiene Raleigh N C Personal communication
- 78 YORK C J 1951 Aspects of control in bovine leptospirosis Proc 55th Ann Meeting U S Livestock San Assoc 295-300 Kansas City Kans
- 79 YORK C J & J A BAKER 1953 Vaccination for bovine leptospirosis Am J Vet Research 14 5-8
- 80 YORK C J & V JOHNSTON & V B ROBINSON 1955 The use of vaccine in the control of leptospirosis in cattle and swine 92nd Ann Meeting Proc Am Vet Med Assoc 169-171
- 81 BROWN A L A A CRAWFORD & S I SHEIDY 1954 Improved *Leptospira bacterin* for the control of bovine leptospirosis 58th Ann Meeting Proc U S Livestock San Assoc 227-235
- 82 TURNER I W A G WIGGINS & C S FORTY 1957 *Leptospira canicola* infection in cattle Am J Vet Research To be published
- 83 MENGE R W M M GALTON & A D HALL 1958 Diagnosis of leptospirosis from urine specimens by direct culture following bladder tapping J Am Vet Med Assoc 132 53-60
- 84 INADA R Y IDO R HOMI R KANAKO & H ITO 1916 The etiology mode of infection and specific therapy of Weil's disease (*Spirochaeta icterohaemorrhagica*) J Exptl Med 23 377
- 85 FIELD H I & K C SELLERS 1951 *Leptospira sclerosa* infection in the calf Vet Record 62 311-313
- 86 VAN THIEL P H 1948 The Leptospires Universitaire Pers Leiden Netherlands
- 87 FIELD H I & K C SELLERS 1951 *Leptospira sclerosa* infection in piglets Vet Record 63 78-80
- 88 MASON N 1937 Leptospirosis occurring naturally in guinea pigs Lancet 1 564-565
- 89 WINSTON C L JR R TRUB W S GOCHENOUR JR J L SMADEL & W E LANCASTER 1955 Leptospirosis of man and animals in urban rural and jungle areas of southeast Asia Am J Trop Med Hyg 4 29-40
- 90 NOGUCHI H 1917 *Spirochaeta sclerosa* in American wild rats and its relation to the Japanese and European strains J Exptl Med 25 755-763
- 91 YAGER R H & W S GOCHENOUR JR 1952 Leptospirosis in America Am J Trop Med Hyg 1 457-461
- 92 ESVELD H & W A COLLIER 1938 Leptospirosis bij katten te Batavia Meded el Dist Volk gezondheid Ned Indie 27 250
- 93 KRASILNIKOV A P 1957 Natural reservoirs of infection of leptospirosis in the Belo-Russian S S R Zhur Mikrobiol Epidemiol Immunobiol 28 50-54
- 94 WOLFF J W 1954 The Laboratory Diagnosis of Leptospirosis Thomas Springfield Ill
- 95 KMETZ E I PLESKO & E CHYLO 1956 Weitere Ergebnisse der Leptospirosen forschung in der Slowakei Zentr Bakteriell Parasitenk 167 243-253
- 96 BJORG PETERSEN C 1944 *Leptospira ballum* a new serological *Leptospira* type? Acta Pathol Microbiol Scand 21 504-509
- 97 FLETCHER W 1928 Recent work on leptospirosis (tsutsugamushi disease and tropical typhus) in the Federated Malay States Trans Roy Soc Trop Med Hyg 21 265-266
- 98 POPOVA F M & N I AMOSKOVA 1957 Reservoirs of leptospirosis infection in the northwestern regions of the U S S R Results of an investigation on leptospirosis infections of murine rodents Zhur Mikrobiol Epidemiol Immunobiol 28 44-49
- 99 WANNAN J S 1955 Leptospirosis serotypes in Australia Australian Vet J 297-301
- 100 LEMLEY G T 1937 Leptospirosis in Queensland A serological investigation leading to the discovery of distinct serological groups of leptospires causing leptospirosis as it occurs in Northern Queensland with some related observations Med J Australia 24 654-664
- 101 ALEXANDER A D C A GLEISER P MALNATI & H LODER 1957 Observations of the prevalence of leptospirosis in canine populations of the United States Am J Hyg 65 43-56
- 102 CLAYTON C E B E H DERRICK & R CILENTO 1937 The presence of leptospirosis of a mild type (seven day fever) in Queensland Med J Australia 1 647
- 103 ROBERTS S J C J YORK & J W ROBINSON 1952 An outbreak of leptospirosis in horses on eight small farms J Am Vet Med Assoc 121 237

have infected salivary glands been found without concurrent infection in the central nervous system, which indicates an inherent inability of this species to transmit the disease as a symptomless carrier. The transmitting potential of foxes however is great as shown by the fact that daily testing of experi-




FIGURE 1. Percentage of human rabies deaths in the United States by age groups from 1944 through 1954.




FIGURE 2. Principalylvatic rabies areas of the United States.

mentally infected foxes showed that their saliva contained rabies virus for long periods of time in one individual for 14 days

The most recent development in the problem of sylvatic rabies is the finding of rabies in insectivorous bats in the United States. This series of events began with the first isolation of rabies virus from bats in June 1953 when a yellow bat *Dasypterus floridanus* attacked a child near Tampa Fla. This episode aroused great interest prompting surveys and surveillance activities throughout the country. Since the first reported case in Florida 175 cases have been diagnosed from 18 states in widely diverse geographical areas of the United States, 2 from 1 province of Canada and 2 from Yugoslavia. Four species of tree living or solitary bats and 10 species of colonial or cave-dwelling bats have been implicated thus far. All are of the insectivorous variety. The greatest number of rabies virus isolations has been made from the Mexican free tailed bat *Tadarida mexicana* in the southwestern United States.

One of the many problems in the epizootiological puzzle of rabies in insectivorous bats is the question of asymptomatic transmission of the disease. It is well established that their chiropteran cousins, the vampire bats in Latin America, are capable in some instances of transmitting rabies for long periods of time without showing signs of illness. Some have suggested that this symbiotic host-virus relationship may be present in the insectivorous bats. During current studies conducted by the Communicable Disease Center in a cave in the southwestern United States, four infected bats have been found in a random sample of 300 normal bats collected in flight. Rabies virus was recovered from the salivary glands and not from the central nervous system of these four bats which remained clinically normal until sacrificed for testing. The ultimate test for asymptomatic carriers, however, that of demonstrating a continuous yield of virus from saliva of normal insectivorous bats observed over long periods of time, has not yet been done.

Although it has been possible to find the infection in normal bats of the aforementioned cave colony in frequencies ranging from 0.5 to 1.5 per cent, the rate of virus isolations went up to 5.1 per cent in dead and moribund bats collected in the midst of a die-off that lasted about three weeks. During this die-off many of the ill bats exhibited convulsions and spasmodic contractions of the abdominal muscles. Vicious biting of one bat by another was observed in one instance. Monthly sampling for serum neutralizing antibodies in this cave revealed positive antibody rates which increased gradually from 14.7 per cent in the early part of the season to 28.2 per cent late in the season just before the colony left on its southward migration.

Investigations are now being carried out to discover the epidemiological significance of the bat rabies findings. Although as yet there has been no direct and definitive evidence of the natural transmission of rabies from insectivorous bats to man or to lower animals, the public health implications of such findings cannot be minimized. There are now well-documented reports of 35 episodes of bats proven to be rabid that had bitten human beings (about half were unprovoked attacks) and 2 human rabies deaths are attributed to exposure from rabid bats on an anamnestic basis, since neither bat was recovered.

A few observations from the early phases of the current investigations may be of interest here. It has been possible *experimentally* to infect by peripheral routes several species of animals including bats with virus isolated from the brains and salivary glands of naturally infected bats but in most instances this has been done with difficulty (Stamm *et al*, 1956; Enright *et al*, 1955; Burns *et al*, 1956). This phase of the problem including attempts at simulated natural transmission, is described elsewhere in this monograph.

The ultimate solution to the rabies problem is predicated on the control and eventual elimination of the disease from animal populations. This may be accomplished by the establishment of transmission barriers such as animal immunization, elimination of stray dogs and the reduction of excessive numbers of wildlife vectors. Perhaps the greatest advance in the control of rabies in the last decade has been the improvement of animal rabies vaccines and the successful use of canine vaccination. The progress made in this field is discussed elsewhere in this monograph.

References

- BURNS, K. F., C. F. FARINACCI & T. G. MURNANE. 1956. Insectivorous bats naturally infected with rabies in southwestern United States. *J. Am. Public Health* 46: 1083-1091.
- ENRIGHT, J. B., W. W. SADLER, J. E. MOLLTON & D. G. CONSTANTINE. 1955. Isolation of rabies virus from insectivorous bat (*Tadarida mexicana*) in California. *Proc. Soc. Exptl. Biol. Med.* 89: 94-96.
- JOHNSON, H. N. 1948. Methods of rabies control. *Proc. 4th Intern. Congr. Trop. Med.* 1: 581-594.
- STAMM, D. D., R. F. KISSLING & M. E. FIDSON. Experimental rabies infection in insectivorous bats. *J. Infectious Diseases* 98: 10-14.

Discussion of the Paper

OSCAR SUSSMAN (*Bureau of Veterinary Public Health, State of New Jersey, Department of Health, Trenton, N. J.*) It has been contended by some observers when referring to the decrease of animal rabies in the United States during the period from 1944 through 1954 that this decrease has been associated with the use of new live culture avianized rabies virus vaccines. Inference has been made that there is a direct relationship between these reductions and the new live culture vaccines. The decrease shown was indeed substantial; however, it included reductions in areas where avianized vaccine was not used. To attribute this reduction solely to the introduction of a new vaccine is naive and at least a loose use of statistics. If there is any principle in epidemiological analysis it is that the secular movement of any disease is the resultant of many factors (multiple causation) and not merely the behavior of any single variable.

I agree with Ernest Tierkel that vaccination is an excellent procedure around which to conduct a rabies control program. I agree further that the avianized vaccine, where properly administered to dogs, is an efficacious prophylactic agent. In determining the amount of credit, however, that can be attributed to the new live culture vaccines in the decrease of animal rabies, I suggest that we examine the behavior of the disease in earlier times, both for the United States as a whole and for particular geographical subdivisions, with particular

reference to those periods when only phenolized vaccine was available for use and in which it actually was used

(1) In 1937 and 1938 11,500 animals were reported rabid in the United States. In 1942 this number decreased to 7163 animals—a decrease of almost 38 per cent

(2) In the east south-central part of the United States in 1936 2250 animals were reported rabid and by 1939 only 745 were reported rabid—a decrease of 67 per cent

(3) In the Pacific Coast area 2129 animals were reported rabid in 1938. This figure decreased to 515 in 1940—a reduction of 75 per cent

(4) In the state of New Jersey in 1939 619 cases of rabies were reported. This was reduced to 42 cases in 1943—a decrease of approximately 94 per cent

It is true that in the United States as a whole reported rabies increased from 7163 for the calendar year 1942 to 10,870 for the calendar year 1946 and then fell erratically to 3844 for the calendar year 1955. In the east south-central portion of the United States the number of reported cases increased from 145 in 1939 to 2066 by 1945, dropped to 1476 by 1951, only to increase to 2119 by 1953 and then to fall to 973 by 1955. On the Pacific Coast reported rabies decreased from 2129 in 1938 to 515 in 1940, only to increase to 916 by 1944, then to drop to 75 by 1951 and then to increase to 425 by 1955.

All of this demonstrates that reported rabies behaves in a cyclic fashion. If one were to choose any particular interval of time in a particular geographical area, a case could be made out for the use of killed vaccine for the control of stray dogs or for no effort whatsoever.

It must be admitted in all fairness that Ernest Tierkel's citation of the changing nature of the rabies problem from one in which dogs are the predominant species involved to one in which wildlife seems to be the essential factor in the maintenance of the disease is good supporting evidence that vaccination of pet animals is an excellent prophylactic procedure. I believe however that to ignore other possible factors such as increased awareness on the part of dog owners of their responsibility in disciplining their animals or in ignoring the ordinary behavior of the disease is to be unaware of the subtle interplay of the multiple variables in the natural history of any disease.

It has been said that "An epidemiologist is a disease investigator who slides into glory on the downward slope of an epidemic curve." It appears that attempts have been made to slide the new avianized rabies vaccines to glory on the downward slope of a nationwide cyclic epidemic rabies curve.

Avianized vaccine was discussed at the meeting of the American Public Health Association in New York N. Y. in 1949. A statement was made that in Staten Island N. Y. where there had been a considerable amount of rabies the previous year, no cases of rabies were reported following the use of avianized vaccine in a field trial. The audience was led to believe that this was due to the use of the avianized rabies vaccine. This impression was not corrected until Morris Greenberg of the New York City Department of Health indicated that the safety of the product might be evaluated in this respect but that one could not validly claim that avianized vaccine was responsible for the observed

phenomenon Greenberg pointed out that *while the avianized vaccine was used in Staten Island where there had been rabies and now there was not any the avianized vaccine had not been used in Manhattan where there had been rabies and now there was not any*¹

Phenolized vaccine has been used in many situations throughout the United States, and used successfully. Certainly the previous United States and regional figures referred to in which definite decreases in incidence were observed could not be attributed to the avianized vaccine when it is remembered that in this period only phenolized vaccines were available. These decreases were no less startling than those that occurred following the use of the avianized product in other situations.

Different causal relationships other than the use of vaccines are actually involved in the decrease of rabies incidence. This is particularly true in the case of dog rabies where proper control—a firmer conviction of owner responsibility in supervising the activities of the dogs and the elimination of stray dogs results in a decrease of susceptible populations to attack by wild and predatory infected animals. On the basis of all the evidence available I maintain that the use of any good vaccine aids in reducing the number of susceptibles among owned vaccinated dogs. I have yet to see a stray dog jump on a table and ask to be vaccinated.

In order to show conclusively that avianized vaccine is responsible for the reduction in reported rabies the following evidence must be submitted:

(1) Secular trend lines in rabies prevalence maintain a negative slope.

(2) If the foregoing statement is true then it must be demonstrated that (a) an increasing proportion of dogs vaccinated had been treated with avianized vaccine as compared to other vaccines and (b) that there was no statistical difference in the results between the groups of vaccinated dogs all other factors being equal.

Merlin Kaerberle in his paper "Newer Tools for the Prevention of Rabies in Domestic Animals" pays deserved attention to the duration of immunity experiment conducted jointly by the United States Public Health Service Washington D. C. and Lederle Laboratories Pearl River N. Y. However he failed to mention that the phenolized vaccine and the avianized vaccine both protected 100 per cent of all dogs challenged 24 months after vaccination.

I should like to raise several issues that appear to me to be of paramount importance in future rabies control procedures and in determining products to be used to control the disease in domestic animals.

(1) *Recall immunity*—Do multiple doses of phenolized vaccine administered at yearly intervals produce a more solid immunity than does the single dose of avianized vaccine given every 3 or 4 years? We have some evidence in our work with poliomyelitis and more conclusive evidence in Harald Johnson's² work on the phenolized product to the effect that multiple doses are of value in increasing total immunity. The 39 month study of the United States Public Health Service comparing phenolized and avianized vaccines failed to take into account the annual vaccination system in determining the relative value of multiple doses of killed products and method of administration.

(2) *Shelf life*—Consideration must be given to the shelf life and to the

practitioner's handling of any product. Live-culture vaccines when reconstituted require meticulous handling and may not be mishandled even when desiccated. Their entire value depends on the introduction of sufficient live virus to cause infection and subsequent immunity. This does not imply that phenolized products may not be abused; however, phenolized material can stand more abuse before its potency is lost than can avianized material.

(3) *Safety versus species*. Live vaccines must be used only in those species for which they have been proved safe. Therefore, if there is likelihood of utilization in unauthorized species, a product of greater safety must be used even if some degree of immunity is lost. The use of the avianized product in cats, puppies, cattle, skunks, and foxes has resulted in cases of atypical rabies.

(4) *Sensitivity versus neurological manifestations*. Sensitivity to egg protein may become a hazard when distemper products and other similar vaccines all are made from egg embryo tissue. As to neurological manifestations following the use of phenolized killed products, over 300,000 dogs in the state of New Jersey have been vaccinated over the last 5 years with no definite evidence of neurological manifestations.

(5) *Live rabies virus in nonendemic areas*. Aside from the benefit of recall immunity when annual vaccinations are made, rabies control administrators are faced with the problem of the use of live rabies virus in areas where no cases exist. Should the disease reappear following live vaccine usage, the subsequent public reaction must be taken into account. This consideration may appear inconsequential and a blot on the progress of scientific advancement, but it is a definite problem in areas where the disease has been eliminated.

(6) *Psychological value of annual vaccination*. The psychological impact when annual vaccinations are held must be considered. While in some areas this may be onerous, in others it is considered good procedure for it not only aids in controlling rabies, but it prescribes at least one time in a year when an owner must look closely at his pet and may help to prevent transfer of other animal diseases to the owner.

In conclusion, it is very difficult for one who has had considerable success in the use of the phenolized vaccine to see it buried so rashly before it is really dead. Psychologically, there is some reason for having an annual vaccination of dogs, as has been proved by field experience among those of us who have administered these programs on a continuous basis. Both vaccines certainly are valuable adjuncts in the prevention of rabies. The downward slope of the epidemic curve in the last decade cannot, however, be attributed solely to the introduction of the newer avianized vaccine. As one administrator and epidemiologist involved in the control of rabies in a state that has successfully eliminated the disease by the use of dog control, phenolized vaccine, and the elimination of stray dogs, I cannot accept the definition of an epidemiologist or of a superior vaccine as one that merely slides to glory on the downward slope of an epidemic curve.

Reference

1. JOHNSON, H. 1947. Rabies. *Ann N Y Acad Sci* 48(6): 390.

BAT RABIES EXPERIMENTAL HOST TRANSMISSION STUDIES

By Kenneth F. Burns, Dorothy Fresenius Shelton, and Earl W. Grogan
Fourth United States Army Medical Laboratory, Fort Sam Houston, Texas

The recovery of rabies virus from bats living under natural conditions in the United States indicates the need for evaluating the public health significance of these animals as reservoirs and carriers of this disease. Are rabies infected bats purely of academic interest or do they constitute a true public health hazard? Is it likely that human subjects will be affected directly by the bite of such a bat except as a rare accident, or is it more logical to believe that bats may come to represent an additional sylvatic reservoir in this country contributing to a marked increase of the disease in animals and indirectly in man? These are the more important questions with which public health personnel are concerned.

Originally the recovery of rabies virus from bats infected in nature was thought to be a unique single event localized within a small geographical area in Florida.¹ Within 6 months, however, other rabies isolations were made from bats in Pennsylvania and Texas,^{2,4} and thereafter from other states in the Southwest,⁵ the Midwest, and the West.⁶ From 4 species of solitary and 9 species of colonial bats, approximately 175 isolations of rabies virus have been made since June 1953, and 30 of these isolations have been associated with human exposures.⁷ Further circumstantial evidence concerned with 2 human rabies deaths in Texas would tend to incriminate bats as the vector of these 2 infections.^{8,9}

In attempting to evaluate the public health significance of bat rabies, we studied the ability of laboratory infected and naturally infected indigenous bats to transmit the disease to other animals under conditions of induced association in the laboratory.

Further studies were conducted with other animals to determine their susceptibility to bat rabies viruses and in an attempt to determine whether these agents have assumed some of the characteristics ascribed to the phenomenon of fixation.

MATERIALS AND PROCEDURES

Animals

Bats. *Tadarida mexicana* (Mexican free tailed bats) used in this experiment were obtained from two sources: Bracken Cave, Texas, and Carlsbad Caverns, N. Mex. The group of 400 bats netted at Bracken Cave appeared physically normal at the time of capture and were maintained in captivity for a 21 day period prior to the initiation of the experiment. The group from Carlsbad consisted of 50 encephalitic bats collected from the floor of the caverns during

an epizootic of rabies among *Molossidae* of the cave during August and September 1956*

Twenty five *Myotis pallidus pallidus* (pallid bats) were collected and shipped to us by Bryan P. Class of Oklahoma Agricultural and Mechanical College Stillwater Okla.

Monkeys Five young adult rhesus (*Macaca rhesus*) monkeys were employed. The animals weighed between 12 and 20 lb. and had been imported directly from India by the United States Air Force. They had not been used for any previous studies, having been maintained at this laboratory for 3 months prior to the beginning of this experiment.

Mice Three to four week-old SWLs albino mice weighing 8 to 10 gm. were used throughout the experiment as indicators of infection for virus titrations and for the intracerebral mouse neutralization tests for reidentification of agents employed and serum antibody titrations.

Other animals Fifteen guinea pigs, 16 hamsters, 19 rabbits, 4 Spanish goats, 6 chickens and 1 raccoon were included in the study for biting and induced-confinement experiments and for host susceptibility studies.

Virus Strains

Six rabies strains isolated from the brains of *T. mexicana* bats were employed in these studies. The strains were identified as the virus of rabies by the intracerebral mouse neutralization test. In addition to the bat isolates rabies fixed virus (Challenge Virus Standard) obtained from Lederle Laboratories Pearl River, N. Y. was used for control purposes.

Isolate No. 2915 (third mouse-brain passage) was obtained from a pool of the brains of 2 of the naturally infected Carlsbad Caverns bats collected during the August 1956 rabies epizootic.

Isolate No. 2084 (fifth mouse brain passage) was obtained from a moribund bat collected on the floor of Carlsbad Caverns in September 1955.

Isolates No. 1063 (second mouse brain passage) and No. 1248 (third mouse brain passage) were isolated from individual paralyzed bats collected in the San Antonio, Texas area in July and December 1954.

Isolate No. 1009-13 (fifth mouse brain passage) was obtained from a pool of the brains of 3 apparently normal bats (asymptomatic) collected in Uvalde, Texas in July 1954.

Isolate No. 168 (third mouse brain passage) was obtained from an individual bat exhibiting encephalitic manifestations at Fort Sam Houston, Texas in April 1954.

Diagnostic Criteria and Identification of Rabies Virus

Throughout the studies rabies virus was considered isolated either when Negri bodies could be demonstrated in brain smears from mice that died following a characteristic course of symptoms or when, in the absence of these

*The cooperation of D. G. Constantine of the Veterinary Public Health Section of the Communicable Disease Center, Public Health Service, Atlanta, Ga., is gratefully acknowledged.

identification of the agent could be made by the standard neutralization test. This latter method outlined by the Neurotropic Virus Disease Commission (since renamed the Commission on Viral Infections Armed Forces Epidemiological Board) Walter Reed Army Institute of Research Washington D C as stated by Paul¹⁰ incorporates equal volumes of undiluted inactivated rabies antiserum (horse) and serial tenfold dilutions of virus infected mouse brains.

Rabies Virus Titrations

With all experimental hosts each of the virus strains was titrated by the *intracerebral inoculation of tenfold dilutions into 8 to 10-gm 3 to 4-week-old white Swiss mice*. The total dose inoculated is expressed as mouse intracerebral LD₅₀s (MIC LD₅₀).

Bacterial Sterility Studies

Bacterial sterility studies were employed throughout the experiments on all injectable materials using trypticase soy and thioglycollate culture mediums. After incubation at 37° C and reading at 72 hours all tubes having no clouding or turbidity indicated sterile inocula.

Collection of Saliva Samples

Collections of saliva samples were made by repeated swabbing of the mouths of the bats with 6-inch cotton tipped applicators soaked in 1.0 ml of 10 per cent rabbit serum saline containing 1000 units of penicillin and 2.0 mg of streptomycin. The excess was expressed against the vial wall and after 1 hour of incubation at refrigerator temperature (4° C) with the antibiotics for control of bacterial contamination the mixture was inoculated intracerebrally into mice.

Inoculations

Intracerebral inoculations of mice (0.03 ml) bats (0.02 ml) chicks (0.05 ml) hamsters (0.05 ml) guinea pigs (0.10 ml) and rabbits (0.25 ml) were made using a 27 gauge needle and a 0.25 ml syringe employing sterile technique. In a similar manner but with increased amounts goats and monkeys received an intracerebral dose of 0.5 ml of inoculum using a 1.0-ml syringe with a 22 gauge needle while the animals were under general anesthesia.

Intramuscular inoculations (gluteals) of hamsters (0.05 ml) guinea pigs (0.10 ml) rabbits (0.75 ml) and a raccoon (1.0 ml) were made using a 1.0-ml syringe and a 22 gauge needle. Bats received 0.07 ml into their pectoral muscles with a 0.25 ml syringe and 27 gauge needle.

Subcutaneous inoculation of bats (0.02 ml) and rabbits (1.0 ml) was made in the suprascapular area with 27 gauge needles and with 0.25 ml syringes and 1.0-ml syringes respectively.

Intraperitoneal inoculations of guinea pigs (1.0 ml) and hamsters (0.5 ml) were accomplished with a 1.0-ml syringe and 22 gauge needle.

Serum Collections

Serum specimens were obtained from the monkeys for assay of neutralizing antibody for rabies virus prior to and four months after confinement with bats and again when the challenged animals were moribund with rabies. Sera of all other species of animals including fowl were collected just prior to exposure and either when moribund with clinical rabies or at time of sacrifice of those that remained asymptomatic.

Serum Neutralization Tests

Serum neutralization tests were conducted according to the method quoted by Paul¹⁰. A log neutralization index of 1.7 or greater was considered positive.

INDUCED ASSOCIATION EXPERIMENT INVOLVING INFECTED BATS (*T. mexicana*) AND NORMAL MONKEYS

Separate dog metabolism cages were utilized to confine a number of bats (*T. mexicana*) and 1 monkey each. Three of the cages each contained 100 bats inoculated with 32,000 MIC ID₅₀s of bat rabies isolate No. 2084 intracerebrally, subcutaneously and intramuscularly respectively. One cage contained the 50 naturally infected bats collected from Carlsbad Caverns. One cage contained 100 normal bats as controls. The cages were altered to consist of 2 compartments, upper and lower, separated by a metal grid containing openings large enough to permit passage of the bats. The monkeys were caged in the lower compartments and the bats were placed in the upper compartments at the beginning of the experiment.

Results

Early migration of the inoculated bats from the upper compartments of the dog metabolism cages to the lower compartments occupied by the monkeys was observed. The bats did not appear to return to their own compartments. This was believed to be due to the development of infection from the prodromal period in which the bats were irritable, aggressive and restless to a state of fatal progressive paralysis. No such picture was noted among the asymptomatic bats confined for control purposes. Numerous altercations occurred resulting in bites inflicted on the monkeys by the infected bats. The monkeys' attempts to dismember the confined bats resulted in their being bitten on the fingers. In addition to bite wounds, it appears reasonable to assume that the monkeys received stab wounds in the gums and throat while consuming portions of the bat carcasses. Many fragments of bats were discarded from each of the cages.

Bats inoculated intracerebrally placed with monkey No. 1. Of 100 intracerebrally inoculated bats, all were dead by the eighteenth postinoculation day. Fifty-three bats that were removed between the sixth and eighteenth postinoculation days were examined for rabies infection; the remaining carcasses were discarded due to mutilation by the monkey. These 53 bats were autopsied and tissues were prepared in 13 pools each of brain and salivary gland. Rabies

identification of the agent could be made by the standard neutralization test. This latter method outlined by the Neurotropic Virus Disease Commission (since renamed the Commission on Viral Infections, Armed Forces Epidemiological Board) Walter Reed Army Institute of Research, Washington D C as stated by Paul¹⁰ incorporates equal volumes of undiluted inactivated rabies antiserum (horse) and serial tenfold dilutions of virus infected mouse brains.

Rabies Virus Titrations

With all experimental hosts each of the virus strains was titrated by the intracerebral inoculation of tenfold dilutions into 8 to 10 gm 3 to 4-week-old white Swiss mice. The total dose inoculated is expressed as mouse intracerebral LD₅₀s (MIC LD₅₀s).

Bacterial Sterility Studies

Bacterial sterility studies were employed throughout the experiments on all injectable materials using trypticase soy and thioglycollate culture mediums. After incubation at 37° C and reading at 72 hours all tubes having no clouding or turbidity indicated sterile inocula.

Collection of Saliva Samples

Collections of saliva samples were made by repeated swabbing of the mouths of the bats with 6-inch cotton tipped applicators soaked in 1.0 ml of 10 percent rabbit serum saline containing 1000 units of penicillin and 2.0 mg of streptomycin. The excess was expressed against the vial wall and after 1 hour of incubation at refrigerator temperature (4° C) with the antibiotics for control of bacterial contamination the mixture was inoculated intracerebrally into mice.

Inoculations

Intracerebral inoculations of mice (0.03 ml) bats (0.02 ml) chicks (0.05 ml) hamsters (0.05 ml) guinea pigs (0.10 ml) and rabbits (0.25 ml) were made using a 27 gauge needle and a 0.25 ml syringe employing sterile technique. In a similar manner but with increased amounts goats and monkeys received an intracerebral dose of 0.5 ml of inoculum using a 1.0 ml syringe with a 22 gauge needle while the animals were under general anesthesia.

Intramuscular inoculations (gluteals) of hamsters (0.05 ml) guinea pigs (0.10 ml) rabbits (0.75 ml) and a raccoon (1.0 ml) were made using a 1.0 ml syringe and a 22 gauge needle. Bats received 0.02 ml into their pectoral muscles with a 0.25 ml syringe and 27 gauge needle.

Subcutaneous inoculation of bats (0.05 ml) and rabbits (1.0 ml) was made in the suprascapular area with 26 gauge needles and with 0.25 ml syringes and 1.0-ml syringes respectively.

Intraperitoneal inoculations of guinea pigs (1.0 ml) and hamsters (0.5 ml) were accomplished with a 1.0-ml syringe and 22 gauge needle.

TABLE 1
BAT RABIES EXPERIMENTAL HOST STUDIES
Intracerebral Challenge of Monkeys and Inoculation of Goats

Animal species	Bat brain strain	MIC† LD	Animal No.	Death / inoculation	Post inoculation day	Virus		Remarks
						Blood	Salivary gland	
Monkey (<i>Macaca rhesus</i>)	2084 (Fifth mouse brain passage)	10 ^{2.5}	1 685 595	1/1 1/1 1/1	8 8 10	Pos Pos Pos	Neg Neg Neg	Monkeys challenged 120 days after exposure to laboratory infected rabid bats
	2915 (Third mouse brain passage)	10 ^{2.7}	2	1/1	8	Pos	Neg	Monkey challenged 120 days after exposure to naturally infected rabid bats
	Control CVS	10	697	1/1	7	Pos	Neg	Monkey challenged 120 days after exposure to asymptomatic bats
Goat (caprine sp.)	2084 (Fifth mouse brain passage)	10	1	1/1	12	Pos	Neg	Encephalitis
	1248 (Third mouse brain passage)	10 ⁴	2 3	1/1 1/1	11 13	Pos Pos	Neg Neg	Encephalitis
	1063 (Third mouse brain passage)	10 ¹	4	0/1	50	Neg	Neg	Repeated IC challenge with 1063 (MIC† LD 10 ^{1.5}) on 18th post inoculation day Negative SNI†

Asymptomatic sacrifice

† Serum neutralization index

‡ Mouse intracerebral

isolated from the group of naturally infected bats with which this monkey had been in contact. Monkey 697 whose contact had been with normal bats received 100 000 MIC LD₅₀s of challenge virus standard (CVS). The results are shown in TABLE 1.

A prodromal period was noted in the monkeys as early as the fifth day for the CVS challenged animal and on the sixth to eighth days for those challenged with the bat rabies strains. Fulminating infections developed with weakness of the extremities, grinding of teeth, ease of incitement to convulsive state, moments of mental alertness subsequently drifting into subconsciousness followed by an ascending type paralysis. Deaths occurred on the seventh to tenth postinoculation days.

All monkeys were bled in the moribund state and sera from these bleedings were found to be negative for neutralizing antibodies to rabies. Mice inoculated with brain tissue of all monkeys exhibited positive evidence of rabies while none of the monkey salivary gland tissues yielded a viral agent. Histopathological examination of the monkeys' brains revealed a meningoencephalitis including glial satellitosis, perivascular cuffing and neuronophagia. Sellers stained impressions and Stovall Black stained sections failed to reveal Negri bodies.

virus was recovered and identified from all brain pools and all but 1 salivary gland pool the negative pool being composed of salivary glands of dead bats collected on the sixth postinoculation day

Bats inoculated subcutaneously placed with monkey No 685 Of the 100 subcutaneously inoculated bats 88 were removed dead between the sixth and twenty second postinoculation days Only fragments of the remaining 12 bats could be identified Twenty two pools each of bat brain and salivary gland tissues were prepared and inoculated intracerebrally into mice Five of the 44 pools yielded rabies virus 3 positive brain pools from bats dying the ninth seventeenth and twenty second postinoculation days and 2 positive salivary gland pools from bats dying the fifteenth and twenty second postinoculation days

Bats inoculated intramuscularly placed with monkey No 595 Of the 100 intramuscularly inoculated bats 59 dead were recovered between the sixth and twenty third postinoculation days the remaining 41 bats having been dismembered beyond recognition by the monkey Isolation materials contained in 16 brain and 16 salivary gland pools resulted in recovery of the virus in 5 instances from both brain and salivary glands of bats dying the sixth and thirteenth postinoculation days and from the brain only of bats dying on the twentieth postinoculation day

Bats infected naturally placed with monkey No 2 The 50 naturally infected *T. mexicana* were caged with rhesus monkey No 2 Twenty four hours later 11 bats had died and were removed from the cage Six of the remaining sick bats died on the second postexposure day 3 on the third and 2 on the fourth From the 2 bats dying the fourth postexposure day 2 isolates were obtained (No 2915 Br and No 2915 SG) from pooled tissues of both brain and salivary gland tissue and these were identified as the virus of rabies Twenty eight bats were either eaten or mutilated and could not be utilized

Normal controls placed with monkey No 697 One hundred normal bats were placed with a fifth monkey On the twenty third postexposure day 48 bats were sacrificed for testing and from their tissues 13 brain pools and an equal number of salivary gland pools were prepared Intracerebrally inoculated mice yielded negative results Fifty two bats were mutilated beyond salvage or cannibalized by the monkey

None of the monkeys exhibited clinical symptoms of rabies or other disease during a 120-day period of observation no demonstrable neutralizing antibodies were observed in sera drawn prior to or 120 days after exposure

Challenge

Since all of the monkeys remained asymptomatic during the observation period each was challenged with rabies virus to prove its susceptibility to bat rabies virus

Monkeys Nos 1 685 and 595 were challenged intracerebrally with 6300 MIC ID₅₀s of strain 2084 four months after their contact exposure with bats experimentally infected with that strain Simultaneously monkey No 2 was challenged intracerebrally with 50 000 MIC LD₅₀s of strain 2915 which was

neuronophagii perivascular infiltration glial nodules and Negri inclusion bodies

Strain 1063 was inoculated (approximately 500 000 MIC LD₅₀s) into goat No 4. On the twelfth postinoculation day the animal refused water and food but after a 24 hr period it once again started to eat and drink and assume normal habits. On the eighteenth postinoculation day the goat was challenged intracerebrally with approximately 8000 MIC LD₅₀s of bat rabies virus strain 1063. The animal exhibited no central nervous system symptoms during a 50-day observation period and was sacrificed. Serum specimens collected at time of inoculation and 41 and 50 days postinoculation were negative for neutralizing antibodies for rabies. The mechanism of immunity observed in this animal manifested by its resistance to challenge in the absence of discernible neutralizing antibody cannot be explained.

Rabbits

Nineteen rabbits were utilized in experimental infection studies via intracerebral (I C) intramuscular (I M) and subcutaneous (S C) routes of inoculation resulting in infection and subsequent death of 13 of the group (TABLE 2). Six animals remained asymptomatic and were sacrificed after a 33 to 61 day observation period. All 4 rabbits inoculated intracerebrally and 5 of the 7 inoculated intramuscularly died of rabies the mean incubation period for each of these routes being 17 days. The 2 intramuscularly inoculated survivors were sacrificed on the thirty fifth and sixty first postinoculation days. Of the 8 rabbits inoculated subcutaneously 4 became infected with rabies with a mean incubation period of 22.5 days and 4 of this group remained asymptomatic 3 being sacrificed after 33 days and 1 on the fortieth day.

Rabies virus was recovered from the brains of 11 of the 13 rabbits succumbing to infection with bat rabies virus. No viral agent could be recovered from

TABLE 2
BAT RABIES TRANSMISSION STUDIES IN RABBITS

Bat brain	Route	Dose in LD ₅₀	Deaths / inoculated	Post inoculation day / interval	Neg body	Virus recovered / tested	Neutralizing test / positive / tested
2084 (Fifth mouse brain passage)	I C	10	2/2	13-14	Negative	2/2	2/2
	S C	10	2/4	23-26 (33) (40)	Negative	1/4	3/4
	I M	10*	4/5	14-16 16-23 (61)	Negative	3/3	4/4
1009-13 (Fifth mouse brain passage)	I C	10*	1/1	23 (33) (33)	Positive	1/1	1/1
	S C	10*	0/2	18	Negative	0/2	1/2
	I M	10	1/1		Positive	1/1	1/1
1248 (Third mouse brain passage)	I C	10*	1/1	18	Negative	1/1	ND
	S C	10	2/2	20-21 (35)	Negative	2/2	2/2
	I M	10*	0/1		Negative	0/1	ND

Symbol () all animals terminated after 30 days had remained asymptomatic for the observation period recorded. ND test not performed.

INDUCED-BITING EXPERIMENTS USING LABORATORY INFECTED *A. pallidus*

Fifteen asymptomatic *A. pallidus pallidus* were each inoculated intracerebrally with approximately 20,000 MIC LD₅₀s of bat rabies strain 2084. The bats had been maintained at the laboratory for 11 days prior to inoculation. One bat dying on the second postinoculation day was discarded. Of the remaining 14 mammals, deaths occurred between the eighth and twenty-seventh postinoculation days. 7 of the group dying between the eighth and twelfth days. A period of 24 to 48 hours of marked irritability preceded the deaths. Rabies virus was recovered from the brains of 13 of 14 bats, the single failure being the brain of one of the 4 bats dying on the twelfth postinoculation day. Attempts to isolate rabies virus from the salivary glands were accomplished on 6 of the inoculated bats. Three of these yielded rabies virus. Pooled mouth swabbings collected 7 days postinoculation from 3 bats exhibiting symptoms of a central nervous system infection yielded rabies virus.

Six of the infected bats (including the three that yielded rabies virus from salivary gland) on the tenth, eleventh, twelfth and fourteenth postinoculation days were induced to inflict multiple bites on the ears and abdomens of 6 guinea pigs. The guinea pigs survived a 60-day observation period with no clinical signs of disease and were exsanguinated 60 days following exposure. Their sera failed to exhibit rabies neutralizing antibody. Concurrently, 6 white Swiss mice were placed with and received multiple bite exposures from these bats. The mice survived a 60-day observation period with no signs of clinical illness.

INOCULATIONS OF OTHER ANIMALS AND FOWL

Goats

Selected bat rabies strains were studied in 6-month-old Spanish goats to determine this animal's susceptibility via the intracerebral route of inoculation (TABLE 1).

One caprine (No. 1) inoculated with approximately 800,000 MIC LD₅₀s of bat rabies strain 2084 died on the twelfth postinoculation day, exhibiting typical symptoms of rabies. Rabies virus was recovered from the brain of this animal.

Bat rabies strain 1248 was inoculated (approximately 630,000 MIC LD₅₀s) into 2 goats (Nos. 2 and 3). Goat No. 2 exhibited symptoms of encephalitis on the ninth postinoculation day and died 2 days later. Goat No. 3 was lethargic and incoordinated with a fixed stare on the tenth postinoculation day. 24 hours later a marked grinding of the teeth was noted with progressive ascending type paralysis, and the animal died on the thirteenth postinoculation day. Mice inoculated with goat brain suspensions from these animals were positive for rabies, while the goats' salivary glands failed to reveal infection with the virus on replicate isolation attempts in mice.

Histopathological examination of brain tissue of three goats (Nos. 1, 2, and 3) revealed typical pathological findings including meningoencephalitis.

neuronophagia perivascular infiltration glial nodules and Negri inclusion bodies

Strain 1063 was inoculated (approximately 500 000 MIC I D₅₀s) into goat No 4. On the twelfth postinoculation day the animal refused water and food but after a 24 hr period it once again started to eat and drink and assume normal habits. On the eighteenth postinoculation day the goat was challenged intracerebrally with approximately 8000 MIC I D₅₀s of bat rabies virus strain 1063. The animal exhibited no central nervous system symptoms during a 50-day observation period and was sacrificed. Serum specimens collected at time of inoculation and 41 and 50 days postinoculation were negative for neutralizing antibodies for rabies. The mechanism of immunity observed in this animal manifested by its resistance to challenge in the absence of discernible neutralizing antibody cannot be explained.

Rabbits

Nineteen rabbits were utilized in experimental infection studies via intracerebral (I C) intramuscular (I M) and subcutaneous (S C) routes of inoculation resulting in infection and subsequent death of 13 of the group (TABLE 2). Six animals remained asymptomatic and were sacrificed after a 33 to 61 day observation period. All 4 rabbits inoculated intracerebrally and 5 of the 7 inoculated intramuscularly died of rabies the mean incubation period for each of these routes being 17 days. The 2 intramuscularly inoculated survivors were sacrificed on the thirty fifth and sixty first postinoculation days. Of the 8 rabbits inoculated subcutaneously 4 became infected with rabies with a mean incubation period of 22.5 days and 4 of this group remained asymptomatic 3 being sacrificed after 33 days and 1 on the fortieth day.

Rabies virus was recovered from the brains of 11 of the 13 rabbits succumbing to infection with bat rabies virus. No viral agent could be recovered from

TABLE 2
BAT RABIES TRANSMISSION STUDIES IN RABBITS

Bat brain	Route	Dose mouse LD ₅₀	Death inoculated	Post inoculation day (if infection)	Antibody	Virus recovery brain positive / tests	Neutralization positive / tests
2084 (Fifth mouse brain passage)	I C	10	2/2	13 14	Negative	2/2	2/2
	S C	10	2/4	23 26 (33) (40)	Negative	1/4	3/4
	I M	10*	4/5	14 16 16 23 (61)	Negative	3/5	4/4
1009-13 (Fifth mouse brain passage)	I C	10*	1/1	24	Positive	1/1	1/1
	S C	10*	0/2	(33) (33)	Negative	0/2	1/2
	I M	10	1/1	18	Positive	1/1	1/1
1248 (Third mouse brain passage)	I C	10	1/1	18	Negative	1/1	ND
	S C	10*	2/2	20 21	Negative	2/2	2/2
	I M	10	0/1	(35)	Negative	0/1	ND

Symbols () all animals terminated after 30 days had remained asymptomatic for the observation period recorded. ND test not performed.

the brains of 2 of the rabbits exhibiting clinical rabies (1 I C 1 I M) nor from the brains of 6 asymptomatic rabbits surviving the observation period. Negri bodies were observed in Stovall Black stained sections of 2 of 19 rabbit brains examined (1 I C 1 I M). Rabies virus was recovered from the salivary glands (parotid and submaxillary pool) of only 1 (S C) of 8 rabbits from which tissues were harvested (1 I C 1 I M 6 S C).

Sera collected from 11 of the 13 symptomatic rabbits (2 sera not tested) exhibited positive serum neutralization indices. Sera of 3 of the 6 asymptomatic rabbits drawn the day of sacrifice were positive for rabies neutralizing substance. Two rabbits were negative and no serum specimen was collected from the sixth surviving animal.

Guinea Pigs

Nine guinea pigs (3 each I C I P and I M) received 8000 MIC LD₅₀s of bat rabies strains Nos 768 and 1009-13. The 3 intracerebrally inoculated animals died of rabies with a mean incubation period of 15 days. Rabies virus was recovered from the brains of the 5 animals dying (3 I C 2 I M) and 2 of 4 subjected to histopathological examination of brain tissue revealed the presence of Negri bodies. All 3 of the intraperitoneally inoculated guinea pigs and 1 of the intramuscularly inoculated group survived a 61 day observation period with no clinical signs of disease.

Hamsters

Sixteen hamsters receiving 8000 MIC LD₅₀s of bat rabies virus strain No 768 (5 I P 5 I C and 6 I M) died of rabies with a mean incubation period of 9 days. While rabies virus was recovered from the brains of all of these animals, Negri bodies were observed in only 2 (1 I C 1 I M) of 8 brains examined (4 I C 3 I M 1 I P).

Raccoon

One raccoon inoculated intramuscularly with approximately 630 000 MIC LD₅₀s of bat rabies strain 2084 remained asymptomatic for 62 days at which time it was exsanguinated. The animal's serum was negative for neutralizing antibodies to rabies and the brain yielded no viral agent when inoculated intracerebrally in mice.

One Day Old Chicks

Five of six 1 day old chicks inoculated intracerebrally with bat rabies strains 1009-13 and 1248 survived periods of from 46 to 107 days with no clinical evidence of disease. The sixth bird developed clinical symptoms of rabies including incoordination and leg weakness on the twenty ninth postinoculation day followed by paralysis on the thirtieth day at which time it was sacrificed (TABLE 3). Serum specimens drawn from all 6 inoculated birds on the day of sacrifice were positive for rabies neutralizing antibodies.

Five control chicks inoculated intracerebrally with normal mouse brain suspension were sacrificed after a 63-day observation period with no signs of dis-

TABLE 3
EXPERIMENTAL INFECTION OF 24 HOUR CHICKS WITH BAT RABIES VIRUS

Bat brain strain	IC dose mouse LD	Chick No.	Survival time (days)		Remarks
			Day	Log NIT	
1009-13 (Fifth mouse brain passage)	10 ^{2.4}	1	46 107	4.0 3.5	Asymptomatic
		2	46	4.0	Asymptomatic
		3	30	3.5	Clinical rabies Virus recovered
		4	54	4.0	Asymptomatic
1248 (Third mouse brain passage)	10 ^{2.4}	5	60	4.0	Asymptomatic
		6	60	2.0	Asymptomatic
Controls	NMB†	7-11	63	0.0-0.5	Five normals

Post inoculation day

† Log neutralization index

‡ NMB = normal mouse brain

ease and sera collected at the time of sacrifice were negative for neutralizing antibodies for rabies

Bats (*Introns pallidus pallidus*)

After 8 days acclimatization in the laboratory each of 10 asymptomatic *I. pallidus pallidus* was inoculated intramuscularly with approximately 13 000 MIC LD₅₀s of bat rabies strain 2084. Of the 10 inoculated 1 bat died the first and 2 on the third postinoculation day and were discarded as nonspecific deaths. Six of the remaining 7 bats died between the fifth and ninth postinoculation days. It is noteworthy that 1 inoculated bat survived 10 days before death intervened. Individual attempts to recover the rabies virus from brain and salivary gland tissue of these bats by intracerebral inoculation of mice were futile in spite of the appearance of characteristic pathology other than Negri inclusion bodies. Bacterial sterility controls were also negative.

DISCUSSION

Possible theoretical objection to serial passage of the various bat rabies virus strains in mice on the grounds that such passage may tend to cause some fixation of the virus is recognized. However subsequent experimental infections by the subcutaneous route and isolation of the virus from salivary glands argue against any theory of virus fixation.

Other workers^{11, 12} have demonstrated that rabid hemophagous and fructivorous bats are capable of transmitting rabies and may recover and transmit the disease as asymptomatic carriers. However we have been unsuccessful in

the brains of 2 of the rabbits exhibiting clinical rabies (1 I C, 1 I M) nor from the brains of 6 asymptomatic rabbits surviving the observation period. Negri bodies were observed in Stovall Black stained sections of 2 of 19 rabbit brains examined (1 I C, 1 I M). Rabies virus was recovered from the salivary glands (parotid and submaxillary pool) of only 1 (S C) of 8 rabbits from which tissues were harvested (1 I C, 1 I M, 6 S C).

Sera collected from 11 of the 13 symptomatic rabbits (2 sera not tested) exhibited positive serum neutralization indices. Sera of 3 of the 6 asymptomatic rabbits drawn the day of sacrifice were positive for rabies neutralizing substance. Two rabbits were negative and no serum specimen was collected from the sixth surviving animal.

Guinea Pigs

Nine guinea pigs (3 each I C, I P, and I M) received 8000 MIC LD₅₀s of bat rabies strains Nos. 768 and 1009-13. The 3 intracerebrally inoculated animals died of rabies with a mean incubation period of 15 days. Rabies virus was recovered from the brains of the 5 animals dying (3 I C, 2 I M) and 2 of 4 subjected to histopathological examination of brain tissue revealed the presence of Negri bodies. All 3 of the intraperitoneally inoculated guinea pigs and 1 of the intramuscularly inoculated group survived a 61 day observation period with no clinical signs of disease.

Hamsters

Sixteen hamsters receiving 8000 MIC LD₅₀s of bat rabies virus strain No. 768 (5 I P, 5 I C, and 6 I M) died of rabies with a mean incubation period of 9 days. While rabies virus was recovered from the brains of all of these animals, Negri bodies were observed in only 2 (1 I C, 1 I M) of 8 brains examined (4 I C, 3 I M, 1 I P).

Raccoon

One raccoon inoculated intramuscularly with approximately 630 000 MIC LD₅₀s of bat rabies strain 2084 remained asymptomatic for 62 days at which time it was exsanguinated. The animal's serum was negative for neutralizing antibodies to rabies and the brain yielded no viral agent when inoculated intracerebrally in mice.

One Day Old Chicks

Five of six 1 day old chicks inoculated intracerebrally with bat rabies strains 1009-13 and 1248 survived periods of from 46 to 107 days with no clinical evidence of disease. The sixth bird developed clinical symptoms of rabies including incoordination and leg weakness on the twenty ninth postinoculation day followed by paralysis on the thirtieth day at which time it was sacrificed (TABLE 3). Serum specimens drawn from all 6 inoculated birds on the day of sacrifice were positive for rabies neutralizing antibodies.

Five control chicks inoculated intracerebrally with normal mouse brain suspension were sacrificed after a 63 day observation period with no signs of dis-

While interspecific comparisons are not entirely valid the results obtained in other animals inoculated with the various bat rabies isolates used in the present study do not suggest an attenuation of the virus. Apart from a reduced capacity for pathogenesis there appears to be little difference in our hands between the biological characteristics of bat virus strains studied and those characteristics normally associated with street virus. Despite a tendency toward a negative Negri pattern there is a high degree of infectivity for several laboratory animals and for goats. The fact that 4 of 8 rabbits could be readily infected subcutaneously with a mean incubation period of 22.5 days is of considerable interest. In our hands the incubation period of rabies caused by street virus is from 15 to 30 days for rabbits when the virus is introduced into the brain. Establishment of infection via the relatively less susceptible subcutaneous route of entry by the bat rabies virus attests to its invasiveness and virulence. Also the numerous isolations of the virus from salivary glands of infected animals indicate that it is not a fixed virus.

The failure of infected bats to transmit rabies to confined monkeys and other laboratory animals suggests that a certain inherent mechanical deficit existed that prevented these animals from transmitting the disease. The thickness of the cartilaginous tissues of the ear and the skin layers of the guinea pig appeared to govern the extent of the penetrating wounds that *T. mexicana* and *A. pallidus* were capable of inflicting. In no instance did *A. pallidus* penetrate the skin of guinea pigs in bite experiments to sufficient depth to produce hemorrhage. Only localized edema without extravasation of tissue fluids was noted. This inability to produce a deep penetrating wound may be one of the reasons for our negative transmission studies.

Previously reported⁴ and contrary to the negative host transmission results however is the presence of antirabies neutralizing antibody in large numbers of and the isolation of rabies virus from asymptomatic bats collected in nature. These data suggested that a certain tolerance or diminished virulence in the host virus relationship did exist and suggested further the possibility of infection and recovery among the indigenous bat population. Asymptomatic carriers if capable of transmitting the disease would be a formidable problem from the standpoint of preventive medicine because of the lack of self limiting factors.

Statistical analysis of our unpublished bat serologic data compiled during the last 4 years of study of multiple bat sera pools revealed that the probable estimate of incidence of rabies-neutralizing substances for bat populations of 2 species of Molossidae was 15.4 per cent with 95 per cent confidence limits of 12.5 per cent and 17.9 per cent and for 5 species of Vespertilionidae was 5.7 per cent with 95 per cent confidence limits of 2.0 per cent and 9.4 per cent. Computations were based on normal approximations to the binomial distribution of percentages. It is believed that some importance may be attached to the significantly lessened incidence of rabies neutralizing substances in the Vespertilionidae as compared with the Molossidae. The migratory habits known for *T. mexicana* whose population makes up a large percentage of the Molossidae (composed of 2 genera and 6 species) in contrast with the hibernating behavior exhibited by Vespertilionidae offer an essential physical require

our attempts to prove naturally infected and laboratory infected indigenous insectivorous bats (*T. mexicana* and *A. pallidus pallidus*) capable of transmitting the disease by natural means to monkeys under conditions of induced association or by biting experiments on guinea pigs and mice. The bite of the rabid *T. mexicana* was quite capable of penetrating the skin or ears of the contact animals. Recovery of rabies virus from the brain and salivary gland tissues of the infected bats indicated their potential capabilities for transmission.

In the induced association experiment involving infected bats and normal monkeys, the natural curiosity of monkeys caused them to examine a number of the infected bats. Consequently the monkeys were bitten several times on their hands and fingers before they could relieve themselves of the bats. The reaction on the part of confined monkeys after being bitten was to attack the bats, tear them apart and at times consume them, thereby affording opportunity for further exposure. This type of experience is probably duplicated to a certain extent in nature by predatory animals in their search for food. Indeed, data gathered by Twente¹³ suggests that predators are important limiting factors in the maintenance of *Tadarida* populations. This suggested the possibility that insectivorous bats infected with rabies might be a reservoir for the maintenance of the sylvatic disease. Such a hypothesis could explain some of the bizarre geographical and cyclic disease patterns in wildlife rabies where outbreaks of the disease occur in rapid succession in widely separated regions of the United States.

Results obtained by banding¹³ suggest that the pattern of movement behavior for insectivorous bats is an erratic one in that bats banded in one cavern and released elsewhere will infrequently return to their original habitat but will quite often fly to other caverns not in the direct line of flight from the point of original release. These transient individuals if infected with rabies would appear to offer excellent exposure possibilities for predatory animals that inhabit the same caverns.

However, if a parallel may be drawn between the habits of predatory animals in nature and the action of confined monkeys of the laboratory in attacking and consuming infected bats, ingestion does not appear to contribute to the mode of transmission and perpetuation of the virus in nature.

We experienced no difficulty in infecting bats either intracerebrally or peripherally. This lack of tolerance on the part of the indigenous bats for the bat rabies virus by peripheral inoculation does not suggest that the virus has existed among the Molossidae for a sufficient time to produce an established relationship as manifested by diminished virulence for the host. However, we were unable to recover rabies virus from a group of intramuscularly inoculated *A. pallidus pallidus* that received viable virus based on simultaneous titration of the inoculum in mice that yielded an LD₅₀ titer of $10^{-5.3}$. The absence of the virus in the dead animals may have been due to enzymatic destruction of the virus in the dead cells or to a phenomenon referred to as an autosterilizable neuroinfection, a condition postulated in rabies among other neurotropic virus diseases by Levaditi, Sanchis Bayarri, and Schoen¹⁴ which presupposes that the reaction of the host is so efficient that the virus and the host are destroyed. No rational answer appears to be available immediately.

Guinea pigs and white mice bitten numerous times by bats (*L. pallidus pallidus*) experimentally infected with bat rabies virus failed to contract rabies although isolation of rabies virus from the salivary glands and mouth swabbings of a significant number of the rabid bats was accomplished. The guinea pigs failed to develop demonstrable neutralizing antibodies to rabies virus.

Experimental infections of monkeys, goats, rabbits, guinea pigs, hamsters, mice, and chicks with bat rabies virus were accomplished.

Laboratory infection of 4 of 8 rabbits and numerous bats (*T. mexicana*) by subcutaneous inoculation and isolation of the bat rabies virus from salivary glands of a number of the rabid animals indicate that this studied agent has not assumed the characteristics normally associated with those of fixation. Neither is the virus of indigenous bat rabies considered to have assumed an aberrant form in this unusual host.

Among bats (*T. mexicana* and *L. pallidus pallidus*), goats, rabbits, guinea pigs, hamsters, and mice experimentally inoculated with bat rabies virus, none showed apparent recovery from clinical rabies.

Regardless of our negative results in transmission experiments, the repeated isolation of bat rabies virus from salivary glands and from mouth swabbings of infected bats emphasizes our opinion that persons bitten by bats should consider the incident as a potential rabies exposure and that these cases should be evaluated for vaccine prophylaxis.

Acknowledgment

We express our indebtedness to Lt. Col. John M. Lukeman, MC, Commanding Officer of the Fourth United States Army Medical Laboratory, for his encouragement and cooperation which made it possible to carry out the work described in this paper.

References

1. VENTERS, H. D., W. R. HOFFERT, J. E. SCATTERDAY & A. V. HARDY. 1954. Rabies in bats in Florida. *Am. J. Public Health* 44: 182.
2. WITTE, E. J. 1954. Bat rabies in Pennsylvania. *Am. J. Public Health* 44: 186-187.
3. SULLIVAN, T. D., J. F. GRIMES, R. B. EADS, G. C. MENZIES & J. V. IRONS. 1954. Recovery of rabies virus from colonial bats in Texas. *Public Health Repts. U. S.* 69: 766-768.
4. BURNS, K. F. & C. J. FARINACCI. 1955. Rabies in non sanguivorous bats of Texas. *J. Infectious Diseases* 97: 211-218.
5. BURNS, K. F., C. J. FARINACCI, T. G. MURNANE & D. F. SHELTON. 1956. Insectivorous bats naturally infected with rabies in southwestern United States. *Am. J. Public Health* 46: 1089-1097.
6. ENRIGHT, J. B. 1956. Bats and their relation to rabies. *Ann. Rev. Microbiol.* 10: 369-392.
7. TIERKEL, E. S. 1957. Present status of rabies in insectivorous Chiroptera of the United States. World Health Organization Expert Committee on Rabies. Doc. 92.
8. SCLAIN, S. E. & M. J. GREVE. 1954. Human rabies caused by bat bite. *Texas State J. Med.* 50: 620-621.
9. QUIST, K. D., R. B. FADS & A. CONKLIN. 1957. Studies on bat rabies in Texas. *J. Am. Vet. Med. Assoc.* 130: 66-68.
10. PAUL, J. R. 1944. The filtrable viruses. In *Laboratory Methods of the United States Army*. J. S. Simmons and C. J. Gentzkow, Eds. 579-600. Lea & Febiger, Philadelphia, Pa. Also in *Viral and Rickettsial Infections of Man*, 1952, 2nd ed. T. M. Rivers, Ed. Lippincott, Philadelphia, Pa. 72-77.

ment for interspecies transmission with the known rabies infected vampire bats (*Desmodus sp*) of Mexico. The social behavior of clustering according to species is quite selective for colonial bats, in this respect one might speculate that migrant infected members of the Molossidae, in competing with the indigenous Vespertilionidae in habitat selection, present the necessary opportunity for the chance infection of the latter.

Although positive serologic findings previously reported have suggested that nonfatal infections might occur they are by no means proof that they actually do occur. Certainly the lack of survivors among the laboratory infected bats does not support such a thesis.

Experiments conducted by Pawan^{11, 12} with both laboratory infected and naturally infected bats demonstrated the capabilities of the vampires and fruit eating bats to recover from a clinical infection and continue to be capable of transmitting the virus for prolonged periods.

Among the scores of insectivorous bats that were inoculated by peripheral and intracerebral routes with varying doses of bat rabies virus we have yet to find a single individual exhibiting a frank illness followed by recovery based upon a 70 day observation period.

The possibility of latency of infection should not be dismissed even though we were unable to observe this phenomenon under the conditions described in our experimental methods. To public health veterinarians charged with the responsibility of control of rabies reservoir type animals latency of infection would be of great significance. If indigenous insectivorous bats were true carriers of rabies virus there would be no self limiting mechanism to assist these disease control specialists.

The irregular appearance and low incidence of Negri bodies in infections with the bat rabies strains studied emphasize the danger of relying upon microscopic examination alone in diagnosing bat rabies.

Bat rabies awaits further extensive study and investigation for clarification. Meanwhile in spite of our negative results in transmission experiments repeated isolation of bat rabies virus from salivary glands and from mouth swab bings of infected bats emphasizes our opinion that persons bitten by bats should consider such incidents as potential rabies exposures and these cases should be evaluated for vaccine prophylaxis.

SUMMARY

Six strains of bat rabies virus isolated from naturally infected bats (*T. mexicana*) were used in transmission experiments and in studies of the susceptibility of other animals to the virus.

Monkeys in close confinement with groups of bats (*T. mexicana*) experimentally infected with bat rabies virus and with a naturally infected group of bats did not contract rabies nor develop demonstrable neutralizing antibodies against rabies virus. We observed biting of the monkeys by the rabid bats and in lesion of some of the bats by the monkeys and rabies virus was isolated from the salivary glands of a significant number of the rabid bats. Susceptibility of the monkeys to fatal infection with bat rabies virus was later proved by intracerebral challenge.

NEWER TOOLS FOR THE PREVENTION OF RABIES IN DOMESTIC ANIMALS

By Merlin I. Kaeblerle

College of Veterinary Medicine University of Illinois Urbana Ill

The occurrence of rabies in domestic animals is a problem of major importance in the United States and in many other countries of the world. Of primary concern to the veterinarian and physician is the resulting exposure of many human beings to the dread disease. A large percentage of our human exposures is due to contact with rabid animals classified as domestic. The number of reported cases of rabies in the United States during the ten year period 1947 to 1956 is shown in TABLE 1 which also serves to indicate the high percentage of cases reported in domestic animals as compared to other species. The problem is also an economic one since the number of cattle and other valuable animals that succumb to the disease is substantial. The loss of valuable breeding stock is a serious matter for all too often these animals cannot be replaced even though an indemnification is paid. Consequently prevention of the disease in these animals is of more than minor interest.

The development of prophylactic agents has been essential for once clinical signs of illness appear the animal invariably dies. For this reason ever since inception of the modern era of rabies research scientists have looked to agents that would prevent the disease. Previously particular emphasis was placed upon prophylactic treatment of animals and humans following exposure to the disease rather than upon pre exposure immunization.

Extensive experimentation by Louis Pasteur and his associates in the early 1880s culminated in the immunization of dogs against the disease by 1894. This research employing the use of vaccine in animals led to the successful administration of vaccine to human beings in 1885 and 1886^{23, 24, 25}. The basis for the Pasteur vaccine was a virus modified by serial passage through rabbits until it had become "fixed". The virus was further reduced in virulence by desiccation of the virus infected rabbit spinal cords over caustic potash. During this same period Galtier^{11, 12} was concerning himself with the protection of herbivorous animals against rabies. He subsequently administered street virus intravenously to sheep, challenged them with the homologous strain introduced subcutaneously and reported an immune response. In 1888 Roux and Nocard¹⁶ reported experimental studies in which rabbit brain emulsion containing live rabies virus was administered intravenously to cattle and sheep that were later challenged with street virus by the intra ocular route. Live virus was also administered by Moncet²⁷ and by Krasnitski²⁸ who concluded that it was a safe and efficacious method. Living fixed virus was administered to dogs by Schnurer²⁹ who reported favorable results.

Modifications of the Pasteur vaccine were soon forthcoming as additional workers entered this field of research. Thus in the early 1890s Hognes working at the Pasteur Institute of Budapest Hungary developed a vaccine that bears his name. This vaccine prepared from fixed virus and further

- 11 PAWAN J L 1936 Rabies in the vampire bat of Trinidad with special reference to the clinical course and the latency of infection *Ann Trop Med* 30 401-422
- 12 PAWAN J L 1948 Fruit eating bats and paralytic rabies in Trinidad *Ann Trop Med* 42 143-177
- 13 TWENTE J W 1955 Aspects of a population study of cavern dwelling bats *J Mammalogy* 36 349-390
- 14 LEVADITI C V SANCHIS BAYARRI & R SCHOEN 1978 Neuro-infections autostérilisables (encéphalite herpès rage) *Compt rend soc biol* 98 911

TABLE 2
SUMMARY COMPARATIVE CANINE RABIES VACCINE EXPERIMENTS²⁹

Vaccine	Route	Rabies mortality	Paralysis mortality	Period of immunity	Author
Brain tissue (phenolized) Controls	SC —	6/52 41/52	11.5 78.2	12 months	Johnson ²
Modified live virus (chick embryo) Brain tissue (phenolized) Controls	IM SC —	0/25 3/22 18/25	0 13.6 72	12 months	Kajrowski and Black ³⁰
Modified live virus (chick embryo) Brain tissue (phenolized) Controls	IM SC —	3/25 8/19 21/23	12.0 42.1 91.4	24 months	Kajrowski and Black ³⁰
Modified live virus (chick embryo) Brain tissue (phenolized) Brain tissue (ultraviolet irradiated) Brain tissue (benzene ether treated) Controls	IM IM IM IM —	0/37 0/30 0/31 0/30 18/33	0 0 0 0 54.6	24 months	Tierkel, Kissling, Erlson and Halcl
Modified live virus (chick embryo) Brain tissue (phenolized) Brain tissue (ultraviolet irradiated) Controls	IM IM IM —	0/30 8/34 7/30 31/36	0 23.6 23.3 86.1	30 months	Tierkel, Kissling, Erlson and Halcl

Symbols SC: subcutaneous IM: intramuscular

activated virus vaccines have been proposed and developed their inadequacy has been illustrated by the never ending attempts to develop a better prophylactic agent. The immunizing potential of some of these vaccines has been quite satisfactory and their value in the control of canine rabies has been proved. However other vaccines have been found to be lacking in antigenic activity⁴⁴⁻⁴⁶. There are still other disadvantages to the use of rabies vaccine of nervous-tissue origin. A significant number of nervous type reactions result from the administration of vaccines of this type. This is usually exhibited as a postvaccinal paralysis that may be mild and transient or so severe as to cause the death of the animal. A second disadvantage is the rather limited duration of the immunity conferred by one injection of vaccine containing inactivated virus. A one year period is most commonly accepted although immunity probably persists for a longer period in some animals and for a shorter time in others.

The last decade has been a period of marked progress in the field of rabies immunology and this is due primarily to the development of a modified live virus vaccine (Flury strain) grown in chicken embryos. Leach and Johnson²⁵ isolated this strain of virus from the tissues of a girl who had died of rabies. Johnson subsequently carried the virus through 136 intracerebral passages

TABLE 1
REPORTED CASES OF RABIES IN THE UNITED STATES 1947 to 1956⁴⁰

Year	Domestic animals	Other animals	Man
1947	8192	728	26
1948	7676	819	13
1949	6395	1192	10
1950	6526	1375	9
1951	6621	1387	14
1952	6758	1674	21
1953	7344	1479	14
1954	5577	1697	8
1955	3924	1915	5
1956	3457	2049	10

reduced in virulence by a multiple dilution method has been used extensively for immunizing domestic animals. The results of these immunizations were summarized by Aujeszky¹ who concluded that the vaccine was quite effective. Experimental work conducted by Meissner³⁶ in which fixed virus was administered to calves and sheep by various routes indicates that immunity was induced by several doses of vaccine containing virus from freshly harvested brain tissue material.

A group of vaccines that have been used extensively in the prophylactic antirabies treatment of man and animals are those containing inactivated virus. Since there are many vaccines of this type only a few will be mentioned. Phenolized vaccine first developed by Fermi^{9, 10} as a prophylactic agent for use in man has received wide application in domestic animals. The vaccine was modified by Semple⁴⁷ in 1911 and tested in animals prior to its extensive use in humans. Umeno and Doi⁴⁹ adopted this type of vaccine for use in immunizing dogs by a single injection technique and the results achieved from administering the vaccine to about 15 000 dogs were quite promising. In 1922 Eichorn and Lyon⁶ introduced the method into the United States where they used it experimentally. Based upon evidence presented by these investigators the vaccine was adopted for use by a number of health departments but it was necessary to alter the method of vaccine preparation slightly when Schoening⁴⁴ demonstrated the presence of live virus in it.

As research progressed other agents were employed for the inactivation of rabies virus in the preparation of vaccines. Kelser³ prepared an antirabies vaccine by treating fixed virus with chloroform. The efficacy of this vaccine was demonstrated by Kelser, by Schoening⁴⁵ and by Leach³⁴. A departure from chemical inactivation was made by Hodes, Lavin and Webster¹⁸ in 1937 when they used ultraviolet irradiation to inactivate fixed virus. Habel¹⁷ also prepared a vaccine of this type and reported on its potency as an immunizing agent. This vaccine was incorporated in duration of immunity studies conducted at the Communicable Disease Center Atlanta, Ga. and the results are indicated in TABLE 2. However the irradiated vaccines have never received extensive use in the field.

Although many modifications of the original Pasteur vaccine and the in

TABLE 2
SUMMARY COMPARATIVE CANINE RABIES VACCINE EXPERIMENTS²⁹

Vaccine	Route	Rabies mortality	Percent mortality	Period of incubation	Author
Brain tissue (phenolized) Controls	S.C. —	1/2 41/57	11.5 78.9	12 months	Johnson ²¹
Modified live virus (chick embryo) Brain tissue (phenolized) Controls	I.M. S.C. —	0/25 3/22 18/25	0 13.6 72	12 months	Kojiroski and Black ²⁰
Modified live virus (chick embryo) Brain tissue (phenolized) Controls	I.M. S.C. —	3/25 8/19 21/23	12.0 42.1 91.4	24 months	Koprowski and Black ²⁰
Modified live virus (chick embryo) Brain tissue (phenolized) Brain tissue (ultraviolet irradiated) Brain tissue (benzene ether treated) Controls	I.M. I.M. I.M. I.M. —	0/32 0/30 0/31 0/30 18/33	0 0 0 0 54.6	24 months	Tierkel, Kissling, Ellison and Habel
Modified live virus (chick embryo) Brain tissue (phenolized) Brain tissue (ultraviolet irradiated) Controls	I.M. I.M. I.M. —	0/30 8/34 7/30 31/36	0 23.6 23.3 86.1	39 months	Tierkel, Kissling, Ellison and Habel

Symbols S.C. subcutaneous I.M. intramuscular

activated virus vaccines have been proposed and developed their inadequacy has been illustrated by the never ending attempts to develop a better prophylactic agent. The immunizing potential of some of these vaccines has been quite satisfactory and their value in the control of canine rabies has been proved. However other vaccines have been found to be lacking in antigenic activity.²⁴⁻²⁸ There are still other disadvantages to the use of rabies vaccine of nervous-tissue origin. A significant number of nervous-type reactions result from the administration of vaccines of this type. This is usually exhibited as a postvaccinal paralysis that may be mild and transient or so severe as to cause the death of the animal. A second disadvantage is the rather limited duration of the immunity conferred by one injection of vaccine containing inactivated virus. A one year period is most commonly accepted although immunity probably persists for a longer period in some animals and for a shorter time in others.

The last decade has been a period of marked progress in the field of rabies immunology and this is due primarily to the development of a modified live virus vaccine (Flury strain) grown in chicken embryos. Leach and Johnson²⁵ isolated this strain of virus from the tissues of a girl who had died of rabies. Johnson subsequently carried the virus through 136 intracerebral passages

in day old chicks. After two additional passages in chicks Koprowski and Cox were successful in adapting the virus to the developing chicken embryo.²⁰ The serial passage was then continued in embryonated eggs with a significant change in the virulence of the virus noted between the fortieth and fiftieth egg passage levels. At this stage the virus demonstrated a reduced virulence for hamsters and guinea pigs and a loss of virulence for rabbits and dogs when administered parenterally. Further experimental work was conducted to determine whether or not the virus had retained its antigenicity. A group of dogs inoculated with the virus material and later challenged with virus of the NYC strain demonstrated a good immunogenic response to the virus.²¹ This finding was the basis for the production of vaccine prepared from the fortieth to fiftieth egg passage of the Flury strain virus. It is commonly known as low egg passage (LEP) modified live virus Flury strain and has been extensively tested in the laboratory and in the field.

The development of the Flury strain vaccine initiated extensive experimental activity. As with all new biological products the use of this vaccine both experimentally and in the field raised the usual questions concerning its safety and efficacy. A number of experiments were designed to demonstrate the relative effectiveness of the rabies vaccines commonly used in this country for protecting dogs against rabies. A summary of these studies is given in TABLE 2. In this table are included the results of the important challenge experiments conducted to evaluate rabies vaccines in dogs. A comparison of brain tissue vaccines with modified live virus vaccine Flury strain indicates that all vaccines tested induced some degree of resistance to challenge with street virus. However it appears that the modified live virus is a superior antigen. The 39 month duration of immunity study conducted by Tierkel *et al.*²² at the Communicable Disease Center shows quite conclusively that modified live virus vaccine Flury strain produces a significantly greater degree of immunity than do inactivated virus brain tissue vaccines 39 months after vaccination.

Studies concerning the efficacy of modified live virus rabies vaccine were not confined to the laboratory. Limited field trials had been conducted prior to 1950 when the Expert Committee on Rabies of the World Health Organization (WHO) Geneva Switzerland recommended a demonstration program of rabies control.⁷ This was to include the mass immunization of dogs with chicken embryo vaccine. The demonstration was carried out in the state of Israel where rabies had been a continuous problem for several years preceding 1950 despite control efforts. The mass immunization of dogs with modified live virus vaccine in conjunction with other control measures reduced the number of reported rabies cases to less than ten yearly during 1951, 1952 and 1953.²² The effectiveness of mass immunization of dogs with chicken embryo vaccine was again demonstrated in the Federation of Malaya. Although rabies had existed there since 1924 a federally sponsored rabies control program including mass canine immunization initiated in 1952 practically eliminated the disease from the country.

Modified live rabies virus has been in use in the state of New York since 1950. During that period more than 300,000 dogs have been administered

TABLE 3
EFFECTIVENESS OF RABIES VACCINES IN PUPPIES²⁰

Vaccine	Results of challenge	
	Ratio of challenge	Percentage protected
Modified live virus (LFP)	6/20	30
Brain tissue (phenolized)	6/20	30
Controls	11/11	100

Puppies 2 to 4 months of age at time of vaccination. The time between vaccination and challenge varied from 54 to 242 days.

the vaccine at canine vaccination clinics. The routine investigation of every reported case of canine rabies during that same period has revealed that only four of those dogs later succumbed to rabies despite the enzootic occurrence of the disease in the state.

The immunogenic response of puppies to the antigenic stimulus of rabies vaccine has long been questioned. The minimum age at which dogs might be successfully immunized against the disease was a matter of opinion. Koprowski²¹ conducted studies utilizing two- and three-month-old puppies that were administered LFP Flury strain vaccine but the findings were rather inconclusive. Kissling and his associates⁴ carried out an experiment in which two- to four-month-old puppies from nonimmune dams were administered rabies vaccines and were later challenged. The results of this study are shown in TABLE 3 and indicate that immunity was induced by the vaccines. However, the findings of the experiment hinted that puppies do not respond as well to an antigen as do adult dogs. In 1956 a study was conducted by state and CDC veterinarians in the New York State Department of Health Albany, N. Y., in cooperation with workers at Lederle Laboratories Division American Cyanamid Company, Pearl River, N. Y. Puppies ranging in age from 5 to 16 weeks were administered either LFP Flury or HEI Flury vaccine and five months later they were challenged with NYC strain virus. The results of the challenge are given in TABLE 4 while there was protection against exposure the degree of immunity was not of a level that might be expected in adult animals. In TABLE 5 the test animals are grouped by age at the time of vaccination and the results of challenge are shown. It is thus quite well demonstrated that puppies less than 11 weeks of age do not respond as well to vaccine

TABLE 4
RESPONSE OF YOUNG DOGS TO CHALLENGE

Vaccine	Results of challenge		
	Number tested	Number dying	Percentage
Modified live virus (LFP)	27	14	51.9
Modified live virus (HEI)	23	10	43.4
Controls	28	2	92.8

TABLE 5
EFFECTIVENESS OF RABIES VACCINE IN PUPPIES AT VARIOUS AGE LEVELS

Age at inoculation	Rabid mortality rate	Percent mortality
5 to 7 weeks	11/18	61.0
8 to 10 weeks	10/16	62.5
11 to 16 weeks	3/16	18.7
Control 5 to 16 weeks	26/28	93.0

as do older animals. In young animals there appears to be a maturation factor that must be present to permit the type of antigenic response found in older animals.

The occurrence of rabies in cats and the immunization of this species against the disease also has been a matter of study during the last decade. Information concerning the epidemiology of rabies in cats and the effectiveness of rabies vaccines in protecting these animals against the disease has accumulated gradually. Cats are probably unimportant vectors of the disease in nature due to a number of factors. Recent studies⁵ have shown that cats and kittens are more resistant than dogs and puppies to challenge with street virus by the peripheral route. Also the inherent nature of cats and their behavior probably serves to limit the possibility of their exposure. While this is the case it appears that the great majority of cats that develop rabies exhibit the furious form of the disease. The individual rabid cat therefore is a threat to man and other animals.

Laboratory studies concerned with the response of cats to rabies vaccines have demonstrated the effectiveness of these prophylactic agents. Koprowski⁷ vaccinated cats with IEP Flury vaccine and subjected these animals to challenge 39 days later. The results of that study show that the vaccine induced a significant degree of protection. In an experiment conducted in the state of New York⁸ it was shown that LEP Flury vaccine, HEP Flury vaccine, and phenolized brain tissue vaccine provide an antigenic stimulus to impart a significant degree of immunity 70 days after vaccination (TABLE 6). However, the results would indicate that the nervous tissue vaccine is probably

TABLE 6
EFFECTIVENESS OF ANTIRABIES VACCINES IN PROTECTING CATS AGAINST RABIES

Type	Dose	Route	Number of subjects	Survival	
				Number	Percent
Controls	—	—	26	4	15.4
Modified live virus (IEP)	1.5 ml	I.M.	26	18	69.2
Modified live virus (HFI)	1.5 ml	I.M.	24	18	75.0
Brain tissue (phenolized)	3.0 ml	S.C.	27	26	96.3

Symbl. I.M. intramuscular S.C. subcutaneous

a superior antigen in adult cats. Experiments conducted thus far have not provided data concerning the duration of immunity in cats.

The safety of rabies vaccines when administered to cats has also received consideration. Although the safety of brain tissue vaccines for felines has been little studied, phenol does not appear to be a serious problem.^{3, 4} Accumulated experimental data indicate that cats under six months of age should not receive modified live virus rabies vaccine. Young animals are apparently more susceptible, and an occasional kitten will succumb to LEP virus inoculated peripherally. Although this age factor exists, the maturation point after which greater resistance is demonstrated has not been determined in cats.

The occurrence of rabies in cattle and sheep has been a problem for many years, with the most emphasis placed upon prophylactic treatment of exposed animals. The epizootiology of rabies in cattle was little considered prior to outbreaks of vampire bat rabies in the cattle of Central and South America. Added impetus to these studies was given by the occurrence of bovine rabies in the United States caused by a spillover from the wildlife reservoir. Cattle are apparently quite susceptible to rabies virus. This is confirmed by observations of the high mortality following field exposures, and by the susceptibility to LEP Flury virus^{5, 21} and experimental challenge inoculations.^{14, 21} These findings have increased our knowledge of the disease in this species and brought further progress in rabies prophylaxis.

The greater part of the recent experimental work concerned with the immunization of cattle against rabies has been carried on in connection with the modified live virus vaccines. Since LEP Flury vaccine had proved to be so effective in the immunization of dogs, there existed the possibility of its use in cattle. Gomez Black and Koprowski¹² instituted a study in Colombia to determine the efficacy of LEP and chloroform inactivated rabies vaccine of nervous tissue origin (Kelser method). Extensive serum neutralization tests and challenge procedures were incorporated in this study, which demonstrated the ability of both vaccines to induce immunity in cattle. Shortly thereafter Schroeder and his colleagues¹⁶ conducted an extensive field study in Honduras, Costa Rica, and Guatemala, with particular emphasis on the immunization of cattle against vampire bat rabies. In these studies the immunogenic potency of LEP Flury vaccine was shown quite conclusively. However, field use of the vaccine demonstrated a susceptibility of cattle to this virus, for it was deemed to be the cause of paralysis and even of death in an occasional animal. Since its safety was questionable, the use of the vaccine in the bovine was contraindicated, and its use largely discontinued.

The search for a prophylactic agent that could be used successfully in cattle continued. HEP Flury strain vaccine was the next to be investigated. Continued serial passage of the Flury strain virus in embryonated chicken eggs had resulted in a further change in the virulence of the virus at egg passages 1/6 and 18.⁷ While suckling mice would still succumb to the virus when it was administered intracerebrally, it no longer would kill adult mice. The virus was also found to be apathogenic for rabbits and dogs by the same route. Since there was a possibility of using this for the immunization of cattle, Koprowski, Black, and Johnson²¹ conducted laboratory studies to ascertain the

TABLE 5
EFFECTIVENESS OF RABIES VACCINE IN PUPPIES AT VARIOUS AGE LEVELS

Age at vaccination	Rabid mortality	Percentage mortality
5 to 7 weeks	11/18	61.0
8 to 10 weeks	10/16	62.5
11 to 16 weeks	3/16	18.7
Controls 5 to 16 weeks	26/28	93.0

as do older animals. In young animals there appears to be a maturation factor that must be present to permit the type of antigenic response found in older animals.

The occurrence of rabies in cats and the immunization of this species against the disease also has been a matter of study during the last decade. Information concerning the epidemiology of rabies in cats and the effectiveness of rabies vaccines in protecting these animals against the disease has accumulated gradually. Cats are probably unimportant vectors of the disease in nature due to a number of factors. Recent studies⁵ have shown that cats and kittens are more resistant than dogs and puppies to challenge with street virus by the peripheral route. Also the inherent nature of cats and their behavior probably serves to limit the possibility of their exposure. While this is the case it appears that the great majority of cats that develop rabies exhibit the furious form of the disease. The individual rabid cat therefore is a threat to man and other animals.

Laboratory studies concerned with the response of cats to rabies vaccines have demonstrated the effectiveness of these prophylactic agents. Koprowski⁷ vaccinated cats with LEP Flury vaccine and subjected these animals to challenge 39 days later. The results of that study show that the vaccine induced a significant degree of protection. In an experiment conducted in the state of New York⁸ it was shown that LEP Flury vaccine, HEP Flury vaccine and phenolized brain tissue vaccine provide an antigenic stimulus to impart a significant degree of immunity 10 days after vaccination (TABLE 6). However the results would indicate that the nervous tissue vaccine is probably

TABLE 6
EFFECTIVENESS OF ANTIRABIES VACCINES IN PROTECTING CATS AGAINST RABIES

Type	V		Result		
	Dose	Route	Number tested	Survived	
				Number	Percentage
Controls	—	—	26	4	15.4
Modified live virus (LEP)	1.5 ml	IM	26	18	69.2
Modified live virus (HIP)	1.5 ml	IM	24	18	75.0
Brain tissue (phenolized)	3.0 ml	SC	27	26	96.3

Symbols IM intramuscular SC subcutaneous

in use for several years. Improvements in this method of assay were made to give more consistent results from the testing of inactivated virus vaccine. Among the accomplishments has been the adoption of a standard challenge virus for use in testing vaccine potency. Since the potency mouse test could not be applied to the modified live virus vaccines it was necessary to develop a potency guinea pig test to be used for this purpose. This was accomplished and has now been a standard method for several years. These tests have served to provide better vaccines for the immunization of domestic animals.

The past decade has thus been a period of great progress in the field of rabies prophylaxis in domestic animals. We now have tools that if properly used will serve to reduce greatly the incidence of the disease in this group of animals.

References

1. AJESKY A. 1927. Wutschutzimpfung der Haustiere in Ungarn. Deut. tierärztl. Wochschr. 35: 399. (Cited by Koprowski²⁴.)
2. BABEL V. & M. LEPP. 1939. Recherches sur la vaccination antirabique. Ann. inst. Pasteur 3: 384-390.
3. BAILEY J. 1940. 18th Session of OIE. Rept. No. 167.
4. CARNEIRO V. J. BLACK & H. KOPROWSKI. 1955. Rabies in cattle. V. Immunization of cattle in Brazil against exposure to street virus of vampire bat origin. J. Am. Vet. Med. Assoc. 127: 366-369.
5. DEAN D. J. Observations on anti-rabies vaccines. N. Y. State Health Dept. Unpublished data.
6. EICHORN A. & B. M. LYON. 1927. Prophylactic vaccination of dogs against rabies. J. Am. Vet. Med. Assoc. 61: 38-47.
7. EXPERT COMMITTEE ON RABIES. FIRST REPORT. 1940. World Health Organization. Tech. Rept. Ser. No. 23.
8. EXPERT COMMITTEE ON RABIES. THIRD REPORT. 1951. World Health Organization. Tech. Rept. Ser. No. 121.
9. FERMI C. 1909. Zentr. Bakteriell. Parasitenk. Abt. I. Ref. 49: 457. (Cited by Van Rooyen and Rhodes²⁵.)
10. FERMI C. 1909. Zentr. Bakteriell. Parasitenk. Abt. I. Ref. 52: 236. (Cited by Van Rooyen and Rhodes²⁵.)
11. GALTIER V. 1881. Les injections de virus rabique dans le torrent circulatoire ne provoquent pas l'éclosion de la rage et semblent conférer l'immunité. La rage peut être transmise par l'injection de la matière rabique. Compt. rend. 93: 234. (Cited by Koprowski²⁴.)
12. GALTIER V. 1883. Nouvelles expériences sur l'inoculation antirabique en vue de préserver les animaux herbivores de la rage à la suite des morsures de chiens enragés. Compt. rend. 106: 1189. (Cited by Koprowski²⁴.)
13. GOMEZ C. J. BLACK & H. KOPROWSKI. 1955. Rabies in cattle. III. Comparative studies on vaccination of cattle in Colombia with Flury virus and chloroform inactivatedacci. J. Am. Vet. Med. Assoc. 127: 360-363.
14. HABEL K. 1954. Antiserum in the prophylaxis of rabies. Bull. World Health Organization 10: 81-89.
15. HABEL K. 1945. Seroprophylaxis in experimental rabies. U. S. Public Health Repts. 60: 545-560.
16. HABEL K. & J. T. WRIGHT. 1948. Some factors influencing mouse potency test for rabies vaccine. U. S. Public Health Repts. 63: 44-6.
17. HABEL K. 194. Ultraviolet irradiation in production of potent antirabies vaccines. U. S. Public Health Repts. 62: 91-800.
18. HUNDS H. G. LAYN & L. T. WEBSTER. 1934. Antirabic immunization with culture virus under daylight and ultra violet light. Science 86: 44-445.
19. HUNT A. & M. GURLEY. 1934. Experimental street virus rabies in white mice: studies on passive immunization. Proc. Soc. Exptl. Biol. Med. 37: 454.
20. HUNT A. & M. GURLEY. 1938. Experimental street virus rabies in white mice: studies on passive immunization. Proc. Soc. Exptl. Biol. Med. 33: 40.
21. JOHNSON H. N. 1945. Experimental and field studies of canine rabies vaccination. Proc. U. S. Livestock San. Assoc. 99-107.

immunogenic potency of this virus in cattle. This together with field studies⁴ served to evaluate the vaccine in this species. The value of this antigen for immunizing cattle has thus been shown and the vaccine is now used quite commonly for this purpose in many areas of the world.

The postexposure treatment of domestic animals is a practice that has been carried on for many years. In fact, the earlier research was concerned primarily with the use of prophylactic agents following exposure. This continued to be the case until the advent of mass canine vaccination. Vaccines have been used widely in a number of European countries as post exposure treatment. Although a substantial amount of information has accumulated from the administration of many types of vaccines, it is difficult to draw conclusions due to the nature of the disease and the type of data available.

The possibility of inducing passive immunity by injections of antirabic serum was proposed by Babes and Lepp in 1889. The use of rabies antiserum was not commonly practiced during the ensuing years since its efficacy was questioned. However Habel¹⁴ reports that during this period a number of workers including Marie Fermi Proca and Shortt were investigating its value as a prophylactic agent. The studies of Hoyt and Gorley^{19, 20} and of Habel¹⁵ in which seroprophylaxis was used to immunize mice served to revitalize efforts concerned with this approach. Additional experiments with guinea pigs demonstrated the value of antiserum in protecting laboratory animals. Koprowski has conducted extensive studies with hyperimmune serum in laboratory animals and dogs in the development of a product for human use. Although the preventive effect of antirabies serum has been shown, it is not generally used for treatment of animals. This is due to its cost and to the conditions under which many exposures of domestic animals occur.

The proper handling of dogs and cats exposed to the disease by a known rabid animal has long been a problem. It has been a usual practice to destroy such animals rather than to risk their acquiring the clinical disease and exposing other animals and possibly human beings as well. The alternative was to hold the animal in strict isolation for a period of 6 months. Recognition has now been given to postexposure treatment by using antirabies serum followed by 1 to 3 doses of chicken embryo vaccine or 14 injections of nervous tissue vaccine.⁸ If the animal has been vaccinated within 1 year with nervous tissue or Kelev strain vaccine or within 3 years with LEP Flury vaccine, it is acceptable procedure to revaccinate the animal and place it in quarantine for 30 days. This serves to indicate the faith placed in rabies vaccines.

An inactivated rabies vaccine of duck embryo origin was recently released for use in humans. Clinical and laboratory results reported by Peck, Powell and Culbertson¹⁶ indicate that the antigenic stimulus provided by this vaccine is quite satisfactory. It would seem that a vaccine of this type may have indications for use in domestic animals. Additional experimental work will be required to determine the efficacy of the vaccine in these species.

The development and use of rabies vaccines have been discussed, but marked progress has also been made in the potency testing of vaccines. In the past decade Habel and Wright¹⁷ modified the potency mouse test that had been

in use for several years. Improvements in this method of assay were made to give more consistent results from the testing of inactivated virus vaccine. Among the accomplishments has been the adoption of a standard challenge virus for use in testing vaccine potency. Since the potency mouse test could not be applied to the modified live virus vaccines it was necessary to develop a potency guinea pig test to be used for this purpose. This was accomplished and has now been a standard method for several years. These tests have served to provide better vaccines for the immunization of domestic animals.

The past decade has thus been a period of great progress in the field of rabies prophylaxis in domestic animals. We now have tools that if properly used will serve to reduce greatly the incidence of the disease in this group of animals.

References

1. AUJESKY A. 1927. Wutschutzimpfung der Haustiere in Ungarn. D-ut tierarztl Wochschr 35 399 (Cited by Koprowski¹⁴).
2. BABES V. & M. LEPP. 1889. Recherches sur la vaccination antirabique. Ann inst Pasteur 3 384-390.
3. BAILEY J. 1950. 18th Session of OIE. Rept No 167.
4. CARNEIRO V. J. BLACK & H. KOPROWSKI. 1955. Rabies in cattle. V. Immunization of cattle in Brazil against exposure to street virus of vampire bat origin. J Am Vet Med Assoc 127 366-399.
5. DEAN D. J. Observations on anti rabies vaccines. N. Y. State Health Dept. Unpublished data.
6. FICHORN A. & B. M. LYON. 1922. Prophylactic vaccination of dogs against rabies. J Am Vet Med Assoc 61 38-42.
7. EXPERT COMMITTEE ON RABIES. FIRST REPORT. 1950. World Health Organization. Tech Rept Ser No 28.
8. EXPERT COMMITTEE ON RABIES. THIRD REPORT. 1957. World Health Organization. Tech Rept Ser No 121.
9. FERMI C. 1909. Zentr. Bakteriell. Parasitenk. Abt. I. Ref. 49 452 (Cited by Van Rooyen and Rhodes).
10. FERMI C. 1909. Zentr. Bakteriell. Parasitenk. Abt. I. Ref. 52 536 (Cited by Van Rooyen and Rhodes).
11. GALTIER V. 1881. Les injections de virus rabique dans le torrent circulatoire ne provoquent pas l'eclosion de la rage et semblent conférer l'immunité. La rage peut être transmise par l'ingestion de la matière rabique. Compt. rend. 93 284 (Cited by Koprowski).
12. GALTIER V. 1888. Nouvelles expériences sur l'inoculation antirabique en vac de préserver les animaux herbivores de la rage à la suite des morsures de chiens enragés. Compt. rend. 106 1189 (Cited by Koprowski).
13. GOMEZ C. J. BLACK & H. KOPROWSKI. 1955. Rabies in cattle. III. Comparative studies on vaccination of cattle in Colombia with Flu. J. virus and chlor. form inactivated vaccine. J Am Vet Med Assoc 127 360-363.
14. HABEL K. 1954. Anticrum in the prophylaxis of rabies. Bull. World Health Organ. 10 781-788.
15. HABEL K. 1945. Seroprophylaxis in experimental rabies. U. S. Public Health Repts. 60 545-560.
16. HABEL K. & J. T. WRIGHT. 1948. Some factors influencing mouse potency test for rabies vaccine. U. S. Public Health Repts. 63 44-65.
17. HABEL K. 1947. Ultraviolet radiation in production of potent antirabies vaccines. U. S. Public Health Repts. 62 991-800.
18. HODES H. G. LAVIN L. T. WEBSTER. 1937. Antibody immunization with culture virus rendered avirulent by ultraviolet light. Science 86 447-448.
19. HOYT A. & M. GURLEY. 1937. Experimental street virus rabies in white mice studies on passive immunization. Proc. Soc. Exptl. Biol. Med. 37 454.
20. HOYT A. & M. GURLEY. 1938. Experimental street virus rabies in white mice studies on passive immunization. Proc. Soc. Exptl. Biol. Med. 38 40.
21. JOHNSON H. N. 1945. Experimental and field studies of canine rabies vaccination. Proc. U. S. Livestock San. Assoc. 99-107.

- 22 KAPLAN M Y GOOR & C S TIERKEL 1954 A field demonstration of rabies control using chicken embryo vaccine in dogs Bull World Health Organization 10 743-752
- 23 KELSER R A 1930 Chloroform treated rabies vaccine J Am Vet Med Assoc 77 593-603
- 24 KELSER R A & H W SCHOENING 1943 Manual of Bacteriology 45th ed 508-528 Williams & Wilkins Baltimore Md
- 25 KISSLING R F M E LIDSON D D STAMM & E S TIERKEL Personal communication
- 26 KOPROWSKI H 1925 Rabies in cattle II Review of immunization of herbivorous animals against rabies J Am Vet Med Assoc 127 359-360
- 27 KOPROWSKI H Personal communication
- 28 KOPROWSKI H & J BLACK 1950 Studies on chick embryo adapted rabies virus pathogenicity for dogs and use of egg adapted strains for vaccination purposes J Immunol 64 185-196
- 29 KOPROWSKI H & J BLACK 1952 Studies on chick embryo adapted virus III Duration of immunity in vaccinated dogs Proc Soc Exptl Biol Med 80 410-415
- 30 KOPROWSKI H & J BLACK 1954 Studies on chick embryo adapted rabies virus protection of animals with antiserum and living, attenuated virus after exposure to street strain of rabies virus J Immunol 72 85-93
- 31 KOPROWSKI H J BLACK & W JOHNSON 1955 Rabies in cattle IV Vaccination of cattle with high egg passage chicken embryo adapted rabies virus J Am Vet Med Assoc 127 363-366
- 32 KOPROWSKI H & H K COX 1948 Studies on chick embryo adapted rabies virus I Culture characteristics and pathogenicity J Immunol 60 533-554
- 33 KRASNITSKI V 1902 Immunisation antirabique au moyen des injections intravasculaires du virus rabique Ann inst Pasteur 16 393 (Cited by Koprowski²⁴)
- 34 LEACH C N & H N JOHNSON 1940 Canine rabies vaccination experimental study of efficacy of single subcutaneous injection method with chloroform treated vaccine Am J Hyg 32 14-79
- 35 LEACH C N & H N JOHNSON 1940 Human rabies with special reference to virus distribution and titer Am J Trop Med 20 335-340
- 36 MEISSNER I 1912 Über Tollwutchutzimpfung bei Tieren Zentr Bakteriol Parasitenk 54 13 (Cited by Koprowski²⁴)
- 37 MONCET M 1898 Immunisation des herbivores contre la rage Rev Vet 5 291 (Cited by Koprowski²⁴)
- 38 PASTEUR I 1886 Resultats de l'application de la methode pour prevenir la rage apres morsure Compt rend 102 459-466
- 39 PASTEUR L C F CHAMBERLAND & M ROUX 1884 Nouvelle communication sur la rage Compt rend 98 457-463
- 40 PECK I B H M POWELL & C C CULBERTSON 1955 A new antirabies vaccine for human use clinical and laboratory results using rabies vaccine made from embryonated duck eggs J Lab Clin Med 45 619-683
- 41 ROUX M & M NOCARD 1884 Expériences sur la vaccination des ruminants contre la rage par injections intraveineuses de virus rabique Ann inst Pasteur 2 341 (Cited by Koprowski²⁴)
- 42 SCHREIBY S F J F SKELLEY & H W KULP 1953 Thiolized rabies vaccines in cats J Am Vet Med Assoc 123 330
- 43 SCHNITZER J 1905 Z Hyg Infektionskrankh 61 46 (Cited by Vallee⁴¹)
- 44 SCHOENING H W 1930 Experimental studies with killed canine rabies vaccines J Am Vet Med Assoc 76 25-33
- 45 SCHOENING H W 1925 Studies on the single injection method of vaccination as a prophylactic against rabies in dogs J Agr Research 30 431-439
- 46 SCHROEDER C R J BLACK I I BLUMHART & H KOPROWSKI 1952 Rabies in cattle I Prevention of vampire bat paralytic rabies berriengue by vaccination with chick embryo adapted rabies virus Vet Med 47 507-506
- 47 SMYTH D 1911 Sci Mem Med Sanit Dept India No 44
- 48 TIERKEL E S R F KILG M LIDSON & K HABEL 1953 A brief survey and progress report of controlled comparative experiment in canine rabies immunization Proc Book Am Vet Med Assoc 443-445
- 49 UENO & DOI 1921 Kitasato Arch Exptl Med 4(2) 89 (Cited by Kelsner)

- 50 UNITED STATES DEPARTMENT OF AGRICULTURE 1957 Animal Disease Fra lication Branch Notice Reported Incidence of Ral ies in the United States February 14 1957
- 51 VALLÉE H 1978 La vaccination antirabique chez les animaux Report to the International Rabies Conference Ann in t Pasteur Suppl 41 144-163
- 52 VALLERY RADOT R 1912 (Translated by R L Devonshire) The Life of Pasteur 390-436 Garden City Publ Co Garden City N Y
- 53 VAN ROOYEN C & A RHODES 1948 Virus Diseases of Man 844-875 Nelson & Sons New York N Y
- 54 WEBSTER L T 1939 Immunizing potency of antirabies vaccines critical review Am J Hyg 30 113-134
- 55 WEBSTER L T 1939 Mouse test for measuring immunizing potency of antirabies vaccines J Exptl Med 70 87-106
- 56 WEBSTER L T 1940 Mouse test for measuring immunizing potency of antirabies vaccines J Am Vet Med Assoc 96 62-13

Discussion of the Paper

ERNEST S TIERKEL (*Communicable Disease Center Public Health Service Atlanta Ga*) It seems entirely appropriate at this point to illustrate by means of field experience in rabies control the effects of this research in animal vaccination Two examples of the many successful immunization programs of the last ten years may be cited here Rabies in Memphis Tenn had been at a substantially high simmering level for years when it exploded into epidemic proportions in the early spring of 1948 despite an excellent program for the collection of stray dogs The organization of an intensified mass canine immunization campaign began swiftly In a single week in mid April 23 000 dogs were vaccinated in 70 emergency inoculation clinics strategically located throughout the city and county In the ensuing months positive reports began to drop until the last case of animal rabies and the last human anti rabies treatment were reported in July (TABLE 1) The elimination of rabies from both city and county was a new and refreshing experience for Memphis

A similar campaign was conducted in Houston Texas in the fall of 1954 From 1945 to 1954 Houston had been plagued by an annual average of 300 animal cases with 14 human deaths During 1953 and 1954 the animal cases rose to an average of 40 per month In the last days of September 1954 an intensified 4-day program was carried out in which 45 000 dogs were vaccinated in 91 emergency clinics throughout the city and county Here again the success

TABLE 1
ANIMAL RABIES CASES IN MEMPHIS TENN 1948

January	13
February	24
March	41
April	28—Mass immunization campaign
May	5
June	2
July	2
August	0
September	0
October	0
November	0
December	0

TABLE 2
ANIMAL RABIES CASES IN HOUSTON, TEXAS

1934	September	34—Mass immunization campaign
	October	33
	November	25
	December	17
1935	January	21
	February	15
	March	18
	April	13
	May	9
	June	14
	July	2
	August	3
	September	1
	October	0

of this program can be seen by the gradual decrease in cases from October 1934 to October 1935 (TABLE 2)

Progress achieved in the over all national incidence of rabies during the past decade is reflected in the annual reported cases, as shown in TABLE 3. The total number of cases in 1946 was 10,872. This declined steadily to 5,844 cases for 1955—a decrease of one half. By the same token the human cases dropped from 22 in 1946 to 5 in 1955—a decrease of 77 per cent. Although in most parts of the country the dog still accounts for most of the cases and remains the most important vector of the disease, the striking decline in canine cases from 8,384 in 1946 to 2,657 in 1955 (68 per cent) is eloquent testimony to our efforts to control the disease during the past 10 years.

We now come to measures taken for the prevention of rabies in man. Over the years one of the problems in the specific biological therapy of exposed persons was the ineffectiveness of vaccine in cases of severe bite, especially those involving the region of the head, when the incubation period was too short to allow for the development of immunospecific protection. We have now largely solved this problem by the production of hyperimmune rabies antiserum which can confer sufficient passive immunity during that critical early stage before the vaccine can stimulate active protection. Convincing experimental evidence from challenge tested laboratory animals (Habel 1945; Koprowski *et al.* 1950) and from antibody studies (Atanasiu *et al.* 1956) and natural exposure studies in man (Baltazard and Bahmanyar 1955) have established the value of the antiserum when it is followed by the usual course

TABLE 3
REPORTED CASES OF RABIES IN THE UNITED STATES

Year	Dog	Wild	Cat	Livestock	Man	Total
1946	8,384	956	455	1,055	22	10,872
1955	2,657	1,915	343	924	5	5,844

of vaccine. One serum injection of 0.5 ml/kg of body weight is given as soon as possible after the bite. Since this is a horse serum product precautions should be taken to avoid anaphylactic reactions by testing the patient for sensitivity. Antihistamines, ACTH or cortisone seem to be useful when serum sickness occurs.

The extent of the over all rabies problem in the United States can be measured by the 50 000 human exposures occurring annually each of which requires the expensive painfully long series of 14 to 21 daily vaccine treatments a therapeutic prophylactic regime requiring greater professional medical care than many other infectious diseases. An important phase of this problem is the occasional occurrence in exposed persons of postvaccinal encephalitic, neurologic, and paralytic complications following administration of the prescribed series of rabies vaccine treatments. These reactions are believed to be tissue specific isoallergic phenomena. Apparently they are related not to the rabies antigen present but rather to some encephalitogenic component of the nerve tissue in the vaccine. They occur most frequently after the seventh inoculation and most often in persons who have had previous Pasteur treatment. Reports on the frequency of these complications vary for example 1 in 527 (Cook *et al.* 1955) 1 in 600 (Pait and Pearson 1949) 1 in 7025 (Appelbaum *et al.* 1953) 1 in 7200 (Sellers 1948) 1 in 8500 (McKendrick 1940). Experiments are currently proceeding in order to find new approaches that will lead to an effective vaccination regimen without danger of damage to the central nervous system.

References

- APPLEBAUM E. M. GREENBERG & J. NELSON. 1953. Neurological complications following anti rabies vaccination. *J. Am. Med. Assoc.* 161: 188.
- ATANASIU P. M. BAHMANYAR M. BALTAZARD J. P. FOY K. HABEL M. M. KAPLAN R. F. KISSLING A. KOMAROV H. KOPROWSKI P. LEPIVÉ F. PEREZ GALLARDO & M. SCHAEFFER. 1956. Rabies neutralizing antibody response to different schedules of serum and vaccination inoculations in non exposed persons. *Bull. World Health Organization* 14: 593.
- BALTAZARD M. & M. BAHMANYAR. 1955. Essai pratique du serum antirabique chez les chiens par loups enragés. *Bull. World Health Organization* 13: 747.
- COOK E. B. M. C. STEARNS J. FIELD & J. V. IRONS. 1955. Report on the use of phenolized vaccine in Texas from 1949 through 1953. *Texas Rept. Biol. and Med.* 13: 234-250.
- HABEL K. 1945. Seroprophylaxis in experimental rabies. *U.S. Public Health Repts.* 60: 545.
- KOPROWSKI H. & H. R. COX. 1948. Studies in chick embryo adapted rabies virus. I. Cultural characteristics and pathogenicity. *J. Immunol.* 60: 535-554.
- KOPROWSKI H. J. VAN DER SCHEER & J. BLANK. 1950. Use of hyperimmune serum concentrates in experimental rabies. *Am. J. Med.* 8: 412.
- McKENDRICK A. G. 1940-1941. A ninth analytical review of reports from Pasteur Institutes on the results of anti rabies treatment. *Bull. World Health Organization* 9: 31.
- PAIT C. F. & H. E. PEARSON. 1949. Rabies vaccine encephalomyelitis in relation to the incidence of animal rabies in Los Angeles. *Am. J. Public Health* 39: 375.
- SELLERS T. F. 1948. Rabies the physician's dilemma. *Am. J. Trop. Med.* 28: 453-456.
- TIERKEL E. S. R. E. KISSLING M. C. EDSON & K. HABEL. 1953. A brief survey and progress report of controlled comparative experiments in canine rabies immunization. 90th Ann. Meeting Proc. Am. Vet. Med. Assoc. 443-445.
- TIERKEL E. S. 1955. The present status of rabies control. Editorial. *Am. J. Public Health* 45: 677-680.
- TIERKEL E. S. 1956. Antibody studies in persons vaccinated with HEP Thury vaccine. *World Health Organization Rabies* 94.

TABLE 2
ANIMAL RABIES CASES IN HOUSTON TEXAS

1954	September	34—Mass immunization campaign
	October	33
	November	25
	December	17
1955	January	21
	February	15
	March	18
	April	13
	May	9
	June	14
	July	2
	August	3
	September	1
	October	0

of this program can be seen by the gradual decrease in cases from October 1954 to October 1955 (TABLE 2)

Progress achieved in the over all national incidence of rabies during the past decade is reflected in the annual reported cases as shown in TABLE 3. The total number of cases in 1946 was 10 872. This declined steadily to 5 844 cases for 1955, a decrease of one half. By the same token the human cases dropped from 22 in 1946 to 5 in 1955, a decrease of 77 per cent. Although in most parts of the country the dog still accounts for most of the cases and remains the most important vector of the disease, the striking decline in canine cases from 8384 in 1946 to 2657 in 1955 (68 per cent) is eloquent testimony to our efforts to control the disease during the past 10 years.

We now come to measures taken for the prevention of rabies in man. Over the years one of the problems in the specific biological therapy of exposed persons was the ineffectiveness of vaccine in cases of severe bite especially those involving the region of the head when the incubation period was too short to allow for the development of immunospecific protection. We have now largely solved this problem by the production of hyperimmune rabies antiserum which can confer sufficient passive immunity during that critical early stage before the vaccine can stimulate active protection. Convincing experimental evidence from challenge tested laboratory animals (Habel 1945; Koprowski *et al.* 1950) and from antibody studies (Atanasiu *et al.* 1956) and natural exposure studies in man (Baltazard and Bahmanyar 1955) have established the value of the antiserum when it is followed by the usual course

TABLE 3
REPORTED CASES OF RABIES IN THE UNITED STATES

Y	Dog	Wild	Cat	Livestock	Man	Total
1946	8 384	156	455	1 055	22	10 872
1955	2 657	1 915	343	924	5	5 844

of vaccine. One serum injection of 0.5 ml/kg of body weight is given as soon as possible after the bite. Since this is a horse serum product precautions should be taken to avoid anaphylactic reactions by testing the patient for sensitivity. Antihistamines, ACTH or cortisone seem to be useful when serum sickness occurs.

The extent of the over all rabies problem in the United States can be measured by the 50 000 human exposures occurring annually each of which requires the expensive painfully long series of 14 to 21 daily vaccine treatments a therapeutic prophylactic regime requiring greater professional medical care than many other infectious diseases. An important phase of this problem is the occasional occurrence in exposed persons of postvaccinal encephalitic neurologic and paralytic complications following administration of the prescribed series of rabies vaccine treatments. These reactions are believed to be tissue specific isoallergic phenomena. Apparently they are related not to the rabies antigen present but rather to some encephalitogenic component of the nerve tissue in the vaccine. They occur most frequently after the seventh inoculation and most often in persons who have had previous Pasteur treatment. Reports on the frequency of these complications vary for example 1 in 527 (Cook *et al.* 1955) 1 in 600 (Pait and Pearson 1949) 1 in 2025 (Appelbaum *et al.* 1953) 1 in 1200 (Sellers 1948) 1 in 8500 (McKendrick 1940). Experiments are currently proceeding in order to find new approaches that will lead to an effective vaccination regimen without danger of damage to the central nervous system.

References

- APPELBAUM E. M. GREENBERG & J. NELSON. 1953. Neurological complications following anti rabies vaccination. *J. Am. Med. Assoc.* 161: 188.
- ATANASIU P. M. BAHMANYAR M. BALTAZARD J. P. FOX K. HABEL M. M. KAPLAN R. F. KISSLING A. KOMAROV H. KOPROWSKI P. LEPINE F. PEREZ GALLARDO & M. SCHAEFFER. 1956. Rabies neutralizing antibody response to different schedules of serum and vaccination inoculations in non exposed persons. *Bull. World Health Organization* 14: 593.
- BALTAZARD M. & M. BAHMANYAR. 1955. Essai pratique du serum antirabique chez les moutons par jous enrages. *Bull. World Health Organization* 13: 747.
- COOK E. B. M. C. STEARNS J. FIELD & J. V. IRONS. 1955. Report on the use of phenolized vaccine in Texas from 1949 through 1953. *Texas Rept. Biol. and Med.* 13: 234-250.
- HABEL K. 1945. Seroprophylaxis in experimental rabies. *U. S. Public Health Repts.* 60: 545.
- KOPROWSKI H. & H. R. COX. 1948. Studies in chick embryo adapted rabies virus. I. Cultural characteristics and pathogenicity. *J. Immunol.* 60: 535-554.
- KOPROWSKI H. J. VAN DER SCHEER & J. BLACK. 1950. Use of hyperimmune serum concentrates in experimental rabies. *Am. J. Med.* 8: 412.
- McKENDRICK A. G. 1940-1941. A ninth analytical review of reports from Pasteur Institutes on the results of anti rabies treatment. *Bull. World Health Organization* 9: 31.
- PAIT C. F. & H. E. PEARSON. 1949. Rabies vaccine encephalomyelitis in relation to the incidence of an fatal rabies in Los Angeles. *Am. J. Public Health* 39: 375.
- SELLERS T. F. 1948. Rabies the physician's dilemma. *Am. J. Trop. Med.* 28: 453-456.
- TIERKEL E. S. R. E. KISSLING M. E. FIDSON & K. HABEL. 1953. A brief survey and progress report of controlled comparative experiments in canine rabies immunization. 90th Ann. Meeting Proc. Am. Vet. Med. Assoc. 443-445.
- TIERKEL F. S. 1955. The present status of rabies control. Editorial. *Am. J. Public Health* 45: 677-680.
- TIERKEL F. S. 1956. Antibody studies in persons vaccinated with HFP Flury vaccine. *World Health Organization. Rabies* 94.

PROPHYLAXIS AGAINST RABIES IN HUMANS

By John P. Fox

*Division of Epidemiology Department of Tropical Medicine and Public Health
Tulane University School of Medicine New Orleans La*

Measures intended to prevent rabies in man may be divided into two basic categories—those taken before and those taken after the infecting bite. Historically, since the ancient realization of the relation of dogs to human rabies, the first category has embraced measures to avoid being bitten and, since the early Nineteenth Century, measures to control canine rabies. In the second category, local cleansing treatment of the bite wound has been practiced for centuries, acid treatment since 1830¹. Specific active immunization following suspected exposure has been practiced since the time of Louis Pasteur. With two important exceptions that will be indicated shortly, the foregoing states in broad outline the situation today with respect to the orientation of efforts to prevent human rabies.

Important refinements of course have been introduced in the years since Pasteur, but without change in this basic orientation. These include the development and widespread utilization of canine immunization as an aid to control, improvements in methods of diagnosing animal infections as an aid to defining more certainly the individuals at risk, and development of a reliable method² for measuring vaccine potency to help ensure effective immunization. Despite these important technical advances, completely safe and reliable protection of man against rabies has yet to be achieved.

Three recent developments, however, afford the basis for renewed hope. The first represents another example of technical advance and consists of the possible substitution of vaccine prepared in avian embryos—chicken^{3,4} or duck^{5,7}—for the conventional types of vaccine that are prepared from central nervous system (CNS) tissue of infected animals. Such a vaccine, being essentially free of CNS tissue, should not induce the tissue-specific sensitization postulated as the cause of the demyelinating reactions that too often result from the Pasteur treatment. The remaining two recent developments represent the application of new principles or orientation. One, the introduction of hyperimmune serum⁸, belongs in the after-the-bite category and is intended to protect the recipient by passively conferred antibody against a disease with such a short incubation period that active response would be too long delayed. The other, which technically is still in the developmental stage, falls chiefly in the before-the-bite category and consists of the demonstration that once induced, active immunity resulting from either conventional Pasteur treatment or a primary course of avian embryo vaccine can be recalled by a single booster inoculum^{4,5}. With embryo vaccine this provides the basis for possible safe, long-term immunization initiated well before exposure in high-risk population groups. It also permits abbreviated (single inoculation) and presumably safe postexposure treatment of those who previously have had a course of Pasteur treatment.

This presentation will endeavor to describe briefly and clearly the salient

features of human antirabies prophylaxis at present. The major effort however will be reserved for the three important and most recent developments just mentioned. Those readers desiring a more detailed and fully balanced review of the general problem are referred to an excellent and very recent paper by Karl Habel.⁹

The Situation After the Bite

Presentation of the new information in proper developmental sequence makes it necessary to ignore the obvious logical order and to begin with the situation that confronts us after the bite has been inflicted.

Determination of exposure. Because of cost inconvenience and actual hazard related to unnecessary treatment an essential first step is to determine the likelihood of exposure. The first question is whether the saliva of the animal was in contact with freshly abraded or lacerated skin. To answer this one must determine whether a recent abrasion existed, whether the wound was really a bite rather than a scratch with the paw and if so whether the skin was really broken. Next one must determine whether the saliva actually was infectious. If the animal is not available for observation (that is if it escaped or cannot be identified) the presumption of infectiousness must prevail and specific treatment must be instituted. Otherwise except in instances of very severe exposure that may be associated with a short incubation period specific treatment may await the outcome of diagnostic procedures including observation of the animal.

Since Negri bodies (the detection of which is the most rapid diagnostic procedure) become more numerous and more certainly detectable as disease progresses diagnosis ordinarily is hastened by delaying sacrifice of the animal until signs of rabies are well developed. Mouse inoculation which requires an additional 6 to 14 days is indicated when contact has been well defined and Negri bodies are not found since of all infected animals the brains of about 12 per cent may be Negri negative.¹⁰ While demonstration of Negri bodies or rabies virus in the brain proves that the animal was infected it does not follow necessarily that the saliva at the time of the exposure contact was infectious. Not only does rabies virus completely fail to reach the salivary glands in many instances (in 25 per cent in one series of 28 proved canine infections¹¹) but it is not present in any case for more than a few days before obvious signs of disease appear. The available evidence is undesirably scanty in view of the crucial importance of this point but it suggests that appearance of virus in the saliva is always associated with early signs of illness (that is at least a slight rise in temperature) and is usually followed within no more than 5 days by clearly recognizable illness.¹¹ The practical implication of this is that *if the animal is still well 5 to 7 days after having bitten there was no exposure to rabies* even if disease becomes evident in the next day or two. One important exception to the foregoing relates to bats which should be captured and sacrificed immediately for virologic tests since the outcome of infection is unpredictable in these species.

Treatment of the bite wound. The fundamental consideration here is that

rabies virus may remain localized at the site of introduction for as long as twenty four hours. Treatment is directed at destroying or eliminating the virus and to the extent that it is successful it may completely prevent infection or so reduce the size of the infecting dose that the incubation period is extended. This simple but very important procedure should be carried out in all instances of animal bite without regard to the probability of rabies exposure.

Modern studies agree that the classic method of acid cautery is quite effective, especially when carried out within the first few hours after the bite. However other and less painful and disfiguring methods also are nearly as effective under experimental conditions, these include irrigation and scrubbing with soap or detergent solutions¹²⁻¹³. Recent observations by Perez Gallardo¹³ confirm the earlier work of Habel¹⁴ and indicate that local infiltration of antiserum is of real value.

It must not be forgotten that bites may be infected with agents other than rabies virus and that appropriate measures should be taken to combat these (that is antitetanus and antibacterial precautions).

Passive immunity. Active response to antirabies vaccine requires time in order that effective levels of antibody may be achieved (perhaps 20 days from initiation of treatment) and obviously cannot be expected to prevent a disease that occurs after unusually short periods of incubation as is commonly the case in severely exposed persons. The obvious remedy for this situation is to supply the needed antibody at once by passive means. Perhaps because of the inconclusiveness of early observations¹⁵⁻¹⁶ until recently there has been little sustained interest in this approach. Experimental work demonstrated the high degree of protection afforded animals by systemically and even locally administered antiserum¹³⁻¹⁴. Somewhat later, the importance of early use of serum (within seventy two hours) was demonstrated and it was shown also that the maximum protection of animals requires antiserum supplemented by vaccine⁸⁻¹⁷.

Since 1948 partially purified hyperimmune antirabies serum has been utilized on an experimental basis in man. In Georgia its use was restricted to persons suffering severe and certain exposure less than 72 hours before seeking treatment. By early 1953 68 persons none of whom developed rabies had received antiserum. In each case a course of antirabic vaccine also was begun after injecting the serum¹⁸. In 1954 the opportunity finally occurred to carry out a controlled antiserum trial that had been planned since 1950 by the World Health Organization (WHO) Expert Committee on Rabies. A rabid wolf in Iran attacked and bit 27 persons of whom 17 were severely exposed by bites on the head. These latter were divided into 3 groups for treatment: the groups receiving respectively only vaccine or vaccine plus 1 dose of serum or vaccine plus 2 doses of serum (the second dose given after a 5-day interval). The rabies mortality ratios for these 3 groups were 3.5, 1.6 and 0.6 respectively¹⁹. Sera were collected at frequent intervals from these persons²⁰ and on an experimental basis²¹⁻²² from unexposed volunteers receiving vaccine alone or with serum and have been tested for neutralizing antibody. The resulting data indicate that even a single dose of serum coupled with a full

course of vaccine ordinarily results in the continuous presence of antibody during the 50- or 60 day period of observation after injection of the serum. However especially with marginally effective vaccination active response may be partially or even completely suppressed because of the antiserum this suppressive influence being more pronounced with 2 doses of serum than with a single dose. Hence while the mortality experience in Iran was better (though not significantly so) in the group given 2 doses of serum it would seem safer on theoretical grounds to give but a single dose of serum and to couple it with vaccine of well established antigenic effect.

Antibody solution of equine origin is now generally available and should be used in all instances of severe exposure. In instances where rabies infection of the biting animal is not clearly evident vaccine can be deferred pending the outcome of the usual observation procedures in all other cases serum should be followed promptly by the vaccine. Use of serum in instances of less serious exposure is not mandatory and in my own opinion should be discouraged because of the relatively great frequency of serum sickness observed.

Idiote immunization (Pasteur treatment) If circumstances make it necessary to assume that exposure has occurred the use of antirabic vaccine becomes mandatory. Unique among vaccines in that it is used *after* exposure has occurred its rationale depends on the stimulation of active response in time to prevent the usually slowly incubating infection from involving vital areas of the central nervous system (CNS). Since in the individual instance the incubation period cannot be predicted reliably and may be short rapidity of response is greatly to be desired. Presumably for this reason the course of immunization employed from the time of Pasteur to the present usually has been very intensive although on a largely empirical rather than solidly experimental basis. Recent evidence to be discussed shortly suggests that fewer but better spaced inocula will stimulate a response with at least equal rapidity and magnitude.

All antirabic vaccines consist of relatively crude suspensions of infected tissue. Until recently all were prepared from the CNS tissue of animals (in the United States usually that of rabbits). For the most part the virus in such suspensions is inactivated by phenol (as in Semple type vaccine) or by ultraviolet light. Harris type vaccine containing live fixed virus is produced by one biological firm in the United States and a similar live virus vaccine is distributed by the Georgia State Health Department Atlanta Ga. Since live fixed virus apparently does not multiply in man these live virus and inactivated vaccines may be regarded as essentially similar. They should be effective in proportion to their antigenic content and the immunizing efficiency of the methods by which they are utilized. Also they presumably possess equivalent capacities to cause undesirable complications such as sensitivity to the general animal protein immediate allergic reactions (in persons already sensitive to the animal protein perhaps because of a previous course of treatment) or finally demyelinating reactions of the CNS (or isoallergic encephalitis) due to the development of organ specific sensitivity to the CNS. This latter complication is the most serious and is the major reason for restricting the use of vaccine to cases in which the risk of not using the vaccine is carefully

rabies virus may remain localized at the site of introduction for as long as twenty four hours. Treatment is directed at destroying or eliminating the virus and to the extent that it is successful it may completely prevent infection or so reduce the size of the infecting dose that the incubation period is extended. This simple, but very important, procedure should be carried out in all instances of animal bite without regard to the probability of rabies exposure.

Modern studies agree that the classic method of acid cautery is quite effective especially when carried out within the first few hours after the bite. However other and less painful and disfiguring methods also are nearly as effective under experimental conditions, these include irrigation and scrubbing with soap or detergent solutions.¹⁻¹³ Recent observations by Perez Gallardo¹³ confirm the earlier work of Habel¹⁴ and indicate that local infiltration of antiserum is of real value.

It must not be forgotten that bites may be infected with agents other than rabies virus and that appropriate measures should be taken to combat these (that is antitetanus and antibacterial precautions).

Passive immunity. Active response to antirabies vaccine requires time in order than effective levels of antibody may be achieved (perhaps 20 days from initiation of treatment) and obviously cannot be expected to prevent a disease that occurs after unusually short periods of incubation as is commonly the case in severely exposed persons. The obvious remedy for this situation is to supply the needed antibody at once by passive means. Perhaps because of the inconclusiveness of early observations¹⁵⁻¹⁶ until recently there has been little sustained interest in this approach. Experimental work demonstrated the high degree of protection afforded animals by systemically and even locally administered antiserum.¹³⁻¹⁴ Somewhat later the importance of early use of serum (within seventy two hours) was demonstrated and it was shown also that the maximum protection of animals requires antiserum supplemented by vaccine.⁸⁻¹⁷

Since 1948 partially purified hyperimmune antirabies serum has been utilized on an experimental basis in man. In Georgia its use was restricted to persons suffering severe and certain exposure less than 72 hours before seeking treatment. By early 1953 69 persons none of whom developed rabies had received antiserum. In each case a course of antirabic vaccine also was begun after injecting the serum.¹⁸ In 1954 the opportunity finally occurred to carry out a controlled antiserum trial that had been planned since 1950 by the World Health Organization (WHO) Expert Committee on Rabies. A rabid wolf in Iran attacked and bit 27 persons of whom 17 were severely exposed by bites on the head. These latter were divided into 3 groups for treatment: the groups receiving, respectively only vaccine or vaccine plus 1 dose of serum or vaccine plus 2 doses of serum (the second dose given after a 5 day interval). The rabies mortality ratios for these 3 groups were 3/5, 1/6 and 0/6 respectively.¹⁹ Sera were collected at frequent intervals from these persons⁸ and on an experimental basis⁸⁻²¹ from unexposed volunteers receiving vaccine alone or with serum and have been tested for neutralizing antibody. The resulting data indicate that even a single dose of serum coupled with a full

TABLE 1
TIME OF APPEARANCE OF ANTIBODIES FOLLOWING PRIMARY IMMUNIZATION
WITH HARRIS SEMPLE AND HEP FLURY VACCINES

Type	Inoculation		Interval (day)	Number of people	Number of days after initiation of body						Percentage
	Dose	Formula			0	10	15	20	30	60	
Harris Semple	7.0 ml SC	14	1	9	0	2	9	9	9	9	100
	6.0 ml SC	12	1	10†	0	2	8	9	9	9	90
		4	5	19	0	6	17	19	19	19	100
HEP Flury	12 gm IM	2, 3 or 6	1, 7	14	0	7	8	8	9	12	93
	20 gm IM	5, 10	1 or 2	10	0	7	7	7	8	9	100
	0.99 or 1.32 gm IM	3 or 4	5	18	0	5	9	10	12	13	78 (100)
	2.64 gm IM plus 0.64 gm ID	4	5	8	0	5			8	8	100
	0.12 to 0.36 gm ID	3	5	49	0	23	31	39	38	36	92 (94)
	0.16 to 0.64 gm ID	4	5	19	0	8	14	16	18	18	95

Antibody as indicated by survival of 3 or more of 6 mice inoculated with mixture of 1:4 final dilution of serum and 100 LD₅₀ of virus.

† Based on presence of significant neutralization in at least 1 serum specimen; apparent discrepancies between percentage shown and maximum number positive on any 1 day are due to individual variations in time of relatively transient low level responses. Figures in parentheses are based on retesting sera found negative in the conventional test by means of the long incubation test.

‡ Sera from these 10 persons were obtained through the courtesy of M. M. Kaplan, Veterinary Public Health Section, World Health Organization. These are the same individuals as group K in TABLE 3.

vaccine only in marginally effective amounts it became particularly necessary to determine the most efficient method of rapid immunization.

Numerous small groups of volunteers were given different courses of vaccine including for comparative purposes Semple and Harris type vaccines. In TABLE 1 are presented data as to the time of appearance of neutralizing antibody and the over all proportion responding in some of the groups manifesting the best responses. Although only the 2 most rigorous courses of Flury vaccine (with a total dose of 20 gm. or with combined intramuscular and intradermal inoculations) elicited a readily detected response in 100 per cent of vaccines it is interesting that as measured by rapidity of response (the proportion with antibody by the tenth day) Flury vaccine did better (47 per cent of 118 persons) than Semple and Harris vaccines combined (26 per cent of 38 persons). Also of note is the fact that 4 doses of Semple vaccine spaced at 5 day intervals did at least as well as 12 or 14 daily doses of Semple or Harris vaccine. In TABLE 2 are shown the available data as to titers of neutralizing antibody at 15 and 30 days after initiation of vaccination in some of the same groups of volunteers. The best antibody titers among those given HEP Flury vaccine were in the groups receiving the 20 gm. total dose intramuscularly or a course

evaluated as being greater than the danger of using it. This latter risk is variously estimated²³⁻²⁴ (the maximum estimated frequency is 1 in 600) but is greater in persons who have had one or more previous courses of treatment. Case fatality may be as great as 25 per cent but survivors usually manifest few permanent sequelae.²⁵

The occurrence of these complicating reactions plus the incomplete effectiveness of specific prophylaxis has resulted in recent times in a very vigorous reinvestigation of human immunization. Passive immunity described above appears to afford an answer to the problem posed by the disease that occurs after a short incubation period. Elimination of the CNS specific tissue antigen and reduction in the capacity of the vaccine to induce sensitivity to the animal protein would resolve the problem of complications due to the vaccine. One obvious answer would be to grow the viral antigen in tissue other than CNS and if possible in a tissue poor in sensitizing proteins. Avian embryos provide a suitable host for viral growth from which CNS tissue can be readily eliminated. Furthermore embryonic proteins are very poorly antigenic themselves. However, before vaccines of such different types can be recommended for general use it must be shown that they are no less effective as viral antigens than those currently in use.

Use of these vaccines in man has been studied independently²⁷ and also on a world wide collaborative basis¹⁻²⁶ under the guidance of the WHO Expert Committee on Rabies. This latter study in particular is very comprehensive in scope and has undertaken to make unbiased comparative tests not only of vaccines of different types but also of various inoculation schedules of such vaccines and of the combined use of antiserum plus vaccine as already described. Unfortunately little of the WHO work with avian embryo vaccines has been completed at this time.

The avian vaccine most extensively studied in man is that made from chick embryos infected with the high egg passage (HEI) Flury strain of virus.²⁶⁻²⁷ This vaccine is used as a partially clarified suspension (40 to 80 per cent) of infectious chick embryo (often with penicillin and streptomycin added) with infectivity preserved by lyophilization and is given preferably in doses of 0.1 to 0.2 ml intradermally or 0.5 ml intramuscularly. More than 550 primary immunizations have been given to volunteers with various dose and inoculation schedules and more than 220 recall doses have been given to persons previously immunized with Flury vaccine or by the conventional Pasteur treatment.²⁸ This fairly extensive use provides the basis for considerable confidence as to the primary safety of HEI Flury vaccine. Local reactions especially after intradermal inocula have constituted the chief but still not very considerable postvaccination complaint. Early work³ made two points clear (1) although alive the HEP Flury virus (like fixed rabies virus) is not able to multiply in man since response is directly related to the amount of vaccine inoculated and (2) using closely spaced intramuscular inocula excessive amounts of vaccine (12 to 20 gm total dose of chick embryo) are required to elicit a satisfactorily uniform response. Since immune response is dependent entirely on the viral antigen actually injected and since this apparently is present in the

β propriolactone⁷ This vaccine as of the moment has an apparent advantage over the HEP Flury vaccine in two or three respects First while it has been assumed on theoretical grounds that the Flury vaccine is free of neuro-paralytic factors the duck embryo vaccine actually has been found to be free in adjuvant tests in guinea pigs⁸ Second, the data reported for several lots suggest that in potency tests it consistently surpasses by two or more times the standard vaccine of the National Institutes of Health Public Health Service Bethesda Md whereas for reasons not clear Flury vaccine will not pass this test Third because of the potency test data the duck embryo vaccine has now been licensed for commercial production and is generally available Rather surprisingly however the comparable available data as to antibody response to a course (14 daily doses) of duck embryo vaccine as shown at the bottom of TABLE 2 suggest that the level of human response is little better than response to a 4-dose course of Flury vaccine and considerably lower than that to Semple or Harris vaccines It should be mentioned also that insofar as it can be ascertained from published reports the primary safety of HEP Flury vaccine for man has been tested far more extensively by actual use in man than that of duck embryo vaccine Direct comparisons of HEP Flury and duck embryo vaccines are now in progress

Work with vaccines of CNS origin inactivated with phenol as carried out in the WHO sponsored study¹ is summarized in TABLE 3 with respect to time of appearance of antibody The chief point of interest is that the 1 and 3 dose schedules were grossly less effective than those of 12 or 14 daily doses or even than that of 7 daily doses Generally speaking the results of the schedules with daily doses compared favorably with those of similar schedules with Harris vaccine as already presented However there seems an obvious disparity perhaps related to differences in vaccine potency between the results

TABLE 3

SUMMARY OF OBSERVATIONS AS TO TIME OF APPEARANCE OF NEUTRALIZING ANTIBODY IN PERSONS UNDERGOING PRIMARY IMMUNIZATION WITH PHENOLIZED VACCINES OF ANIMAL CNS ORIGIN IN THE COLLABORATIVE WHO TRIALS

Group	Vaccine	Sgl (ml)	Nb of doses	Interval (days)	Nb of subjects	Number of persons with antibody on day					Overall %
						7	10	14	21	28	
F	20% rabbit brain	3.0	1	—	10	1	3	4	3	4	50
B	20% rabbit brain	0.5	7	daily	10	1	7	8	9	9	90
K†	20% rabbit brain	0.5	12	daily	10	1	2	9	9	9	90
L	5% goat brain	2.0	14	daily	10	NT‡	0	8	10	10	100
N	5% goat brain	2.0	3	5	10	NT	1	1	5	6	60
T	20% rabbit brain	0.5	14	daily	6	NT	1	2	6	6	100

Adapted from papers by Ananasi *et al.*¹⁰

† As in TABLE 1

‡ Group K is the same group as that mentioned in TABLE 1 as receiving 12 doses of Semple vaccine the results shown in TABLE 1 were obtained in tests using 100 LD₅₀ of virus rather than 30 LD₅₀ as employed by the WHO collaborators

§ NT = not tested

TABLE 2

OBSERVATIONS ON LEVEL OF NEUTRALIZING ANTIBODY FOLLOWING PRIMARY IMMUNIZATION WITH HARRIS SEMPLE AND HFP FLURY VACCINES AND FOR COMPARISON DUCK EMBRYO VACCINE†

Type of vaccine	Total dose administered	Number of inoculations	Intervals (days)	Total number of persons	Days after first inoculation	Number of persons with antibody titer in following range						
						<4	4-7	8-31	32-127	128-511	+512	+512
Harris	7.0 ml SC	14	daily	9 9	15 30			2	4 2	3 3		4
Semple	2.0 ml SC	4	5	19 19	15 30	2		1 1	7 4	7 9	2 5	
HLP Flury	20.0 gm IM	5-10	1-2	6 6	15 30	2	1		3 3			
	0.99 to 2.64 gm IM	3 or 4	5	16 16	15 30	6 2	6 8	4 4	1 1	1 1		
	2.64 gm IM + 0.32 gm ID	4	5	8 8	15 30	1	1	2 3	3 5	1		
	0.12 to 0.96 gm ID	3	5	45 45	15 30	19 11	16 14	8 13	5	2 2		
	0.16 to 0.64 gm ID	4	5	19 19	15 30	5 1	7 8	6 6	1 2	2		
Duck embryo	7.0 ml SC	14	daily	21 17	15 30	1 1	7 2	13 9	5			

As measured by testing serial twofold dilutions of sera in groups of 6 mice per dilution using 100 LD₅₀ of challenge virus per intracerebral mouse inoculum.

† The duck embryo data abstracted from the paper by Peck *et al.*⁷ appear comparable in that the same methods of determining neutralizing antibody were employed.

of 4 intradermal doses alone or together with intramuscular doses with 5 day intervals. While the range of titers observed after the administration of Flury vaccine clearly overlaps those resulting from Harris or Semple vaccine the fact remains that no course of Flury vaccine so far employed has induced response fully equivalent to that following Harris or Semple vaccine with respect to both uniformity and antibody level. A further point to note is that the 4-dose course of Semple vaccine induced antibody titers in essentially the same range as did the 14-dose course of Harris vaccine. Unfortunately no comparable group was given a 14-dose course of the same lot of Semple vaccine.

The growth of fixed virus in duck embryos⁸ provided the basis for the more recent type of avian embryo vaccine. Although first used as a live virus vaccine⁸ the product as recently licensed is inactivated by treatment with

β propriolactone⁷ This vaccine as of the moment has an apparent advantage over the HEI Flury vaccine in two or three respects First while it has been assumed on theoretical grounds that the Flury vaccine is free of neuro-paralytic factors the duck embryo vaccine actually has been found to be free in adjuvant tests in guinea pigs⁸ Second the data reported for several lots suggest that in potency tests it consistently surpasses by two or more times the standard vaccine of the National Institutes of Health Public Health Service Bethesda Md whereas for reasons not clear Flury vaccine will not pass this test Third because of the potency test data the duck-embryo vaccine has now been licensed for commercial production and is generally available Rather surprisingly however the comparable available data as to antibody response to a course (14 daily doses) of duck embryo vaccine as shown at the bottom of TABLE 2 suggest that the level of human response is little better than response to a 4-dose course of Flury vaccine and considerably lower than that to Semple or Harris vaccines It should be mentioned also that insofar as it can be ascertained from published reports the primary safety of HEP Flury vaccine for man has been tested far more extensively by actual use in man than that of duck embryo vaccine Direct comparisons of HEI Flury and duck embryo vaccines are now in progress

Work with vaccines of CNS origin inactivated with phenol, as carried out in the WHO sponsored study²¹⁻²² is summarized in TABLE 3 with respect to time of appearance of antibody The chief point of interest is that the 1 and 3 dose schedules were grossly less effective than those of 12 or 14 daily doses or even than that of 7 daily doses Generally speaking the results of the schedules with daily doses compared favorably with those of similar schedules with Harris vaccine as already presented However there seems an obvious disparity perhaps related to differences in vaccine potency between the results

TABLE 3
SUMMARY OF OBSERVATIONS AS TO TIME OF APPEARANCE OF NEUTRALIZING ANTIBODY IN PERSONS UNDERGOING PRIMARY IMMUNIZATION WITH PHENOLIZED VACCINES OF ANIMAL CNS ORIGIN IN THE COLLABORATIVE WHO TRIALS

Group designation	Vaccine	Single dose (ml)	Number of doses	Interval (days)	Number of persons	Number of persons with antibody per day					On percent response at
						7	10	14	21	28	
E	20% rabbit brain	3.5	1	—	10	1	3	4	3	4	50
B	20% rabbit brain	0.5	7	daily	10	1	2	8	9	9	90
K†	20% rabbit brain	0.5	12	daily	10	1	2	9	9	9	90
L	5% goat brain	2.0	14	daily	10	NT‡	0	8	10	10	100
N	5% goat brain	2.0	3	5	10	NT	1	1	5	6	60
T	20% rabbit brain	0.5	14	daily	6	NT	1	2	6	6	100

Adapted from papers by Atanasiu *et al.*²³

† As in TABLE 1

‡ Group K is the same group as that mentioned in TABLE 1 as receiving 12 doses of Semple vaccine; the results shown in TABLE 1 were obtained in tests using 100 LD₅₀ of virus rather than 30 LD₅₀ as employed by the WHO collaborators

§ NT = not tested

TABLE 2

OBSERVATIONS ON LEVEL OF NEUTRALIZING ANTIBODY FOLLOWING PRIMARY IMMUNIZATION WITH HARRIS SEMPLE AND HEP FLURY VACCINES AND FOR COMPARISON DUCK EMBRYO VACCINE†

Type of vaccine	Total dose	Number of animals	Interval (day)	Total number of pups	Days after infection	Number of mice positive in first					
						<4	4-7	8-31	3-127	128-511	+512
Harris	7.0 ml SC	14	daily	9 9	15 30			2	4 2	3 3	4
Semple	2.0 ml SC	4	5	19 19	15 30	?		1 1	7 4	7 9	2 5
HEP Flury	20.0 gm IM	5-10	1-2	6 6	15 30	2	1		3 3		
	0.99 to 2.64 gm IM	3 or 4	5	16 16	15 30	6 2	6 8	4 4	1 1	1	
	2.64 gm IM + 0.32 gm ID	4	5	8 8	15 30	1	1	2 3	3 5	1	
	0.12 to 0.96 gm ID	3	5	45 45	15 30	19 11	16 14	8 13	5	2 ?	
	0.16 to 0.64 gm ID	4	5	19 19	15 30	5 1	7 8	6 6	1 2	2	
Duck embryo	7.0 ml SC	14	daily	21 17	15 30	1 1	7 2	13 9	5		

As measured by testing serial twofold dilutions of sera in groups of 6 mice per dilution using 100 LD₅₀ of challenge virus per intracerebral mouse inoculum

† The duck embryo data abstracted from the paper by Peck *et al.*⁷ appear comparable in that the same methods of determining neutralizing antibody were employed

of 4 intradermal doses alone or together with intramuscular doses with 5 day intervals. While the range of titers observed after the administration of Flury vaccine clearly overlaps those resulting from Harris or Semple vaccine the fact remains that no course of Flury vaccine so far employed has induced response fully equivalent to that following Harris or Semple vaccine with respect to both uniformity and antibody level. A further point to note is that the 4-dose course of Semple vaccine induced antibody titers in essentially the same range as did the 14 dose course of Harris vaccine. Unfortunately no comparable group was given a 14 dose course of the same lot of Semple vaccine.

The growth of fixed virus in duck embryos⁸ provided the basis for the more recent type of avian embryo vaccine. Although first used as a live virus vaccine⁶ the product as recently licensed is inactivated by treatment with

β propiolactone⁷ This vaccine as of the moment has an apparent advantage over the HFI Flury vaccine in two or three respects First while it has been assumed on theoretical grounds that the Flury vaccine is free of neuro-paralytic factors the duck embryo vaccine actually has been found to be free in adjuvant tests in guinea pigs¹⁰ Second the data reported for several lots suggest that in potency tests it consistently surpasses by two or more times the standard vaccine of the National Institutes of Health Public Health Service Bethesda Md whereas for reasons not clear Flury vaccine will not pass this test Third because of the potency test data the duck embryo vaccine has now been licensed for commercial production and is generally available Rather surprisingly however the comparable available data as to antibody response to a course (14 daily doses) of duck embryo vaccine as shown at the bottom of TABLE 2 suggest that the level of human response is little better than response to a 4-dose course of Flury vaccine and considerably lower than that to Semple or Harris vaccines It should be mentioned also that insofar as it can be ascertained from published reports the primary safety of HEP Flury vaccine for man has been tested far more extensively by actual use in man than that of duck embryo vaccine Direct comparisons of HFI Flury and duck embryo vaccines are now in progress

Work with vaccines of CNS origin inactivated with phenol as carried out in the WHO-sponsored study^{1, 2} is summarized in TABLE 3 with respect to time of appearance of antibody The chief point of interest is that the 1 and 3 dose schedules were grossly less effective than those of 12 or 14 daily doses or even than that of 1 daily dose Generally speaking the results of the schedules with daily doses compared favorably with those of similar schedules with Harris vaccine as already presented However there seems an obvious disparity perhaps related to differences in vaccine potency between the results

TABLE 3
SUMMARY OF OBSERVATIONS AS TO TIME OF APPEARANCE OF NEUTRALIZING ANTIBODY IN PERSONS UNDERGOING PRIMARY IMMUNIZATION WITH PHENOLIZED VACCINES OF ANIMAL CNS ORIGIN IN THE COLLABORATIVE WHO TRIALS

Group designation	Vaccine	Vaccine dose (ml)	No. of subjects	Interval (days)	No. of subjects	No. of subjects with antibody					Overall percentage
						7	10	14	21	28	
E	20% rabbit brain	3.5	1	—	10	1	3	4	3	4	50
B	20% rabbit brain	0.5	7	daily	10	1	2	8	9	9	90
K†	20% rabbit brain	0.5	12	daily	10	1	2	9	9	9	90
L	5% goat brain	2.0	14	daily	10	NT‡	0	8	10	10	100
N	5% goat brain	2.0	3	5	10	NT	1	1	5	6	70
T	20% rabbit brain	0.5	14	daily	6	NT	1	2	6	6	100

Adapted from papers by Ananasiu et al.¹¹

† As in TABLE 1

‡ Group K is the same group as that mentioned in TABLE 1 as receiving 12 doses of Semple vaccine; the results shown in TABLE 1 were obtained in tests using 100 LD₅₀ of virus rather than 30 LD₅₀ as employed by the WHO collaborators

§ NT = not tested

with 3 doses in TABLE 3 and those with 4 doses of Semple vaccine shown in TABLE 1 in each case with a 5 day interval

To recapitulate briefly, when exposure must be presumed, active antirabic immunization is mandatory. A major hazard is isoallergic encephalitis to avoid which embryo vaccines may provide the answer. Duck embryo vaccine has been licensed whereas HEP Flury vaccine thus far has not been licensed. Although promising results have been observed after modified dose schedules these must be confirmed before the usual 14 daily dose treatment is abandoned. Management of exposure in persons who previously have had Pasteur treatment is a special situation now to be described.

The Situation Before the Bite

Broadly speaking the preventive measures that should be taken before the bite fall into one or another of three main categories as follows: those that help to minimize contact between potentially infected animals and man; those directed at controlling rabies in domestic and wild animals; and those that may afford man specific protection in advance of exposure. I or lack of space I shall pass over the first two categories remarking only upon two relatively recent developments. The first of these is the availability and apparently superior usefulness of canine Flury vaccine³ which because of the long duration of the resulting immunity is a potentially great adjunct to the control of canine rabies. The second discussed at length elsewhere in these pages by K. F. Burns is the emergence of extensive rabies infection in herbivorous and insectivorous bats as a possibly important new factor in wildlife rabies.

Measures falling into the third category—the specific protection of man in advance of exposure—have been possible in theory for a long time but have been rarely practiced because of the risk associated with receiving antirabic vaccine of the types heretofore available. The advent of the presumably safe avian embryo vaccines makes possible a very great change in this situation and considerable work has been done already with the HEP Flury vaccine^{4, 5} as I shall try to indicate in the following summary. The observations in question suggest that primary immunization with HEP Flury vaccine or by means of Pasteur treatment (presumably including that with duck embryo vaccine) provides a foundation immunity that conditions the recipient into responding rapidly to single stimuli given as much twenty or more years later. Upon such a basis one can build a long term immunity by means of periodic booster inocula.

About 64 persons who had received a primary course of HEP Flury vaccine were given single booster inocula of the same vaccine 5 to 48 months after the primary course; the results are summarized in TABLE 4. It is of interest that small intradermal inocula elicited responses just as effectively as did the larger ones given intramuscularly. Although definite failures of response (in 3 cases only) were observed only among those whose primary response had failed or was uncertain, there was observed a definite correlation between the maximum antibody titer after the primary course and that after the booster. This is shown in TABLE 5 and it suggests that in terms of antibody level at

TABLE 4

SUMMARY OF ANTIBODY RESPONSES OBSERVED IN PERSONS IMMUNIZED EXPERIMENTALLY WITH DIFFERING PRIMARY COURSES OF HEP FLURY VACCINE AND GIVEN SINGLE BOOSTER INOCULA OF THE SAME VACCINE AT FROM 5 TO 48 MONTHS AFTER PRIMARY IMMUNIZATION

HEP Flury booster inoculum		Response primary	Number booster	Number exposed	Range (minimum-maximum) titer (1:1)
Dose (g/ml)	Route				
0.001-0.16	ID	None Dubious Yes	1 5 19	1 3 18(19)	3? >32-96 <4-256
0.33	IM	Dubious Yes	2 12	2 7(12)	>4 >4-3?
2.0	IM	None Dubious Yes	4 11 10	2(3) 11 10	<4->756 <4-96 >4->1024

Figures in parentheses include instances in which response was likely but because of inadequate serum specimens the conclusive quantitative test could not be done

† Results of <4 mean some mice were protected in the lowest serum dilution tested (1:4 final) but not enough to give a 1:4 titer; use of the > sign means that because of inadequate serum specimens retests to determine the definite end point could not be done

least best results with boosters are obtained after the more effective primary courses. However without regard to the level of primary response nearly three fourths of the 42 persons responded to the booster with titers of 1:32 or better. Some idea of the maintenance of antibody levels after primary and booster immunizations is given in TABLE 6. With respect to the primary course it seems clear that except for the less common individual with a primary response titer of 1:32 or better antibody persistence at a readily demonstrated level is likely to be brief. While the same relation exists in the case of the postbooster persistence the pattern differs significantly in that the maximum titer after booster was usually 1:32 or better. In practice therefore it would seem that adequate levels persist in most instances for at least two years after

TABLE 5

CORRELATION OF MAXIMUM ANTIBODY TITER AFTER HEP FLURY BOOSTER AND MAXIMUM TITER FOLLOWING PRIMARY COURSE OF VACCINE

Maximum antibody titer (1:1) following primary course	Total number exposed	Number (percentage) of maximum antibody titer (1:1) following booster inoculum				
		0	<4	4-7	8-31	32+
3?+	5					5
8-31	5				1	4
4-7	6		1		1	4
<4	19	1	2	1		15
0	7	3	1		1	2
All	47	4	4	1	3	30

Symbols <4 as in TABLE 4

with 3 doses in TABLE 3 and those with 4 doses of Semple vaccine shown in TABLE 1 in each case with a 5 day interval

To recapitulate briefly, when exposure must be presumed, active antirabic immunization is mandatory. A major hazard is isoallergic encephalitis to avoid which embryo vaccines may provide the answer. Duck embryo vaccine has been licensed whereas HEP Flury vaccine thus far has not been licensed. Although promising results have been observed after modified dose schedules these must be confirmed before the usual 14 daily dose treatment is abandoned. Management of exposure in persons who previously have had Pasteur treatment is a special situation now to be described.

The Situation Before the Bite

Broadly speaking the preventive measures that should be taken before the bite fall into one or another of three main categories as follows: those that help to minimize contact between potentially infected animals and man; those directed at controlling rabies in domestic and wild animals, and those that may afford man specific protection in advance of exposure. For lack of space I shall pass over the first two categories remarking only upon two relatively recent developments. The first of these is the availability and apparently superior usefulness of canine Flury vaccine³⁰ which because of the long duration of the resulting immunity, is a potentially great adjunct to the control of canine rabies. The second discussed at length elsewhere in these pages by K. F. Burns is the emergence of extensive rabies infection in herbivorous and insectivorous bats as a possibly important new factor in wildlife rabies.

Measures falling into the third category—the specific protection of man in advance of exposure—have been possible in theory for a long time but have been rarely practiced because of the risk associated with receiving antirabic vaccine of the types heretofore available. The advent of the presumably safe avian embryo vaccines makes possible a very great change in this situation and considerable work has been done already with the HEP Flury vaccine^{4, 5} as I shall try to indicate in the following summary. The observations in question suggest that primary immunization with HEP Flury vaccine or by means of Pasteur treatment (presumably including that with duck embryo vaccine) provides a foundation immunity that conditions the recipient into responding rapidly to single stimuli given as much twenty or more years later. Upon such a basis one can build a long term immunity by means of periodic booster inocula.

About 64 persons who had received a primary course of HEP Flury vaccine were given single booster inocula of the same vaccine 5 to 48 months after the primary course; the results are summarized in TABLE 4. It is of interest that small intradermal inocula elicited responses just as effectively as did the larger ones given intramuscularly. Although definite failures of response (in 3 cases only) were observed only among those whose primary response had failed or was uncertain, there was observed a definite correlation between the maximum antibody titer after the primary course and that after the booster. This is shown in TABLE 5 and it suggests that in terms of antibody level at

with respect to the management of treatment on the occasion of any subsequent new exposure to rabies. Deliberate use of Pasteur treatment as the foundation for maintenance immunity would not be recommended obviously because of the associated risk; therefore this aspect has not been explored in detail. However, when treatment has been required because of exposure, it is possible that profit may be taken from the misfortune and immunity maintained thereafter as suggested above. Indeed, proof that this may be done and guidance as to timing may be derived from the following data, which bear more directly on managing the treatment of a new exposure.

A total of 136 persons with histories of previous Pasteur treatment, chiefly veterinarians, but including some prisoner volunteers, were persuaded to submit to a single inoculation of HEI Flury vaccine, usually by the intradermal route, and to one preinoculation and several postinoculation bleedings. The results are summarized in TABLE 7. Definite response was demonstrated in 86 per cent of the total or, excluding 9 persons for whom the tests were not determinative (see footnotes to TABLE 7), in 92 per cent (11 of 12) of those properly tested. Furthermore, of the 10 definite failures, 6 were among 17 persons last treated 20 or more years previously. Although some variation

TABLE 8
SUMMARY OF OBSERVATIONS AS TO PERSISTENCE OF
NEUTRALIZING ANTIBODY AFTER PASTEUR TREATMENT

Number of treatments received	Antibody level (1:1)	Number of persons observed at indicated years last treatment									
		<1	1-2	3-4	5-6	7-8	9-11	12-15	16-20	21-25	35-40
Only 1	32+	1		3		2	3	1	3		
	8-31	2	3	2	2			2	1	1	
	4-7						1				
	<4 or I LT 0	2	7	1	1	3	1	3	4	1	1
		2	2	3	5	1	4	5	7	6	3
	All pos	3	10	6	3	5	5	6	8	2	1
	All	5	12	9	8	6	9	11	15	8	4
Two or more	32+	1	6	4	4	6	1	1			
	8-31		1	1			1	1	2		
	4					2	2				
	<4 or I LT 0					1	1	1	2	1	
									2		
	All pos	1	7	5	4	8	4	5	4	0	
	All	1	7	5	4	9	5	5	6	1	
Only 1	Per cent positive	60	84	67	38	84	56	55	53	25	25
		13		57		54		25			
100		100	100	100	89	80	100	67	0		
100		89		18		—					
Two or more											

P.L.T. mean positive in qualitative long incubation test in most such instances no quantitative tests were made. Also <4 as in TABLE 4.

TABLE 6

RELATION BETWEEN MAXIMUM ANTIBODY TITER AFTER PRIMARY COURSE AND AFTER BOOSTER OF HFP FLURY VACCINE AND PRESENCE OF ANTIBODY AT SUBSEQUENT INTERVALS

Times of follow-up	Maximum titer (1:)	Presence of antibody at designated intervals								
		5 Months			12 Months			24 or more months		
		Total N	Number with titer		Total No.	Number with titer		Total N	Number with titer	
			<4	4 >		<4	4 >		<4	4 >
Primary course	32+	2	1	1	2		2	4		4
	8-31	3	2	1	4	1	3	1	1	
	4-7	7	7		2	1	1	3	3	
	<4	8	7	1	16	15	1	6	6	
Booster	32+				17	2	15	8	1	1
	8-31				1	1		0		
	4-7				4	4		2	1	1
	<4				1	1		1	1	

The symbol <4 in this case may include sera devoid of antibody since many were negative in 1:4 final dilution and not tested in lower dilution

a booster. Second, third, and even fourth (in one instance only) boosters have been given but thus far to an insufficient number of persons to permit definition of the optimum spacing.

As stated previously, Pasteur treatment itself provides a foundation immunity upon which a program of maintenance immunity also may be built and which in any event carries important and long persisting implications.

TABLE 7

SUMMARY OF ANTIBODY RESPONSES TO HFP FLURY BOOSTER INOCULA GIVEN TO PERSONS WITH HISTORIES OF PREVIOUS PASTEUR TREATMENT

Year last treated	Number tested	Persons responded		Range of maximum response titer (1:)
		Number	Percent	
unknown	5	4	80	>32->1 074
35-40	4	1	25	128
20-26	13	9 (10)	69 (71)	<4 >4 100
15-19	21	20 (21)	95 (100)	<4 1 500
10-14	20	20	100	32-2 048
5-9	34	31 (33)	91 (97)	4-16 400
3-4	15	14 (15)	93 (100)	4-3 640
1-2	18	16 (17)	89 (94)	38->4 096
<1	6	2 (5)	33 (83)	18-282
All	136	111 (126)	86 (93)	<4 16 400

Inocula usually intra-dermal and in amounts ranging from 0.08 to 0.16 gm. in 15 persons; however, intramuscular inocula of 2.0 gm. were given.

† Use of parentheses has same meaning as in TABLE 4.

‡ Use of < and > has same meaning as in TABLE 4.

with respect to the management of treatment on the occasion of any subsequent new exposure to rabies. Deliberate use of Pasteur treatment as the foundation for maintenance immunity would not be recommended obviously because of the associated risk; therefore this aspect has not been explored in detail. However when treatment has been required because of exposure it is possible that profit may be taken from the misfortune and immunity maintained thereafter as suggested above. Indeed proof that this may be done and guidance as to timing may be derived from the following data which bear more directly on managing the treatment of a new exposure.

A total of 136 persons with histories of previous Pasteur treatment chiefly veterinarians but including some prisoner volunteers were persuaded to submit to a single inoculation of HEI Flury vaccine usually by the intradermal route and to one preinoculation and several postinoculation bleedings. The results are summarized in TABLE 7. Definite response was demonstrated in 86 per cent of the total or excluding 9 persons for whom the tests were not determinative (see footnotes to TABLE 7) in 92 per cent (117 of 127) of those properly tested. Furthermore of the 10 definite failures 6 were among 17 persons last treated 20 or more years previously. Although some variation

TABLE 8
SUMMARY OF OBSERVATIONS AS TO PERSISTENCE OF
NEUTRALIZING ANTIBODY AFTER PASTEUR TREATMENT

Number of treatments	Antibody titer	Number of persons bled at stated years									
		<1	1-2	3-4	5-6	7-8	9-11	12-15	16-20	21-25	35-40
Only 1	32+	1		3		2	3	1	3		
	8-31	2	3	2	?			2	1	1	
	4-7		7	1	1	3	1	3	4	1	1
	<4 or P LT 0	2	2	3	5	1	4	5	1	6	3
	All pos	3	10	6	3	5	5	6	8	2	1
	All	5	12	9	8	6	9	11	15	8	4
Two or more	32+	1	6	4	4	6	1	1			
	8-31		1	1			1	1	2		
	4-7					2	2				
	<4 or P LT 0					1	1	1	2	1	
	All pos	1	7	5	4	8	4	5	4	0	
	All	1	7	5	4	9	5	5	6	1	
Only 1	Per cent positive	60	84	67	38	84	56	55	53	25	25
		13			57			54		25	
		100	100	100	100	89	80	100	67	0	
		100			89			18		—	
Two or more											

P LT means positive qualitative long incubation test in most such instances no quantitative tests were made. All <4 as in TABLE 4.

TABLE 6
RELATION BETWEEN MAXIMUM ANTIBODY TITER AFTER PRIMARY COURSE AND
AFTER BOOSTER OF HFP FLURY VACCINE AND PRESENCE OF ANTIBODY
AT SUBSEQUENT INTERVALS

Titer flow g	Ma ximum titer (1°)	Pre s e n c e o f a n t i b o d y t i t e r a t i n t e r v a l s									
		5 Months				12 M o n t h s				24 m o n t h	
		T t a l N o	Numb w i t h t i t e r 1		T t a l N	Numb w i t h t i t e r 1		T t a l N o	N m b w i t h t i t e r 1		
			<4	4 o r >		<4	4 o r >		<4	4 o r >	
Primary course	32+	2	1	1	2		2	4		4	
	8-31	3	2	1	4	1	3	1	1		
	4-7	7	7		2	1	1	3	3		
	<4*	8	7	1	16	15	1	6	6		
Booster	32+				17	2	15	8	1	7	
	8-31				1	1		0			
	4-7				4	4		2	1	1	
	<4				1	1		1	1		

The symbol <4 in this case may include sera devoid of antibody since many were negative in 1:4 final dilution and not tested in lower dilution

a booster. Second, third and even fourth (in one instance only) boosters have been given but thus far to an insufficient number of persons to permit definition of the optimum spacing.

As stated previously, Pasteur treatment itself provides a foundation immunity upon which a program of maintenance immunity also may be built and which in any event carries important and long persisting implications.

TABLE 7
SUMMARY OF ANTIBODY RESPONSES TO HFP FLURY BOOSTER INOCULA GIVEN
TO PERSONS WITH HISTORIES OF PREVIOUS PASTEUR TREATMENT

Years since last treatment	Number of persons	Persons positive		Range of positive titer (1°)
		Number	Percent	
unknown	5	4	80	>32 >1:074
35-40	4	1	25	128
20-26	13	9 (10)	69 (77)	<4->4:100
15-19	21	20 (71)	95 (100)	<4-1:500
10-14	20	20	100	32-2:048
5-9	34	31 (33)	91 (97)	4-16:400
3-4	15	14 (15)	93 (100)	45-3:640
1-2	18	16 (17)	89 (94)	38->4:096
<1	6	2 (5)	33 (83)	18:282
All	136	117 (126)	86 (93)	<4:16:400

Inocula usually intra dermal and in amounts ranging from 0.08 to 0.16 gm. in 15 persons; however intramuscular inocula of 2.0 gm. were given.

† Use of parentheses has same meaning as in TABLE 4.

‡ Use of < and > has same meaning as in TABLE 4.

with respect to the management of treatment on the occasion of any subsequent new exposure to rabies. Deliberate use of Pasteur treatment as the foundation for maintenance immunity would not be recommended obviously because of the associated risk (therefore this aspect has not been explored in detail). However when treatment has been required because of exposure it is possible that profit may be taken from the misfortune and immunity maintained thereafter as suggested above. Indeed proof that this may be done and guidance as to timing may be derived from the following data which bear more directly on managing the treatment of a new exposure.

A total of 136 persons with histories of previous Pasteur treatment chiefly veterinarians but including some prisoner volunteers were persuaded to submit to a single inoculation of HFI Flury vaccine usually by the intradermal route and to one preinoculation and several postinoculation bleedings. The results are summarized in TABLE 7. Definite response was demonstrated in 86 per cent of the total or excluding 9 persons for whom the tests were not determinative (see footnotes to TABLE 7) in 92 per cent (117 of 124) of those properly tested. Furthermore of the 10 definite failures 6 were among 17 persons last treated 20 or more years previously. Although some variation

TABLE 8
SUMMARY OF OBSERVATIONS AS TO PERSISTENCE OF
NEUTRALIZING ANTIBODY AFTER PASTEUR TREATMENT

Number of inoculations	Antibody level (t)	Number of persons by titer titer									
		<1	1-2	3-4	5-6	7-8	9-11	12-15	16-20	21-25	35-40
Only 1	32+	1		3		2	3	1	3		
	8-31	2	3	2	2			2	1	1	
	4-7						1				
	<4 or P L T		7	1	1	3	1	3	4	1	1
	0	2	2	3	5	1	4	5	7	6	3
	All pos	3	10	6	3	5	5	6	8	2	1
Two or more	All	5	12	9	8	6	9	11	15	8	4
	32+	1	6	4	4	6	1	1			
	8-31		1	1			1	1	2		
	4-7						2	2			
	<4 or P I T					2	2	1	2	1	
	0					1	1		2		
	All pos	1	7	5	4	8	4	5	4	0	
	All	1	7	5	4	9	5	5	6	1	
Only 1		60	84	67	38	84	56	55	53	25	25
	Per cent post		73			57		54		25	
	tive										
Two or more		100	100	100	100	89	80	100	67	0	
			100			89		18		—	

P L T mean post in qualit tivel g i cubation test in most such instances no quantitative tests were made Als <4 as in TABLE 4

in response was noted, peak responses (usually by the fifteenth day) tended to surpass those following a full primary course of Pasteur treatment. Although no sera were collected earlier than 15 days after the booster the common observation of maximum response titers in these 15 day sera indicates that the booster response usually was more rapid as well as greater than that to primary immunization. It should be noted that 43 (or 32 per cent) of the group admitted having received 2 or more previous courses of treatment. An analysis not here presented of the frequency and degree of response showed no differences as between those treated once and those given multiple treatments.

Data as to duration of post treatment immunity became available as the prebooster sera were tested. These are summarized separately in TABLE 8 for those previously treated only once and for those with histories of multiple courses. In general remarkable antibody persistence was observed but the differences related to number of previous courses are of interest. After but a single course more than one half of the subjects retained antibody through 20 years and nearly three fourths for 5 years. However if 2 or more courses had been administered essentially 80 per cent had antibody for 20 years and all possessed it through 5 years. From these data one might suggest the following: (1) for maintenance immunity a booster dose (preferably of avian embryo vaccine) might be given at from 1 to 2 years after Pasteur treatment with subsequent boosters at 5 to 6 year intervals (or upon new exposure) and (2) for new exposures in previously treated persons a single booster inoculum (again preferably of avian embryo vaccine) is probably adequate if the most recent treatment occurred no longer than 20 years previously.

Comment

The heart of the problem confronting the physician in the after the bite situation is the weighing of the evidence that true exposure has taken place against the risk known to be associated with conventional antirabic vaccination. This has been delineated in classic manner by T. F. Sellers²¹ in a paper aptly entitled "Rabies: the physician's dilemma" which closed with the caution that for exposures other than actual bites the danger of treatment complications far exceeds that of either rabies or rabophobia. These words were published in 1948. At present although these words of caution must be borne in mind at least a little longer use of the avian embryo vaccines seems to ease the problem considerably especially in the case of those who already have undergone one course of Pasteur treatment. Meanwhile to ensure the best possible response in the emergency situation efforts should continue to increase the antigenic potency of the embryo vaccines and to improve the methods for their use. The other major problem posed by instances of severe exposure with probability of disease after a short incubation period has been greatly eased by the advent of antirabies hyperimmune serum. However room for important improvements remains in the direction of increasing the potency and degree of purification in order to reduce the volume of the inoculation the average size of which is now 40 to 50 ml per adult and the likelihood of serum sickness reactions.

For small but selected high risk segments of the population of the United States including veterinarians dog catchers and postmen the observations as to the long persistence after Pasteur treatment both of antibody and of conditioning to later booster inocula have special implications with respect to management of new exposures. Often no inoculation may be needed (see TABLE 8 for guidance) and in any case one inoculum preferably of embryo vaccine should be sufficient. For these same groups plus children resident in regions of high rabies endemicity the embryo vaccines would seem to offer a safe mechanism for laying a foundation immunity and for maintaining such a basic immunity including that induced by Pasteur treatment indefinitely by means of periodic booster doses. The optimum spacing of such doses remains to be worked out but if the duration of postbooster seroimmunity is at all similar to that following Pasteur treatment in previously treated persons (again see TABLE 8) a possible schedule would have the foundation course followed within 1 or 2 years by the first booster and subsequent boosters coming every 5 or 6 years thereafter or as with tetanus toxoid upon the occasion of suspected exposure.

A final note perhaps should be added. The foregoing presentation and discussion have been based on the possibly debatable assumption that immunity to rabies and the readily demonstrable presence of neutralizing antibody are synonymous. I am familiar with evidence (largely unpublished) that highly susceptible animals possessing antibody may succumb to challenge with rabies street virus and that other animals may resist it even though antibody can not be demonstrated. Because species susceptibility to rabies may vary significantly one cannot translate the experimental work in animals directly to man nor can one evaluate seroimmunity in man by the method of challenge. As a consequence one must have a reasoned faith in the importance of neutralizing antibody the degree of faith obviously varying directly with the antibody titer.

References

1. JOHNSON, H. N. 1957. Viral and Rickettsial Infections of Man. Chap. 12. 2nd ed. T. M. Rivers, Ed. Lippincott, Philadelphia, Pa.
2. HABEL, K. 1940. Evaluation of a mouse test for the standardization of the immunizing power of anti-rabies vaccines. Public Health Repts U. S. 55: 1473.
3. SCHWAB, M. I., J. P. FOX, D. P. CONWELL, & T. A. ROBINSON. 1954. A standardized rabies virus vaccination in man. Bull. World Health Organization 10: 873.
4. FOX, J. P., D. P. CONWELL, & I. GERHARDT. 1956. Anti-rabies vaccination of man with HEP Flury virus. Bull. Tulane Univ. Med. Fac. 16: 1.
5. FOX, J. P., H. KOPROWSKI, D. P. CONWELL, J. BLACK, & H. M. GELFAND. 1957. Study of anti-rabies immunization of man. Observations with HEP Flury vaccine and other vaccines and with hyperimmune serum in primary and recall immunizations. Bull. World Health Organization. In press.
6. PECK, F. B., JR., H. M. POWELL, & C. G. CULBERTSON. 1955. New anti-rabies vaccine for human use. J. Lab. Clin. Med. 45: 619.
7. PECK, F. B., JR., H. M. POWELL, & C. G. CULBERTSON. 1956. Duck embryo rabies vaccine. Study of fixed virus vaccine grown in embryonated duck eggs and killed with beta-propiolactone (BPL). J. Am. Med. Assoc. 162: 1373.
8. HABEL, K. 1954. Antiserum in the prophylaxis of rabies. Bull. World Health Organization 10: 781.
9. HABEL, K. 1957. Rabies prophylaxis in man. Pediatrics 19: 923.
10. JOHNSON, H. N. 1942. The significance of the Negri body in the diagnosis and epidemiology of rabies. Ill. Med. J. 81: 387.

in response was noted peak responses (usually by the fifteenth day) tended to surpass those following a full primary course of Pasteur treatment. Although *no sera were collected earlier than 15 days after the booster* the common observation of maximum response titers in these 15 day sera indicates that the booster response usually was more rapid as well as greater than that to primary immunization. It should be noted that 43 (or 32 per cent) of the group admitted having received 2 or more previous courses of treatment. An analysis not here presented of the frequency and degree of response showed no differences as between those treated once and those given multiple treatments.

Data as to duration of post treatment immunity became available as the prebooster sera were tested. These are summarized separately in TABLE 8 for those previously treated only once and for those with histories of multiple courses. In general remarkable antibody persistence was observed but the differences related to number of previous courses are of interest. After but a single course more than one half of the subjects retained antibody through 20 years and nearly three fourths for 5 years. However if 2 or more courses had been administered essentially 80 per cent had antibody for 20 years and all possessed it through 5 years. From these data one might suggest the following (1) for maintenance immunity a booster dose (preferably of avian embryo vaccine) might be given at from 1 to 2 years after Pasteur treatment with subsequent boosters at 5 to 6 year intervals (or upon new exposure) and (2) for new exposures in previously treated persons a single booster inoculum (again preferably of avian embryo vaccine) is probably adequate if the most recent treatment occurred no longer than 20 years previously.

Comment

The heart of the problem confronting the physician in the after the bite situation is the weighing of the evidence that true exposure has taken place against the risk known to be associated with conventional antirabic vaccination. This has been delineated in classic manner by T. F. Sellers²¹ in a paper aptly entitled *Rabies: the physician's dilemma* which closed with the caution that for exposures other than actual bites the danger of treatment complications far exceeds that of either rabies or rabophobia. These words were published in 1948. At present although these words of caution must be borne in mind at least a little longer use of the avian embryo vaccines seems to ease the problem considerably especially in the case of those who already have undergone one course of Pasteur treatment. Meanwhile to ensure the best possible response in the emergency situation efforts should continue to increase the antigenic potency of the embryo vaccines and to improve the methods for their use. The other major problem posed by instances of severe exposure with probability of disease after a short incubation period has been greatly eased by the advent of antirabies hyperimmune serum. However room for important improvements remains in the direction of increasing the potency and degree of purification in order to reduce the volume of the inoculation the average size of which is now 40 to 50 ml per adult and the likelihood of serum sickness reactions.

Part V Ornithosis

ORNITHOSIS IN DOMESTIC FOWL LOWER FINDINGS IN TURKEYS

By John P. Delaplane*

*Department of Veterinary Microbiology Agricultural and Mechanical College of Texas and
Texas Agricultural Experiment Station College Station Texas*

The public health aspect of ornithosis infection in turkeys has been of interest since 1951 when Irons Sullivan and Rowen¹ published serologic and epidemiologic studies on an outbreak that occurred among poultry dressing plant workers in Giddings Texas in 1948. No additional infections involving such workers were recognized until November and December 1951 and in April 1952 when infections in poultry plant workers again were observed in the same plant that had experienced the 1948 outbreak reported by Irons *et al*^{1, 2}

As a result of the outbreaks in 1951 and 1952 attempts were made to recover ornithosis virus from turkeys showing lower respiratory involvements because it was suspected that ornithosis might be masked by the lower form of infectious sinusitis. In 1952 Bonev *et al*³ isolated an agent from turkeys that was later identified as ornithosis virus.

Much interest was stimulated in ornithosis infection in turkeys in 1952 when a number of laboratory workers inadvertently became infected with ornithosis of turkey origin thus demonstrating the high pathogenicity of this agent for man. No further outbreaks of ornithosis were recognized in turkeys or processing plant workers in Texas until April May and June of 1954 when approximately 200 nonfatal cases occurred in five processing plants in Texas. A number of isolations of virus were made from turkey flocks in that state by Pate Bonev and Delaplane⁴.

Following recognition of the disease in Texas in 1952 and 1954 infections traceable to turkeys have been reported from New Jersey⁵ and Oregon^{6, 7}. According to Andrews⁸ ornithosis virus has been isolated from turkeys in seven states.

Since 1955 a comprehensive research program on ornithosis as it affects turkeys has been under way at the Texas Agricultural Experiment Station. Financial support for the research has been provided by the Department of Agriculture Agricultural Research Service Washington D C the Division of Research Grants National Institutes of Health Bethesda Md American Cyanamid Co Research Division Pearl River N Y Merck & Co Rahway N J and Hess and Clark Inc Ashland Ohio.

The following phases of work have received special emphasis (1) the gross lesions of the disease in turkeys from one through twenty four weeks (2) the effect of chlortetracycline in the feed on the course of the infection in turkeys on a prophylactic basis and as therapy following infection (3) the possibility

Deceased

- 11 JOHNSON H N & T F SELLERS 1948 Rabies *In Diagnostic Procedures for Virus and Rickettsial Diseases* 1st ed Thomas Francis Jr Chairman Committee on Diagnostic Procedures for Virus and Rickettsial Diseases Am Public Health Assoc
- 12 SHAUGHNESSY H J & J ZICHIS 1954 Treatment of wounds inflicted by rabid animals Bull World Health Organization 10 805
- 13 PEREZ GALLARDO F 1951 Bull World Health Organization To be published
- 14 HABEL K 1945 Seroprophylaxis in experimental rabies Public Health Repts U S 60 545
- 15 SHORTT H E J P MCGUIRE A G BROOKS & E D STEPHENS 1955 Anti-rabic immunization probable lines of progress in improvement of methods Indian J Med Research 22 537
- 16 PROCA G & S BABES 1940 Anti-rabic immunization living vaccines and killed vaccines Bull World Health Organization 9 79
- 17 KOPROWSKI H J VAN DER SCHEER & J BLACK 1950 Use of hyperimmune anti-rabies serum concentrates in experimental rabies Am J Med 8 412
- 18 SELLERS T F 1953 Rabies—Immunization and management of human exposures Okla Acad Gen Pract Oklahoma City Okla
- 19 BALTAZARD M & M BAHMANYAR 1955 Essai pratique du sérum antirabique chez les mordus par loups enragés Bull World Health Organization 13 747
- 20 HABEL K & H KOPROWSKI 1955 Laboratory data supporting the clinical trial of antirabies serum in persons bitten by a rabid wolf Bull World Health Organization 13 773
- 21 ATANASIU P M BALTAZARD J P FOX K HABEL M KAPLAN I F KISSLING A KOMAROV H KOPROWSKI I LÉPINE & I PEREZ GALLARDO 1956 Rabies neutralizing antibody response to different schedules of serum and vaccine inoculations in non-exposed persons Bull World Health Organization 14 593
- 22 ATANASIU P M BALTAZARD J P FOX K HABEL M KAPLAN R F KISSLING A KOMAROV H KOPROWSKI P LÉPINE & F PEREZ GALLARDO 1957 Bull World Health Organization To be published
- 23 SELLERS T F 1947 Complications of antirabic treatment J Med Assoc Georgia 36 30
- 24 PAIT C F & H E PEARSON 1949 Rabies vaccine encephalomyelitis in relation to the incidence of animal rabies in Los Angeles Am J Public Health 39 875
- 25 MCKENDRICK A G 1940 A ninth analytical review of reports from Pasteur Institutes on the results of anti-rabies treatment Bull World Health Organization 9 31
- 26 KOPROWSKI H & H R COX 1948 Studies on chick embryo-adapted rabies virus I Culture and characteristics and pathogenicity J Immunol 60 533
- 27 KOPROWSKI H J BLACK & D J NELSON 1954 Studies on chick embryo-adapted rabies virus VI Further changes in pathogenic properties following prolonged cultivation in the developing chick embryo J Immunol 72 94
- 28 POWELL H M & C C CLBERTSON 1950 Cultivation of fixed rabies virus in embryonated duck eggs Public Health Repts U S 65 400
- 29 MACFARLANE J O & C G CLBERTSON 1954 Attempted production of allergic encephalomyelitis with duck embryo suspensions and vaccines Can J Public Health 45 28
- 30 JOHNSON H N 1954 Experimental studies on the duration of immunity in dogs vaccinated against rabies Bull World Health Organization 10 725
- 31 SELLERS T F 1948 Rabies the physician's dilemma Am J Trop Med 28 453

ORNITHOSIS IN DOMESTIC FOWL AND FINDINGS IN TURKEYS

By John P. Delaplane*

*Department of Veterinary Microbiology and Immunology and Mechanical College of Texas and
Texas Agricultural Experiment Station College Station Texas*

The public health aspect of ornithosis infection in turkeys has been of interest since 1951 when Irons, Sullivan, and Rowen¹ published serologic and epidemiologic studies on an outbreak that occurred among poultry-dressing plant workers in Cuddings, Texas, in 1948. No additional infections involving such workers were recognized until November and December 1951 and in April 1952 when infections in poultry plant workers again were observed in the same plant that had experienced the 1948 outbreak reported by Irons *et al.*^{1, 2}

As a result of the outbreaks in 1951 and 1952 attempts were made to recover ornithosis virus from turkeys showing lower respiratory involvement because it was suspected that ornithosis might be masked by the lower form of infectious sinusitis. In 1952 Boney *et al.*³ isolated an agent from turkeys that was later identified as ornithosis virus.

Much interest was stimulated in ornithosis infection in turkeys in 1952 when a number of laboratory workers inadvertently became infected with ornithosis of turkey origin, thus demonstrating the high pathogenicity of this agent for man. No further outbreaks of ornithosis were recognized in turkeys or processing plant workers in Texas until April, May, and June of 1954 when approximately 200 nonfatal cases occurred in five processing plants in Texas. A number of isolations of virus were made from turkey flocks in that state by Pate, Boney, and Delaplane.⁴

Following recognition of the disease in Texas in 1952 and 1954, infections traceable to turkeys have been reported from New Jersey⁵ and Oregon.^{6, 7} According to Andrews, ornithosis virus has been isolated from turkeys in seven states.

Since 1955 a comprehensive research program on ornithosis as it affects turkeys has been under way at the Texas Agricultural Experiment Station. Financial support for the research has been provided by the Department of Agriculture, Agricultural Research Service, Washington, D. C.; the Division of Research Grants, National Institutes of Health, Bethesda, Md.; American Cyanamid Co., Research Division, Pearl River, N. Y.; Merck & Co., Rahway, N. J.; and Hess and Clark, Inc., Ashland, Ohio.

The following phases of work have received special emphasis: (1) the gross lesions of the disease in turkeys from one through twenty-four weeks; (2) the effect of chlortetracycline in the feed on the course of the infection in turkeys on a prophylactic basis and as therapy following infection; (3) the possibility

of egg transmission of the virus to the newly hatched poult (4) serology studies using the indirect complement fixation and other serologic tests to study sera from turkeys under field and experimental conditions (5) potential reservoirs of virus in nature through the screening of representative free flying birds and rodents from six turkey farms under study

More recently specimens of wading birds have been obtained from the rookeries along the coast of the Gulf of Mexico

Gross Lesions

Extensive studies of the gross lesions of ornithosis in turkeys by Davis *et al*⁹ confirmed the preliminary observations of Boney *et al*² and of Pate Boney and Delaplane⁴ and showed that the major lesions consisted of a high incidence of caseofibrinous pericarditis perihepatitis occurring as a fibrinous coating or plastic exudate biliary stasis caseofibrinous peritonitis caseofibrinous exudates on the air sacs pneumonia and splenomegaly In experimental infections the mortality⁹ varied with the age of the turkey It was almost 100 per cent in 1 week old poults but declined to approximately 20 to 25 per cent in the 12 to 24 week old birds The incubation period varied from 4 days in young poults to 7 days in older turkeys

Microscopic studies of the lesions in tissues collected from the gross pathology phase of the study are in progress

In limited studies⁹ evidence of airborne transmission of ornithosis in turkeys could not be proved

Prophylaxis and Therapy

Prophylactic therapy using chlortetracycline at the rate of 100 200 400 600 and 800 gm per ton of feed as reported by Davis and Delaplane^{9 11} showed that all levels were effective in preventing mortality in 3 week old experimentally infected poults over a 2 week period of observation Extensive symptoms and lesions were observed in the inoculated poults on the 100-gm level of antibiotic virus was easily recovered from these poults as well as from the untreated infected controls Only slight lesions were observed in the poults receiving 200 gm of antibiotic per ton of feed and virus was recovered only from the poults with lesions No symptoms or lesions were observed in the poults receiving 400 600 and 800 gm of antibiotic per ton of feed and virus was not recovered from any of these poults

Recently Davis and Watkins¹² studied furazolidone at the rate of 100 200 and 300 gm per ton of feed in the prevention of ornithosis using similar procedures as for the chlortetracycline studies and they found it to be ineffective

The therapy studies of Davis and Delaplane¹ using chlortetracycline at the rate of 100 200 400 and 800 gm per ton of feed for 2 weeks in 3 week-old infected turkeys showed that such levels did not result in the elimination of virus When similar treatment was given for 3 weeks 100 gm was ineffective but 200 gm and more suppressed the infection

Infected adult turkeys receiving 200 and 400 gm of chlortetracycline per ton of feed for 2 or 3 weeks were able to overcome the virus Ornithosis virus

was recovered from untreated infected adults 24 days after inoculation but not after 31 days

In the fall of 1956 Davis Watkins and Delaplane¹⁴ reported on the treatment of naturally occurring ornithosis in a farm flock. Attempted therapeutic measures by the owner had not been successful. Later supervised therapy using 200 gm chlortetracycline per ton of feed for 3 weeks suppressed the infection. The flock was processed at a federally inspected plant and ornithosis virus was not recovered. One hundred sick turkeys were removed from the farm and placed under study at the Texas Agricultural Experiment Station Research Farm and given 200 gm of chlortetracycline per ton of feed for 3 weeks. Virus was present in the experimental flock at the end of 2 weeks but not after 3 weeks.

Egg Transmission

Preliminary studies on the role of turkey eggs in the transmission of ornithosis by Davis and Delaplane¹⁵ revealed no evidence of transmission. More extensive studies were undertaken by these investigators¹⁶ and no virus was recovered from 2954 eggs laid by breeder turkeys experimentally infected with the Jo strain of ornithosis although 300 of the eggs were laid during the time at which virus could be recovered from the blood stream of representative birds. When fresh turkey eggs were inoculated with ornithosis virus and incubated the virus was not recovered after the ninth day of incubation. When 1 day-old embryonating turkey eggs were inoculated the embryo mortality rate was high and the virus was not recovered beyond the sixteenth day in the few that survived.

Further studies showed that the virus was not recovered from eggs laid by hens in the viremic stage of ornithosis when isolation attempts were made on freshly laid and 24-hour incubated eggs. Ornithosis virus failed to live longer than 3 days on turkey egg shells or longer than 4 days on cotton swabs under egg incubator conditions.

Serologic Findings

Since 1955 turkey serum samples obtained from the pullorum testing laboratories representing Texas turkey breeding flocks have been tested using the indirect complement fixation (ICF) test. Approximately 50 per cent of the flocks showed reactions at 1:2, 1:4, 1:8 and 1:16 but the history of such flocks did not indicate the presence of ornithosis infection. Recently Neal and Davis¹⁷ compared the efficiency of the direct complement fixation (DCF) of Benedict^{18, 19} and an agglutination test with the ICF tests on the same serum samples of experimental and farm source of flocks. The agglutination test as used was of no value however there was about 85 per cent correlation between the ICF and the Benedict DCF test results.

During and immediately following the age of maturity of one of the experimental groups on study approximately 30 per cent of these known laboratory reared ornithosis free turkeys showed reactions much like those experienced under farm flock conditions. Later representative birds from the ornithosis-

of egg transmission of the virus to the newly hatched poult (4) serology studies using the indirect complement fixation and other serologic tests to study sera from turkeys under field and experimental conditions (5) potential reservoirs of virus in nature through the screening of representative free flying birds and rodents from six turkey farms under study

More recently specimens of wading birds have been obtained from the rookeries along the coast of the Gulf of Mexico

Gross Lesions

Extensive studies of the gross lesions of ornithosis in turkeys by Davis *et al*³ confirmed the preliminary observations of Boney *et al*³ and of Pate Boney and Delaplane⁴ and showed that the major lesions consisted of a high incidence of caseofibrinous pericarditis perihepatitis occurring as a fibrinous coating or plastic exudate biliary stasis caseofibrinous peritonitis caseofibrinous exudates on the air sacs pneumonia and splenomegaly. In experimental infections the mortality³ varied with the age of the turkey. It was almost 100 per cent in 1 week old poults but declined to approximately 20 to 25 per cent in the 12 to 24 week old birds. The incubation period varied from 4 days in young poults to 7 days in older turkeys.

Microscopic studies of the lesions in tissues collected from the gross pathology phase of the study are in progress.

In limited studies³ evidence of airborne transmission of ornithosis in turkeys could not be proved.

Prophylaxis and Therapy

Prophylactic therapy using chlortetracycline at the rate of 100 200 400 600 and 800 gm per ton of feed as reported by Davis and Delaplane^{9 11} showed that all levels were effective in preventing mortality in 3 week old experimentally infected poults over a 2 week period of observation. Extensive symptoms and lesions were observed in the inoculated poults on the 100 gm level of antibiotic virus was easily recovered from these poults as well as from the untreated infected controls. Only slight lesions were observed in the poults receiving 200 gm of antibiotic per ton of feed and virus was recovered only from the poults with lesions. No symptoms or lesions were observed in the poults receiving 400 600 and 800 gm of antibiotic per ton of feed and virus was not recovered from any of these poults.

Recently Davis and Watkins¹³ studied furazolidone at the rate of 100 200 and 300 gm per ton of feed in the prevention of ornithosis using similar procedures as for the chlortetracycline studies and they found it to be ineffective.

The therapy studies of Davis and Delaplane¹ using chlortetracycline at the rate of 100 200 400 and 800 gm per ton of feed for 2 weeks in 3 week old infected turkeys showed that such levels did not result in the elimination of virus. When similar treatment was given for 3 weeks 100 gm was ineffective but 200 gm and more suppressed the infection.

Infected adult turkeys receiving 200 and 400 gm of chlortetracycline per ton of feed for 2 or 3 weeks were able to overcome the virus. Ornithosis virus

order of effectiveness as tetracycline oxytetracycline erythromycin chlortetracycline and penicillin Dihydrostreptomycin had no apparent effect on the virus

Roth and Delaplane⁴ reported that oxymycin was ineffective against 6 BC psittacosis virus employing chicken embryos Davis Watkins and Delaplane⁵ showed that synnemetin B was ineffective except possibly at very high dosage levels using chicken embryos

References

- 1 IRONS J V T D SULLIVAN & J FOWEN 1951 Outbreak of psittacosis (ornithosis) from working with turkeys & chickens *Am J Public Health* 41 931
- 2 IRONS J V M L DENLEY & F D SULLIVAN 1955 Psittacosis in Turkeys and Poultry as a Source of Human Infection Psittacosis Diagnosis Epidemiology and Control F R Beaudette Ed Rutgers Univ Press New Brunswick N J
- 3 BONEY W A I C GRUMBLES J P DELAPLANE J V IRONS & T D SULLIVAN 1952 Another agent causing air sac involvement in turkeys *J Am Vet Med Assoc* 121 269
- 4 PATE D D W A BONEY & J I DELAPLANE 1954 Turkey ornithosis I A report of natural and experimental infections *J Am Vet Med Assoc* 125 4 6-478
- 5 BEAUBETTE F R C B HUDON & V K KASCHULA 1956 An outbreak of psittacosis in turkeys *Vet Med* 51 436
- 6 O GOOD S B M A HOLMES D M MASON G CAPLAN K F MEYER & B LIDDE 1956 Ornithosis in Oregon A report of outbreaks in turkeys and man Oregon State Board of Health U S Public Health Service Communicable Disease Center and The George Williams Hooper Foundation Univ Calif Med Center San Francisco Calif
- 7 DICKINSON F M W L BABCOCK & J C KILIAN 1957 Ornithosis in Oregon turkeys *J Am Vet Med Assoc* 130 117-119
- 8 ANDREWS J M 1957 The importance of psittacosis in the United States *J Am Vet Med Assoc* 130 109-116
- 9 DAVIS D E & J P DELAPLANE 1956 The lesions of ornithosis in turkeys *In Psittacosis* F R Beaudette Ed Rutgers Univ Press New Brunswick N J In press
- 10 DAVIS D F & J P DELAPLANE 1956 Turkey ornithosis therapy and control *Proc U S Livestock Sanitary Assoc* 59th Ann Rept 250
- 11 DAVIS D F & J P DELAPLANE 1956 Turkey ornithosis prophylaxis using chlorotetracycline *In Psittacosis* F R Beaudette Ed Rutgers Univ Press New Brunswick N J In press
- 12 DAVIS D E & J P DELAPLANE 1957 The treatment of ornithosis with chlorotetracycline *Am J Vet Research* In press
- 13 DAVIS D F & J R WATKINS 1957 Furazolidone as a prophylactic agent against turkey ornithosis *Avian Diseases* In press
- 14 DAVIS D E J R WATKINS & J P DELAPLANE 1957 The treatment of turkey ornithosis in a farm flock *Southwestern Vet* 3 223
- 15 DAVIS D E & J P DELAPLANE 1955 Ornithosis in turkeys *Proc Book Am Vet Med Assoc* 296-301
- 16 DAVIS D F & J P DELAPLANE 1957 The role of turkey eggs in the transmission of ornithosis *Am J Vet Research* 18 409-413
- 17 NEAL J E & D L DAVIS 1957 A comparison of the indirect complement fixation test the direct complement fixation test and the microscopic agglutination test for ornithosis antiserum in turkey sera *Am J Vet Research* To be published
- 18 BENEDICT A A & C McFARLAND 1956 Direct complement fixation test for diagnosis of ornithosis in turkeys *Proc Soc Exptl Biol Med* 92 168
- 19 BENEDICT A A & E O BRIEN 1956 Antigenic studies on the psittacosis lymphogranuloma cereum group of viruses II Characterization of complement fixing antigens extracted with sodium lauryl sulfate *J Immunol* 76 293
- 20 BENEDICT A A F O BRIEN & C McFARLAND 1956 Antigenic studies on the psittacosis lymphogranuloma cereum group of viruses III Detection of ornithosis hypersemitivity in experimentally infected chickens *J Am Vet Research* *Abstr Soc Exptl Biol Med* 94 146
- 21 BRUMFIELD H P & B S POMEROY 1957 Direct complement fixation by turkey and chicken serum in viral systems *Proc Soc Exptl Biol Med* 94 146

free groups gave negative reactions after having shown the type of reaction mentioned. This indicated that factors possibly unrelated to ornithosis can account for positive reactions to the ICF test. To determine whether Benedict's DCF test would have also picked positive reacting serums, 22 of the positive reacting samples were tested by Benedict. 8 of these were positive 1/4 or 1/8 by this method. The Benedict skin allergy test⁶ gave negative findings in representative birds. Twenty one of the same positive turkey samples were tested by Brumfield, using her DCF test¹ and 10 of the samples showed positive reactions. These results indicate that some unknown factors other than ornithosis account for nonspecific reactions using these various testing procedures. Work is now in progress to determine further something of the nature of the explanation for these reactions. Volkert *et al*² have shown that a bacterial antigen made from an organism identified by them as *Bacterium antitratum* possesses a high degree of activity against human psittacosis antisera; possibly something of this nature is involved in some nonspecific reactions.

Potential Virus Reservoirs

In the study of reservoirs of possible sources of ornithosis virus in nature six farms in the area in which the original virus was isolated in 1952 were selected for routine observation and recording of the various species of free flying birds, rodents, and animals having contact with the turkeys on range. Approximately 2500 birds and mammals have been screened for virus to date. So far as possible, serum samples have been collected for serologic studies. All sick and dead birds are kept frozen on the farms and are submitted to the laboratory once each week for examination. To date ornithosis virus has not been isolated from any of the specimens. In two ornithosis outbreaks in Texas in 1956 it was observed that in both instances the turkeys had access to or were near bodies of water that attracted water fowl such as herons and egrets. During the spring and summer of 1957 specimens sick and dead and screened for ornithosis virus with negative results have been collected. E. M. Dickinson, Oregon State College, Corvallis, Ore. (personal communication) found ornithosis virus in sea gulls from an affected farm in Oregon. It is hoped that eventually this phase of the study may yield findings of importance regarding reservoirs of ornithosis virus.

Pathogenicity Studies

Studies involving a comparison of the symptoms and lesions of various strains of psittacosis and ornithosis virus for various age groups of turkeys are in progress. The isolates from turkeys appear to be much more pathogenic than those from parakeets and parrots. It would appear from the studies to date that the occurrence of ornithosis in turkeys is sporadic and accidental in nature and that the strain of virus involved is of a highly pathogenic nature at the time it becomes introduced into a flock.

Antibiotic Activity

Studies by Davis²³ involving the screening of six antibiotics for activity against the Jo strain of ornithosis virus using chicken embryos indicate the

order of effectiveness as tetracycline oxytetracycline erythromycin chlortetracycline and penicillin Dihydrostreptomycin had no apparent effect on the virus

Koth and Delaplane⁴ reported that oxytetracycline was ineffective against 6 BC psittacosis virus employing chicken embryos Davis Watkins and Delaplane³ showed that synnemetin B was ineffective except possibly at very high dosage level using chicken embryos

References

- 1 IRONS J V T D SULLIVAN & J ROWEN 1951 Outbreak of psittacosis (ornithosis) from working with turkeys or chickens *Am J Public Health* **41** 931
- 2 IRON J V M L DENLEY & T D SULLIVAN 1955 Psittacosis in Turkeys and Fowl as a Source of Human Infection Psittacosis Diagnosis Epidemiology and Control F K Beaudette Ed Rutgers Univ Press New Brunswick N J
- 3 BONEY W A I C GRUMBLES J P DELAPLANE J V IRONS & T D SULLIVAN 1952 Another agent causing air sac involvement in turkeys *J Am Vet Med Assoc* **121** 769
- 4 PATE D D W A BONEY & J P DELAPLANE 1954 Turkey ornithosis I A report of natural and experimental infections *J Am Vet Med Assoc* **125** 476-478
- 5 BEAUDETTE F R C B HUDON & V K KASCHULA 1956 An outbreak of psittacosis in turkeys *Vet Med* **51** 496
- 6 O-GOOD S B M A HOLMES D M MASON C CAPLAN K F MEYER & B EDDIE 1956 Ornithosis in Oregon A report of outbreaks in turkeys and man Oregon State Board of Health U S Public Health Service Communicable Disease Center and The George Williams Hooper Foundation Univ Calif Med Center San Francisco Calif
- 7 DICKINSON E M W E BARCOCK & J G KILIAN 1957 Ornithosis in Oregon turkeys *J Am Vet Med Assoc* **130** 117-119
- 8 ANDREWS J M 1957 The importance of psittacosis in the United States *J Am Vet Med Assoc* **130** 109-116
- 9 DAVIS D E & J P DELAPLANE 1956 The lesions of ornithosis in turkeys In Psittacosis F R Beaudette Ed Rutgers Univ Press New Brunswick N J In press
- 10 DAVIS D E & J P DELAPLANE 1956 Turkey ornithosis therapy and control Proc U S Livestock Sanitary Assoc 9th Ann. Rept 250
- 11 DAVIS D E & J P DELAPLANE 1956 Turkey ornithosis prophylaxis using chlor tetracycline In Psittacosis F R Beaudette Ed Rutgers Univ Press New Brunswick N J In press
- 12 DAVIS D E & J P DELAPLANE 1957 The treatment of ornithosis with chlortetracycline *Am J Vet Research* In press
- 13 DAVIS D E & J K WATKINS 1957 Fusidic acid as a prophylactic agent against turkey ornithosis *Avian Diseases* In press
- 14 DAVIS D E J R WATKINS & J P DELAPLANE 1957 The treatment of turkey ornithosis in a farm flock *Southwestern Vet* **3** 223
- 15 DAVIS D E & J P DELAPLANE 1955 Ornithosis in turkeys Proc Book Am Vet Med Assoc 226-301
- 16 DAVIS D E & J P DELAPLANE 1957 The role of turkey eggs in the transmission of ornithosis *Am J Vet Research* **18** 409-413
- 17 NEAL J E & D E DAVIS 1957 A comparison of the indirect complement fixation test the direct complement fixation test and the macroscopic agglutination test for ornithosis antiserum in turkey sera *Am J Vet Research* To be published
- 18 BENEDICT A A & C MCFARLAND 1956 Direct complement fixation test for diagnosis of ornithosis in turkeys Proc Soc Exptl Biol Med **92** 68
- 19 BENEDICT A A & L O'BRIEN 1956 Antigenic studies on the psittacosis lymphogranuloma venereum group of viruses II Characterization of complement fixing antigens extracted with sodium lauryl sulfate *J Immunol* **76** 293
- 20 BENEDICT A A F O'BRIEN & C MCFARLAND 1956 Antigenic studies on the psittacosis lymphogranuloma venereum group of viruses III Detection of ornithosis hypersensitivity in experimentally infected chickens *J Am Vet Research Also Soc Exptl Biol Med* **94** 146
- 21 BRUMFIELD H P & B S POMEROY 1957 Direct complement fixation by turkey and chicken serum in viral systems Proc Soc Exptl Biol Med **94** 146

free groups gave negative reactions after having shown the type of reaction mentioned. This indicated that factors possibly unrelated to ornithosis can account for positive reactions to the ICF test. To determine whether Benedict's DCF test would have also picked positive reacting serums 22 of the positive reacting samples were tested by Benedict, 8 of these were positive 14 or 16 by this method. The Benedict skin allergy test⁹ gave negative findings in representative birds. Twenty one of the same positive turkey samples were tested by Brumfield, using her DCF test¹¹ and 10 of the samples showed positive reactions. These results indicate that some unknown factors other than ornithosis account for nonspecific reactions using these various testing procedures. Work is now in progress to determine further something of the nature of the explanation for these reactions. Volkert *et al.* have shown that a bacterial antigen made from an organism identified by them as *Bacterium antitratum* possesses a high degree of activity against human psittacosis anti sera possibly something of this nature is involved in some nonspecific reactions.

Potential Virus Reservoirs

In the study of reservoirs of possible sources of ornithosis virus in nature, six farms in the area in which the original virus was isolated in 1952 were selected for routine observation and recording of the various species of free flying birds, rodents, and animals having contact with the turkeys on range. Approximately 2500 birds and mammals have been screened for virus to date. So far as possible serum samples have been collected for serologic studies. All sick and dead birds are kept frozen on the farms and are submitted to the laboratory once each week for examination. To date ornithosis virus has not been isolated from any of the specimens. In two ornithosis outbreaks in Texas in 1956 it was observed that in both instances the turkeys had access to or were near bodies of water that attracted water fowl such as herons and egrets. During the spring and summer of 1957 specimens sick and dead and screened for ornithosis virus with negative results have been collected. E. M. Dickinson, Oregon State College, Corvallis, Ore. (personal communication) found ornithosis virus in sea gulls from an affected farm in Oregon. It is hoped that eventually this phase of the study may yield findings of importance regarding reservoirs of ornithosis virus.

Pathogenicity Studies

Studies involving a comparison of the symptoms and lesions of various strains of psittacosis and ornithosis virus for various age groups of turkeys are in progress. The isolates from turkeys appear to be much more pathogenic than those from parakeets and parrots. It would appear from the studies to date that the occurrence of ornithosis in turkeys is sporadic and accidental in nature and that the strain of virus involved is of a highly pathogenic nature at the time it becomes introduced into a flock.

Antibiotic Activity

Studies by Davis¹ involving the screening of six antibiotics for activity against the Jo strain of ornithosis virus using chicken embryos indicate the

order of effectiveness as tetracycline oxytetracycline erythromycin chlortetracycline and penicillin Dihydrostreptomycin had no apparent effect on the virus

Roth and Delaplane⁴ reported that oxamycin was ineffective against 6 BC psittacosis virus employing chicken embryos Davis Watkins and Delaplane⁵ showed that synnemetin B was ineffective except possibly at very high dosage levels using chicken embryos

References

- 1 IRONS J V T D SULLIVAN & J ROWEN 1951 Outbreak of psittacosis (ornithosis) from working with turkeys or chickens *Am J Public Health* 41 931
- 2 IRONS J V M I DENSEY & T D SULLIVAN 1953 Psittacosis in Turkey and Fowls as a Source of Human Infection *Psittacosis Diagnosis Epidemiology and Control* F K Beaullette Ed Rutgers Univ Press New Brunswick N J
- 3 BONEY W A I C GRUMBLES J I DELAPLANE J V IRONS & T D SULLIVAN 1952 Another agent causing air-borne involvement in turkeys *J Am Vet Med Assoc* 121 269
- 4 PATE D D W A BONEY & J I DELAPLANE 1954 Turkey ornithosis I A report of natural and experimental infections *J Am Vet Med Assoc* 125 476-478
- 5 BEAULETTE F K C B HEDSON & V K KASCHULA 1956 An outbreak of psittacosis in turkeys *Vet Med* 51 496
- 6 OGDEN S B M A HOLMES D M MASON G CAPLAN K F MEYER & B EDDIE 1956 Ornithosis in Oregon A report of outbreaks in turkeys and man Oregon State Board of Health U S Public Health Service Communicable Disease Center and The George Williams Hooper Foundation Univ Calif Med Center San Francisco Calif
- 7 DICKINSON F M W I BABCOCK & J C KELTAN 1957 Ornithosis in Oregon turkeys *J Am Vet Med Assoc* 130 117-119
- 8 ANDREWS J M 1957 The importance of psittacosis in the United States *J Am Vet Med Assoc* 130 109-116
- 9 DAVIS D F & J I DELAPLANE 1956 The lesions of ornithosis in turkeys *In Psittacosis* F K Beaullette Ed Rutgers Univ Press New Brunswick N J In press
- 10 DAVIS D F & J P DELAPLANE 1956 Turkey ornithosis therapy and control *Proc U S Livestock Sanitary Assoc 59th Ann Rept* 250
- 11 DAVIS D F & J I DELAPLANE 1956 Turkey ornithosis prophylaxis using chlortetracycline *In Psittacosis* F K Beaullette Ed Rutgers Univ Press New Brunswick N J In press
- 12 DAVIS D F & J P DELAPLANE 1957 The treatment of ornithosis with chlortetracycline *Am J Vet Research* In press
- 13 DAVIS D F & J R WATKINS 1957 Furazolidone as a prophylactic agent against turkey ornithosis *Avian Diseases* In press
- 14 DAVIS D F J R WATKINS & J P DELAPLANE 1957 The treatment of turkey ornithosis in a farm flock *Southwestern Vet* 3 273
- 15 DAVIS D F & J P DELAPLANE 1955 Ornithosis in turkeys *Proc Book Am Vet Med Assoc* 296-301
- 16 DAVIS D F & J P DELAPLANE 1957 The role of turkey eggs in the transmission of ornithosis *Am J Vet Research* 18 409-413
- 17 NEAL J E & D L DAVIS 1957 A comparison of the indirect complement fixation test the direct complement fixation test and the macroscopic agglutination test for ornithosis antibodies in turkey sera *Am J Vet Research* To be published
- 18 BEAULETTE F K & C McFARLAND 1956 Direct complement fixation test for diagnosis of ornithosis in turkeys *Proc Soc Exptl Biol Med* 92 168
- 19 BEAULETTE F K & L O'BRIEN 1956 Antigenic studies on the psittacosis lymphogranuloma virus group of viruses II Characterization of complement fixing antigens extracted with sodium lauryl sulfate *J Immunol* 76 293
- 20 BEAULETTE F K A E O'BRIEN & C McFARLAND 1956 Antigenic studies on the psittacosis lymphogranuloma virus group of viruses III Detection of ornithosis hyper-sensitivity in experimentally infected chicken *J Am Vet Research Assoc Soc Exptl Biol Med* 91 146
- 21 BRUMFIELD H T & B S I JMEROV 1957 Direct complement fixation by turkey and chicken serum in viral systems *Proc Soc Exptl Biol Med* 94 146

free groups gave negative reactions after having shown the type of reaction mentioned. This indicated that factors possibly unrelated to ornithosis can account for positive reactions to the ICF test. To determine whether Benedict's DCF test would have also picked positive reacting serums 22 of the positive reacting samples were tested by Benedict. 8 of these were positive 14 or 18 by this method. The Benedict skin allergy test⁹ gave negative findings in representative birds. Twenty one of the same positive turkey samples were tested by Brumfield, using her DCF test²¹ and 10 of the samples showed positive reactions. These results indicate that some unknown factors other than ornithosis account for nonspecific reactions using these various testing procedures. Work is now in progress to determine further something of the nature of the explanation for these reactions. Volkert *et al*²² have shown that a bacterial antigen made from an organism identified by them as *Bacterium antitratum* possesses a high degree of activity against human psittacosis anti sera possibly something of this nature is involved in some nonspecific reactions.

Potential Virus Reservoirs

In the study of reservoirs of possible sources of ornithosis virus in nature six farms in the area in which the original virus was isolated in 1952 were selected for routine observation and recording of the various species of free flying birds, rodents and animals having contact with the turkeys on range. Approximately 2500 birds and mammals have been screened for virus to date. So far as possible serum samples have been collected for serologic studies. All sick and dead birds are kept frozen on the farms and are submitted to the laboratory once each week for examination. To date ornithosis virus has not been isolated from any of the specimens. In two ornithosis outbreaks in Texas in 1956 it was observed that in both instances the turkeys had access to or were near bodies of water that attracted water fowl such as herons and egrets. During the spring and summer of 1957 specimens sick and dead and screened for ornithosis virus with negative results have been collected. E. M. Dickinson, Oregon State College, Corvallis, Ore. (personal communication) found ornithosis virus in sea gulls from an affected farm in Oregon. It is hoped that eventually this phase of the study may yield findings of importance regarding reservoirs of ornithosis virus.

Pathogenicity Studies

Studies involving a comparison of the symptoms and lesions of various strains of psittacosis and ornithosis virus for various age groups of turkeys are in progress. The isolates from turkeys appear to be much more pathogenic than those from parakeets and parrots. It would appear from the studies to date that the occurrence of ornithosis in turkeys is sporadic and accidental in nature and that the strain of virus involved is of a highly pathogenic nature at the time it becomes introduced into a flock.

Antibiotic Activity

Studies by Davis²³ involving the screening of six antibiotics for activity against the Jo strain of ornithosis virus using chicken embryos indicate the

order of effectiveness as tetracycline oxytetracycline erythromycin chlortetracycline and penicillin Dihydro streptomycin had no apparent effect on the virus

Roth and Delaplane⁴ reported that oxamycin was ineffective against 6 BC psittacosis virus employing chicken embryos Davis Watkins and Delaplane⁵ showed that synnemetin B was ineffective except possibly at very high dosage levels using chicken embryos

References

- 1 IRONS J V T D SULLIVAN & J KOWEN 1951 Outbreak of psittacosis (ornithosis) from working with turkeys or chickens *Am J Public Health* 41 931
- 2 IRONS J V M L DENLEY & T D SULLIVAN 1955 Psittacosis in Turkeys and Fowls as a Source of Human Infection Psittacosis Diagnosis Epidemiology and Control F R Beaudette Ed Rutgers Univ Press New Brunswick N J
- 3 BONEY W A I C GRUMBLES J I DELAPLANE J V IRONS & T D SULLIVAN 1952 Another agent causing air sac involvement in turkeys *J Am Vet Med Assoc* 121 269
- 4 PATE D D W A BONEY & J I DELAPLANE 1954 Turkey ornithosis I A report of natural and experimental infections *J Am Vet Med Assoc* 125 476-478
- 5 BEAUDETTE F R C B HUDSON & A K KASCHULA 1956 An outbreak of psittacosis in turkeys *Vet Med* 51 496
- 6 OSGOOD S B M A HOLMES D M MASON C CAPLAN K F MEYER & B FODIE 1956 Ornithosis in Oregon A report of outbreaks in turkeys and man Oregon State Board of Health U S Public Health Service Communicable Disease Center and The George Williams Hooper Foundation Univ Calif Med Center San Francisco Calif
- 7 DICKINSON F M W F BABCOCK & J C KILIAN 1957 Ornithosis in Oregon turkeys *J Am Vet Med Assoc* 130 117-119
- 8 ANDREWS J M 1951 The importance of psittacosis in the United States *J Am Vet Med Assoc* 130 109-116
- 9 DAVIS D E & J P DELAPLANE 1956 The lesions of ornithosis in turkeys *In Psittacosis* F R Beaudette Ed Rutgers Univ Press New Brunswick N J In press
- 10 DAVIS D F & J P DELAPLANE 1956 Turkey ornithosis therapy and control *Proc U S Livestock Sanitary Assoc 59th Ann Rept* 250
- 11 DAVIS D E & J I DELAPLANE 1956 Turkey ornithosis prophylaxis using chlor tetracycline *In Psittacosis* F R Beaudette Ed Rutgers Univ Press New Brunswick N J In press
- 12 DAVIS D F & J P DELAPLANE 1957 The treatment of ornithosis with chlortetracycline *Am J Vet Research* In press
- 13 DAVIS D F & J R WATKINS 1957 Furazolidone as a prophylactic agent against turkey ornithosis *American Diseases* In press
- 14 DAVIS D F J R WATKINS & J P DELAPLANE 1957 The treatment of turkey ornithosis in a farm flock *Southwestern Vet* 3 223
- 15 DAVIS D F & J P DELAPLANE 1955 Ornithosis in turkeys *Proc Book Am Vet Med Assoc* 296-301
- 16 DAVIS D F & J P DELAPLANE 1957 The role of turkey eggs in the transmission of ornithosis *Am J Vet Research* 18 409-413
- 17 NEAL J E & D F DAVIS 1957 A comparison of the indirect complement fixation test the direct complement fixation test and the macroscopic agglutination test for ornithosis antibodies in turkey sera *Am J Vet Research* To be published
- 18 BENEDICT A A & C MCFARLAND 1956 Direct complement fixation test for diagnosis of ornithosis in turkeys *Proc Soc Exptl Biol Med* 92 768
- 19 BENEDICT A A & F O BRIEN 1956 Antigenic studies on the psittacosis lymphogranuloma venereum group of viruses II Characterization of complement fixing antigens extracted with sodium lauryl sulfate *J Immunol* 76 223
- 20 BENEDICT A A F O BRIEN & C MCFARLAND 1956 Antigenic studies on the psittacosis-lymphogranuloma venereum group of viruses III Detection of ornithosis hypersensitivity in experimentally infected chickens *J Am Vet Research Assoc Soc Exptl Biol Med* 94 146
- 21 BRUMFIELD H P & B S POMEROY 1957 Direct complement fixation by turkey and chicken serum in viral systems *Proc Soc Exptl Biol Med* 94 146

- 22 VOLBERT M & M MATTHIJSSEN 1956 An ornithosis related antigen from a coccid bacterium *Acta Pathol Microbiol Scand* 39 114
- 23 DAVIS D I 1957 The activity of several antilotics against turkey ornithosis virus in chicken embryos *Avian Diseases* In press
- 24 ROTH F E & J P DELAPLANE 1955 Dosage level and activity of oxytetracycline against psittacosis virus in chicken embryos *Southwestern Vet* 8 154
- 25 DAVIS D L J P WATKINS & J P DELAPLANE 1956 The activity of salmotin against ornithosis virus *Southwestern Vet* 10 47

Discussion of the Paper

OSCAR SLESSMAN (*Bureau of Veterinary Public Health, State of New Jersey Department of Health Trenton N J*) Raymond Fagan and I have recently completed a study based on the treatment of known infected turkeys with 160 mg of oxytetracycline given intramuscularly and into the hanging wattles of the birds. Our report is not yet ready for publication, but I can state that the injection of this amount of antibiotic into birds in the midst of an epidemic resulted in isolation of no virus in birds killed 3 weeks 4 to 7 weeks and as long as 8 weeks thereafter. Spleens were examined and isolation attempts were made on all of the flock after treatment. Virus could be obtained only from birds that had not received treatment. The material used was an intramuscular compound diluted with physiological saline solution to 80 mg/cc. The mineral oil suspension* was one of 80 mg/cc. It should be noted that the intramuscular material discolored the injected area of the musculature and it was found that injection into the wattle appeared to be as effective and did not injure any of the salable portion of the bird.

Supplied by Chas Pfizer & Co Inc Brooklyn N Y

NEWER METHODS FOR DETECTION OF AVIAN ORNITHOSIS*

By Albert A. Benedict† and Clarence McFarland†
*Department of Preventive Medicine and Public Health, University of
Texas Medical Branch, Galveston, Texas*

The increased awareness of the public health importance of turkey ornithosis has stimulated re-examination of the methods used for detection of this infection in domestic fowls. The detection of exposure to ornithosis may be considered to have 3 principal objectives: (1) for diagnostic purposes, (2) to explore the epidemiological facets of the infection chain which ultimately will lead to (3) the development of control measures. Obviously definitive diagnosis is made by isolation and identification of the agent and as evidenced by recent epizootics in turkeys virus isolation is made frequently and easily. In severe epizootics the appearance of high titer circulating antibody in a large number of birds can be considered as direct evidence of infection. Diagnosis is also justified by a significant rise in titer when paired serum samples are available.

The significance of the presence of a few serologic reactors with low antibody titers among large flocks is not clear. There exist several possibilities that might explain these situations such as: (1) the sign of initial ornithosis activity in a flock, (2) the residue of ornithosis infection that was originally manifested with or without clinical ornithosis, (3) the presence of a cross-reacting antibody to an organism similar to the bacterium isolated by Volkert and Matthiesen¹ and (4) the presence of unknown nonspecific serologic reactions. The control of ornithosis will be aided by the accurate interpretation of the significance of these reactions.

Until recently the serologic technique used exclusively for revealing circulating antibody in domestic fowl has been the indirect complement fixation test (ICF) developed by Karrer, Meyer and Eddie. The complement fixation (CF) test was not applied for such studies since among other avian sera Rice² reported the failure of antipullorum turkey sera to fix complement (C) with *S. pullorum* and Eddie and Francis³ observed fixation at only low dilutions with a small number of turkey anti-ornithosis sera. The presence of a specific inhibitor, presumably an incomplete type antibody, was demonstrated in avian noncomplement fixing sera^{2, 3} and thus the ICF was devised for detection of such antibodies^{2, 3, 4}.

It was generally believed therefore that turkeys and chickens do not produce the usual CF antibodies until recent studies showed that detection of turkey CF antibody depended partially upon the type of antigen employed.⁵ Moreover in 1924 Bushnell and Hudson⁶ reported that chicken antipullorum sera fixed C with homologous antigen but that this property was destroyed

This work reported in this paper was supported in part by funds provided under Contract AF 41(657) 93 with the School of Aviation Medicine, USAF Randolph Air Force Base, Texas and in part by Research Grant E 1217 from the National Biological Institute, Public Health Service, Bethesda, Md.

† Present address: Department of Bacteriology, University of Kansas, Lawrence, Kansas.

by heat inactivation of the sera. This early observation was reinvestigated by our laboratory and by Brumfield and Pomeroy,⁹ and it revealed that the CF ability of turkey and chicken ornithosis antibody was destroyed by heat inactivation of the sera if virus particle antigens were employed.

A variety of serologic tests have been made available; however, only the ICF and the direct complement fixation test (DCF) will be considered in this report. In addition, the intradermal allergic test was developed as an ancillary epidemiological tool.¹⁰⁻¹¹ At present there is an urgent need to evaluate and to compare the various methods that have been proposed so that a standard technique may be developed. With this in view, this report presents the development and evaluation of the DCF and allergic tests.

Direct Complement Fixation Using Heat Inactivated Serum

Technique. We have described the extraction of a soluble psittacosis CF antigen with a detergent;¹² this antigen fixed C consistently with turkey anti-ornithosis sera.⁷ Fixation occurred with sera that had been inactivated by heating either at 56° C for 30 min. or at 60° C for 20 min. For this test 4 units of the detergent extracted antigen were used. In studies that compared antigen activities complement was titrated in the presence of antigens. Preliminary incubation was carried out at 37° C for 75 min. and after the addition of sensitized sheep erythrocytes a secondary incubation at 37° C for 30 min. was done. The method used in performing the ICF test was similar to that described by Karrer, Meyer, and Eddie. Titers represent the reciprocal of the highest dilution of serum showing either a 1+ or 2+ reaction in the ICF (partial inhibition) or a 4+ reaction (complete fixation) in the DCF.

Comparison of the DCF and ICF methods. A comparison of these techniques was made with naturally infected and experimentally infected turkeys. Forty-seven turkeys were selected at random from a flock that had undergone a severe epizootic several months prior to bleeding. TABLE 1 shows that 41 sera in this group (A) were positive by both the ICF and DCF and that 3 sera were positive in low dilutions in the DCF and negative in the ICF. Also shown in TABLE 1 are results with sera from 2 flocks (B and C) in which there was no evidence of infection at the time of bleeding. One serum from flock B was negative by the ICF and had a DCF titer of 1:4. Sera obtained from 12 turkeys that had undergone a natural infection 18 months prior to the test were all positive to the DCF and only 5 sera had positive ICF titers. A direct correlation analysis on the basis of the log distribution of the ICF and DCF titers of group A was found to be 0.72, which was significant at the 1 per cent level. Although there is no direct evidence as yet that the same antibody was being measured by the 2 tests, the DCF titers correlated well with the ICF titers of these sera.

In search of a reliable, simple, and economical serologic technique for survey purposes, Neil and Davis¹³ compared the ICF, DCF (detergent antigen) and a macrophage agglutination test¹⁴ that employed stained elementary bodies as antigen. The ICF and DCF agreed in 85.1 per cent of 923 turkey sera tested. This is similar to the findings shown in TABLE 1 in which the 2

TABLE 1
COMPARISON OF THE DIRECT AND INDIRECT COMPLEMENT FIXATION TESTS
FOR THE DETECTION OF TURKEY ORNITHOSIS

Group	History	Number tested	DCF pos ICF pos.	DCF pos ICF neg.	DCF neg ICF pos.	DCF neg ICF neg.	Percent positive		Percent agreement
							DCF	ICF	
A	Natural infection for 3 months	47	41	3	0	3	93.6	87.3	93.6
B	No evidence of in- fection	61	1	1	0	59	3.2	1.6	98.3
C	No evidence of in- fection	50	2	0	0	48	4.0	4.0	100.0
D	Natural infection 18 months be- fore	12	7	5	0	0	100.0	58.3	58.3
F	Experimentally in- fected 2 months	21	15	1	3	2	61.9	85.7	80.9
F	Experimentally in- fected 3 months	12	8	0	0	4	66.6	66.6	100.0
Total		203	64	10	3	116	41.3	37.9	88.5

Direct complement fixation

† Indirect complement fixation

tests agree in 88.5 per cent of 203 turkey sera tested. On the other hand the correlation between the results of the agglutination and ICF tests was 42.3 per cent and between the agglutination and DCF tests the correlation was 40.4 per cent. Many nonspecific reactions were given by the agglutination test by lower serum dilutions, an observation that we had also made relative to an ornithosis precipitin reaction with turkey and chicken sera.¹⁴ On the basis of these comparisons further study with this DCF test was considered justified.¹²

Onset of DCF antibody. It was felt that if incomplete CF antibodies were formed in turkey sera they might have been predominant during the early stages of the infection. On the contrary, however, positive DCFs were demonstrated one to two weeks following infection (TABLE 2). The earliest appearance of the DCF antibodies paralleled the appearance of ICF antibodies.

Activity of antigens. Numerous attempts were made to yield CF reactions with heated turkey sera and antigens made from either yolk sac or allantoic fluid virus suspensions. For example, a heated phenolized psittacosis (strain 6 BC) yolk sac antigen, a viable concentrated meningopneumonitis virus suspension, and the detergent extract were each titrated against 4 units each of rabbit and turkey antipsittacosis sera. The soluble antigen had the same titer when tested against rabbit and turkey sera (TABLE 3). The viral suspensions were considerably less active when titrated with turkey antiserum than with rabbit antiserum. It was obvious that viral suspensions were unsuitable as antigens with heat inactivated sera. Direct fixation with turkey sera would not usually be obtained if the virus suspension antigens were standardized with nonavian or pigeon sera, and if then the usual 4 units of antigen were used in the direct test with turkey sera. When tested with rabbit antiserum, 4 units of the allantoic fluid and yolk sac antigens shown in TABLE 3 represent

by heat inactivation of the sera. This early observation was reinvestigated by our laboratory and by Brumfield and Pomeroy⁹ and it revealed that the CF ability of turkey and chicken ornithosis antibody was destroyed by heat inactivation of the sera if virus particle antigens were employed.

A variety of serologic tests have been made available; however only the ICF and the direct complement fixation test (DCF) will be considered in this report. In addition the intradermal allergic test was developed as an ancillary epidemiological tool^{10, 11}. At present there is an urgent need to evaluate and to compare the various methods that have been proposed so that a standard technique may be developed. With this in view this report presents the development and evaluation of the DCF and allergic tests.

Direct Complement Fixation Using Heat Inactivated Serum

Technique. We have described the extraction of a soluble psittacosis CF antigen with a detergent¹²; this antigen fixed C consistently with turkey anti ornithosis sera⁷. Fixation occurred with sera that had been inactivated by heating either at 56° C for 30 min. or at 60° C for 20 min. For this test 4 units of the detergent extracted antigen were used. In studies that compared antigen activities complement was titrated in the presence of antigens. Preliminary incubation was carried out at 37° C for 75 min. and after the addition of sensitized sheep erythrocytes a secondary incubation at 37° C for 30 min. was done. The method used in performing the ICF test was similar to that described by Karrer, Meyer and Eddie. Titers represent the reciprocal of the highest dilution of serum showing either a 1+ or 2+ reaction in the ICF (partial inhibition) or a 4+ reaction (complete fixation) in the DCF.

Comparison of the DCF and ICF methods. A comparison of these techniques was made with naturally infected and experimentally infected turkeys. Forty seven turkeys were selected at random from a flock that had undergone a severe epizootic several months prior to bleeding. TABLE 1 shows that 41 sera in this group (A) were positive by both the ICF and DCF and that 3 sera were positive in low dilutions in the DCF and negative in the ICF. Also shown in TABLE 1 are results with sera from 2 flocks (B and C) in which there was no evidence of infection at the time of bleeding. One serum from flock B was negative by the ICF and had a DCI titer of 1/4. Sera obtained from 12 turkeys that had undergone a natural infection 18 months prior to the test were all positive to the DCI and only 3 sera had positive ICF titers. A direct correlation analysis on the basis of the log distribution of the ICF and DCF titers of group A was found to be 0.72 which was significant at the 1 per cent level. Although there is no direct evidence as yet that the same antibody was being measured by the 2 tests the DCF titers correlated well with the ICF titers of these sera.

In search of a reliable, simple and economical serologic technique for survey purposes Neal and Davis¹³ compared the ICF, DCF (detergent antigen) and a macroscopic agglutination test¹⁴ that employed stained elementary bodies as antigen. The ICF and DCF agreed in 85.1 per cent of 923 turkey sera tested. These results were similar to the findings shown in TABLE 1 in which the 2

TABLE 1
COMPARISON OF THE DIRECT AND INDIRECT COMPLEMENT FIXATION TESTS
FOR THE DETECTION OF TURKEY ORNITHOSIS

Case	History	Number of sera tested	DCF pos ICF pos	DCF pos ICF neg	DCF neg ICF pos	DCF neg ICF neg	Percent fixation		Percent agglutination
							DCF	ICF	
A	Natural infection for 3 months	47	41	3	0	3	93.6	87.3	93.6
B	No evidence of in- fection	61	1	1	0	59	3.2	1.6	98.3
C	No evidence of in- fection	50	2	0	0	48	4.0	4.0	100.0
D	Natural infection 18 months be- fore	12	1	5	0	0	100.0	58.3	58.3
E	Experimentally in- fected 2 months	21	15	1	3	2	76.1	85.7	80.9
F	Experimentally in- fected 3 months	12	8	0	0	4	66.6	66.6	100.0
Total		203	74	10	3	116	41.3	37.9	88.5

Direct complement fixation

† Indirect complement fixation

tests agree in 84.5 per cent of 203 turkey sera tested. On the other hand the correlation between the results of the agglutination and ICF tests was 42.3 per cent and between the agglutination and DCF tests the correlation was 40.4 per cent. Many nonspecific reactions were given by the agglutination test by lower serum dilutions, an observation that we had also made relative to an ornithosis precipitin reaction with turkey and chicken sera.¹³ On the basis of these comparisons further study with this DCF test was considered justified.¹²

Onset of DCF antibody. It was felt that if incomplete CF antibodies were formed in turkey sera they might have been predominant during the early stages of the infection. On the contrary however positive DCFs were demonstrated one to two weeks following infection (TABLE 2). The earliest appearance of the DCF antibodies paralleled the appearance of ICF antibodies.

Activity of antigens. Numerous attempts were made to yield CF reactions with heated turkey sera and antigens made from either yolk sac or allantoic fluid virus suspensions. For example a heated phenolized psittacosis (strain 6 BC) yolk sac antigen, a viable concentrated meningo-pneumonia virus suspension and the detergent extract were each titrated against 4 units each of rabbit and turkey anti-psittacosis sera. The soluble antigen had the same titer when tested against rabbit and turkey sera (TABLE 3). The viral suspensions were considerably less active when titrated with turkey antiserum than with rabbit antiserum. It was obvious that viral suspensions were unsuitable as antigens with heat inactivated sera. Direct fixation with turkey sera could not usually be obtained if the virus suspension antigens were standardized with nonavian or pigeon sera and if then the usual 4 units of antigen were used in the direct test with turkey sera. When tested with rabbit antiserum 4 units of the allantoic fluid and yolk sac antigens shown in TABLE 3 represent

TABLE 2
THE APPEARANCE OF COMPLEMENT FIXING ANTIBODIES
IN EXPERIMENTALLY INFECTED TURKEYS

Turk y	D ys aft fe t u n				
	0	8	10	14	108
1	0	2 (3+)		64	
2	0	16		32	
3	0	2 (3+)		8	
4	0	2		64	
5	0	4		8	
1977	0		64		128
1852	0		128		<4
1915	0		128		32
1849	0		256		<4
1984	0		256		<4
1943	0		128		4
1967	0		512		64

Negative at 1:2 dilution

titers of 32 and 64 respectively these concentrations were insufficient to fix C with turkey sera

The detergent extract apparently consisted of an antigen that differed either quantitatively or qualitatively from that found in the particle suspensions. To test this hypothesis the antigen was purified by acid precipitation and fractionated with phenol¹. Phenol fractionation produces soluble and insoluble CF antigens that differ chemically. TABLE 3 summarizes the results of titrating these fractions against 4 units each of guinea pig, human, and turkey

TABLE 3
COMPLEMENT FIXING ACTIVITY OF ANTIGENS WHEN TESTED WITH
MAMMALIAN AND TURKEY ANTIPSITTACOSIS SERA

A t g	P s i t t a t r u m			
	G e a i g	H m a	R b b t	T k y
Living meningo-pneumonitis allantoic fluid suspension			128†	16
Killed psittacosis yolk sac suspension			256	8
Crude detergent extract			64	64
Phenol insoluble fraction of detergent extract				
1	16			16
2	16			16
3		32		32
Phenol soluble fraction of detergent extract				
1	8			0
2	32			0
3		16		0

Four units of each antiserum
† CF titer of antigens

antisera. The phenol soluble fraction reacted similarly to the particle antigens and failed to fix C with heated turkey antibody, whereas the phenol insoluble antigen had the same titer when tested against turkey, human and guinea pig sera. Possibly this latter fraction represents a subsurface antigen that was released by extraction. However, this hypothesis is not yet tenable in view of the fact that the absorption of heated turkey sera with virus particles also absorbs the antibody to the phenol insoluble fraction.

When we employed the detergent extracted antigen, our attempts to obtain direct fixation of C with heat inactivated chicken antisera met with repeated failures. Nevertheless, this antigen combined with heated chicken antibody and gave an ICI reaction.

Direct Complement Fixation Using Unheated Serum

Techniques. As previously mentioned, Bushnell and Hudson⁴ observed that chicken and turkey antibody failed to fix C with heat inactivated sera. However, CF tests with unheated sera are unsuitable because of strong anticomplementary (AC) properties. Brumfield and Pomeroy⁵ proposed a method in which the C was increased to overcome the AC activity of unheated fowl sera. In this test, yolk sac antigens were employed and complement was titrated in the presence of 10 unheated sera diluted 1:8. Sera to be tested were diluted 1:8 to minimize AC effects and lytic action due to the fowl C. There was agreement between the ICI and DCF in more than 94 per cent of 125 turkey sera tested.

A second method might be one that selectively removes the AC substances from unheated sera, thereby avoiding use of excessive amounts of C and possibly permitting tests with lower serum dilutions. With this in mind, the following procedures were attempted to remove chicken and turkey sera AC activity: (1) chloroform extraction, (2) hypertonicization with NaCl¹⁶, (3) acid precipitation¹⁷, (4) adsorption with activated charcoal, (5) adsorption with a mouse liver preparation¹⁸ and (6) adsorption with kaolin. Kaolin adsorption proved to be the most effective. Complete details of the procedure will be given elsewhere and are given in brief form as follows: A 1:2 dilution of 1.2 ml. of antiserum is adsorbed with 40 to 50 mg. of kaolin for 60 min. The kaolin is removed by centrifugation at 1500 rpm for 5 min. Under these conditions AC activity was effectively removed (TABLE 4). Also, the serum was decomplemented and prozones were avoided. Antibody was rarely adsorbed from turkey sera with 40 to 50 mg. of kaolin, but some chicken sera showed decreased ICF and DCF titers following adsorption. Both DCF and ICF chicken antibodies were completely removed from 1.2 ml. of a 1:4 dilution of serum adsorbed with the above quantities of kaolin. Nikkila and Oker-Blom¹⁹ observed, however, that adsorption of human serum with clay minerals resulted in partially selective removal of lipids without loss of gamma globulin and Clarke and Casals⁹ successfully used the clay mineral bentonite to remove hemagglutination inhibitor from antisera.

Effect of storage on DCF antibody. The ability of the detergent extracted antigen to fix C with heated turkey serum and the inability of the particle antigens to exert the same degree of activity suggested that these antigens

TABLE 2
THE APPEARANCE OF COMPLEMENT FIXING ANTIBODIES
IN EXPERIMENTALLY INFECTED TURKEYS

Turkey	Days of infection				
	0	8	10	14	108
1	0*	2 (3+)		64	
2	0	16		32	
3	0	2 (3+)		8	
4	0	2		64	
5	0	4		8	
1977	0		64		128
1857	0		178		<4
1915	0		128		32
1849	0		256		<4
1984	0		256		<4
1943	0		128		4
1962	0		512		64

Negative at 1:2 dilution

titers of 32 and 64 respectively, these concentrations were insufficient to fix C with turkey sera

The detergent extract apparently consisted of an antigen that differed either quantitatively or qualitatively from that found in the particle suspensions. To test this hypothesis the antigen was purified by acid precipitation and fractionated with phenol.¹ Phenol fractionation produces soluble and insoluble CI antigens that differ chemically. TABLE 3 summarizes the results of titrating these fractions against 4 units each of guinea pig, human, and turkey

TABLE 3
COMPLEMENT FIXING ACTIVITY OF ANTIGENS WHEN TESTED WITH
MAMMALIAN AND TURKEY ANTIPSITTACOSIS SERA

Antigen	Potency of serum			
	Guinea pig	Human	Rabbit	Turkey
Living meningopneumonitis allantoic fluid suspension			128†	16
Killed psittacosis yolk sac suspension			256	8
Crude detergent extract			64	64
Phenol insoluble fraction of detergent extract				
1	16			16
2	16			16
3		32		32
Phenol soluble fraction of detergent extract				
1	8			0
2	32			0
3		16		0

Four units of each antiserum
† CF titer of antigens

TABLE 6
COMPLEMENT FIXATION TITERS OF UNHEATED KAOLIN ADSORBED
TURKEY ANTISERA STORED AT 4 C

T key	Days stored									
	0		4		8		16		32	
	D	YS†	D	YS	D	YS	D	YS	D	YS
12	16	0 ₄								
149	64	16								
42	8	0								
14	2	0								
109	8	2			8	0				
101	32	8			32	2				
29	32	2			32	4				
127	16	8			16	0				
16	32	8			32	16				
49	8	0			8	0				
148	16	8	16	2	16	0				
167	16	4					16	0		
13	16	2					16	0		
106	32	4					32	0		
155	(-)	4					<64	0		
29	64*	8					32	0		
46	16	4							16	0
8	16	4							16	4
28	8	2							4	0
9	4	(-)							2	0
27	16	2							16	0

Symbol (-) not tested

Detergent antigen

† Yolk sac antigen

‡ Less than 1:2

presents the results of titrating these unheated kaolin adsorbed sera. The titers were lower with the yolk sac antigen than with the soluble antigen and they decreased following refrigeration for 13 days. Studies with turkey sera are being continued with only the soluble antigen since CF antibodies of unheated turkey sera are potentially labile to the particle antigens. Unheated chicken antibody however loses the ability to fix C following storage at 4 C to both the soluble and particle antigens.

Detection of chicken DCF antibody. An interesting study performed by N. T. Briggs in our laboratory showed good correlation between the kaolin treated serum method and the ICF test. To simulate a natural infection by the oral route the following experiment was performed. Fifteen white leghorn chickens were infected with psittacosis virus (Texas turkey). Beginning 48 hours after infection feces were collected daily from these donors for 25 days. Collections were pooled, added to 200 cc of distilled water and homogenized in a Waring Blendor and a 1:4 dilution was made. Forced feeding of this suspension to recipient chickens was accomplished by introducing a blunted pipette directly into the esophagus thereby passing the tracheal opening. All daily feedings were made using stock inocula prepared from feces passed

TABLE 4
EFFECT OF CONCENTRATION OF KAOLIN AND TIME OF ADSORPTION ON
REMOVAL OF ANTICOMPLEMENTARY ACTIVITY FROM CHICKEN SERUM

Mg kaolin	Time of adsorption minutes			
	15	30	45	60
10	4	4	4	4
20	4	4	4	4
30	(-)	3	4	4
40	4	4	3	±
50	3	±	0	0

Anticomplementary activity of 1:2 dilution of serum

might also have different activities with unheated sera. Turkey sera that had been stored at -20°C for 11 months and had been repeatedly frozen and thawed during this time were titrated (unheated and adsorbed with kaolin) with 4 units each of the detergent antigen and a yolk sac antigen. The serum titers obtained with the yolk sac antigen were significantly lower, and the titers obtained with the soluble antigen were the same as titers obtained when tested 11 months before by the heat inactivation method. Upon further storage of these sera at 4°C the titers continued to drop when tested with the yolk sac antigen.

Fresh turkey antisera had the same titers to the 2 antigens (TABLE 5). Following storage at 4°C for 4 days these sera showed decreased titers when tested with the yolk sac antigen (TABLE 5). According to these results and to those mentioned by Brumfield and Pomeroy,⁹ caution must be exercised when using the viral suspension antigens for the testing of unheated fowl sera that had not been frozen soon after procurement. This fact was further borne out by testing turkey sera that had been mailed to us unrefrigerated. TABLE 6

TABLE 5
COMPLEMENT FIXATION TITERS OF UNHEATED KAOLIN ADSORBED
TURKEY ANTISERA STORED AT 4°C

Turkey	Days			
	0		4	
	D	YS†	D	YS
1849	256‡	256	256	256
1852	128	128	128	<32
1984	256	256	256	32
1962	256	256	512	<32
1977	64	64	32	16
1406	0	0	0	0

Detergent antigen

† Yolk sac suspension

‡ CF titer of sera

TABLE 6
COMPLEMENT FIXATION TITERS OF UNHEATED KAOLIN ADSORBED
TURKEY ANTISERA STORED AT 4 C.

Turkey	D y t e d									
	0		4		5		10		13	
	D	YS†	D	YS	D	YS	D	YS	D	YS
12	16	0 ₄								
149	64	16								
42	8	0								
14	2	0								
109	8	2			8	0				
101	32	8			32	2				
29	32	2			32	4				
127	16	8			16	0				
16	32	8			32	16				
49	8	0			8	0				
148	16	8	16	2	16	0				
167	16	4					16	0		
13	16	2					16	0		
106	32	4					32	0		
155	(-)	4					<64	0		
29	64*	8					32	0		
46	16	4							16	0
8	16	4							16	4
28	8	2							4	0
9	4	(-)							2	0
27	16	2							16	0

Symbol (-) not tested

Detergent antigen

† Yolk sac antigen

‡ Less than 1/2

presents the results of titrating these unheated kaolin adsorbed sera. The titers were lower with the yolk sac antigen than with the soluble antigen and they decreased following refrigeration for 13 days. Studies with turkey sera are being continued with only the soluble antigen since CF antibodies of unheated turkey sera are potentially labile to the particle antigens. Unheated chicken antibody however loses the ability to fix C following storage at 4 C. to both the soluble and particle antigens.

Detection of chicken DCF antibody. An interesting study performed by N. T. Briggs in our laboratory showed good correlation between the kaolin treated serum method and the ICF test. To simulate a natural infection by the oral route the following experiment was performed. Fifteen white leghorn chickens were infected with psittacosis virus (Texas turkey). Beginning 48 hours after infection feces were collected daily from these donors for 25 days. Collections were pooled, added to 200 cc. of distilled water and homogenized in a Waring Blendor and a 1/4 dilution was made. Forced feeding of this suspension to recipient chickens was accomplished by introducing a blunted pipette directly into the esophagus thereby passing the tracheal opening. All daily feedings were made using stock inocula prepared from feces passed

TABLE 4
EFFECT OF CONCENTRATION OF KAOLIN AND TIME OF ADSORPTION ON
REMOVAL OF ANTICOMPLEMENTARY ACTIVITY FROM CHICKEN SERUM

Mg kaolin	Time of ad r p t on m t s			
	15	30	4	60
10	4*	4	4	4
20	4	4	4	4
30	(-)	3	4	4
40	4	4	3	±
50	3	±	0	0

Anticomplementary activity of 1:2 dilution of serum

might also have different activities with unheated sera. Turkey sera that had been stored at -20°C for 11 months and had been repeatedly frozen and thawed during this time were titrated (unheated and adsorbed with kaolin) with 4 units each of the detergent antigen and a yolk sac antigen. The serum titers obtained with the yolk sac antigen were significantly lower and the titers obtained with the soluble antigen were the same as titers obtained when tested 11 months before by the heat inactivation method. Upon further storage of these sera at 4°C the titers continued to drop when tested with the yolk sac antigen.

Irish turkey antisera had the same titers to the 2 antigens (TABLE 5). Following storage at 4°C for 4 days these sera showed decreased titers when tested with the yolk sac antigen (TABLE 5). According to these results and to those mentioned by Brumfield and Pomeroy,⁹ caution must be exercised when using the viral suspension antigens for the testing of unheated fowl sera that had not been frozen soon after procurement. This fact was further borne out by testing turkey sera that had been mailed to us unrefrigerated. TABLE 6

TABLE 5
COMPLEMENT FIXATION TITERS OF UNHEATED KAOLIN ADSORBED
TURKEY ANTISERA STORED AT 4°C

Turk y	D y stor d			
	0		4	
	D	Yst	D	YS
1849	256†	256	256	256
1857	128	128	128	<32
1984	256	256	256	32
1962	256	256	512	<32
1971	64	64	32	16
1406	0	0	0	0

Detergent antigen

† Yolk sac suspension

‡ CF titer of sera

TABLE 6
COMPLEMENT FIXATION TITERS OF UNHEATED KAOLIN ADSORBED
TURKEY ANTISERA STORED AT 4 C

Titer	D y t e d									
	0		4		8		16		32	
	D	YS†	D	YS	D	YS	D	YS	D	YS
12	16	0								
149	64	16								
42	8	0								
14	2	0								
109	8	2			8	0				
101	32	8			32	2				
29	32	2			32	4				
127	16	8			16	0				
16	32	8			32	16				
49	8	0			8	0				
148	16	8	16	2	16	0				
167	16	1					16	0		
13	16	2					16	0		
106	32	4					32	0		
155	(-)	4					<64	0		
29	64*	8					32	0		
46	16	4							16	0
8	16	4							16	4
28	8	2							4	0
9	4	(-)							2	0
27	16	2							16	0

Syrrhol (-) not tested

Detergent antigen

† Yolk sac antigen

* Less than 1:2

presents the results of titrating these unheated kaolin adsorbed sera. The titers were lower with the yolk sac antigen than with the soluble antigen and they decreased following refrigeration for 13 days. Studies with turkey sera are being continued with only the soluble antigen since CF antibodies of unheated turkey sera are potentially labile to the particle antigens. Unheated chicken antibody however loses the ability to fix C following storage at 4 C to both the soluble and particle antigens.

Detection of chicken DCF antibody. An interesting study performed by N. T. Briggs in our laboratory showed good correlation between the kaolin treated serum method and the ICF test. To simulate a natural infection by the oral route the following experiment was performed. Fifteen white leghorn chickens were infected with psittacosis virus (Texas turkey). Beginning 48 hours after infection feces were collected daily from these donors for 25 days. Collections were pooled, added to 200 cc of distilled water and homogenized in a Waring Blendor and a 1:4 dilution was made. Forced feeding of this suspension to recipient chickens was accomplished by introducing a blunted pipette directly into the esophagus thereby passing the tracheal opening. All daily feedings were made using stock inocula prepared from feces passed

TABLE 7
COMPARISON OF INDIRECT AND DIRECT COMPLEMENT FIXING ANTIBODY
RESPONSES IN ORALLY INFECTED CHICKENS

No. days d. nos. infected	Chick	Titer by indirect complement fixation test recipients								No. pos.
		14		21		28		40		
		DCF	ICF†	DCF	ICF	DCF	ICF	DCF	ICF	
2-6	1	16	16	4	8	4	16	4	8	3/4
	2	4	4	0	8	8	8	4	16	
	3	2	0	8	8	16	8	16	8	
	4	0	0	0	0	0	0	0	0	
	5 Cont	0	0	0	0	0	0	0	0	
7-11	6	0	0	16	32	0	2	sacrificed		3/5
	7	16	16	32	64	16	8	sacrificed		
	8	0	0	0	0	0	0	0	0	
	9	0	0	0	0	8	8	16	4	
	10	0	0	0	0	0	0	0	0	
	11 Cont	0	0	0	0	0	0	0	0	
12-16	12	0	0	0	(-)	0	0	0	0	1/5
	13	0	0	0	(-)	0	0	0	0	
	14	32	64	32	(-)	8	32	sacrificed†		
	15	0	0	0	(-)	0	0	0	0	
	16	0	0	0	(-)	0	0	0	0	
	17 Cont	0	0	0	(-)	0	0	0	0	
17-21	18	0	(-)	0	0	0	(-)	0	(-)	0/5
	19	0	(-)	0	0	0	(-)	0	(-)	
	20	0	(-)	0	0	0	(-)	0	(-)	
	21	0	(-)	0	0	0	(-)	0	(-)	
	22	0	(-)	0	0	0	(-)	0	(-)	
	23 Cont	0	(-)	0	0	0	(-)	0	(-)	
22-26	24	0	(-)	0	0	0	(-)	0	(-)	0/3
	25	0	(-)	0	0	0	(-)	0	(-)	
	26	0	(-)	0	0	0	(-)	0	(-)	
	27 Cont	0	(-)	0	0	0	(-)	0	(-)	

Symbol (-) not tested

Direct complement fixation test with unheated sera that had been adsorbed with kaolin

† Indirect complement fixation

‡ Ornitho is isolated from spleen

during the previous 24 hour period. Inocula prepared daily from donor feces passed on days 2 through 6 were fed (1 ml. per bird per day) to a group of recipients. Four other groups similarly treated received respectively feces passed by donors 7 to 11, 12 to 16, 17 to 21, and 22 to 26 days following infection. All birds were therefore given 5 daily feedings. Birds within a given group were housed in the same cage along with an uninfected control bird. The birds were bled on the days following the first feeding as shown in TABLE 7. Sera were frozen within 24 hours after bleedings. All birds were serologically negative on the first day of feeding. The DCF and ICF tests were equally sensitive for detection of serologic reactors and the serologic response was interpreted as a sign of infection. Apparently the feces were most infective

2 to 11 days following infection of the donor birds 6 of 9 recipient birds produced antibody Chicken No 14 was sacrificed 29 days following first feeding and the spleen yielded ornithosis virus on mouse passage Three blind passages of the spleens and livers of chickens No 6 and No 7 failed to reveal virus Working with the bovine encephalomyelitis virus Matumoto *et al*²¹ indicated that the serologic response might be as efficient as direct isolation procedures for the detection of virus in infective materials

The Intradermal Test

Although the tuberculin test for detection of fowl tuberculosis²² has had limited usage the presence of ornithosis hypersensitivity in chickens that had been experimentally infected suggested that the skin test might serve as an added epidemiological method for rapid detection of endemic foci^{10 11 23} Sensitivity was detected as early as four weeks and as late as one year following infection¹¹ Circulating antibody appeared earlier than cutaneous hypersensitivity during the course of the infection however the serologic and lin



FIGURE 1. Positive development of response in an experimentally infected turkey following 24-hour reaction.

TABLE 7
COMPARISON OF INDIRECT AND DIRECT COMPLEMENT FIXING ANTIBODY
RESPONSES IN ORALLY INFECTED CHICKENS

No. days died	Chick	Titers of feces to recipients								No. pos.
		14		21		28		40		
		DCF	ICF†	DCF	ICF	DCF	ICF	DCF	ICF	
2-6	1	16	16	4	8	4	16	4	8	3/4
	2	4	4	0	8	8	8	4	16	
	3	2	0	8	8	16	8	16	8	
	4	0	0	0	0	0	0	0	0	
	5 Cont	0	0	0	0	0	0	0	0	
7-11	6	0	0	16	32	0	2	sacrificed		3/3
	7	16	16	32	64	16	8	sacrificed		
	8	0	0	0	0	0	0	0	0	
	9	0	0	0	0	8	8	16	4	
	10	0	0	0	0	0	0	0	0	
	11 Cont	0	0	0	0	0	0	0	0	
12-16	12	0	0	0	(-)	0	0	0	0	1/5
	13	0	0	0	(-)	0	0	0	0	
	14	32	64	32	(-)	8	32	sacrificed†		
	15	0	0	0	(-)	0	0	0	0	
	16	0	0	0	(-)	0	0	0	0	
	17 Cont	0	0	0	(-)	0	0	0	0	
17-21	18	0	(-)	0	0	0	(-)	0	(-)	0/5
	19	0	(-)	0	0	0	(-)	0	(-)	
	20	0	(-)	0	0	0	(-)	0	(-)	
	21	0	(-)	0	0	0	(-)	0	(-)	
	22	0	(-)	0	0	0	(-)	0	(-)	
	23 Cont	0	(-)	0	0	0	(-)	0	(-)	
22-26	24	0	(-)	0	0	0	(-)	0	(-)	0/3
	25	0	(-)	0	0	0	(-)	0	(-)	
	26	0	(-)	0	0	0	(-)	0	(-)	
	27 Cont	0	(-)	0	0	0	(-)	0	(-)	

Symbols: (-) not tested

Direct complement fixation test with unheated sera that had been adsorbed with kaolin

† Indirect complement fixation

* Ornithosis isolated from spleen

during the previous 24 hour period. Inocula prepared daily from donor feces passed on days 2 through 6 were fed (1 ml per bird per day) to a group of recipients. Four other groups similarly treated received respectively feces passed by donors 7 to 11, 12 to 16, 17 to 21, and 22 to 26 days following infection. All birds were therefore given 5 daily feedings. Birds within a given group were housed in the same cage along with an uninfected control bird. The birds were bled on the days following the first feeding as shown in TABLE 7. Sera were frozen within 24 hours after bleedings. All birds were serologically negative on the first day of feeding. The DCF and ICF tests were equally sensitive for detection of serologic reactors and the serologic response was interpreted as a sign of infection. Apparently the feces were most infective

TABLE 8

DETECTION OF ALLERGIC AND SEROLOGIC REACTORS AMONG HEALTHY TURKEYS AND TURKEYS NATURALLY INFECTED WITH ORNITHOSIS

Flock	History	Skin test results			Serologic results		
		N tested	N pos.	Per cent pos.	N tested	No pos.	Per cent pos.
SK ₁	1 yr-olds natural infection 3 to 4 months	237	165	71	47	40	85
SK ₂	2 yr-olds close contact with infected birds SK ₁	34	12	35	(-)	(-)	(-)
SK	2 to 5 yr olds close contact with infected birds SK ₁	53	6	11	52	4	8†
J	1 yr olds natural infection 3 to 4 months	255	196	78	190	157	83
H ₁	1 yr olds healthy	87	0	0	(-)	(-)	(-)
H ₂	1 yr olds healthy	(-)	(-)	(-)	109	5	5
R	1 yr olds healthy	242	0	0	10	19	11
A & M	8 to 10 months old healthy	110	1	0.9	98	0	0†
Colorado	14 wks old healthy	1049	0	0	10	0	0†
Total		2057	380	19	837	225	27

Symbol (-) not tested

Indirect complement fixation test

† Direct complement fixation test heat inactivated sera

The results of testing these birds for hypersensitivity and for the presence of serum antibody are summarized in TABLE 8. Seventy one and 78 per cent respectively of the 1 year-old SK and J flocks were skin reactors. The percentages of serologic reactors from these flocks were essentially the same as the allergic reactor tests. In contrast to the large number of reactors among the 1 year-old birds, only 35 per cent of the 2 year old and 11 per cent of the 2 to 5 year old birds of the SK flock were skin reactors. Similarly, only 8 per cent of the oldest birds were serologically positive. In the SK flock the breeder observed that the infection was not as severe and that the mortality rate was less in the toms than in the hens²⁴. Such differences were not noted in the J flock. The number of skin reactors among the toms concurred with this observation. Of the 232 SK birds tested, 24 were toms and only 3, or 12.2 per cent, were positive. In the J flock (255 birds), 5, or 71 per cent of 7 toms were positive reactors.

Further suggesting the application of the intradermal test for epidemiological purposes, it was noted by the breeder that during the epizootic the mortality rate among older birds on ranch SK was lower than among the younger birds. The small number of allergic and serologic reactors attest to this observation.

Flocks H and R, although in close proximity to flocks SK and J, showed no clinical ornithosis at the time of testing and no skin reactors were found. However, of 110 birds from flock R tested for ICF antibody, 19 showed titers of 1:2 or above. Among 109 birds tested from flock H, 4.6 per cent were positive.

test methods were equally sensitive for detection of reactors among birds infected for four weeks or longer. To further evaluate the intradermal test we tested naturally infected turkeys under field conditions.²¹

Technique In the early experiments intradermal inoculation of turkeys with 0.1 ml of antigen was made in the dewlap and on the lateral surface of the thigh. The dewlap was finally selected as the inoculation site since reactions at this site were easier to judge and developed earlier than the thigh reaction. The reactions were judged by palpation of the dewlap for edema 24 hours following skin test. Positive dewlap reactions are shown in FIGURES 1 and 2. Since the antigen was more likely to yield false negative rather than false positive reactions, detection of minimal edema was considered positive.

Comparison of intradermal and serologic tests Turkeys were obtained from the following sources: (1) normal broad breasted bronze (BBB) and Beltsville (BV) turkeys and BV turkeys that had been experimentally infected at the Agricultural and Mechanical College of Texas, College Station, Texas; (2) 1 to 5 year old BBB turkeys that had experienced an ornithosis epizootic in Oregon approximately 3 to 4 months prior to testing (flocks SK and J); (3) 2 flocks of 1 year old BBB turkeys (flocks H and R) from the same vicinity as SK and J flocks but without evidence of clinical ornithosis infection when tested; and (4) 2 flocks of healthy 14 week old BBB turkeys located in east Colorado.



FIGURE 2. Positive dewlap allergic response in a naturally infected turkey. Twenty-four hour reaction.

TABLE 10
COMPARISON OF THE SKIN TEST AND SEROLOGIC METHODS FOR THE DETECTION OF ORNITHOSIS
IN NATURALLY AND EXPERIMENTALLY INFECTED TURKEYS

T key	N tested	ST pos. sero f pos	ST neg. sero pos.	ST pos. sero neg.	ST neg. sero neg.	P e a t s		P e r c e n t both tests agreed
						ST	Sero	
Natural infection for 3 to 4 months SK flock	48	39	5	2	2	85	92‡	85
Natural infection for 3 to 4 months J flock	45	27	3	8	7	73	67‡	76

Skin test

† Serologic

‡ Complement fixation test heat inactivated sera

§ In direct complement fixation test

by the ICF test. It should be pointed out that the birds skin tested from flock H were considered prize birds and had been separated from the flock because of their robust appearance whereas the serologically positive birds were from the culled flock. It was known that the sensitivity of the intradermal test was limited during early infection. Studies have shown that allergic sensitization developed later than circulating antibody.¹¹ Subsequent to this study an ornithosislike agent was isolated from flock H and the number of serum reactors from both H and R flocks had increased.⁴ It would appear therefore that the infection in these flocks was initiated shortly prior to the time of skin testing and that serum antibody appeared prior to the development of allergy. In such a situation the skin test might yield supporting data on the initiation of ornithosis activity in this flock. It should be emphasized that the intradermal test is not suitable for diagnostic purposes but is suitable for mass surveys to determine the extent of past infections.

The ICF DCF (heat inactivated sera) and skin test methods were compared in 48 randomly selected birds from flock SK. the results given in TABLE 9 show that the allergic test compared favorably with the two serologic techniques. The results further summarized in TABLE 10 show that the serologic and skin tests agreed in 85 and 76 per cent of the naturally infected birds.

TABLE 11
ABILITY OF DIFFERENT ANTIGENS TO FIX COMPLEMENT WITH
TURKEY AND CHICKEN ANTI-ORNITHOSIS SERA

A n t i g e n	T u r k e y s e r u m		C h i c k e n s e r u m	
	U n h e a t e d	H e a t e d	U n h e a t e d	H e a t e d
Virus suspension	+	-	+	-
Detergent extract	+	+	+	-
Phenol-soluble fraction of extract	+	-	+	-
Phenol-insoluble fraction of extract	+	+	+	+

TABLE 9
COMPARISON OF THE INTRADERMAL AND SEROLOGIC REACTIONS IN
TURKEYS NATURALLY INFECTED WITH ORNITHOSIS

Tu l y	T t of sera		Sk test
	D ect complemen t fixat o	Ind ct complemen t fixat on	
3	16	32	neg
12	4	0	neg
20	0	0	neg
23	8	4	neg
35	256	128	neg
49	0	0	neg
50	128	128	neg
1	8	256	+
2	32	128	+
4	128	256	+
5	(-)	0	+
66	256	256	+
7	16	(-)	+
8	32	64	+
9	256	32	+
10	256	64	+
11	32	8	+
13	128	128	+
14	128	256	+
15	64	8	+
16	256	128	+
17	256	128	+
18	32	16	+
19	128	32	+
21	64	128	+
22	64	128	+
24	128	32	+
25	8	16	+
26	256	256	+
27	8	8	+
28	32	32	+
29	64	64	+
30	16	4	+
31	8	8	+
32	64	8	+
33	512	256	+
34	16	4	+
36	4	0	+
37	128	32	+
38	128	64	+
39	128	16	+
40	16	16	+
41	16	0	+
42	0	0	+
43	64	256	+
45	16	4	+
47	16	8	+
48	8	256	+
Per cent positive	92	85	85

Symbol (-) not tested
Heat inactivated sera

- 3 RICE C F 1947 Can J Comp Med Vet Sci 11 236
- 4 IIDDIF B & T FRANCIS 1942 Proc Soc Exptl Biol Med 50 291
- 5 RICE C F 1948 J Immunol 69 3f
- 6 KARRER H K F MEYER & B IIDDIF 1950 J Infectious Diseases 87 24
- 7 BENEDICT A A & C MCFARLAND 1956 Proc Soc Exptl Biol Med 92 768
- 8 BUSINFELL I D & C B HILDON 1927 J Infectious Diseases 41 388
- 9 BRUMFIELD H I & B S POMEROY 1954 Proc Soc Exptl Biol Med 94 146
- 10 BENEDICT A A I O'BRIEN & C MCFARLAND 1956 Am J Vet Research 17 543
- 11 BENEDICT A A & C MCFARLAND 1956 J Immunol 77 165
- 12 BENEDICT A A & E O'BRIEN 1956 J Immunol 76 293
- 13 NEAL J F & D I DAVIS 1957 Am J Vet Research In press
- 14 LABROFFSKY N A 1946 J Infectious Diseases 79 96
- 15 BENEDICT A A & C MCFARLAND Unpublished data
- 16 NORDBERG B K & K SCHJERNING-THIESEN 1956 J Infectious Diseases 98 266
- 17 SACHS H 1921 Berlin klin Wochschr 58 1075
- 18 KAPP F G M GNESU & I GORDON 1955 Proc Soc Exptl Biol Med 90 335
- 19 NIKKILA F & N OKER-BLOM 1952 Science 116 685
- 20 CLARKE D & J CASALS 1955 Proc Soc Exptl Biol Med 88 96
- 21 MATSUMOTO M T OMORI T MORIMOTO K HARADA Y INABA & S ISHII 1955 Japan J Exptl Med 25 223
- 22 VAN ES I & A E SCHALK 1914 N Dakota Agr Expt Sta Bul No 108 3
- 23 BENEDICT A A 1957 Am J Hyg 65 245
- 24 ORNITHOSIS IN OREGON 1956 Report Oregon State Board of Health U S Public Health Service and The George W Hooper Foundation Laboratory Unpublished

This was in good agreement with the 80 per cent correlation previously obtained with experimentally infected chickens¹¹

Conclusions

The results of the experiments with turkey and chicken antibody indicate that their abilities to fix C' with homologous antigen differ. These two species probably cannot be grouped as producing qualitatively similar CF antibodies. The reactive group (or groups) on the antibody molecules associated with fixation of C' are apparently heat labile but the degree of lability is different for turkey and chicken antibody as evidenced by the fixation of C with heated turkey serum and the soluble antigen. Turkey and chicken antibody are dissimilar in at least this one respect. A summary of these reactions is given in TABLE 11. According to our present knowledge of the ornithosis system the animal species can be listed in decreasing order of their ability to fix C as follows: mammals (rabbit, guinea pig, man), psittacines, pigeon, turkey, and chicken. Whether duck antibody is similar to turkey or chicken antibody remains to be determined. Work is in progress to determine whether ICF antibody represents the DCF antibody and has lost CF ability because of inactivation of reactive sites.

In regard to the intradermal test, the results of skin testing naturally infected ornithotic turkeys confirm previous observations made on cutaneous hypersensitivity in experimentally infected chickens¹¹. The good correlation between the serologic and allergic reactor rates suggests the usefulness of the intradermal test for detection of ornithosis exposure. From a practical viewpoint, skin testing large numbers of turkeys in the field was found to be a less cumbersome procedure than bleeding the birds and then processing the blood samples. According to our experience in testing more than 2000 turkeys in the field, 5 or 6 birds were skin tested for every bird bled. Of all the tests proposed, it is felt that the intradermal test is the most practical for extensive ornithosis survey studies.

Acknowledgments

We express our appreciation to John Delaplane and Donald F. Davis and their colleagues at the Agricultural and Mechanical College of Texas, College Station, Texas, for their generous cooperation. We are indebted to Samuel Osgood of the Public Health Service and the Oregon State Board of Health, Portland, Ore.; Donald Mason of the Public Health Service and The George Williams Hooper Foundation, University of California Medical Center, San Francisco, Calif., for their kind assistance in studies with the 1956 Oregon epizootic; and to John Clayton of the Public Health Service, Denver, Colo., and Martin Baum, Colorado State Department of Health, Denver, for their aid in the Colorado survey. Grateful acknowledgment is made to Bernice Eddie of The George Williams Hooper Foundation for some of the ICF data given in this report.

References

1. VOLKERT, M. & M. MATTHIJSSEN. 1956. *Acta Pathol. Microbiol. Scand.* **39**: 117.
2. KARRER, H., K. F. MEYER & B. LIDDIE. 1950. *J. Infectious Diseases* **87**: 13.




FIGURE 1 Annual total of reported murine typhus fever cases in the United States between 1941 and 1956

The period 1941 to 1944 might be considered the culmination of the classic era of murine typhus in the United States which began in 1913 with Paullin's description. This period might be characterized as follows:

- (1) There was a steady increase in the reported cases of murine typhus from 2784 in 1941 to 5401 in 1944 as shown in FIGURE 1
- (2) There was an increasing number of deaths ascribed to murine typhus as shown in TABLE 1 108 in 1941 and 193 in 1944
- (3) There were sizable populations of oriental rat fleas in many parts of the southeastern United States. Davis (1945) reported that the index of oriental rat fleas varied from as low as 3 to as high as 25 per rat in San Antonio Texas in 1944. Data from Public Health Service surveys of seaport towns made before 1940 and after 1944 before the initiation of DDT dusting suggest that in many parts of the southeastern United States the oriental rat flea index was rather high from 1941 to 1944. For example Cole and Koepke (1947) have published data showing an oriental rat flea index in Savannah Ga. varying between 1 and 12 fleas per rat during the years 1932 and 1933 while Ludwig and Nicholson (1947) showed that in May and June of 1945 the index varied from 5 to as high as 57 fleas per rat depending on the type of habitat and date of trapping. It is reasonable to assume that in the intervening years 1941 to 1944 the oriental rat flea population in Savannah was at comparable levels.
- (4) There was a rather high index of murine typhus infections in domestic rats. As one example Davis (1948) has written that the blood of 35 per cent of the roof rats and 51 per cent of the Norway rats in San Antonio Texas gave

Part VI Other Contemporary Disease Problems

THE CHANGING PICTURE OF MURINE TYPHUS IN THE UNITED STATES

By Harry D. Pratt

Communicable Disease Center Public Health Service Atlanta Ga

The picture of murine typhus first appeared in the United States in 1913 when J. E. Paullin reported six cases in Atlanta, Ga. The epidemiological studies of Maxcy (1926 and 1929) in the southeastern part of the country showed conclusively that endemic typhus was a milder disease than epidemic typhus—that it occurred in individuals who were not infested with body lice—that cases occurred more commonly in summer and fall than in winter and early spring—as in the case of the classic louse-borne typhus, and that there might be a rodent reservoir of the disease. The investigations of Dyer, Rumberg, and Badger (1931) showed that the oriental rat flea (*Xenopsylla cheopis* Rothschild) was the vector of murine typhus. In this same year Mosser, Castaneda, and Zinsser (1931) established the fact that the domestic rat was the reservoir of this disease. In the 1930s the Weil-Felix agglutination test was used increasingly in the laboratory to confirm the clinical diagnosis of murine typhus. The number of cases of murine typhus reported annually increased from 511 in 1930 in four wavelike surges with peaks of 2070 in 1933, 2393 in 1937, 2996 in 1939, and 5401 in 1944 when the greatest number of cases was reported. Bengtson (1941) and Bengtson and Topping (1941) reported that the complement fixation test was specific for endemic typhus fever. This was followed by a later paper of Brigham and Bengtson (1945).

In 1944 and 1945 a new approach to the control of murine typhus—the use of DDT to control the oriental rat fleas that transmit the murine typhus rickettsiae from the rodent reservoir to man—was begun in two widely separated areas. Savannah, Ga., where Ludwig and Nicholson (1947) carried out much of the research basic to the cooperative CDC State Health Department murine typhus control program begun in the last half of 1945, and San Antonio, Texas, where Davis (1945) carried out equally significant experiments on the control of rat fleas with DDT.

In 1949 chloramphenicol (Chloromycetin) came into the picture as the first of a new series of antibiotics that were highly effective in controlling the rickettsial diseases (Smadel *et al.* 1949). The decline in the number of cases and deaths due to murine typhus coincides with these last three significant discoveries: the DDT program to control oriental rat fleas, thus breaking the transmission cycle from rodents to man; the complement fixation test for laboratory confirmation of suspected cases; and the widespread use of antibiotics for the treatment of human cases.

It is the purpose of this paper to show the changing picture of murine typhus in the United States by comparing two four-year periods: 1941 to 1944 and 1953 to 1956.




FIGURE 2. Counties in 11 southern states reporting cases of endemic typhus fever between 1922 and 1947.

ported from Coffee County in 1943 but in a door-to-door survey 211 persons said they had had the disease. Blood specimens were obtained from 177 of the 211 reputed cases and 135 revealed positive results in complement fixation and/or Weil Felix tests. Therefore more than twice as much typhus occurred in Coffee County in 1943 than was reported. Davis and Pollard (1946) made a careful complement fixation survey of 4219 persons in San Antonio, Texas, between February and June of 1945. They estimated that about 700 persons had been infected each year during the ten years from 1935 to 1945, although the peak number of cases reported by the San Antonio Health Department was 91 in 1944.

C. L. Williams (1949) has summarized the thoughts of many people interested in murine typhus control during this period in his discussion of the typhus control program initiated in 1946 by the Georgia State Health Department and the Communicable Disease Center in Brooks, Thomas, Grady, and Decatur counties in southwestern Georgia. He wrote: "Our workers first made a house-to-house survey in these counties to determine the number of cases of murine typhus that had occurred during the one year period immediately preceding the survey. This study revealed that about one third the number of cases had been reported by physicians. If this figure is true for the whole typhus area, it would indicate that there actually occur three times as many cases as are reported. In 1944 this would have been about 10,000."

TABLE 1
REPORTED DEATHS DUE TO TYPHUS IN THE UNITED STATES 1940 TO 1955

Year	Mu typh s	Un pecu d†
1940	84	37
1941	108	39
1942	149	60
1943	154	58
1944	193	80
1945	173	75
1946	128	35
1947	88	31
1948	68	15
1949	2	23
1950	—	13
1951	—	16
1952	—	6
1953	—	6
1954	—	7
1955	—	5

* Source: *Vital Statistics of the United States* the years 1940 to 1944 published by the Department of Commerce Bureau of Census the years 1945 through 1949 by the Federal Security Agency Public Health Service National Office of Vital Statistics and the years 1950 through 1955 by the Department of Health Education and Welfare Public Health Service National Office of Vital Statistics Bethesda Md

† Excludes Brill's disease and other rickettsial diseases

a positive complement fixation test for murine typhus in 1944 to 1945. He wrote an analysis of presence of antibodies for typhus shows that [rats in] grain mills had a significantly higher per cent than stores or residences. This localization of typhus in such places as granaries, mills and along railroad sidings was noted by many investigators.

(5) There were small epidemics of murine typhus in cities and fairly wide spread infections in rural areas in the southeastern United States. Meleney (1941) has given a most interesting summary of the cases of endemic typhus occurring in cities such as Savannah, Macon and Atlanta, Ga. for the years 1922 to 1939 including an outbreak of 15 cases in Nashville, Tenn. in 1939. This trend continued from 1939 up through 1944 for Chatham County (Savannah) and Fulton County (Atlanta), Ga. had 139 and 69 cases respectively in 1944 (Communicable Disease Center 1947). Coffee County, Ala. may have had 500 cases in 1943 according to Hill and Ingraham (1947). Meleney's maps are similar to those shown in FIGURE 2.

(6) Typhus may have been under reported in the southeastern United States from 1941 through 1944. There was a general feeling among health authorities that considerably more murine typhus existed than was reported to health departments. During the war years it was not possible to conduct an accurate survey to determine the status of typhus because of the shortage of manpower and the emphasis on malaria control. However the survey made by Hill and Ingraham (1947) in 1943 in Coffee County, Ala. may be considered typical of the typhus situation at that time in parts of the Southeast. These investigators wrote that only 61 cases of murine typhus had been officially re-




FIGURE 4 Reported cases of murine typhus in the southeastern United States during 1954




FIGURE 5 Reported cases of murine typhus in the southeastern United States during 1955

(7) There were no effective drugs for the treatment of clinical cases of murine typhus. The authoritative *Textbook of Medicine* edited by Cecil (1940) which was widely used during the years between 1941 and 1944 stated simply:

There is no specific treatment for typhus. Treatment should be directed toward supporting the patient and eliminating as far as possible the source of exhaustion until spontaneous recovery can occur. Continuous rest in bed and nursing are most essential.

(8) Control of murine typhus was based on good general sanitation to limit rodent and oriental rat flea populations supplemented by fumigation, rat trapping, ratproofing of buildings and rat poisoning with relatively ineffective rodenticides such as red squill.

From 1953 through 1956 quite a different picture of murine typhus may be seen in the United States as indicated by the following data:

(1) There was a steady decrease in the reported cases of murine typhus—from 221 in 1953 to 98 in 1956 (FIGURES 1, 3, 4, 5, and 6). This is the first year since 1924 in which fewer than 100 cases of murine typhus have been officially reported to the Public Health Service.

(2) There have been no officially recorded deaths due to murine typhus since 1950 although a number of deaths due to unspecified typhus have been registered (TABLE 1).

(3) The population of oriental rat fleas has been highly variable throughout the southeastern United States. In a study made in Atlanta, Ga. in August 1957, a total of 482 oriental fleas was collected from 205 rats, an average of slightly more than 2 fleas per rat. The data for a series of rat, rat flea, and




FIGURE 3

Reported cases of murine typhus in the southeastern United States during 1953

TABLE 2
DATA ON RATS TRAPPED RAT SERA TESTED FOR TYPHUS AND NUMBER OF *LEVOPISILLAE*
CHIEPIS FOUND ON RATS TRAPPED IN FULTON COUNTY GA FROM 1945 TO 1957

Year	Number rats trapped	Number rats tested for typh	Number rat sera positive for typh	Number rats with <i>Levopisillae</i>	Number <i>X</i> <i>k p</i>
1945	150	35	23	84	761
1946	507	416	63	14	433
1947	1298	1243	255	257	1496
1948	657	641	80	35	61
1949	347	329	19	99	396
1950	190	190	7	59	132
1957	205	195	4	74	482

Twenty-one sera anticomplementary

cases were appraised as not typhus. In Georgia McCroan *et al* (1955) were able to confirm only 13 of 41 reported cases of typhus in 1953. They wrote

One of the confirmed typhus cases was reported as Rocky Mountain spotted fever. Five of the 29 unconfirmed cases were classified as presumptive but 24 of these cases remained diagnostic problems even after prolonged study.

(1) There are effective drugs for the treatment of murine typhus such as chloramphenicol and the tetracycline agents such as Aureomycin and Terramycin. However these new antibiotics have had an important effect on the reporting of rickettsial infections to health departments. According to McCroan *et al* (1955) The rickettsial infections are so well masked by antibiotics that many cases are aborted and diagnosis with complete laboratory confirmation is a rarity. Development of complement fixation antibodies is delayed sometimes for 6 or more months. Confirmation of diagnosis in recent years has been achieved largely as the result of epidemiological follow up of cases by public health agencies.

(8) Control of murine typhus today is based on good general sanitation rat trapping ratproofing and rat poisoning with red squill and a number of new rodenticides such as ANTU 1080 and particularly the anticoagulants such as warfarin and pival which give effective rodent control thus limiting the flea populations. Since 1944 a variety of new organic insecticides such as DDT dieldrin and heptachlor have been developed. These are cheap and effective and they have a long residual action in controlling oriental rat fleas.

Between 1926 and 1945 the majority of the cases of murine typhus were reported from the southeastern and the Gulf Coast states. In fact the distribution of cases in this area was so pronounced that the Communicable Disease Center manual (1949) included the statement "Over 90% of the cases occur south of a line drawn from Charleston South Carolina to Dallas Texas." Data on cases of murine typhus reported in this area for the 7 southeastern states of South Carolina Georgia Florida Alabama Mississippi Louisiana and Texas are presented in TABLES 3 and 4. It will be noted that in the years 1941 to 1944 from 90 to 93 per cent of the cases of typhus were reported from these states while in the later period from 1953 to 1956 82 to 86 per

rat blood surveys are summarized in TABLE 2. These figures indicate that, while typhus infection rates in Norway rats have not changed significantly since 1950, the number of oriental rat fleas particularly the average number of fleas per infested rat increased in Atlanta in 1957.

(4) There was a rather low index of murine typhus infection in rats in some parts of the southeastern United States and a considerably higher index in other parts. In laboratory tests of 195 rat sera collected in Atlanta, Ga. during August and September 1957, only 4 sera gave indications of typhus infections by complement fixation and the titers were low in all these specimens in the range of 1:4 to 1:16.

(5) There have been no murine typhus epidemics in cities or in rural areas of the magnitude seen in the 1930s or early 1940s. Quinby and Schubert (1953) made a special effort to 'detect association of proved cases in time, place, and persons thus defining areas considered as outbreaks.' This was successful only as follows:

Brundidge, Ala.	1950	2 cases
Samson, Ala.	1951	2 cases
Santa Rosa County, Fla.	1951	2 cases
Columbia, S. C.	1951	6 cases

(6) Typhus may have been over reported in the southeastern United States from 1953 through 1956. This trend seems to have become apparent beginning in 1948. Quinby and Schubert wrote: 'From 1948 through 1951 351 reported and 99 unreported cases of murine typhus were appraised in 8 southern states. After complement fixation testing 58 per cent of the reported cases and 33 of the unreported cases were confirmed. At least 25 per cent of the reported




FIGURE 6. Reported cases of murine typhus in the southeastern United States during 1956.

ments It is probable that the widespread use of many of the newer rodenticides such as warfarin pival 1080 or ANTU has had an over all influence in reducing the number of older rats with typhus infections thus reducing the reservoir of infected rodents Since 1949 the increasing use of new antibiotics such as chloramphenicol Terramycin and Aureomycin may have affected the reporting of cases by delaying the development of murine typhus antibodies in the blood and making positive laboratory confirmation of this disease difficult or hopeless

It is impossible to predict whether murine typhus may eventually be eradicated from the United States as seems possible with malaria or whether the disease will recur with increasing frequency The Atlanta survey made during the summer of 1951 did indicate that the average rat flea index was high enough (an average of 2.4 oriental rat fleas per rat) for the transmission of murine typhus rickettsiae from the rodent reservoir to man but that the level of typhus infection in rats was very low For this reason very few typhus cases should be expected in Atlanta this year It is worth noting however that the few typhus infections found in rats in Atlanta in 1957 occurred in typical typhus areas namely in low income housing areas in old business districts and around milling companies In some of these areas there were between 30 and 40 oriental rat fleas per rat a dangerously high index if the typhus infection rate should increase in rats in these areas In this connection it is interesting to note that the *Morbidity and Mortality Weekly Report* for August 23 1957 reported 16 cases of murine typhus for the period January 1 to August 1 1957 as compared with 10 for the comparable period in 1956 It is therefore possible that more cases of murine typhus will occur in 1957 than in 1956 and that murine typhus will become a more severe problem than it has been in the last several years

Summary

The changing picture of murine typhus in the United States is discussed by comparing two 4-year periods 1941 to 1944 and 1953 to 1956

In the first period the number of cases of murine typhus showed a steady increase from 2184 reported in 1941 to 5401 in 1944 in the latter period the number of cases of this disease decreased from 221 in 1953 to 98 in 1956 There has also been a decrease in the number of deaths ascribed to murine typhus The number of deaths varied from 108 in 1941 to 193 in 1944 Since 1950 not one death has been officially recorded as due to murine typhus fever although a number of deaths due to unspecified typhus have been registered

In the first period 90 to 93 per cent of all the reported cases in the United States occurred in seven states South Carolina Georgia Florida Alabama Mississippi Louisiana and Texas in the second period 82 to 86 per cent were concentrated in this area

In the first period treatment of murine typhus was largely symptomatic today there are broad spectrum antibiotics such as Chloromycetin Aureomycin and Terramycin which give quick and effective cures

These new drugs however mask all types of rickettsial infection so that diagnosis with laboratory confirmation is difficult or rare The use of these

TABLE 3

REPORTED CASES OF MURINE TYPHUS IN SEVEN SOUTHEASTERN STATES 1941 TO 1944

	1941	1942	1943	1944
South Carolina	105	192	193	178
Georgia	946	1154	1256	1182
Florida	196	313	314	484
Alabama	299	380	629	892
Mississippi	66	58	133	178
Louisiana	200	165	232	283
Texas	733	1204	1452	1740
Subtotal	2545	3466	4209	4937
U S total	2784	3722	4528	5401
Percentage in 7 south eastern states	90%	93%	93%	90%

cent of the cases were reported from this area. In 1956 51 cases or more than one half of the total for the entire United States were reported from Texas. 17 of them were confirmed by laboratory diagnosis.

Many people have speculated on the reasons for this decline in murine typhus since 1944. A number of public health officials feel that the extensive DDT dusting program carried out by state and local health departments in cooperation with the Communicable Disease Center played an important part in the initial lowering of the typhus rate from 1944 to 1950. There was a considerably larger reduction in the number of cases of typhus in urban than in rural areas. Undoubtedly the re emphasis on basic sanitation has played some part in reducing rat and rat flea populations in cities since the end of World War II, particularly programs for new municipal housing, slum clearance, ratproofing of business and food handling establishments, better garbage collections and sanitary landfills. In addition there has been an ever increasing reliance by private industry on exterminators for the control of rodents and all types of insects including fleas in business and food handling establish-

TABLE 4

REPORTED CASES OF MURINE TYPHUS IN SEVEN SOUTHEASTERN STATES 1953 TO 1956

	1953	1954	1955	1956
South Carolina	11	11	6	7
Georgia	41	32	15	9
Florida	10	5	11	6
Alabama	19	23	20	10
Mississippi	3	—	3	0
Louisiana	7	3	1	2
Texas	92	64	52	51
Subtotal	183	138	114	85
U S total	221	163	135	98
Percentage in 7 south eastern states	82%	84%	84%	86%

- DIER R I A RUMFICH & I F BADGER 1931 Typhus fever a virus of the typhus type derived from fleas collected from wild rats Public Health Repts U S 46 334-338
- HILL I L & S C INGRAHAM 1947 A study of murine typhus fever in Coffee County Alabama Public Health Repts U S 62(24) 875-881
- LUDWIG R G & H P NICHOLSON 1947 The control of rat ectoparasites with DDT Public Health Repts U S 62 77-84
- MAXCY K I 1926 An epidemiological study of endemic typhus (Brill's disease) in the southeastern United States Public Health Repts U S 41 2967-2995
- MAXCY K I 1929 Typhus fever in the United States Public Health Repts U S 44 (29) 1735-1747
- MCCROAN J F R L RAMAN W J MURPHY & L S DICK 1955 The status of Rocky Mountain spotted fever in the southeastern United States Public Health Repts U S 70(3) 319-325
- MFLÉNEY H F 1941 Recent extension of endemic typhus fever in the southern United States Am J Public Health 31(3) 219-227
- MOORE H M K CASTANEDA & H ZINSSER 1931 Rats as carriers of Mexican typhus fever J Am Med Assoc 97 231-237
- MORBIDITY AND MORTALITY WEEKLY REPORT 1957 National Office of Vital Statistics Washington D C 6(33) 1
- PAULLIN J F 1913 Typhus fever with a report of cases Southern Med J 6(1) 36-43
- QUINBY G E & J H SCHUBERT 1953 Epidemiologic and serologic appraisal of murine typhus in the United States 1948-1951 Am J Public Health 43(2) 160-164
- SMADAL J F E B JACKSON H L ILEY & R LEWTHWAITE 1949 Comparison of synthetic and fermentation chloramphenicol (chloromycetin) in rickettsial and viral infections Proc Soc Exptl Biol Med 70(2) 191-194
- WILLIAMS C L 1949 The control of murine typhus with DDT Military Surgeon 104(3) 163-167
- WORTH C B & E R RICKARD 1951 Evaluation of the efficiency of common rat ectoparasites in the transmission of murine typhus Am J Trop Med 31(3) 295-298

new drugs has therefore probably obscured the true incidence of this disease today

In the first period the prevention of murine typhus was based on controlling infected rat fleas by fumigation rat trapping rat poisoning with relatively ineffective rodenticides such as red squill ratproofing and sanitation Today in addition to these methods there are powerful insecticides such as DDT dieldrin and heptachlor which give excellent control of infected rat fleas for weeks or months Many new rodenticides have been developed such as ANTU 1080 and particularly the anticoagulants such as warfarin pival and fumarin which give effective rodent control and thus limit the flea population

Finally one can speculate that improved sanitation programs in cities throughout the United States and the increasing reliance of private industry on commercial pest control services to control rodents and all types of insects have had a long term influence on the steady and continued decrease of murine typhus in the United States

Acknowledgments

The writer acknowledges the assistance of Ida Sherman Sarah Welch and Paul White of the Epidemiology Branch Communicable Disease Center in making available important data with regard to murine typhus cases and deaths of Joseph Schubert and Pamela Peltier of the Laboratory Branch of the Center in testing about 205 rat sera for murine typhus antibodies, of A J Brooks B F Bjornson and W H Johnson of the Training Branch of the Center and of John Grennor W Moye and E Walker of the Rodent and Insect Division Fulton County Health Department Atlanta Ga for conducting the murine typhus survey in Atlanta during August and September 1957

References

- BENGSTON I A 1941 Complement fixation in endemic typhus Public Health Repts U S 56 649-653
- BENGSTON I A & N H TOPPING 1941 The specificity of the complement fixation test in endemic typhus fever using a rickettsial antigen Public Health Repts U S 56(35) 1723-1727
- BRIGHAM G D & I A BENGSTON 1945 A study of the complement fixation and Weil Felix reactions in wild rats as related to the isolation of the virus of endemic typhus Public Health Repts U S 60(2) 29-46
- CECIL R L 1940 A Textbook of Medicine xx 1-1744 Saunders Philadelphia Pa
- COLE L C & J A KOEPKE 1947 A study of rodent ectoparasites in Savannah Georgia Public Health Repts U S Suppl 207 42-59
- COMMUNICABLE DISEASE CENTER 1947 Evaluation data for murine typhus fever control activities 1946 CDC Engineering Division 1-53 U S Public Health Service Atlanta Ga
- COMMUNICABLE DISEASE CENTER 1949 Rat borne disease prevention and control 1 292 Fed Sec Agency Atlanta Ga
- DAVIS D L 1945 The control of rat fleas (*Xenopsylla cheopis*) by DDT Public Health Repts U S 60(18) 485-499
- DAVIS D F 1948 Observations on rats and typhus fever in San Antonio Texas Public Health Repts U S 63 183-190
- DAVIS D E & M POLLARD 1946 Irrelevance of typhus complement fixing antibodies in human serums in San Antonio Texas Public Health Repts U S 61 928-931

- DIER, R. F., A. RUMFICH & I. F. HABER. 1931. Typhus fever: a virus of the typhus type derived from fleas collected from wild rats. *Public Health Repts. U. S.* 46: 334-339.
- HILL, F. L. & S. C. INGRAM. 1947. A study of murine typhus fever in Coffee County, Alabama. *Public Health Repts. U. S.* 62(23): 85-88.
- LUDWIG, R. C. & H. I. NICHOLSON. 1947. The control of rat ectoparasites with DDT. *Public Health Repts. U. S.* 62: 77-84.
- MAXCY, A. F. 1926. An epidemiological study of endemic typhus (Brill's disease) in the southern United States. *Public Health Repts. U. S.* 41: 296, 299.
- MAXCY, A. F. 1929. Typhus fever in the United States. *Public Health Repts. U. S.* 44(29): 1735-1743.
- MCCROAN, J. F., R. L. RAWSON, W. J. MURPHY & L. S. DICK. 1955. The status of Rocky Mountain spotted fever in the southern United States. *Public Health Repts. U. S.* 70(3): 319-325.
- MELÉNEY, H. F. 1941. Recent extension of endemic typhus fever in the southern United States. *Am. J. Public Health* 31(3): 219-221.
- MOOSER, H. W., R. C. TANEDA & H. ZINSER. 1931. Rats as carriers of Mexican typhus fever. *J. Am. Med. Assoc.* 97: 11-23.
- MORBIDITY AND MORTALITY WEEKLY REPORT. 1957. National Office of Vital Statistics. Washington, D. C. 6(33): 1.
- JACULLIN, J. L. 1913. Typhus fever with a report of cases. *Southern Med. J.* 6(1): 36-43.
- QUINBY, G. E. & J. H. SCHUBERT. 1953. Epidemiologic and serologic appraisal of murine typhus in the United States, 1949-1951. *Am. J. Public Health* 43(2): 160-164.
- SMADAL, J. F., F. B. JACKSON, H. L. IFFY & K. LEUTHWAITE. 1949. Comparison of steam autoclave and fermentation chloramphenicol (chloromycin) in tick-borne and viral infections. *Proc. Soc. Exptl. Biol. Med.* 70(2): 191-194.
- WILLIAMS, C. L. 1949. The control of murine typhus with DDT. *Military Surgeon* 104(5): 163-167.
- WORTH, C. I. & E. R. RICKARD. 1951. Evaluation of the efficiency of common rat ectoparasites in the transmission of murine typhus. *Am. J. Trop. Med.* 31(3): 295-298.

AIR BORNE TRANSMISSION OF Q FEVER THE ROLE OF PARTURITION IN THE GENERATION OF INFECTIVE AEROSOLS*

By Hartwell H. Welsh, Edwin H. Lennette, Francis R. Abinanti† and John F. Winn‡

Viral and Rickettsial Disease Laboratory, California State Department of Public Health, Berkeley, Calif.

Prior to 1944-1945 Q fever as a disease of humans was a clinical rarity seen as a naturally occurring illness only in Australia. In the past thirteen years the disease has shown itself on all the continents and the British Isles in circumstances varying from sharply defined and explosive outbreaks to smoldering and continuous endemics. It now appears that it can be found in any part of the world where an effort is made to detect it. Whether the disease has spread from Australia to the rest of the world in the years since its original description by Derrick¹ in 1937 or whether propitious circumstances led to its recognition first in Australia is a question that cannot be answered at present. In any case Q fever has become a disease of world wide importance and the considerable effort expended in attempts to define the natural history of this disease has been highly productive especially during the last decade.

Derrick² and Stoker and Marmion³ have presented very fine reviews of the findings relative to the epidemiology of the disease in general beyond a brief synopsis of this subject the present paper will deal rather with data concerning the actual generation and persistence of infective sources of the agent.

THE EPIDEMIOLOGY OF Q FEVER

Sources of the agent in nature. *Coxiella burnetii* has been isolated from at least 22 species of ticks⁴⁻³¹, 1 insect³², 8 mammals³³⁻⁴⁹ and at least 1 bird⁵⁰. Many of the ticks found naturally infected have been shown capable under laboratory conditions of transmitting the infection by bite. In some species stage-to-stage transfer of the agent occurs and in others the agent is transmitted transovarially. The tissues and excreta of such infected ticks have been shown to be potential sources of large numbers of the organism.

Among domestic animals cattle, sheep and goats would appear to be the most important sources of the organism because of their world wide distribution and large numbers. *C. burnetii* has been isolated from the tissues of cattle and sheep and from the secretions and excretions of all three species. The placenta is an especially rich source of the agent. In the case of sheep it has been shown that the wool itself¹⁻⁶ may be heavily contaminated with the organism. This is presumably true of cattle and goat hides as well although this has not as yet been firmly established.

The work on which these observations are based was supported by Grant F 195 from the National Institute of Allergy and Infectious Diseases, Public Health Service, Bethesda, Md.

† Present address: National Institute of Allergy and Infectious Diseases, Public Health Service, Bethesda, Md.

‡ Present address: Communicable Disease Center, Public Health Service, Atlanta, Ga.

Other domestic animals and wild or domestic birds or fowl are perhaps potential sources of *C burnetii* but are less frequently involved in the disease cycle than are the three major domestic animal species.

Man has on occasion been the source of rickettsiae³³ ³⁴ in tissues from fatal cases and from sputum and urine during the acute illness. Person-to-person transmission of the agent has been of importance only in isolated instances³⁵ and does not seem to be of consequence in spread of the disease.

Modes of transmission to man or to domestic animals. Transmission of the infective agent from natural sources to man or to domestic animals may occur in a number of ways depending upon the situation. In the case of man the association between infection and exposure to domestic animal or their products is very striking; exposure may be occupational or residential in type and intense or merely casual in nature. In the case of domestic animals a similar association may exist but it is considerably more difficult to establish because in most instances the occasions on which domestic animals can be kept from close association with other livestock are far less common.

Definite epidemiological evidence exists that *C burnetii* may be carried over considerable distances on inanimate objects such as clothing³⁶ wool³⁷ hair³⁸ straw³⁹ packing materials⁴⁰ and dust⁴¹. More direct evidence exists in the case of wool.

As mentioned previously the possibility of transmission by the bite of infected ticks is a potential method utilized by the agent for transfer from one host to another. However the evidence that such a method is of any real consequence is scant. On the other hand the tissues and excreta of infected ticks have undoubtedly been responsible for some human infections brought about by handling of items thus contaminated.

In the same manner that is through manipulation or agitation of contaminated surfaces the ground and soil of areas housing infected livestock may be through the generation of dusts containing infectious particles a very important method for transmission of the agent to susceptible hosts⁴² ⁴³.

Routes of infection for man and animals. Several portals of entry into the susceptible host exist in the case of *C burnetii*. Thus laboratory animals may be infected through the unbroken (probably abraded) skin by subcutaneous intradermal or intraperitoneal inoculation through ingestion of infective materials by placement of infective materials on the conjunctiva by intranasal instillation or through inhalation of artificially generated infective aerosols. It is assumed that the same potential routes of infection exist in the case of larger animals and man as a matter of fact some of these routes have been shown epidemiologically to be of importance in producing infection under natural conditions.

With respect to man at least a general survey of the reported epidemiological findings indicates that the majority of cases of Q fever may well have resulted from contact with air borne rickettsiae taken into the body either through inhalation or through deposition onto the ocular conjunctivae or other mucous membranes. Of these possibilities the respiratory route constitutes the most probable means of ingress for the agent.

AIR BORNE TRANSMISSION OF Q FEVER THE ROLE OF PARTURITION IN THE GENERATION OF INFECTIVE AEROSOLS*

By Hartwell H. Welsh, Edwin H. Lennette, Francis R. Abinanti†
and John F. Winn‡

*Typhoid and Rickettsial Disease Laboratory, California State
Department of Public Health, Berkeley, Calif.*

Prior to 1944-1945 Q fever as a disease of humans was a clinical rarity seen as a naturally occurring illness only in Australia. In the past thirteen years the disease has shown itself on all the continents and the British Isles in circumstances varying from sharply defined and explosive outbreaks to smoldering and continuous endemics. It now appears that it can be found in any part of the world where an effort is made to detect it. Whether the disease has spread from Australia to the rest of the world in the years since its original description by Derrick¹ in 1937 or whether propitious circumstances led to its recognition first in Australia is a question that cannot be answered at present. In any case Q fever has become a disease of world wide importance and the considerable effort expended in attempts to define the natural history of this disease has been highly productive especially during the last decade.

Derrick² and Stoker and Marmion³ have presented very fine reviews of the findings relative to the epidemiology of the disease in general, beyond a brief synopsis of this subject the present paper will deal rather with data concerning the actual generation and persistence of infective sources of the agent.

THE EPIDEMIOLOGY OF Q FEVER

Sources of the agent in nature. *Coxiella burnetii* has been isolated from at least 22 species of ticks^{4,5} 1 insect²⁷ 8 mammals^{28,29} and at least 1 bird³⁰. Many of the ticks found naturally infected have been shown capable under laboratory conditions of transmitting the infection by bite. In some species stage to stage transfer of the agent occurs and in others the agent is transmitted transovarially. The tissues and excreta of such infected ticks have been shown to be potential sources of large numbers of the organism.

Among domestic animals cattle, sheep and goats would appear to be the most important sources of the organism because of their world wide distribution and large numbers. *C. burnetii* has been isolated from the tissues of cattle and sheep and from the secretions and excretions of all three species. The placenta is an especially rich source of the agent. In the case of sheep it has been shown that the wool itself^{31,32} may be heavily contaminated with the organism. This is presumably true of cattle and goat hides as well although this has not as yet been firmly established.

The work on which these observations are based was supported by Grant F 195 from the National Institute of Allergy and Infectious Diseases, Public Health Service, Bethesda, Md.

† Present address: National Institute of Allergy and Infectious Diseases, Public Health Service, Bethesda, Md.

‡ Present address: Communicable Disease Center, Public Health Service, Atlanta, Ga.

Other domestic animals and wild or domestic birds or fowl are perhaps potential sources of *C burnetii* but are less frequently involved in the disease cycle than are the three major domestic animal species.

Man has on occasion been the source of rickettsiae⁵²⁻⁵⁴ in tissues from fatal cases and from sputum and urine during the acute illness. Person to-person transmission of the agent has been of importance only in isolated instances⁵⁵ and does not seem to be of consequence in spread of the disease.

Modes of transmission to man or to domestic animals Transmission of the infective agent from natural sources to man or to domestic animals may occur in a number of ways depending upon the situation. In the case of man the association between infection and exposure to domestic animal or their products is very striking; exposure may be occupational or residential in type and intense or merely casual in nature. In the case of domestic animals a similar association may exist but it is considerably more difficult to establish because in most instances the occasions on which domestic animals can be kept from close association with other livestock are far less common.

Definite epidemiological evidence exists that *C burnetii* may be carried over considerable distances on inanimate objects such as clothing⁵⁶ wool⁵⁷ hair⁵⁸ straw⁵⁹ packing materials⁶⁰ and dust⁶¹. More direct evidence exists in the case of wool.

As mentioned previously the possibility of transmission by the bite of infected ticks is a potential method utilized by the agent for transfer from one host to another. However the evidence that such a method is of any real consequence is scant. On the other hand the tissues and excreta of infected ticks have undoubtedly been responsible for some human infections brought about by handling of items thus contaminated.

In the same manner that is through manipulation or agitation of contaminated surfaces the ground and soil of areas housing infected livestock may be through the generation of dusts containing infectious particles a very important method for transmission of the agent to susceptible hosts⁶²⁻⁶⁴.

Routes of infection for man and animals Several portals of entry into the susceptible host exist in the case of *C burnetii*. Thus laboratory animals may be infected through the unbroken (probably abraded) skin by subcutaneous intradermal or intraperitoneal inoculation through ingestion of infective materials by placement of infective materials on the conjunctiva by intranasal instillation or through inhalation of artificially generated infective aerosols. It is assumed that the same potential routes of infection exist in the case of larger animals and man as a matter of fact some of these routes have been shown epidemiologically to be of importance in producing infection under natural conditions.

With respect to man at least a general survey of the reported epidemiological findings indicates that the majority of cases of Q fever may well have resulted from contact with air borne rickettsiae taken into the body either through inhalation or through deposition onto the ocular conjunctivae or other mucous membranes. Of these possibilities the respiratory route constitutes the most probable means of ingress for the agent.

AIR BORNE TRANSMISSION OF Q FEVER THE ROLE OF PARTURITION IN THE GENERATION OF INFECTIVE AEROSOLS*

By Hartwell H. Welsh, Edwin H. Lennette, Francis R. Abinanti†
and John F. Winn‡

*Isral and Rickettsial Disease Laboratory, California State
Department of Public Health, Berkeley, Calif.*

Prior to 1944-1945 Q fever as a disease of humans was a clinical rarity seen as a naturally occurring illness only in Australia. In the past thirteen years the disease has shown itself on all the continents and the British Isles in circumstances varying from sharply defined and explosive outbreaks to smoldering and continuous endemics. It now appears that it can be found in any part of the world where an effort is made to detect it. Whether the disease has spread from Australia to the rest of the world in the years since its original description by Derrick¹ in 1937 or whether propitious circumstances led to its recognition first in Australia is a question that cannot be answered at present. In any case Q fever has become a disease of world wide importance, and the considerable effort expended in attempts to define the natural history of this disease has been highly productive especially during the last decade.

Derrick² and Stoker and Marmion³ have presented very fine reviews of the findings relative to the epidemiology of the disease in general beyond a brief synopsis of this subject the present paper will deal rather with data concerning the actual generation and persistence of infective sources of the agent.

THE EPIDEMIOLOGY OF Q FEVER

Sources of the agent in nature. *Coxiella burnetii* has been isolated from at least 22 species of ticks,^{4, 21} 1 insect,²² 8 mammals^{23, 49} and at least 1 bird.⁵⁰ Many of the ticks found naturally infected have been shown capable under laboratory conditions of transmitting the infection by bite. In some species stage to stage transfer of the agent occurs and in others the agent is transmitted transovarially. The tissues and excreta of such infected ticks have been shown to be potential sources of large numbers of the organism.

Among domestic animals cattle, sheep and goats would appear to be the most important sources of the organism because of their world wide distribution and large numbers. *C. burnetii* has been isolated from the tissues of cattle and sheep and from the secretions and excretions of all three species. The placenta is an especially rich source of the agent. In the case of sheep it has been shown that the wool itself^{51, 5} may be heavily contaminated with the organism. This is presumably true of cattle and goat hides as well although this has not as yet been firmly established.

The work on which these observations are based was supported by Grant F 195 from the National Institute of Allergy and Infectious Diseases, Public Health Service, Bethesda, Md.

† Present address: National Institute of Allergy and Infectious Diseases, Public Health Service, Bethesda, Md.

‡ Present address: Communicable Disease Center, Public Health Service, Atlanta, Ga.

Other domestic animals and wild or domestic birds or fowl are perhaps potential sources of *C burnetii* but are less frequently involved in the disease cycle than are the three major domestic animal species.

Man has on occasion been the source of rickettsiae⁵³⁻⁵⁴ in tissues from fatal cases and from sputum and urine during the acute illness. Person to-person transmission of the agent has been of importance only in isolated instances⁵⁵ and does not seem to be of consequence in spread of the disease.

Modes of transmission to man or to domestic animals—Transmission of the infective agent from natural sources to man or to domestic animals may occur in a number of ways depending upon the situation. In the case of man the association between infection and exposure to domestic animal or their products is very striking; exposure may be occupational or residential in type and intense or merely casual in nature. In the case of domestic animals a similar association may exist but it is considerably more difficult to establish because in most instances the occasions on which domestic animals can be kept from close association with other livestock are far less common.

Definite epidemiological evidence exists that *C burnetii* may be carried over considerable distances on inanimate objects such as clothing⁵⁶ wool⁵⁷ hair⁵⁸ straw⁵⁹ packing materials⁶⁰ and dust⁶¹. More direct evidence exists in the case of wool.

As mentioned previously the possibility of transmission by the bite of infected ticks is a potential method utilized by the agent for transfer from one host to another. However the evidence that such a method is of any real consequence is scant. On the other hand the tissues and excreta of infected ticks have undoubtedly been responsible for some human infections brought about by handling of items thus contaminated.

In the same manner that is through manipulation or agitation of contaminated surfaces the ground and soil of areas housing infected livestock may be through the generation of dusts containing infectious particles a very important method for transmission of the agent to susceptible hosts⁶²⁻⁶⁴.

Routes of infection for man and animals—Several portals of entry into the susceptible host exist in the case of *C burnetii*. Thus laboratory animals may be infected through the unbroken (probably abraded) skin by subcutaneous intradermal or intraperitoneal inoculation through ingestion of infective materials by placement of infective materials on the conjunctiva by intranasal instillation or through inhalation of artificially generated infective aerosols. It is assumed that the same potential routes of infection exist in the case of larger animals and man as a matter of fact some of these routes have been shown epidemiologically to be of importance in producing infection under natural conditions.

With respect to man at least a general survey of the reported epidemiological findings indicates that the majority of cases of Q fever may well have resulted from contact with air borne rickettsiae taken into the body either through inhalation or through deposition onto the ocular conjunctivae or other mucous membranes. Of these possibilities the respiratory route constitutes the most probable means of ingress for the agent.

Infective aerosols may be classified in several ways and for present purposes they may be classified as either primary or secondary in type. A primary aerosol may be defined as one generated directly by or in direct association with an infected animal for example, an infective aerosol arising during parturition of an infected animal. A secondary aerosol then is one produced by some factor or action or incident temporally or spatially removed from those giving rise to primary aerosols. Secondary aerosols are produced during the handling or processing of contaminated materials such as wool and hides at some distance in time and/or space from for example the site of the original parturition.

Infection with *C. burnetii* can obviously be produced by both primary and secondary aerosols. The production of secondary aerosols is relatively easy indeed, many instances of infection recorded in the literature appear to have been unavoidable. The handling of contaminated materials in plants or places far removed from the actual animal sources responsible for the contamination has on occasion produced outbreaks of Q fever in wool plants⁵⁷ laundries⁵⁸ and in such unlikely places as an art school⁵⁹ and a locksmith shop.⁶⁰ Such outbreaks appear to be possible whenever objects contaminated by infected animals are handled. In the immediate environs of infected live stock such infective aerosols are readily produced by a mere agitation of the ground surface.⁶¹

Secondary aerosols are highly important epidemiologically being a major factor in the dissemination of the disease to new areas and populations and perhaps they play a greater role in the production of human illness than do the primary aerosols nevertheless the circumstances surrounding the production and persistence of primary aerosols are of more than passing interest. It is the circumstances surrounding the production of these primary aerosols with which we shall be concerned in the remainder of this report.

THE PRODUCTION OF AEROSOLS OF *C. BURNETII* BY INFECTED ANIMALS

Many prior observations cited above had shown that a large percentage of human illness due to Q fever occurred in individuals exposed to domestic livestock either by occupation or residence. In attempts to clarify this relationship we found that the rickettsia is excreted by infected animals during parturition in the placenta and birth fluids and afterwards in the feces. Furthermore it appeared likely that human and perhaps also animal infection could be acquired by exposure to aerosolized infectious materials. Consequently air sampling studies were undertaken to determine the factors related to the appearance and persistence if any of *C. burnetii* in the air of the environs of known infected animals. A group of pregnant sheep was inoculated intravenously with varying doses of *C. burnetii*. Six animals were used in the studies and the air sampling trials using a cotton plug sampler were conducted over a period encompassing the ante partum parturient and post partum states of each infected sheep.

Methods and Materials

Inoculation of animals Six sheep in 2 groups of 3 animals each were inoculated intravenously via the external jugular vein. Each animal of the first group received 1×10^9 hamster infective doses (intrapertoneal) and each animal of the second group received 1×10^4 hamster infective doses of *C. burnetii*. The inoculum consisted of 10 ml of infected yolk sac diluted to the appropriate concentration with sterile skimmed milk.

Housing of infected animals The sheep used in these studies were housed in isolation cubicles of the following dimensions: width 3 feet, length 8 feet, and height 8 feet; total volume 192 cubic feet. Each cubicle was completely enclosed by wooden tongue and groove walls and ceilings. The floors were of cement. A solid plywood door at one end of the cubicle opened onto a common interior alleyway. The doorstep was raised 8 inches from the concrete floor. At the other end of the cubicle a door half glass gave access to an outer alleyway open to the exterior but covered by a roof.

Ventilation in each cubicle was accomplished by natural air flow through a 6×18 inch screened opening near the bottom of the door to the outer alleyway. There were no other gross openings in the cubicles. Traffic between the cubicles was confined to the inner alleyway; the usual isolation techniques were carried out at all times. However, the possibility that traffic or minute openings between the cubicles could result in cross contamination cannot be ruled out.

The sheep were placed in these pens prior to inoculation and were kept there throughout the entire study.

Once each week up to the time of parturition the pens were cleaned and fresh wood shavings added to a depth of approximately 3 inches. Pens were not cleaned for 2 weeks following parturition. Additional shavings were added as necessary, however, to provide a clean, dry surface for the comfort of the animals. Fresh water and feed were provided daily.

Types of air sampling equipment and preparation for use Cotton plug samplers of the type described by Rosebury⁶⁴ and as modified by S. Division, Camp Detrick, Md., were used throughout these studies.

Clean glass sampler shells were sterilized in an autoclave at 18 lb. pressure for 30 min. and then packed with wafers punched from a single roll, thickness of sterile No. 1 absorbent cotton. The wafers were cut with a three-fourths inch diameter leather punch and were thus similar with respect to thickness, diameter, and fiber length. The wafers were forced into the glass shell against a stainless steel retaining screen previously placed in position. Sufficient cotton wafers were placed in each sampler shell to provide resistance to air flow equal to displacement of 28 to 32 mm. of mercury at a flow rate of 20 l./min.

After packing and calibration the samplers were placed in cotton stoppered glass tubes and sterilized by dry heat (350° F. for 60 min.). Tests indicated that the resistance and the general physical characteristics of the samplers were not affected by this procedure. The samplers were kept in closed containers until used in the sampling trials.

Infective aerosols may be classified in several ways and for present purposes they may be classified as either primary or secondary in type. A primary aerosol may be defined as one generated directly by, or in direct association with, an infected animal for example, an infective aerosol arising during parturition of an infected animal. A secondary aerosol then is one produced by some factor, or action or incident temporally or spatially removed from those giving rise to primary aerosols. Secondary aerosols are produced during the handling or processing of contaminated materials such as wool and hides at some distance in time and/or space from, for example the site of the original parturition.

Infection with *C burnetii* can obviously be produced by both primary and secondary aerosols. The production of secondary aerosols is relatively easy indeed many instances of infection recorded in the literature appear to have been unavoidable. The handling of contaminated materials in plants or places far removed from the actual animal sources responsible for the contamination has on occasion produced outbreaks of Q fever in wool plants⁵⁷ laundries⁵⁸ and in such unlikely places as an art school⁶⁰ and a locksmith shop.⁵⁹ Such outbreaks appear to be possible whenever objects contaminated by infected animals are handled. In the immediate environs of infected live stock such infective aerosols are readily produced by a mere agitation of the ground surface.^{6 63}

Secondary aerosols are highly important epidemiologically being a major factor in the dissemination of the disease to new areas and populations and perhaps they play a greater role in the production of human illness than do the primary aerosols nevertheless the circumstances surrounding the production and persistence of primary aerosols are of more than passing interest. It is the circumstances surrounding the production of these primary aerosols with which we shall be concerned in the remainder of this report.

THE PRODUCTION OF AEROSOLS OF *C. burnetii* BY INFECTED ANIMALS

Many prior observations cited above had shown that a large percentage of human illness due to Q fever occurred in individuals exposed to domestic livestock either by occupation or residence. In attempts to clarify this relationship we found that the rickettsia is excreted by infected animals during parturition in the placenta and birth fluids and afterwards in the feces. Furthermore it appeared likely that human and perhaps also animal infection could be acquired by exposure to aerosolized infectious materials. Consequently air sampling studies were undertaken to determine the factors related to the appearance and persistence if any of *C. burnetii* in the air of the environs of known infected animals. A group of pregnant sheep was inoculated intravenously with varying doses of *C. burnetii*. Six animals were used in the studies and the air sampling trials using a cotton plug sampler were conducted over a period encompassing the ante partum parturient and post partum states of each infected sheep.

TABLE 1
RESULTS OBTAINED FROM AIR SAMPLING CARRIED OUT IN PENS HOLDING PREGNANT SHEEP ARTIFICIALLY INFECTED WITH *COXIELLA BURNETII* SAMPLING DONE PRIOR TO PARTURITION

Dose Pia ta	P 9 Sheep 12		P 10 Sheep 13		P 11 Sheep 2		P 12 Sheep 7		P 13 Sheep 4		P 14 Sheep 38	
	1 X 10 ⁶ O t		1 X 10 ⁶ Inf		1 X 10 ⁶ Out		1 X 10 ⁶ I		1 X 10 ⁶ O t		1 X 10 ⁶ I	
D t	P	CS	P	CS	P	CS	P	CS	P	CS	P	CS
January 16	-14	-									-20	-
17	-13	-									-19	-
18	-12	-									-18	-
19	-11	NC									-17	NC
20	-10	NC									-16	NC
21	-9	-									-15	-
22	-8	-									-14	-
23	-7	-									-13	-
24	-6	-									-12	-
25	-5	NC									-11	NC

Placenta removed from pen immediately postpartum

† Placenta left in pen throughout entire sampling period

I = Days before (-) or days after (+) parturition

CS = Results of hamster inoculation of the cotton plug air sample taken on a given day

+ = *C. burnetii* present in air sample

- = *C. burnetii* not present in sample

NC = No sample collected

On January 28 parturition occurred in sheep No 13 in pen 10 and it will be noted from TABLE 2 that the rickettsiae were recovered from the air of this pen on this day and that they were present therein for at least 14 days thereafter. The air sample taken in this same pen on the day before parturition did not contain the rickettsiae. It will also be observed that on the day prior to parturition in sheep No 12 pen 9 rickettsiae were present in the air of the pen and presumably represented contamination from the adjacent pen 10. It is possible that contamination of the atmosphere in the other pens may have occurred but since sampling in pens 11 to 13 was not done no statement can be made. However no rickettsiae were recovered from the air of pen 14 which was the farthest removed from pens 9 and 10. Parturition in sheep No 12 on January 30 was also associated with the presence of the rickettsiae in the air of the pen (pen 9) in which it lambled and the organisms were present continuously for at least 11 days after the parturition.

Essentially the same situation was observed in the cases of sheep Nos 2, 4, 7 and 38 that is parturition was accompanied by the presence of rickettsiae in the air during an appreciable period of time. Some cross contamination presumably occurred between the pens in several instances since the rickettsiae could occasionally be detected in pens in which the animals had not yet lambled. This sporadic appearance in these pens is in contrast to the essentially continuous presence of the organism in the same pen after the occupant had lambled. These observations on the appearance of rickettsiae in the air

After use in the trials the cotton plug portion of the sampler was removed aseptically from the glass shell placed in a sterile screw capped jar and stored at -20°C until it was prepared for testing in laboratory animals. Prior to testing the jar containing the cotton plug was placed in cold running tap water until the sampler was completely thawed. The cotton plug was then broken up with the use of sterile forceps and to each sample was added 25 ml of a 0.4 per cent solution of Triton X 100* in distilled water. The mixture was agitated for 20 min and 5 ml of the resulting supernatant fluid was pipetted off. Penicillin was added to a final concentration of 400 units/ml and the mixture allowed to incubate for 4 hours at 4°C prior to inoculation into animals.

Inoculation of laboratory animals Hamsters six to eight weeks of age were used as test animals in these studies. Each mixture was inoculated into four hamsters each animal receiving 1 ml intraperitoneally. Six weeks after inoculation the animals were bled by cardiac puncture and their sera were examined for the presence of antibodies to *C. burnetii* by the complement fixation technique. The presence of antibody in the serum of one or more animals of a test group of four was regarded as evidence of infection and as an indication that the specimen inoculated had contained *C. burnetii* (large scale studies had indicated that the hamster population in use had had no previous experience with the organism). During the six week test period uninoculated control animals were left in the same quarters as the inoculated hamsters. Complement fixation tests done on sera drawn from these animals after this period were negative indicating that the antibody present in the sera of inoculated test animals was not elicited by cross contaminations.

Conditions of air sampling Samplers were placed 4 feet above the surface level of the sheep pens through a small port cut into the interior door. The sampling rate was set at 10 l/min by means of calibrated limiting orifices set into the sampling lines. Samples were taken for periods of one half to one hour daily starting with a date approximately 1 week prior to the expected parturition in each pen and continuing to a date 2 weeks postpartum in each case.

Results

The results of the air sampling trials together with certain ancillary data are presented in TABLES 1 and 2. Air sampling trials were conducted prior to the first parturition in this group of 6 animals sampling being done in pens 9 and 14. The original intention was to determine for each pen whether the ante partum atmosphere was free of the organism. The long period of sampling shown for pens 9 and 14 was unintentional and it arose because parturition was not as imminent as was believed at the time the sampling was initiated. However the prolonged observations made on these 2 pens show quite well that no rickettsiae at least in detectable amounts were present in the air during the period between inoculation of the animals and the occurrence of the first parturition in the group a period of 12 days.

TABLE 3
INFECTIVITY OF PLACENTA BIRTH FLUIDS AND FECES OF PARTURIENT
SHEEP INFECTED WITH *COXIELLA BURNETII*

Sheep No.	Infected Material			Feces collected post partum per day	Time of inoculation day
	Placenta	Birthing fluid	Feces		
12 (Pen 9)	1×10^1	NC	1×10^4 (12 hr)	14	1×10^2 (14)
13 (Pen 10)	1×10^2	1×10^2	1×10^2 (10 min)	15	1×10^2 (14)
2 (Pen 11)	1×10^1	1×10^2	1×10^1 (1 hr)	18	1×10^2 (14)
7 (Pen 12)	1×10^2	1×10^2	1×10^1 (10 min)	11	Negative (14)
4 (Pen 13)	1×10^1	NC	1×10^1 (4 hr)	14	1×10^1 (10)
38 (Pen 14)	1×10^1	NC	1×10^2 (4 hr)	14	1×10^1 (14)

Infecitivity titers of material expressed as 50 per cent hamster infective doses per gm or ml

† First postpartum specimen collected 10 min to 12 hr postpartum as indicated

the stools is open to conjecture. It is possible that the presence of the organism in the stool represents true excretion in the sense that the organisms were released from some focus in the body and were excreted into the intestinal tract or that multiplication may have occurred in the intestinal mucosa. On the other hand the organisms in the stool may represent mere passive contamination due to the ingestion of contaminated food or inhalation of infectious materials subsequently swallowed. Experimental work directed to a firmer understanding of these points is highly desirable.

That the act of parturition itself is responsible for the initial contamination of the air is well illustrated in TABLE 4 which gives the chronological sequence of events at parturition as they relate to the first appearance of the rickettsiae in the air. In the case of sheep No. 38 the last sample of air taken before parturition was obtained approximately 18 hours before the birth of the lamb. No rickettsiae were detectable in this specimen. However a sample taken at an interval of about 3 to 5 hours after the parturition did contain demonstrable numbers of the organism. In the case of sheep No. 2 rickettsiae were detected in air samples taken 1 to 2 hours after parturition. The air of this same pen had not contained the rickettsiae 24 hours prior to parturition. The most interesting findings pertain to sheep No. 13. As can be seen from TABLE 4 air sampling was being conducted for 55 minutes before and for 5 minutes after the first lamb of a set of twins was born. The specimen of air collected during this 1 hour period was negative for rickettsiae even though sampling was carried out for about 5 minutes after the birth. Air sampling was resumed using fresh filters at 2:54 P.M. and continued for 10 minutes being discontinued just as the second lamb was born. This air sample contained rickettsiae indicating that the organisms appeared in the air within 5 to 24 minutes after the birth of the first lamb (worthy of remark is the fact that the air samplers were in the normal position as described earlier—that is, no attempt was made to place the samplers in especially favorable locations near the animal during the actual parturition). Offhand it is difficult to see how more direct evidence implicating parturition in environmental

TABLE 2
RELATIONSHIP BETWEEN PARTURITION OF SHEEP INFECTED WITH *COXIELLA BURNETII* AND
APPEARANCE OF THE RICKETTSIA IN THE AERIAL ENVIRONMENT

D sag Pl c ta	P n 9 She p 12		Pe 10 Sheep 13		P 11 She p 2		P 12 Sh p 7		P 13 Sh p 4		Pe 14 Sh p 35	
	1 X 10 Out		1 X 10 I		1 X 10 Out		1 X 10 ⁴ In		1 X 10 ⁴ O t		1 X 10 In	
D t	P	CS	P	CS	P	CS	P	CS	P	CS	P	CS
January 26	-4	-									-10	-
27	-3	-	-1	-							-9	NC
28	-2	-	0	+							-8	-
29	-1	+	+1	+							-7	-
30	0	+	+2	+							-6	+
31	+1	+	+3	+							-5	NC
February 1	+2	+	+4	+							-4	+
2	+3	+	+5	+	-1	-			0	+	-3	NC
3	+4	NC	+6	NC	0	+			+1	+	-2	+
4	+5	+	+7	+	+1	+	-4	-	+2	+	-1	-
5	+6	+	+8	+	+2	+	-3	+	+3	+	0	-
6	+7	+	+9	-	+3	+	-2	-	+4	+	+1	+
7	+8	+	+10	+	+4	-	-1	-	+5	-	+2	+
8	+9	+	+11	+	+5	+	0	+	+6	+	+3	-
9	+10	+	+12	+	+6	+	+1	+	+7	+	+4	+
10	+11	+	+13	+	+7	+	+2	+	+8	+	+5	+
11	+12	-	+14	+	+8	+	+3	+	+9	+	+6	+
12	+13	+			+9	+	+4	+	+10	-	+7	+
13	+14	-			+10	-	+5	+	+11	+	+8	+
14					+11	-	+6	+	+12	+	+9	+
15					+12	-	+7	-	+13	-	+10	+
16					+13	-	+8	+	+14	-	+11	-
17					+14	-	+9	+			+12	+
18							+10	-			+13	+
19							+11	-			+14	-
20							+12	-				
21							+13	-				
22							+14	-				

Refer to TABLE 1 for explanation of symbols

of pens housing preparturient animals coincidentally with and following parturition in infected animals in adjacent pens agree well with other studies done in this laboratory

From TABLE 3 it will be noted that both the placenta and the birth fluids contained rickettsiae and that the rickettsial content of the former was tremendous. It might be assumed therefore that the initial and most important source of rickettsiae was the placenta and to a lesser extent the birth fluids. Nevertheless it would appear that the long continued excretion of the organism in the stool postpartum also must have contributed to the contamination of the environment. On the whole the rickettsial content of the stools was considerably less than that of the placenta. The true origin of the rickettsiae in

which the organism could be detected in the air. It should be kept in mind however that even though the placenta was removed there still remained the possibility that an appreciable amount of contamination could have arisen from spillage of the infected birth fluids. In view of the relatively low rickettsial content of these fluids their ability to give rise to long continuous contamination of the air can be questioned. The possible role of infected feces in providing a constantly renewed source of the organism cannot be overlooked. The rickettsial content of the feces was in some instances quite high but here the question of cause and effect must be considered that is whether the infected stools were the result of true excretion or of only passive contamination having as its source the infected environment itself.

The relative importance of these various materials in determining the time span of aerial contamination with *C. burnetii* will be shown only when it is determined whether replenishment of this nature is necessary. It is entirely possible that some proportion of the total concentration of rickettsiae initially introduced into the air at parturition will remain airborne in detectable amounts for a considerable period especially under interior conditions.

In attempting to elucidate some of the problems concerning the original sources of *C. burnetii* in the environment of infected animals we have raised other questions with respect to the persistence of the rickettsiae in such environments. The observations made thus far have shown that the rickettsiae are not normally present in the aerial environment of artificially infected pregnant sheep until parturition occurs. It is also clear that the act of parturition itself is responsible in some manner for the generation of the primary aerosols of this organism but that the persistence of the agent in the air over periods up to 14 days postpartum may depend on other factors.

References

- 1 DERRICK E. H. 1937. Q fever a new fever entity clinical features diagnosis and laboratory investigation. *Med J Australia* 2: 281.
- 2 DERRICK E. H. 1933. The epidemiology of Q fever a review. *Med J Australia* 1: 245.
- 3 STOKER M. G. P. & B. P. MARMION. 1955. The spread of Q fever from animal to man. *Bull. World Health Organization* 13: 781.
- 4 BLANC G. & J. BRUNEAU. 1949. Un réservoir de virus de la Q. fever en Algérie la tique *Hyalomma marginatum*. *Bull. Acad. Natl. Med.* 133: 115.
- 5 BLANC G., J. BRUNEAU & D. CHABAUD. 1950. Présence de *Rickettsia (Coxiella) burnetii* chez des tiques de l'espèce *Amblyomma tigris* recueillies en Afrique Equatoriale Française. *Compt. rend.* 230: 474.
- 6 BLANC C., L. A. MARTIN & A. MAURICE. 1947. Le mélon (*Mersinella*) de la région de Goulmine est un réservoir de virus de la Q. fever marocaine. *Compt. rend.* 224: 1673.
- 7 BLANC G., J. BRUNEAU, R. POITROT & B. DELAGE. 1948. Quelques données sur la Q. fever (maladie de Derrick-Burnet) expérimentale. *Bull. Acad. Med.* 132: 243.
- 8 CAPLEY J. G. & J. H. POPE. 1933. The isolation of *C. burnetii* from the tick *Ixodes holocyti* in Queensland. *Australian J. Exptl. Biol. Med. Sci.* 31: 613.
- 9 COX H. R. 1940. *Rickettsia diapausa* and American Q fever. *Am. J. Trop. Med.* 20: 463.
- 10 DAVIS G. E. 1943. American Q fever experimental transmission by the *Argasid* ticks *Ornithodoros* *microti* and *O. hermsi*. *U. S. Public Health Repts.* 58: 984.
- 11 DAVIS G. E. & H. R. COX. 1938. A filter passing infectious agent isolated from ticks *I. solomonensis* from *Dermacentor* a *dermatitis* reactions in animals and filtration experiments. *U. S. Public Health Repts.* 63: 2259.

TABLE 4
RELATIONSHIP BETWEEN PARTURITION OF ARTIFICIALLY INFECTED SHEEP
AND CONTAMINATION OF THE ENVIRONMENT WITH *COXIELLA*
BURNETII AS SHOWN BY AIR SAMPLING TRIALS

Sheep No	Date	Sampling Period		Sample Result & Its	Remark
		Start	Stop		
38 (Pen 14)	February 4	12 20 P M	1 20 P M	-	17-19 hr prepartal
	5	10 30 A M	11 30 A M	+	3-5 hr postpartum
	6	11 50 A M	12 50 P M	+	30-32 hr postpartum
2 (Pen 11)	2	11 20 A M	11 50 A M	-	24 hr prepartal
	3	12 00 M	1 00 I M	+	1-2 hr postpartum
	4	2 50 P M	3 50 P M	+	28 hr postpartum
13 (Pen 10)	January 28	1 45 I M	2 45 P M	-	(Bag of waters protruding broke as first lamb born at 2 40 P M)
		2 54 P M	3 04 P M	+	(Labor—second lamb born 3 04 P M)
		4 30 P M	5 30 P M	+	1½ to 2½ hr postpartum

contamination could be obtained. With regard to the environment of the other 3 animals involved it will be recalled from TABLE 2 that parturition of the animals in the respective pens resulted in either initial or renewed evidence of aerial contamination although the time sequence could not be as closely followed.

In discussing TABLE 2 it was seen that the rickettsiae persisted in the air of the pens for several days after parturition. The concentration of the inoculum originally administered to the animals apparently had no influence on the length of persistence. The air of pens 9, 10, and 11 holding those sheep that received 1×10^9 hamster infective doses contained rickettsiae in detectable amounts for 13, 14, and 9 days respectively, or an average of 12 days postpartum. Pens 12, 13, and 14 housing the animals that received 1×10^4 doses contained rickettsiae for 9, 12, and 13 days for an average persistence in these cases of 11.33 days postpartum.

Because of the tremendous concentrations of rickettsiae present in the placental tissues it seemed that removal of this highly infected material might affect the extent to which rickettsiae would continue to be found in the air. However, in this small series no obvious differences were noted regardless of whether the placenta was removed from the pen or remained there during the entire sampling period.

DISCUSSION

In these studies it has been possible to show rather conclusively that the act of parturition is responsible for the generation of primary aerosols of *C. burnetii*. The actual mechanisms involved are admittedly obscure.

In this present series the removal of the highly infective placenta from the pen immediately postpartum had no effect on the length of time during

les animaux domestiques vis à vis des antigènes rickettsiens en milieu contaminé d
fièvre Q. Comp. rend 231 86

- 38 HUEBNER R J & J A BELL. 1951 Q fever studies in Southern California. Summary of current results and a discussion of possible control measures. J Am Med Assoc 145 301
- 39 HUEBNER R J W I JELLI ON M D BECK R K PARKER & C C SHEPARD. 1948 Q fever studies in Southern California. I Recovery of *Rickettsia burnetii* from raw milk. U S Public Health Repts 63 214
- 40 JELLI ON W I K ORM REF M D BECK R J HUEBNER R R PARKER & F J BELL. 1948 Q fever studies in Southern California. V Natural infection in a dairy herd. U S Public Health Repts 63 1611
- 41 JELLI ON W I H H WELSH B F ELSON & I J HUEBNER. 1950 Q fever studies in Southern California. VI Recovery of *Coxiella burnetii* from milk of sheep. U S Public Health Repts 65 395
- 42 KILSCHNIGER C & F WIESMAN. 1949 Mortus Epidemic bei Ziegen bedingt durch *Rickettsia burnetii*. Schweiz. Arch. Tierheilk 91 553
- 43 KITAHARA M & K TAKANO. 1952 Complement fixation of Q fever. Comparative evaluation of several techniques. Japan J Med Sci Biol 5 493
- 44 LENNETTE F H B H DEAN F R ABINANTI W H CLARK J F WINN & M A HOLME. 1951 Q fever in California. V Serologic survey of sheep goats and cattle in three epidemiological categories from several geographic areas. Am J Hyg 64 1
- 45 LUTTO I & R J HUEBNER. 1950 Q fever studies in Southern California. IX Isolation of Q fever organisms from parturient placentas of naturally infected dairy cows. U S Public Health Repts 65 541
- 46 MANTOVANI A & I BENAZZI. 1953 The isolation of *Coxiella burnetii* from *Rhipicephalus sanguineus* on naturally infected dogs. J Am Vet Med Assoc 122 117
- 47 PEREZ GALLARDO F G CLAVERO & S FERNANDEZ FERNANDEZ. 1952 Epidemiological studies of Q fever in Syrian Mountain rabbits as intermediate reservoirs of infection of *Coxiella burnetii*. Rev. sanit. hig. publ. Madrid 26 81
- 48 WELSH H H F H LENNETTE F R ABINANTI & J F WINN. 1951 Q fever in California. IV Occurrence of *Coxiella burnetii* in the placenta of naturally infected sheep. U S Public Health Repts 66 1473
- 49 WINN J F F H LENNETTE H H WELSH & F R ABINANTI. 1953 Q fever studies. VII Presence of *Coxiella burnetii* in the feces of naturally infected sheep. Am J Hyg 68 183
- 50 BABUDIERI B & C MO CONICI. 1952 Experimental and natural infection of birds by *Coxiella burnetii*. Nature 169 195
- 51 ABINANTI F R H H WELSH J F WINN & F H LENNETTE. 1955 Q fever studies. XIX Presence and epidemiological significance of *Coxiella burnetii* in sheep wool. Am J Hyg 61 362
- 52 STOKER M C I R D BROWN F J I KETT P C COLLINGS & B P MARMION. 1955 Q fever in Britain. Isolation of *Rickettsia burnetii* from placenta and wool of sheep in an endemic area. J Hyg 53 313
- 53 SPICKNALL C C R J HUEBNER J A TINGER & W P BLOCKER. 1947 Report of an outbreak of Q fever at the National Institutes of Health. Ann Internal Med 27 28
- 54 BROWN D C I A KNIGHT & W L JELLI ON. 1948 A fatal case of Q fever in Southern California. Clin Med 69 3
- 55 SIEGERT R W SIMROCK & U STRODER. 1950 Über eine epidemischen Ausbruch von Q-Fieber in einem Krankenhaus. Z Tropenmed u Parasitol 2 1
- 56 OLIPHANT J W D A GORDON A MEIS & R R PARKER. 1949 Q fever in laundry work is presumably transmitted from contaminated clothing. Am J Hyg 49 16
- 57 SIGFL M M T I McN SCOTT W HENLY & O H JANTON. 1950 Q fever in a wool and hair processing plant. Am J Public Health 40 524
- 58 BEHRENS E A. 1950 Q fever—an epidemiological note. U S Public Health Repts 65 88
- 59 WEFMAN F. 1948 Über die Q-Fever (Queenslandfieber) Epidemie in Graubünden. Sch. et med. Wochschr 78 529
- 60 HARFAN M S C B FORBES & B P MARMION. 1951 An outbreak of Q fever in the United Kingdom. J Hyg 51 152
- 61 CLARK W H A S BLOCKER F H LENNETTE B H DEAN & J R WALKER. 1951

- 12 DE PRADA J, B GAY & A LLORENTE 1950 Primer caso de fiebre Q humana en España Med Colon 15 131 also Abstr Trop Dis cases Bull 19 0 47 533
- 13 FREYCHIE M J & Z DEUTSCHMAN 1950 Human rickettsioses in Africa Epidemiological and vital statistics Rept World Health Organization 3 160
- 14 GIROLD P 1950 Les rickettsioses en Afrique équatoriale Bull World Health Organization 4 535
- 15 HENGEL VON R G A KALSCHIE & E SHERIS 1950 Über zwei dorflche Q fieber epidemien in Baden Deut med Wochschr 75 1505
- 16 INSTITUT PASTEUR DU MAROC 1948 Rapport sur le fonctionnement Rickettsioses 3
- 17 JADIN J 1950 Les rickettsioses du Congo Belge et du Ruanda Urundi I ouvain Abstr in Trop Diseases Bull 1951 48 879
- 18 JELLISON W L F J BELL R J HUEBNER R R PARAFR & H H WELSH 1948 Q fever studies in Southern California IV Occurrence of *Coxiella burnetii* in the Spi nose ear tick *Otobius megnini* U S Public Health Repts 63 1483
- 19 KOHLS G M 1948 Vectors of rickettsial diseases In Rickettsial Dis cases of Man 83-96 Am Assoc Advance Sci Washington D C
- 20 PARKER R R & G E DAVIS 1938 A filter passing infectious agent isolated from ticks II Transmission by *Dermacentor andersoni* U S Public Health Repts 63 2267
- 21 PARKER R R & G M KOHLS 1943 American Q fever the occurrence of *Rickettsia diaporica* in *Amblyomma americanum* in Eastern Texas U S Public Health Repts 58 1510
- 22 PARKER R R & O SLESSMAN 1949 Spontaneous infection of the brown dog tick *Rhipicephalus sanguineus* with *Coxiella burnetii* U S Public Health Repts 64 1159
- 23 PARKER R R J DE PRADA E J BELL & D B LACKMAN 1949 Recovery of *C burnetii* from *H. savignyi* collected in Spain U S Public Health Repts 64 1616
- 24 PAYZIN S & N AKKAY 1952 Natural occurrence of *Coxiella burnetii* in *Ornithodoros lahorensis* ticks Turk Ijen Tetrubi Biol Dergisi 12 7 Abstr in Trop Diseases Bull 1953 60 103
- 25 PEREZ GALLARDO F G CLAVERO & S HERNANDEZ FERNANDEZ 1949 Hallazgo en España de la *Rickettsia burnetii* agente etiologico de la fiebre Q Rev sanidad hig publ (Madrid) 23 489
- 26 SMITH D J W 1940 Studies in the epidemiology of Q fever 3 The transmission of Q fever by the tick *Haemaphysalis humerosa* Australian J Exptl Biol Med 18 103
- 27 SMITH D J W 1941 Studies in the epidemiology of Q fever 8 The transmission of Q fever by the tick *Rhipicephalus sanguineus* Australian J Exptl Biol Med 19 133
- 28 SMITH D J W 1942 Studies in the epidemiology of Q fever 10 The transmission of Q fever by the tick *Ixodes holocyclus* Australian J Exptl Biol Med 20 213
- 29 SMITH D J W & E H DERRICK 1940 Studies in the epidemiology of Q fever I The isolation of six strains of *Rickettsia burnetii* from the tick *Haemaphysalis lumsrosa* Australian J Exptl Biol Med 18 1
- 30 STONER M G P & B P MARMION 1955 Q fever in Britain Isolation of *Rickettsia burnetii* from the tick *Haemaphysalis punctata* J Hyg 63 322
- 31 TAYLOR K M R A MOUNT H HOOGSTRAAL & H R DRESSLER 1952 The presence of *Coxiella burnetii* (Q fever) in Egypt J Egypt Public Health Assoc 27 123
- 32 GIROLD P & J JADIN 1950 Le pou dans la fièvre Q au Ruanda Urundi (Congo Belge) Conservation naturelle de l'antigène de la fièvre Q sur le pou Bull soc pathol exotique 43 674
- 33 ABINANTI I K H H WELSH F H LENNETT & O BRUNETTI 1953 Q fever studies XVI Some aspects of the experimental infection induced in sheep by the intratracheal route of inoculation Am J Hyg 67 140
- 34 ANTONETTI F 1952 Ricerche sperimentali sulla rickettsiosi *burnetii* del cane Z o profilassi 6 249
- 35 CAMINOPETROS J I 1948 La Q fever en Grece Le lait source de l'infection pour l'homme et les animaux Ann parasit humaine et comparée 23 107
- 36 DERRICK F H & D J W SMITH 1940 Studies in the epidemiology of Q fever 2 The isolation of three strains of *Rickettsia burnetii* from the tan hooat *Ixodrom torosus* Australian J Exptl Biol Med 18 99
- 37 GIROLD P & J JADIN 1950 Médecine expérimentale Comportement sérologique

- les animaux domestiques vis à vis les arthropodes rickettsiens en milieu contaminé de fièvre Q. *Comp rend* 231 96
- 38 HUEBNER K J & J A BELL. 1951. Q fever studies in Southern California. Summary of current results and a discussion of possible control measures. *J Am Med Assoc* 145 301
 - 39 HUEBNER K J, W L JELLINE, M D BECK, R R TARKER & C C SHEPARD. 1949. Q fever studies in Southern California. I. Recovery of *Coxiella burnetii* from raw milk. *U S Public Health Repts* 63 214
 - 40 JELLINE, W L, R ORMSBEE, M D BECK, R J HUEBNER, R R TARKER & F J BELL. 1949. Q fever studies in Southern California. V. Natural infection in a dairy herd. *U S Public Health Repts* 63 1611
 - 41 JELLINE, W L, H H WELSH, B F ELSON & R J HUEBNER. 1950. Q fever studies in Southern California. VI. Recovery of *Coxiella burnetii* from milk of sheep. *U S Public Health Repts* 65 395
 - 42 KILCHERFFER C & F WISMAN. 1949. Abortus Epileptici bei Ziegen bedingt durch *Rickettsia burnetii*. *Schweiz Arch Tierhilk* 91 553
 - 43 KITAHARA M & K TAKANO. 1952. Complement fixation in Q fever. Comparative evaluation of several techniques. *Japan J Med Sci Biol* 5 493
 - 44 LENNETTE E H, B H DEAN, F R ABINANTI, W H CLARK, J F WINN & M A HULME. 1951. Q fever in California. V. Serologic survey of sheep, goats and cattle in three epidemiological categories from several geographic areas. *Am J Hyg* 64 1
 - 45 LOTO L & R J HUEBNER. 1950. Q fever studies in Southern California. IX. Isolation of Q fever organisms from parturient placentas of naturally infected dairy cows. *U S Public Health Repts* 65 541
 - 46 MANTOVANI A & P BENAZZI. 1953. The isolation of *Coxiella burnetii* from *Rhipicephalus sanguineus* on naturally infected dogs. *J Am Vet Med Assoc* 122 117
 - 47 PEREZ GALLARDO F, C CLAVERO & S ESPANDEZ FERNANDEZ. 1952. Epidemiological studies of Q fever in Spain. Mountain riffs and domestic reservoirs of infection of *Coxiella burnetii*. *Rev sanitad hig publ Mat* 26 81
 - 48 WELSH H H, F H LENNETTE, F R ABINANTI & J F WINN. 1951. Q fever in California. IV. Occurrence of *Coxiella burnetii* in the placentas of naturally infected sheep. *U S Public Health Repts* 66 1473
 - 49 WINN J F, F H LENNETTE, H H WELSH & F R ABINANTI. 1953. Q fever studies. XVII. Presence of *Coxiella burnetii* in the feces of naturally infected sheep. *Am J Hyg* 68 183
 - 50 BARUZZI B & C MOSCOWITZ. 1952. Experimental and natural infections of *Coxiella burnetii*. *Nature* 169 195
 - 51 ABINANTI F, R H H WELSH, J F WINN & F H LENNETTE. 1955. Q fever studies. XIX. Presence and epidemiologic significance of *Coxiella burnetii* in sheep wool. *Am J Hyg* 61 362
 - 52 STOKER M, C P R D BROWN, F J J KEFF, I C CALINGS & B I MARMON. 1955. Q fever in Britain. Isolation of *Rickettsia burnetii* from placenta and wool of sheep in an endemic area. *J Hyg* 53 313
 - 53 SPICKALL C, R J HUEBNER, J A HINCHER & W I BLOCKER. 1947. Report of a outbreak of Q fever at the National Institutes of Health. *Ann Internal Med* 27 28
 - 54 BROWN D C, L A KNIGHT & W I JELLINE. 1948. A fatal case of Q fever in Southern California. *Calif Med* 69 3
 - 55 SIGBERT R, W SIMROCK & U STRÖDER. 1950. Über einen epidemischen Ausbruch von Q-Fieber in einem Ferienhaus in Tropen. *Monatsschr Parasitol* 2 1
 - 56 OLIPHANT J, W D A CORDON, A MEIS & R R TARKER. 1949. Q fever in London. Work is preliminary transmitted from contaminated clothing. *Am J Hyg* 49 76
 - 57 SIGEL M, M T F MCN SCOTT, W HENEF & O H JANTON. 1950. Q fever in a wool mill handling processing plant. *Am J Public Health* 40 524
 - 58 BYRN S L A. 1950. Q fever—an epidemiological note. *U S Public Health Repts* 65 88
 - 59 WEFMANN I. 1948. Über eine Q fever (Queenslandfieber) Epidemie in Craulund. *Schweiz med Wochschr* 78 52
 - 60 HARVEY M S, G B F BRUCE & B I MARMON. 1951. An outbreak of Q fever in East Kent. *Lancet* 2 1152
 - 61 CLARK W H, A S BOGUCKI, F H LENNETTE, B H DEAN & J R WALKER. 1951

- Q fever in California VI Description of an epidemic occurring at Davis California in 1948 Am J Hyg 54 15
- 62 DELAY P D E H LENNETTE & K B DEOME 1950 Q fever in California II Recovery of *Coxiella burnetii* from naturally infected air borne dust J Immunol 65 211
- 63 LENNETTE F H & H H WELSH 1951 Q fever in California V Recovery of *Coxiella burnetii* from the air of premises harboring infected goats Am J Hyg 64 44
- 64 ROEBURN T 1947 Experimental Air Borne Infection Williams & Wilkins Baltimore Md

HUMAN BRUCELLOSIS WITH SPECIAL REFERENCE TO THE DISEASE IN THE UNITED STATES

By Norman B. McCullough

*Laboratory of Clinical Investigation, National Institute of Allergy and
Infectious Diseases, Public Health Service, Bethesda, Md.*

To evaluate the problems encountered in brucellosis it is necessary to understand the nature of the disease and some of the characteristics of the causative organism.

Brucella melitensis, *Brucella abortus*, and *Brucella suis* are closely related organisms that cause similar diseases in animals and man. However, there are differences that are reflected in the epidemiology, the reported incidence, the laboratory diagnosis, the clinical aspects, and the therapy of brucellosis. In actuality we are dealing with three diseases. A few of the characteristics of *Brucella* that bear on the problems at hand will be considered.

Br. melitensis and *Br. suis* are less exacting in their growth requirements than is *Br. abortus*. The former grow better on the older laboratory media, require no added carbon dioxide, and can be isolated much more readily from man. Prior to the development of adequate media and techniques, *Br. abortus* was seldom isolated from human cases of the disease. Even now, the average laboratory routine seldom succeeds in isolating this species. Since the majority of cases of brucellosis in the United States have been due to *Br. abortus*, these facts have influenced the reported vital statistics of the disease.

Another characteristic of *Brucella* that has been of great significance is the tendency for most strains of the organism to undergo dissociation when grown on laboratory media. Until recent years, most of the experimental work was done with dissociated cultures. With progressive dissociation, there is first loss in virulence and then change in antigenicity. Antigen for the agglutination test made from dissociated cultures does not perform satisfactorily. Surveys^{1, 2} in which the same sera were examined in different laboratories have shown great discrepancies in test results, even to the degree that sera containing agglutinins in high titer were reported as negative in some laboratories. Unsatisfactory antigens and nonstandard techniques are still too widely in use.

The difficulty in isolating the organism from the patient, together with the lack of dependability and uniformity in the results of the agglutination test, not only has led (and still leads) to inaccuracy of diagnosis, but also to a distorted concept of the disease. Since they were forced to rely mainly on a clinical impression, many physicians ceased to employ laboratory results or disregarded them, and made diagnoses of brucellosis in the absence of any objective laboratory evidence of the disease. This is not a problem confined to the past, but one that is currently still of considerable magnitude. Its bearing on the reported incidence of the disease and on concepts of therapy should be apparent.

A consideration of the nature of the infection is of great help in understanding the problems in diagnosis and therapy of the disease. Brucellosis in both domestic animals and man probably has existed since the dawn of history. The organism has become well adapted to life in these hosts and accordingly

it usually produces a subacute or chronic disease. Man tolerates the infection less well than do domestic animals and acute disease may result. However this is the exception rather than the rule. In exposed populations a few individuals develop serious acute and chronic disease, the rest have subclinical infections or brief or low grade illnesses that escape diagnosis. In fact *Brucella* has been cultured from the blood streams of persons with no symptoms of illness, who subsequently remained well. Overt disease may range from a brief illness of a few days to a severe febrile illness lasting for weeks or months with one or more relapses. The onset can be abrupt or insidious. Long term chronic illness may follow either type of onset. In acute cases there is a septicemia or an intermittent bacteremia. The organisms localize in the tissues particularly in the reticuloendothelial system and viscera and have been recovered from local lesions in practically every tissue of the body. In the tissues the organisms are found predominantly intracellularly which gives them protection from the action of antibodies and most drugs. Pathologically there is nonspecific hyperplasia of the reticuloendothelial tissue and focal necrosis, microabscesses and granulomas throughout the viscera. The granulomas are usually minute and consist primarily of collections of epithelioid cells and lymphocytes. However large abscesses and extensive granulomatous inflammation may occur. The formation of granulomas probably operates further to protect the organism from the defense mechanisms of the body and from administered drugs. In persons recovering from acute brucellosis or in those who never have had an acute phase, localized lesions may persist for years and produce chronic or recurrent ill health.

With localization numerous complications may occur. Perhaps the most common is osteomyelitis^{2,5}. The author has seen cases with this complication caused by all three species of *Brucella*. *Brucella* spondylitis with destruction of an intervertebral disc and adjacent vertebral bodies is severely painful and incapacitating. Osteomyelitis of any bone may occur; the spine, pelvis, ribs and long bones are the most frequent sites. Suppurative arthritis usually monoarticular may occur in any joint, usually the hip or knee. This may arise during the course of systemic infection or may be the presenting complaint without other evidence of illness. Other complications are orchitis, epididymitis, subacute endocarditis, cholecystitis, chronic pyelonephritis, hepatic pericholecystic and subdiaphragmatic abscesses and localized abscesses in any part of the body, acute or chronic meningoencephalitis, peripheral neuropathies with the auditory and optic nerves most commonly attacked, granulomatous lesions of the eye and a severe hepatitis leading to cirrhosis of the liver⁶. With local infection in the brain, organic brain damage results leaving a variety of sequelae. Evidence has been presented that long term chronic infection may be accompanied by organic brain disease and deterioration^{7,8}. Finally, neuropsychiatric disorders occur so commonly in brucellosis that they must be regarded as part of the manifestations of the disease.

The disease then presents many aspects ranging from mild or severe acute illness to recurrent or chronic disease persisting for as long as twenty five years. The multiple manifestations may require the practice of all the medical specialties in diagnosis and treatment.

Incidence

The incidence of brucellosis in man is difficult to determine. One index is the number of cases reported through official channels of the various state departments of health. In interpreting such data it must be borne in mind that cases are notified on the basis of a physician's diagnosis. How truly a reported incidence may represent the real incidence of any disease is dependent upon the criteria used for diagnosis as well as upon the completeness of reporting. With brucellosis more than most diseases there has been great variation in the criteria used for diagnosis from rigid proof by isolation of the organism from the patient to clinical diagnosis unsupported by laboratory tests. It is obvious that the reported figures by themselves without knowledge of the criteria used are most difficult to interpret.

In the two decades following the recognition that *Br. abortus* and *Br. suis* cause disease in man laboratory techniques were not sufficiently dependable to provide cultural proof of the diagnosis and nonstandard and undependable antigens for the agglutination test were in use as already mentioned. Of necessity the physician was forced to rely mainly on his clinical judgment with the result that nearly any vague illness came to be called brucellosis. In later years diagnostic criteria used by conservative physicians have become more rigid but even today many physicians arrive at a clinical diagnosis of brucellosis in spite of entirely negative laboratory tests.

Difficulties in diagnosis failure to suspect brucellosis and the occurrence of mild cases that go unrecognized all add to the unreliability of the reported statistics of the disease.

Nevertheless if one excepts salmonellosis deriving from animal sources brucellosis in man in the United States is still of higher reported incidence than any other disease of animals transmitted to man. FIGURE 1 presents the annual incidence derived from notified cases for the period 1921 to 1956*. There was a steady increase in the number of notified cases up to 1947 reaching a peak of 6321. Since that date the notified incidence has decreased sharply becoming currently a little more than 1000 cases per year. Since there was no really significant increase in brucellosis of domestic animals prior to 1947 it is presumed that the increasing number of reported cases during this period was due to a growing awareness of the disease by physicians rather than to a real increase in incidence. In recent years other factors have become important (1) the more widespread practice of pasteurization of milk (2) the introduction and indiscriminate use of the broad spectrum antibiotics and (3) the intensification of the control and eradication program for bovine brucellosis. Disease due to milk borne brucellosis is now practically limited to the rural areas. Some drop in incidence may be due to this factor. The broad-spectrum antibiotics are effective in suppressing clinical brucellosis. The indiscriminate use of these antibiotics has probably obscured the diagnosis of brucellosis in many instances. Such cases do not appear in our vital statistics. The drop in incidence of the disease in recent years may be more apparent than real.

In the last ten years great strides have been made in the control and eradication of bovine brucellosis. Programs in some states have been more active




FIGURE 1 Notified cases of brucellosis in the United States from 1927 to 1956 (84,154 cases)

and successful than in others. It might be expected that a corresponding relative decrease in the reported incidence of brucellosis in man would result in those states that have achieved the greatest decrease in incidence of bovine brucellosis. The annual incidence of brucellosis in man by region and by state for the years 1947 to 1956 is given in TABLE 1.

In 1947 brucellosis was reported from every state. Seventeen states reported more than 100 cases each and accounted for 78.5 per cent of all cases. These same 17 states in 1956 accounted for 67.7 per cent of all reported cases. In 1947 over one half of the total reported cases, 3332 in number or 52.7 per cent, were from the following 7 states: Iowa, Illinois, Texas, Wisconsin, Minnesota, Michigan, and California. In 1956 52.2 per cent of the total cases were from the following 7 states: Iowa, Illinois, Minnesota, Michigan, Kansas, Georgia, and Missouri. In 4 states there was an absolute increase in reported cases: Kentucky (5 to 22), New Mexico (3 to 4), North Dakota (3 to 23), and Wyoming (1 to 4).

The total number of reported cases for 1956 was 1080. This is 1.1 per cent of the total number reported for the peak year 1947 (6321 cases). The percentage of the total contributed by individual states showed an increase in 20 states and a decrease in 26 others.

TABLE 2 presents a comparison of the incidence and distribution of notified cases in 1947 and 1956 assuming a uniform rate of reduction for the country.

TABLE 1
ANNUAL INCIDENCE OF NOTIFIED CASES OF BRUCellosis IN THE UNITED STATES
FOR THE YEAR 1947 TO 1956

	1947	1948	1949	1950	1951	1952	1953	1954	1955	1956	1947 to 1956
New England	464	203	132	91	86	76	68	48	22	40	1 230
Maine	32	12	12	12	4	8	4	1	2	1	88
New Hampshire	24	8	3	3	1	0	0	1	1	1	42
Vermont	136	36	3	7	45	32	20	13	4	14	310
Massachusetts	76	39	31	22	10	18	15	13	9	15	249
Rhode Island	18	10	5	4	4	3	2	3	1	4	54
Connecticut	178	98	78	43	22	15	27	17	5	5	488
Middle Atlantic	470	397	296	263	180	135	76	66	42	24	1 894
New York	261	248	146	149	89	77	51	47	23	8	1 094
New Jersey	45	49	35	35	13	16	10	6	1	2	212
Pennsylvania	114	95	115	79	78	42	15	18	18	14	588
East North Central	1 515	1 295	1 127	803	694	505	444	452	287	247	7 364
Ohio	100	191	137	41	23	11	15	9	10	9	546
Indiana	115	70	44	42	19	10	27	13	9	8	357
Illinois	554	513	514	442	445	294	243	203	140	133	3 481
Michigan	302	219	193	93	64	52	68	111	60	57	1 222
Wisconsin	444	307	237	185	140	138	91	116	68	3	1 738
West North Central	1 765	1 103	1 098	1 109	1 150	1 016	831	666	660	407	9 490
Minnesota	386	307	355	283	189	139	133	151	116	65	2 119
Iowa	907	412	371	49	767	724	556	351	405	179	5 222
Missouri	141	82	115	80	67	45	44	54	39	40	707
North Dakota	6	5	29	38	20	7	13	15	20	73	176
South Dakota	80	67	46	10	32	73	23	21	49	33	334
Nebraska	95	89	46	15	10	3	3	32	8	16	317
Kansas	155	146	120	134	65	75	59	42	23	46	865
South Atlantic	419	307	392	280	272	168	163	119	92	117	2 419
Delaware	7	2	2	0	2	0	0	1	0	0	14
Maryland	52	57	45	44	39	12	13	6	0	7	276
District of Columbia	0	3	0	0	4	1	1	0	2	0	11
Virginia	74	52	73	64	81	47	58	46	30	35	593
West Virginia	14	8	7	8	13	7	5	6	9	3	80
North Carolina	21	16	25	21	28	21	5	3	2	11	153
South Carolina	48	22	31	9	11	6	8	3	2	5	145
Georgia	136	133	122	9	84	64	63	45	35	44	821
Florida	64	74	86	36	10	10	10	9	12	12	376
East South Central	313	226	204	174	183	136	115	106	88	73	1 618
Kentucky	15	21	23	19	17	12	11	15	27	22	177
Tennessee	92	10	39	51	44	41	35	38	34	29	473
Alabama	174	74	70	43	55	37	29	18	11	11	478
Mississippi	87	61	60	61	64	46	40	35	21	11	490

TABLE 1—Continued

	1947	1948	1949	1950	1951	1952	1953	1954	1955	1956	1947 & 1956
West South Central	640	778	545	456	279	241	207	213	137	107	3 553
Arkansas	47	44	30	34	37	32	47	40	26	30	367
Louisiana	44	37	30	32	48	23	18	49	29	28	338
Oklahoma	89	86	144	102	71	68	38	40	31	17	686
Texas	460	561	341	288	123	118	104	84	51	32	2 162
Mountain	415	401	254	132	124	80	48	84	44	29	1 611
Montana	6	4	13	18	9	9	8	7	6	1	81
Idaho	33	30	29	24	30	14	9	13	6	5	193
Wyoming	1	11	8	5	8	4	10	4	3	4	58
Colorado	268	249	99	55	39	20	8	17	12	5	772
New Mexico	3	6	8	1	0	3	2	12	3	4	42
Arizona	21	18	18	16	8	12	1	14	6	3	117
Utah	82	87	75	9	27	18	10	16	8	7	334
Nevada	1	1	4	4	3	0	0	1	0	0	14
Pacific	370	246	197	207	171	180	80	69	72	46	1 633
Washington	66	43	26	37	62	58	5	11	6	8	372
Oregon	20	39	45	45	9	7	7	10	10	3	195
California	284	164	176	120	100	115	68	48	56	35	1 116
Total	6 321.4	991.4	235.3	510.3	139.2	537.2	032.1	823.1	444.1	080	31 112

* Subject to slight change on final verification of cases

at large. The comparison indicates that the reduction in numbers of cases was not uniform for all of the states as evidenced by a calculated χ^2 of 518.93 and a theoretical χ^2 of 56.50 with 48 degrees of freedom at the probability level of 0.01. Of those states in which the expected numbers of cases are less than the actual reported numbers the deviations for Arkansas, Louisiana, Kentucky, and North Dakota appear significant.

Since the greatest decrease in incidence of bovine brucellosis has occurred in the last 5 years, the incidence and distribution of the total cases by states are compared with the data for the preceding 5 years in TABLE 3. Here again it is evident that the reduction in numbers of cases was not uniform as attested by a calculated χ^2 of 2088.47 and a theoretical χ^2 of 76.15 with 48 degrees of freedom at a probability level of 0.01. From this analysis of states reporting less than the expected number of cases the deviations may be significant for the following states: Wisconsin, Texas, New York, Pennsylvania, Connecticut, Indiana, Nebraska, Colorado, Utah, Ohio, Florida, Maryland, Oregon, New Jersey, South Carolina, and New Hampshire. Wisconsin, Connecticut, and New Hampshire are accredited under the brucellosis eradication program. Oregon, Pennsylvania, and Utah are expected to receive accreditation during 1958. Although the control and eradication program was organized in Texas in 1957 and an effective program is not yet operative, the relative decrease in

TABLE 2

A COMPARISON OF THE INCIDENCE AND DISTRIBUTION OF NOTIFIED CASES OF BRUCELLOSIS IN THE UNITED STATES IN 1941 AND 1956

State	1941	1956	Expected number of cases
	Number of cases	Number of cases	
1 Iowa	902	149	15
2 Illinois	554	133	95
3 Texas	460	32	19
4 Wisconsin	444	35	76
5 Minnesota	387	65	66
6 Michigan	307	51	52
7 California	284	35	49
8 Colorado	268	5	46
9 New York	261	8	45
10 Connecticut	178	5	30
11 Kansas	155	47	26
12 Missouri	141	40	24
13 Georgia	136	44	23
14 Vermont	136	14	23
15 Alabama	124	11	21
16 Indiana	115	8	20
17 Pennsylvania	114	14	19
18 Ohio	100	9	17
19 Nebraska	95	16	16
20 Tennessee	92	29	16
21 Oklahoma	88	11	15
22 Mississippi	82	11	14
23 Utah	82	7	14
24 South Dakota	80	33	14
25 Massachusetts	76	15	13
26 Virginia	74	35	13
27 Florida	67	12	11
28 Washington	66	8	11
29 South Carolina	48	5	8
30 Arkansas	47	30	8
31 New Jersey	45	2	8
32 Louisiana	44	28	7
33 Maryland	52	7	9
34 Idaho	33	5	5
35 Maine	32	1	5
36 New Hampshire	24	1	4
37 North Carolina	21	11	4
38 Arizona	21	3	4
39 Oregon	20	3	3
40 Rhode Island	18	4	3
41 Kentucky	15	27	3
42 West Virginia	14	3	3
43 Delaware	7	0	1
44 Montana	6	1	1
45 North Dakota	6	23	1
46 New Mexico	3	4	1
47 Wyoming	1	4	0
48 Nevada	1	0	0
49 District of Columbia	0	0	0
Totals	6 321	1 080	1 072

Based on the assumption of a uniform rate of reduction

TABLE 3

A COMPARISON OF THE INCIDENCE AND DISTRIBUTION OF NOTIFIED CASES OF BRUCELLOSIS FOR THE PERIODS 1947 TO 1951 AND 1952 TO 1956

State	1947 to 1951	1952 to 1956	
	Number of cases	Number of cases	Expected number of cases
1 Iowa	3 007	2 215	1 208
2 Illinois	2 468	1 013	991
3 Minnesota	1 515	604	609
4 Wisconsin	1 310	448	526
5 Texas	1 773	389	712
6 Michigan	874	348	351
7 California	794	322	319
8 Georgia	510	251	230
9 Kansas	670	245	250
10 Missouri	485	222	195
11 Virginia	377	216	151
12 New York	893	201	359
13 Oklahoma	497	194	198
14 Tennessee	296	177	119
15 Arkansas	192	175	77
16 Mississippi	337	153	135
17 South Dakota	235	149	94
18 Louisiana	191	141	77
19 Pennsylvania	481	107	193
20 Alabama	372	106	149
21 Washington	234	88	94
22 Vermont	227	83	91
23 Kentucky	95	82	38
24 North Dakota	98	78	39
25 Massachusetts	178	70	72
26 Connecticut	419	69	168
27 Indiana	290	61	116
28 Nebraska	255	62	102
29 Colorado	710	62	285
30 Utah	275	59	110
31 Ohio	492	54	198
32 Florida	273	53	110
33 Idaho	146	47	59
34 North Carolina	111	42	45
35 Maryland	238	38	96
36 Oregon	158	37	63
37 Arizona	81	36	33
38 New Jersey	177	35	71
39 Montana	50	31	20
40 West Virginia	50	30	20
41 Wyoming	33	25	13
42 New Mexico	18	24	7
43 South Carolina	121	24	49
44 Maine	72	17	22
45 Rhode Island	41	13	16
46 District of Columbia	7	4	3
47 New Hampshire	39	3	16
48 Nevada	13	1	5
49 Delaware	13	1	5
Totals	22 196	8 916	8 716

Based on the assumption of a uniform rate of reduction

notified cases in Texas is greater than in Wisconsin. For states reporting more than the expected numbers of cases the deviations may be significant in the following instances: Iowa, Virginia, Tennessee, Arkansas, South Dakota, North Dakota, Alabama, Louisiana, Kentucky, and New Mexico. No states in this list have received accreditation. New Mexico is expected to be accredited during 1958. The data for Iowa probably reflect the relative importance of swine brucellosis in the epidemiology of the human disease in that state. In Minnesota, which has been accredited, and Michigan, which is probably due for accreditation in 1958, the reported cases are approximately the numbers expected. States not previously mentioned that have received accreditation are North Carolina, Maine, Washington, Delaware, and Vermont; those probably soon to be accredited are Rhode Island and Montana.

As yet, no clear relationship is apparent between the reduction in incidence of notified cases of brucellosis and the degree of control and eradication achieved in the various states. Perhaps in the next several years a real effect will be noted. It should be mentioned that in states with the more effective programs greater numbers of infected cattle are moved to slaughter. This may result in a temporary increase in exposure of occupational groups handling these animals.

As already stated, the notified incidence of the disease may be entirely too low. Epidemiological studies offer another means of evaluating the incidence. *Brucella* infection usually results in specific dermal hypersensitivity to *Brucella* antigen. Repeated injection of *Brucella* antigen into control subjects does not produce this sensitivity^{10, 11} nor does the ingestion of dead organisms.^{1, 12} Presumably growth of the organism within the host is necessary for sensitization to occur. Persons exhibiting dermal hypersensitivity to *Brucella* antigen are regarded as having been infected at some time. From 10 to 25 per cent of the general population in the United States, varying in different areas of the country, exhibit specific allergy for *Brucella*; therefore they must be assumed at one time to have been infected with this agent. If one considers this incidence in the light of the notified incidence of brucellosis in man, it is difficult to reconcile the two. Probably many of these individuals remain asymptomatic and never develop clinical disease. Even so, one cannot avoid the conclusion that a great deal of clinical illness due to *Brucella* escapes recognition and does not appear in the reports of notified cases. In recognition of this discrepancy, there have been a number of attempts to estimate a more realistic incidence than that reported. Difficulties arise because of the duration of the disease. One estimate placed the number of persons in the United States suffering from *Brucella* infection during any particular year at between 40,000 and 4,000,000 (4000 cases were officially reported).¹⁴ A conservative estimate for the year 1949, based upon data derived from a few centers regarded as providing reliable information by isolation of the organism from a high proportion of patients and by employment of an adequate serologic test extrapolated on a national basis, placed the actual incidence at approximately 2½ times the notified incidence.¹⁵

Epidemiology

Brucellosis in man is mainly an endemic disease of sporadic occurrence. Man contracts the disease by consuming unpasteurized milk or other dairy products derived from infected animals or by direct contact with infected animals or their tissues. The only substantiated cases of transmission from human to human are from mother to nursing infant (M. K. Castaneda Hospital General Mexico D. F. Mexico personal communication).

While brucellosis has been described in many animal species, those of primary epidemiological importance for man are cattle, swine, goats, and sheep. *Br. abortus* is found primarily in cattle, *Br. suis* in swine, and *Br. melitensis* in goats and sheep. However, these species are not confined strictly to their respective main hosts. Cost-borne disease due to *Br. melitensis* has existed in the United States for many years in the states along the Mexican border.^{18, 19} There have been few other foci of infected goats, the most noteworthy in Colorado in 1946,¹⁹ where a considerable number of human cases occurred. The infected goats were destroyed and, at present, *Br. melitensis* infection in goats is believed to occur only in the southwestern states in low incidence. The disease in sheep has been nonexistent in the United States, perhaps because we do not produce milking sheep. The only important sources of the disease for man are cattle and swine. Human disease derived from infected cattle has been present throughout the entire country. Disease contracted from swine has been more limited in geographical extent and has been confined mainly to hog raising and hog slaughtering areas.

FIGURE 2 presents the total reported cases by states for the period 1927 to 1956. Six states accounted for the bulk of the reported cases. There is a marked correlation between the numbers of cases, the sites of large packing plants, and the distribution of farm animals.¹⁹ Since most milk now sold in trade is pasteurized, milk-borne brucellosis is largely confined to the rural areas. Brucellosis in the United States has become almost entirely an occupational disease,^{19, 20} of those who contact infected animals or their tissues, that is, farmers, dairymen, packing plant workers, butchers, veterinarians, and laboratory scientists. Epidemiological studies done in 1949 show that in 75 per cent of proved cases of brucellosis there was a history of direct contact with infected animals or their tissues.²¹ Only 25 per cent of cases were judged to be due to drinking unpasteurized milk. Spink²² states that only 10 per cent of his current cases fail to give a history of direct contact and are presumed to be due to the consumption of unpasteurized dairy products.

While brucellosis in cattle is due mainly to *Br. abortus*, the other species may infect this host and be transmitted to man. *Br. melitensis* has been isolated but seldom from cattle in the United States. *Br. suis* however has been found in infected cattle, and many of the reported milk-borne epidemics have been due to this species.^{23, 24} *Br. suis* infection does not seem to spread readily in cattle and, currently, bovine brucellosis due to this species is believed to be of little epidemiological importance.

With the reduction in milk-borne disease, swine have assumed relatively




FIGURE 2 Cumulative notified cases of brucellosis in the United States from 1977 to 1986

greater importance in the epidemiology of human brucellosis. While most clinical disease in swine is attributable to *Br suis*, both of the other species infect these animals. In 1946 *Br melitensis* was found infecting hogs in Iowa and cases of human brucellosis due to this species were attributed to this source.¹⁷ There have been continued reports of cases from Iowa, Minnesota, and elsewhere. The clinical disease in swine resembles that caused by *Br suis*. In culturing the mandibular lymph glands of 5000 marketed hogs, we isolated both *Br melitensis* and *Br abortus* as frequently as *Br suis*.²⁰⁻²² This was the first report of the recovery of *Br abortus* from naturally infected hogs. Meat inspectors who worked exclusively on the hog kill and who developed brucellosis were found to be infected with *Br abortus* and they presumably contracted this infection from hogs. Similarly human cases of *Br melitensis* infection were found in persons working only on the hog kill. Spink also reports epidemiological evidence incriminating the hog as a source of *Br abortus* infection.

of man³⁰ The infection in swine due to *Br abortus* seems limited in clinical manifestations and in infectivity for other swine However epidemiologically it is of importance since persons who come in contact with the tissues of infected swine may develop infection caused by this species

While the control and eradication program for bovine brucellosis has brought about a marked reduction in the number of infected cattle in the country and while further reduction is anticipated it should be realized that even when all the states receive accreditation some infection in cattle will still exist and occupational groups will still be exposed Many factors seem to be involved in determining whether or not man contracts this disease Perhaps the host factors are the most important Even though the number of infected animals may be less persons working in exposed environments will still come in contact with such animals and their tissues repeatedly those individuals who are susceptible may be offered adequate exposure to contract the disease It is anticipated that brucellosis in man due to *Br abortus* contracted from cattle will decline to a low figure and then remain static Furthermore there has been no serious effort to eradicate swine brucellosis and this source of infection for man continues to be of importance At present we cannot foresee the day when brucellosis in man will cease to occur in the United States

Diagnosis

In view of the vagaries of the organism and of the disease the diagnosis of brucellosis in man can be most difficult In the acute disease the symptoms resemble those of many other acute infections in the chronic phase they may be multiple vague and varied There are no pathognomonic signs The clinician cannot make a clinical diagnosis with any degree of assurance a definitive diagnosis is dependent upon the results of laboratory tests Factors relative to culture of the organism and application of the agglutination test have already been discussed With the development of better culture techniques and an increased understanding of the organism and of the nature and course of the infection in man sound principles of diagnosis have emerged^{31 32} These have been well summarized by the Committee on Brucellosis of the National Research Council Washington D C In the second report of this committee Diagnostic Criteria for Human Brucellosis³² it was recognized that the only proof of the diagnosis lies in the isolation of the organism from the patient and that in most acute cases of the disease this is possible In the absence of cultural proof it was considered that an agglutination titer of 1:320 or higher or a rising titer offered presumptive confirmation of a clinical diagnosis It was further recognized that tests to determine dermal sensitivity to *Brucella* antigen provide no information useful to the clinician in diagnosis and it was recommended that they should not be used for this purpose

Since bacteremia is often intermittent success in isolating *Brucella* from human blood may necessitate a series of cultures The organism often may be isolated from bone marrow or lymph nodes removed for culture

When attempts at culture fail to yield the organism dependence must be placed on the results of an agglutination test Since the advent of the broad

spectrum antibiotics many patients have been treated with these drugs prior to the consideration of the possibility of brucellosis or before referral to a diagnostic center. The disease is partially suppressed and the organism less readily isolated from the blood. the agglutination test assumes increasing importance in diagnosis.

If dependence must be placed on a serologic test it is evident that a standard antigen and a standard technique for performing the test are most necessary to provide accuracy and uniformity in diagnosis. The need for and choice of a standard antigen were considered by the National Research Council Committee and these deliberations led to Report No. 3, A Standardized Antigen and Agglutination Technic for Human Brucellosis.²⁴ An antigen made from *Br. abortus* Strain 1119 prepared according to the method used at the Animal Disease Station, Agricultural Research Service, United States Department of Agriculture, Beltsville, Md. for preparation of the antigen used in veterinary medicine was recommended as the standard. A tube test with incubation at 37° C. for 48 hours was specified. The technique and reading of the test were described in detail.

The development of these criteria for diagnosis and of a standard agglutination test are important accomplishments and should promote greater accuracy and more uniformity in the diagnosis and reporting of brucellosis.

The greatest remaining problem in diagnosis is posed by chronic brucellosis. There are two situations that are very difficult: one presented by the patient who first comes to medical attention during the chronic stage of the disease when laboratory tests are less helpful; the other by the patient with a past history of proved brucellosis in whom all signs of the disease have cleared but who continues to have subjective complaints for months or years.

Therapy

The principles of therapy in brucellosis have emerged slowly and painfully. Prior to the days of antibiotics dozens of remedies were enthusiastically proposed and later found wanting. Among these were various vaccines that are still widely used although convincing evidence of their usefulness has not been furnished. The specific therapy of the disease dates from 1946 with the introduction of a combination of streptomycin and sulfadiazine.²⁵⁻²⁷ While this combination was effective and constituted a marked advance in therapy it proved fairly toxic²⁸ and the relapse rate was considerable. This treatment fell into disrepute with the introduction of the broad spectrum antibiotics. These drugs were found to suppress the clinical manifestations of the disease but the relapse rate was high. A combination of dihydrostreptomycin and chlortetracycline²⁹⁻³⁰ has proved effective with or without the addition of sulfadiazine or triple sulfonamide.³¹⁻³³ I prefer this latter combination. In growth inhibition, bactericidal and manometric studies these drugs have been shown to act synergistically.³⁴ They are highly effective against *Brucella* *in vitro* but when the organisms are contained inside cell their action becomes mainly suppressive rather than bactericidal.³⁵ Any of the broad-spectrum antibiotics may replace chlortetracycline in this combination.

of man³⁰ The infection in swine due to *Br abortus* seems limited in clinical manifestations and in infectivity for other swine However, epidemiologically it is of importance since persons who come in contact with the tissues of infected swine may develop infection caused by this species

While the control and eradication program for bovine brucellosis has brought about a marked reduction in the number of infected cattle in the country and while further reduction is anticipated, it should be realized that even when all the states receive accreditation some infection in cattle will still exist and occupational groups will still be exposed Many factors seem to be involved in determining whether or not man contracts this disease Perhaps the host factors are the most important Even though the number of infected animals may be less persons working in exposed environments will still come in contact with such animals and their tissues repeatedly, those individuals who are susceptible may be offered adequate exposure to contract the disease It is anticipated that brucellosis in man due to *Br abortus* contracted from cattle will decline to a low figure and then remain static Furthermore there has been no serious effort to eradicate swine brucellosis and this source of infection for man continues to be of importance At present we cannot foresee the day when brucellosis in man will cease to occur in the United States

Diagnosis

In view of the vagaries of the organism and of the disease the diagnosis of brucellosis in man can be most difficult In the acute disease the symptoms resemble those of many other acute infections in the chronic phase they may be multiple vague and varied There are no pathognomonic signs The clinician cannot make a clinical diagnosis with any degree of assurance a definitive diagnosis is dependent upon the results of laboratory tests Factors relative to culture of the organism and application of the agglutination test have already been discussed With the development of better culture techniques and an increased understanding of the organism and of the nature and course of the infection in man sound principles of diagnosis have emerged³¹ These have been well summarized by the Committee on Brucellosis of the National Research Council Washington D C In the second report of this committee Diagnostic Criteria for Human Brucellosis³² it was recognized that the only proof of the diagnosis lies in the isolation of the organism from the patient and that in most acute cases of the disease this is possible In the absence of cultural proof it was considered that an agglutination titer of 1:320 or higher or a rising titer offered presumptive confirmation of a clinical diagnosis It was further recognized that tests to determine dermal sensitivity to *Brucella* antigen provide no information useful to the clinician in diagnosis and it was recommended that they should not be used for this purpose

Since bacteremia is often intermittent success in isolating *Brucella* from human blood may necessitate a series of cultures The organism often may be isolated from bone marrow or lymph nodes removed for culture

When attempts at culture fail to yield the organism dependence must be placed on the results of an agglutination test Since the advent of the broad

tion of human beings with a mixed heat killed vaccine of *B. typhosus*, *B. abortus* and *B. melitensis*. J Infectious Diseases 67:258

- 11 McCULLOUGH N B Unpublished data
- 12 McCULLOUGH N B & C W EISELE & C A BEAL 1949 Oral administration of killed *B. ucella* to man. Public Health Repts U S 64:1613
- 13 BRADFORD A I D GILB & D ANDERSON 1949 Formation of antibodies in human subjects after the ingestion of heat killed *B. ucella abortus*. J Lab Clin Med 34:44
- 14 EVANS A C 1947 Brucellosis in the United States. Am J Public Health 37:139
- 15 STIFFLE J H & L O FINK 1940 Brucellosis incidence in the United States. In Third Inter American Congress on Brucellosis. Washington D C. Pan Am San Bur 28
- 16 CENTRY E R & T L FERENBAUGH 1911 Femic Malta (Mediterranean) fever in Texas with isolation of the *Micrococcus melitensis* from two patients. J Am Med Assoc 67:889
- 17 LAKE G C 1922 Malta fever in southwestern United States with special reference to a recent outbreak in Phoenix Ariz. Public Health Repts U S 37:2895
- 18 STILES G W 1950 Controlling brucellosis in Colorado goats. In Third Inter American Congress on Brucellosis. Washington D C. Pan Am San Bur 276
- 19 JORDAN C F 1950 The epidemiology of brucellosis. In Brucellosis a Symposium. Washington D C. Am Assoc Advance Sci 98
- 20 McCULLOUGH N B 1950 Brucellosis—a packing plant problem. In Third Inter American Congress on Brucellosis. Washington D C. Pan Am San Bur 16
- 21 MAGOFFIN R L P KASLER W W SPINK & D S FLEMING 1949 An epidemiologic study of brucellosis in Minnesota. Public Health Repts U S 64:1021
- 22 SPINK W W 1956 The Nature of Brucellosis. 109 Univ Minn Press Minneapolis Minn.
- 23 BEATTIE C P & K M RICE 1934 Endulant fever due to *B. ucella* of the porcine type—*B. ucella suis*. J Am Med Assoc 102:160
- 24 HORNING B G 1935 Outbreak of undulant fever due to *B. ucella suis*. J Am Med Assoc 105:1918
- 25 BORTS I H D M HARRIS M F JOYNT J R JENNINGS & C F JORDAN 1943 A milk borne epidemic of brucellosis. J Am Med Assoc 121:319
- 26 BORTS I H 1947 Transmission of Brucella in milk. J Am Med Assoc 134:394
- 27 BORTS I H S H MCNITT & C F JORDAN 1946 *B. ucella melitensis* isolated from swine tissues in Iowa. J Am Med Assoc 130:966
- 28 McCULLOUGH N B C W EISELE & E PAVELCHIK 1949 Isolation of *B. ucella abortus* from hogs. Public Health Repts U S 64:537
- 29 McCULLOUGH N B C W EISELE & E PAVELCHIK 1951 Survey of brucellosis in slaughtered hogs. Public Health Repts U S 66:205
- 30 SPINK W W 1956 The Nature of Brucellosis. 101 Univ Minn Press Minneapolis Minn.
- 31 McCULLOUGH N B 1949 Laboratory tests in the diagnosis of brucellosis. Am J Public Health 39:866
- 32 McCULLOUGH N B 1950 Laboratory tests in brucellosis. In Brucellosis a Symposium. Washington D C. Am Assoc Advance Sci 116
- 33 SPINK W W N B McCULLOUGH L M HUTCHINGS & C K MINGLE 1952 Diagnostic criteria for human brucellosis. J Am Med Assoc 149:805
- 34 SPINK W W N B McCULLOUGH L M HUTCHINGS & C K MINGLE 1954 A standardized antigen and agglutination technique for human brucellosis. Am J Clin Pathol 24:496
- 35 PULSART E J & W H AMPECHER 1947 Streptomycin therapy in brucellosis. Bull U S Army Med Dept 7:271
- 36 EISELE C W & N B McCULLOUGH 1947 Combined streptomycin and sulfadiazine treatment in brucellosis. J Am Med Assoc 135:1053
- 37 SPINK W W W H HALL J B SHAFER & A I BRADY 1948 Human brucellosis its specific treatment with a combination of streptomycin and sulfadiazine. J Am Med Assoc 136:387
- 38 McCULLOUGH N B & C W EISELE 1949 Toxic reactions during streptomycin sulfadiazine therapy of brucellosis. J Am Med Assoc 139:80
- 39 HERRELL W E & T E BARBER 1950 A new method for the treatment of brucellosis. J Am Med Assoc 144:519
- 40 HERRELL W F & T E BARBER 1952 Treatment of brucellosis with Aureomycin or Terramycin combined with dihydrostreptomycin. Postgrad Med 11:46

Since 1949 I have used dihydrostreptomycin (2 gm per day) chlortetracycline or currently tetracycline (2 gm per day) and sulfadiazine or triple sulfonamide in a dosage to yield a blood level of 10 mg per cent in the treatment of proved cases of brucellosis. No case has relapsed or failed to recover from the disease. In the uncomplicated case treatment is continued for 3 weeks and in chronic cases or in those with complications, for a longer period.

In formulating a logical approach to the therapy of brucellosis it should be realized that in most patients it is a self limiting disease with spontaneous recovery occurring within a few weeks or months. Evaluation of the effectiveness of therapy can be most difficult. The more important principles of treatment may be summarized as follows:

- (1) There should be prompt and accurate diagnosis
- (2) Bed rest is always indicated, many patients will recover on bed rest alone
- (3) The physician should search for local lesions that may necessitate longer medical treatment or definitive surgical treatment, in any case of more than a few weeks duration a Roentgen survey of the skeleton should be made to detect osteomyelitic lesions
- (4) Appropriate specific antibiotic therapy should be employed
- (5) As the best current therapy is mainly suppressive treatment should be continued for a sufficient period to allow the defenses of the body to overcome the infection
- (6) The patient should have graded activity and should undergo careful observation during convalescence to avoid relapses
- (7) Sympathetic and intelligent attention should be given to the psychiatric aspects of the disease since many patients develop depressions suicidal tendencies and even psychoses
- (8) There should be adequate diet and supportive measures as indicated
- (9) The physician should have a strong reluctance to make a diagnosis of psychoneurosis when the possibility of brucellosis exists

References

- 1 FISELE C W N B McCULLOUGH & G A BEAL 1947 Discrepancies in the agglutination test for brucellosis as performed with various antigens and as reported from different laboratories J Lab Clin Med 32 847
- 2 CRIGGS J F & L W CASE 1948 Variations in *Brucella* agglutination reactions in different laboratories Am J Clin Pathol 18 506
- 3 LOWE G H JR & P R LIPSCOMB 1947 Brucellosis osteomyelitis Surgery 22 524
- 4 LOWMEER L 1949 Brucellic osteomyelitis of man and animal Proc Staff Meeting Hillcrest Mem Hosp Tulsa Okla 6 1
- 5 SPINK W W 1956 The Nature of Brucellosis 180 Univ Minn Press Minneapolis Minn
- 6 McCULLOUGH N B & C W FISELE 1951 *Brucella* hepatitis leading to cirrhosis of the liver Arch Internal Med 88 193
- 7 APTER N S W C HALSTEAD C W FISELE & N B McCULLOUGH 1948 Impaired cerebral functions in chronic brucellosis Am J Psychiat 105 361
- 8 APTER N S C W FISELE N B McCULLOUGH & W C HALSTEAD 1948 Cerebral pathology in brucellosis Trans Am Neurol Assoc 73 3
- 9 PUBLIC HEALTH SERVICE NATIONAL OFFICE OF VITAL STATISTICS 1927 1956 Reported Incidence of Selected Notifiable Diseases
- 10 KOLMER J A A BONDI JR & A M RUFF 1940 The results of active immuniza-

BRUCELLOSIS IN LIVESTOCK CONTROL AND ERADICATION

By C. K. Mingle

Agricultural Research Service United States Department of Agriculture Washington D. C.

Brucellosis of livestock has been recognized as a serious problem in the United States for almost a half century. However, it was not until 1934 that organized efforts were made to combat the disease in cattle. In that year the co-operative state federal bovine brucellosis eradication program was launched as part of a cattle reduction plan that was brought about as a result of drought conditions in many sections. In the beginning it was estimated that approximately 10 per cent of the adult cattle in the United States were affected with brucellosis.

In the years prior to 1941 the program was greatly retarded by a lack of uniformity in the procedures employed in different parts of the country. As a means of correcting this deficiency, the former Bureau of Animal Industry, Washington, D. C., took steps to encourage the establishment of uniform control and eradication practices. In time sufficient interest was developed along these lines to bring about the adoption of recommended uniform bovine brucellosis eradication procedures at the December 1941 meeting of the United States Livestock Sanitary Association held in Omaha, Nebr. These recommendations were designed to provide the flexibility necessary for handling brucellosis problems under varying herd conditions and still maintain a standard level of uniformity. The resulting Uniform Methods and Rules have contributed much to the progress achieved in recent years. Basically, the current program is built around 4 control and eradication plans: Plan A, test and slaughter with or without calf vaccination; Plan B, test calf vaccination, temporary retention of reactors; Plan C, calf vaccination without test of any part of the herd; and Plan D, adult vaccination.

Plans B, C, and D are considered temporary procedures to be used only in herds where heavy infection exists until Plan A can be put into operation without serious economic shock. Plan A, on the other hand, has eradication as its immediate goal and is the method of initial choice wherever possible.

The Uniform Methods and Rules also provide for the establishment and maintenance of Modified Certified Areas in which the animal and herd infection rates do not exceed 1 per cent and 5 per cent, respectively. It has been shown conclusively that the increased protection afforded by operating a program of this type on an area basis is essential in maintaining the gains made through field operations.

Bovine Brucellosis Eradication—Progress from 1934 to 1954

During the first 20 years of the bovine brucellosis eradication program a total of nearly 11 million herd-representing approximately 127 million cattle—were blood tested for brucellosis. The indicated animal infection rate over this same period was reduced from about 10 per cent to 2.6 per cent. In the fiscal year 1954 infection was disclosed in 14.2 per cent of the herds tested.

- 41 FISELE C W & N B McCULLOUGH 1950 Discussion of Herrell W E and T E Barber ²⁹
- 42 FISELE C W & N B McCULLOUGH 1950 Treatment of brucellosis (undulant fever) *In Current Therapy* 19 0 5 Saunders Philadelphia Pa
- 43 McCULLOUGH N B 1955 Brucellosis in children *Pediat Clin N Am* **2** 73
- 44 McCULLOUGH N B & C A BEAL 1957 Antimetabolic action of sulfadiazine and certain antibiotics for *Brucella* *J Infectious Diseases* **90** 196
- 45 MAGOFFIN R I & W W SPINK 1951 The protection of intracellular *Brucella* against streptomycin alone and in combination with other antibiotics *J Lab Clin Med* **37** 924




FIGURE 3. Official calf vaccinations. Symbol \bar{M} = million. Reproduced by permission of the Agricultural Research Service.

ties carried out during the first three fiscal years of the accelerated program (1955 through 1957)

Blood testing Over this period a total of 46.8 million cattle in 3.3 million herds were blood tested for brucellosis. Of this number 2.2 per cent of the cattle and 12.8 per cent of the herds tested showed evidence of infection. In comparison with the preceding 3-year period this represents an increase of 63.2 per cent in the number of herds tested, a 92.4 per cent increase in cattle tested, and a reduction of 3 per cent and 1.2 per cent in the respective infection rates.

Brucellosis ring testing Remarkable increases also have occurred in milk and cream ring testing during the past 3 years, and the reduced percentages of ring-suspected herds disclosed are significant of the progress being made. Over this period a total of 4.7 million herds were tested, of which 15.6 per cent were classed as suspicious. These figures represent an increase of 133.1 per cent in the number of herds tested and a reduction of 11.4 per cent in the number of suspicious herds disclosed as compared with the preceding 3-year period. Although the brucellosis ring test has recognized imperfections, there is increasing evidence that it is proving to be one of the most valuable tools we possess for combating brucellosis in dairy sections of the country.

Vaccination In common with testing operations, vaccination has increased also each year during the accelerated phase of the program. For the fiscal years 1955, 1956, and 1957 a total of 14.6 million official vaccinations was recorded. This compares with 10.8 million reported for the preceding 3 years and represents an increase of 34.9 per cent. It has been interesting to note the extremely low incidence of infection detected in areas where vaccination has been practiced on a wide scale over the past few years. Many vaccinated areas in which high infection rates were known to have existed now show surprisingly low levels of infection.

Area certifications Strong encouragement has been given in the accelerated campaign to the development of Modified Certified Brucellosis-Free Areas. As explained before, this designation signifies that the infection rates in such

Although Strain 19 vaccination was not officially adopted until 1941 there were more than 21 million calves vaccinated during the following 14 years. The use of vaccine has increased consistently each year since it became an integral part of the program. While it is impossible to evaluate accurately the benefits derived from vaccination there is increasing evidence that it has played a very important role in reducing the previously high incidence of infection in certain areas of this country. The importance of providing a serviceable degree of protection in otherwise susceptible cattle will continue until complete eradication of the disease is accomplished.

The Accelerated Bovine Brucellosis Eradication Program

Through additional funds made available by the Eighty third Congress it was possible in 1954 to expand the cooperative state federal campaign to eradicate brucellosis from cattle. All phases of the program were vastly increased over the previous levels. FIGURES 1, 2 and 3 graphically illustrate the activity




FIGURE 1 Bovine brucellosis Official blood serum agglutination tests Symbol \bar{M} = million Reproduced by permission of the Agricultural Research Service




FIGURE 2 Official brucellosis ring tests Symbol \bar{M} = million Reproduced by permission of the Agricultural Research Service




FIGURE 5 Brucellosis eradication program area certification status as of June 30, 1957. Reproduced by permission of the Agricultural Research Service.




FIGURE 6 Brucellosis eradication program certified and area work counties in the United States, Puerto Rico, and the Virgin Islands as of June 30, 1957. Reproduced by permission of the Agricultural Research Service.

Many states have set 1960 as the goal for certification. Although this objective will not be attained in all cases, it does not appear nearly as visionary as when that date was designated a few years ago. The enthusiasm and support for the eradication of brucellosis on the part of the livestock industry has never been greater than at present. If the momentum generated over the past few years can be maintained by adequate financing for another five or six years all of the United States should be certified by that time.

A progress toward certification is accelerated additional attention must be given to other known reservoirs of infection, namely swine, goats and sheep. At present we are considering plans that it is hoped will lead to pilot field




FIGURE 4 Brucellosis eradication program area certification status as of September 30 1954. Reproduced by permission of the Agricultural Research Service.

areas do not exceed 1 per cent for animals and 5 per cent for herds. FIGURES 4, 5, and 6 depict the advances made during the past 3 years in qualifying such areas. As illustrated in the FIGURE 4 map, there was a total of only 334 counties, including 3 entire states, recognized as Modified Certified Areas at the time the accelerated program became operative on October 1, 1954. In contrast, FIGURE 5 shows a total of 735 counties, including 7 complete states, qualified as certified areas on June 20, 1957. It also depicts an additional 712 counties operating on a complete area basis leading directly toward early certification. During the past fiscal year, new initial county certifications have been recorded at the rate of about 20 each month. As indicated on FIGURE 6, there were 735 certified counties and 112 other counties conducting complete area work as of June 30, 1957. Thus, of a total of 3150 counties in the United States, Puerto Rico, and the Virgin Islands, 46 per cent were either currently certified or rapidly approaching that status.

Future Outlook

Since the time data were assembled for the fiscal year 1957, the entire states of Vermont and Connecticut and 13 counties elsewhere have been added to the list of certified areas. This makes a total of 808 counties, including 9 complete states, so qualified as of September 1, 1957. It is expected that at least 800 additional counties, including 5 more states, will achieve certification during the current fiscal year. This would result in a total of 14 states and about 50 per cent of all domestic and territorial counties being certified by June 30, 1958. It is anticipated also that by the same date most if not all states will be conducting complete area programs designed to achieve early certifications.

all interested groups By maintaining the pressures developed over the past few years through adequate financing from state and federal sources we should reduce the incidence of bovine brucellosis throughout the United States to 1 per cent or less within the next five years and the brucellosis threat from this source eventually should be eliminated

Discussion of the Paper

OSCAR SUSSMAN (*Bureau of Veterinary Public Health State of New Jersey Department of Health Trenton N J*) C K Mingle in his use of the editorial we undoubtedly meant to include the public health authorities local state and national who were of considerable help in the preparation of the effective control of brucellosis or even of its eradication He indicated that it was gratifying to him to note that in the last few years *Brucella* infected cattle were moving out of the farm herds and into the slaughterhouses where they could no longer infect other cattle

In this connection it is worthy of mention that the action of the Board of Health of Chicago Ill gave considerable incentive to the State of Wisconsin in its determination to do an effective job in its attempt to eliminate brucellosis Several years ago the health authorities of Chicago indicated that they would not continue the purchase of milk from areas in which the cattle were infected with this disease even if the milk were pasteurized Subsequently similar action was taken by the public health authorities of several other areas including the State of New Jersey Department of Health which set a deadline beyond which the importation of milk from such areas would be prohibited thereafter no milk could be sold for human consumption except from animals that were known to be free of brucellosis

Regulations statutes and codes of this type have not been and could not be introduced and enforced by the agricultural authorities until such time as the public health authorities gave them the needed administrative support and created an economic incentive to the dairy farmer to cooperate if he wished to sell his product

Mingle has indicated that the agricultural authorities were unable to move *Brucella* reacting cattle off the farms after they had been identified by branding However of recent years these cattle have been moving to the slaughterhouses as already noted I submit that this exodus from the farms was initiated and sustained primarily by the public health authorities This experience repeats the history of the milk control laws that gave impetus to the tuberculosis-eradication campaign I am quite certain that Mingle and others in departments of agriculture realize that the cooperative efforts of the state and local health departments have resulted in considerable aid to farmers and to the national economy through health regulations that implement the desires of the informed segment of the agricultural interests

My remarks are made not in disagreement with Mingle but merely as additional evidence for the reader that there exists a considerable amount of valuable cooperative enterprise between the public health and agricultural authorities in their mutual concern for the control of animal diseases as they relate to human health

studies designed to determine the adequacy of existing recommended procedures for eradicating *Brucella suis* infection from swine on an area basis

Based on the available knowledge of *Brucella suis* and *Brucella melitensis* infections as they exist in the United States the control and eradication of these types of infection should not be too difficult to achieve Records assembled by the Agricultural Research Service over the past 6 years indicate the following probable incidence of brucellosis in swine and goats Of 501 160 swine tested in 57 666 herds 26 425 (or 5.3 per cent) were classified as reactors In the case of goats 1.8 per cent of 306 696 animals tested in 13 273 herds were likewise recorded as reactors Inasmuch as these data were accumulated from voluntary testing and represent a significant number of problem herds it is probable that the indicated infection rates are higher than would be true for the United States as a whole

It is important, of course to recognize the fact that certification alone does not represent complete eradication Obviously the last vestiges of infection will be the most difficult to eliminate However with continued effort and vigilance there is no apparent reason to believe that the goal of final eradication cannot be attained in time The resulting economic and public health benefits will fully justify and repay the effort and expense involved

Summary

Over the past 23 years the Cooperative State Federal Brucellosis Program has been responsible for reducing the incidence of bovine brucellosis in the United States from approximately 10 per cent in 1935 to 1.8 per cent in 1957

The first three years of the accelerated brucellosis eradication campaign undertaken in 1954 has resulted in more substantial progress than was accomplished in any similar period in the past

During the 3 year period ending June 30 1957 a total of 46.8 million cattle in 3.3 million herds were officially tested for brucellosis Of these 1.8 per cent of the cattle and 10.6 per cent of the herds showed evidence of infection This compares with 2.6 per cent animal and 14.2 per cent herd infection disclosed during the year preceding the inauguration of the expanded program

Extensive field experience has shown conclusively that the intelligent use of a standardized blood serum agglutination test the brucellosis ring test and Strain 19 vaccine can be employed effectively to combat bovine brucellosis on an area basis

The establishment of Modified Certified Brucellosis Free Areas is progressing at a faster rate than expected As of September 1 1957 there were 808 certified counties and 800 other counties that were actively working toward early qualification New county certifications are being recorded at an average rate of 20 per month

Although the certification of areas is an essential part of the program it is important to bear in mind that this designation is only an advanced step toward complete eradication With the recognized economic and public health benefits to be derived from this status we should accept nothing less

The bovine brucellosis eradication campaign now has strong support from

apply to the larval or adult stages or to both or to neither. Thus liver flukes have a strict larval specificity but little adult specificity. *H. schrefferi bancrofti* on the other hand has a strict adult specificity but a low larval one occurring in a great many mosquitoes. The human malarial parasites are specific to man and to anopheline mosquitoes. The trichina worm has little if any host specificity among the mammals.

Parasites in general are least likely to be acquired from animals that are very distantly related to man phylogenetically or that are physiologically quite different from him. He has acquired few parasites from herbivores and none from horses but he shares many with both dogs and pigs. Also it is obvious that the closer the physical association of an animal with man the better will be the opportunity of infection by a suitable mutant. In general domestic animals are more important potential sources of parasites than are wild animals.

The specialized external environmental requirements of parasites determine their geographical distribution. Some cannot exist in certain climates because of unsuitable temperatures or humidity or because of lack of a suitable intermediate host. Moreover some cannot gain entrance to hosts otherwise suitable because the habits of the host do not expose it to infection. Thus for example the custom of cooking food prevents the infection of man with a great many species of parasites. Herbivores fail to become naturally infected with the trichina worm only because they do not eat meat just as man does not often become infected with the fluke *Fasciola hepatica* because he does not eat grass.

Man is a recently evolved animal who in modern times has been recorded as host for about 200 species of parasites. A great many of these are of purely academic interest some are plain errors of identification but all have been or are being acquired from other animals. A few were acquired from prehumans the malarial parasites the pinworms and the lice all came from man's ancestors. Some such as *Bertiella* a common tapeworm of primates that is still occasionally recorded in humans and is carried by certain species of mites man may well have lost as a regular parasite after he gave up a purely herbivorous diet and took to an omnivorous one that is at about the same time that he acquired his *Taenias*. *Physaloptera*, *Terndens* and *Oesophagostomum apistomum* all common nematodes parasitizing monkeys in Africa are also in the process of being lost by man. Some parasites were acquired quite early in human history from dogs swine and other animals that man domesticated others were acquired directly or indirectly in the past from wild animals and still others man continues to acquire from both of these sources.

Protozoa

The relationship between the intestinal protozoa of man and those of other animals is still very obscure. The chief potential pathogen of man is *Entamoeba histolytica* a form that is usually nonpathogenic living an apparently saprozoic life in the large bowel. Under circumstances that are still far from clear this parasite enters the mucosa and becomes extremely pathogenic. Under

PARASITES OF ANIMALS AND HUMAN DISEASE

By Thomas W. M. Cameron

*McGill University, Montreal, P. Q. and the Institute of Parasitology,
Macdonald College, P. Q., Canada*

Parasitism is an extremely ancient mode of life and we have evidence of it from as long ago as the Devonian period. It is important to realize, however, that parasitism has no necessary connection with disease. In fact, if a species of parasite is to continue to exist, it is essential that it should not produce severe pathogenic effects in its host—at least not until sufficient opportunity has been given to the next generation of the parasite to gain a foothold in another host. When disease is produced, it is generally accidental, resulting from superparasitism or from some abnormality in the normal life cycle of the host or of the parasite.

Superparasitism is the result of the presence of too many parasites in a single host and it occurs when the parasite can reproduce itself without serious inhibition until overwhelming numbers prove too much for the defense mechanisms of the host. This is seen particularly among the protozoan parasites such as the hemoflagellates or *falciparum* malarial parasites.

This condition is seen among the helminths under quite different circumstances. The worms cannot multiply within a single host and their progeny must escape to the exterior, there to undergo some essential development before they can re-enter the host. Consequently, superparasitism and parasitic disease depend on the numbers of parasites that enter the body. Occasionally, as in trichinosis, a large number can do so simultaneously and cause an explosive, often fatal result. As a general rule, however, the accumulation of large numbers of parasites in a host is the consequence of continued exposure to infection. There is a very important and distinct difference between helminthic infection and helminthic disease.

Disease is not always due to excessive numbers of parasites; sometimes some abnormality in the life cycle causes serious symptoms. Thus the migrations of a single ascarid may occlude an important duct. Similarly, the larval stage of a parasite in an abnormal host may be able to survive but not develop and, in surviving, may produce serious lesions. In general, larval stages tend to have a greater pathogenicity than do the corresponding adult stages and this is particularly the case when the parasites are in an abnormal host.

Parasites not only have existed for a long time but they have evolved along with their hosts and, in evolving, they have become more and more specialized in their environmental requirements, internal as well as external.

Their specialized internal environmental requirements have led to a considerable degree of host specificity that varies in type and degree among the different groups of parasites. Some are so host specific that they can live only in a single species of host animal; others can exist in a group of closely related animals or in a group of animals having some common physiological characteristics; and still others have little host specificity. This specificity may

America for example *Metorchis conjunctus* is a common parasite of the liver of fish eating mammals and when it occurs in numbers it causes a serious morbidity and may cause death. It can and occasionally does occur in man but as its second intermediate host is the Common Sucker a fish rarely eaten by man it is not an important human parasite even though the fish is common and often heavily infected in the northeastern part of North America.

The blood flukes of man now collectively form one of the most important causes of human disease and it is estimated that they infect about one fifth of the total population of the world. The Far Eastern species has been known for many years to have numerous animal hosts and although I was able to show that *S. mansoni* in the West Indies had a sylvatic host in the naturalized African green monkeys there (Cameron 1928) it was only recently demonstrated that both African species occur in a wide variety of native animals (Kuntz 1935). Man however is still the main source of his own infection both in Africa and in South America. Blood flukes have existed as invaders of man for centuries but there is little doubt that irrigation has done much to increase their incidence in modern times.

Blood flukes capable of developing to sexual maturity in man are absent from North America but blood flukes of birds and rodents which have a developmental cycle similar to that of the common human pathogens have larvae capable of penetrating human skin that it reaches by way of infected water. These larvae die almost immediately but in dying they produce a sensitivity which when excited by subsequent exposure causes a sometimes violent and usually irritating dermatitis that is often called Swimmers Itch. Such schistosome dermatitis is a common condition throughout North America and it can be contracted even in salt water.

Tapeuorms

The two human species of *Taenia* occur as adults in man only but although the larvae normally are found in food animals the cysticercus of *Taenia solium* can develop aberrantly in man as well as in the pig. Growing as it does in an abnormal host it often develops in an abnormal situation. In the pig it is confined to the striated muscles in man it is often found in the central nervous system where it can cause serious and often fatal cerebral symptoms. This parasite is absent from the more northerly part of North America at present but it is still common in Central and South America as well as in parts of Europe and Asia.

Dipylidium although a normal parasite of urban dogs and cats is found not infrequently in human beings often in young children. As its normal intermediate host is a flea or a biting louse it is a little difficult to explain how humans become infected even when it is obvious that the original source of the infection was a pet dog or cat.

Hymenolepis is shared by man with domestic rodents and is not uncommon in the warmer parts of the continent. Its ability to dispense with an intermediate host facilitates infection of man by contamination of his food by mouse or rat feces.

In its adult stage the genus *Felinoecoccus* is confined to the Canidae and to

these circumstances its normal biology is at least partly suspended and it seems probable that this phenomenon can be correlated with the relatively recent appearance of this protozoon as a parasite in man. It is common and usually nonpathogenic in Asian monkeys but it can infect South American monkeys and cause a serious morbidity. Morphologically identical species occur in a variety of other mammals and it would seem that this genus which also occurs in reptiles is a widespread parasite of vertebrates probably of long standing. This is probably true also of most of the other parasitic amoebae. There is no evidence, however, that man normally becomes infected from other animals except perhaps in India, where monkeys may form a reservoir of potential human infection.

The position is even less clear with regard to the flagellate *Giardia*, a universal and common parasite of man. What may be the same species is also common in dogs and other animals in whom, as in man, it is quite often associated with diarrhea. There is, however, no evidence as yet that man becomes infected naturally from a canine source although opportunities for such infection are undoubtedly common.

Perhaps the most puzzling of the intestinal protozoa are the coccidia. The genus *Eimeria* occurs in a large variety of animals but it does not affect man whereas the genus *Isospora* which has a much more restricted host range is occasionally reported from him.

The role of the hemoflagellates like that of the intestinal protozoa is also not well understood. Insofar as the two varieties of African trypanosomes that infect man are concerned there is no doubt that they were originally mutants of *Trypanosoma brucei*, a parasite of antelopes that is able to live in man. There is also little doubt that they are now exclusively human parasites and although attempts have been made to implicate an antelope reservoir none has been completely successful.

The *Leishmania* group of which *Trypanosoma cruzi* should be considered a member is, I believe, of South American origin although it has spread to many other parts of the world. *T. cruzi* has not left the Americas although it has spread as far north as Texas. It is definitely a parasite of native South American animals (edentates and opossums) that only occasionally infects man and some of his domestic animals. The genus *Leishmania* with its various varieties is less well understood although it is known to occur in dogs and rodents and possibly in other animals in certain places. Some strains, however, appear to have become restricted to human beings.

The human malarias, some varieties of which have been experimentally transmitted to African anthropoid apes, are restricted to man and probably existed as such in prehumanoids.

Trematodes

The Opisthorchid trematodes of mammals and the Heterophyid trematodes of both mammals and birds appear to have little host specificity in their adult stages. Although normal to fish-eating hosts (including dogs and cats) in many parts of the world they occur in human beings only when fish is eaten raw or considerably undercooked (Cameron, 1945). In northern North

America for example *Metorchis conjunctus* is a common parasite of the liver of fish eating mammals and when it occurs in numbers it causes a serious morbidity and may cause death. It can and occasionally does occur in man but as its second intermediate host is the Common Sucker a fish rarely eaten by man it is not an important human parasite even though the fish is common and often heavily infected in the northeastern part of North America.

The blood flukes of man now collectively form one of the most important causes of human disease and it is estimated that they infect about one fifth of the total population of the world. The Far Eastern species has been known for many years to have numerous animal hosts and although I was able to show that *S. mansoni* in the West Indies had a sylvatic host in the naturalized African green monkeys there (Cameron 1978) it was only recently demonstrated that both African species occur in a wide variety of native animals (Kuntz 1955). Man however is still the main source of his own infection both in Africa and in South America. Blood flukes have existed as invaders of man for centuries but there is little doubt that irrigation has done much to increase their incidence in modern times.

Blood flukes capable of developing to sexual maturity in man are absent from North America but blood flukes of birds and rodents which have a developmental cycle similar to that of the common human pathogens have larvae capable of penetrating human skin that it reaches by way of infected water. These larvae die almost immediately but in dying they produce a sensitivity which when excited by subsequent exposure causes a sometimes violent and usually irritating dermatitis that is often called Swimmers Itch. Such schistosome dermatitis is a common condition throughout North America and it can be contracted even in salt water.

Tapeuorms

The two human species of *Taenia* occur as adults in man only but although the larvae normally are found in food animals the *cysticercus* of *Taenia solium* can develop aberrantly in man as well as in the pig. Growing as it does in an abnormal host it often develops in an abnormal situation. In the pig it is confined to the striated muscles in man it is often found in the central nervous system where it can cause serious and often fatal cerebral symptoms. This parasite is absent from the more northerly part of North America at present but it is still common in Central and South America as well as in parts of Europe and Asia.

Dipylidium although a normal parasite of urban dogs and cats is found not infrequently in human beings often in young children. As its normal intermediate host is a flea or a biting louse it is a little difficult to explain how humans become infected even when it is obvious that the original source of the infection was a pet dog or cat.

Hymenolepis is shared by man with domestic rodents and is not uncommon in the warmer parts of the continent. Its ability to dispense with an intermediate host facilitates infection of man by contamination of his food by mouse or rat feces.

In its adult stage the genus *Echinococcus* is confined to the Canidae and to

these circumstances its normal biology is at least partly suspended, and it seems probable that this phenomenon can be correlated with the relatively recent appearance of this protozoon as a parasite in man. It is common and usually nonpathogenic in Asian monkeys, but it can infect South American monkeys and cause a serious morbidity. Morphologically identical species occur in a variety of other mammals, and it would seem that this genus which also occurs in reptiles is a widespread parasite of vertebrates, probably of long standing. This is probably true also of most of the other parasitic amoebae. There is no evidence, however, that man normally becomes infected from other animals except perhaps in India, where monkeys may form a reservoir of potential human infection.

The position is even less clear with regard to the flagellate *Giardia*, a universal and common parasite of man. What may be the same species is also common in dogs and other animals in whom, as in man, it is quite often associated with diarrhea. There is, however, no evidence as yet that man becomes infected naturally from a canine source, although opportunities for such infection are undoubtedly common.

Perhaps the most puzzling of the intestinal protozoa are the coccidia. The genus *Eimeria* occurs in a large variety of animals, but it does not affect man, whereas the genus *Isospora*, which has a much more restricted host range, is occasionally reported from him.

The role of the hemoflagellates like that of the intestinal protozoa is also not well understood. Insofar as the two varieties of African trypanosomes that infect man are concerned, there is no doubt that they were originally mutants of *Trypanosoma brucei*, a parasite of antelopes that is able to live in man. There is also little doubt that they are now exclusively human parasites and although attempts have been made to implicate an antelope reservoir, none has been completely successful.

The *Leishmania* group of which *Trypanosoma cruzi* should be considered a member, is I believe of South American origin, although it has spread to many other parts of the world. *T. cruzi* has not left the Americas, although it has spread as far north as Texas. It is definitely a parasite of native South American animals (edentates and opossums) that only occasionally infects man and some of his domestic animals. The genus *Leishmania*, with its various varieties, is less well understood, although it is known to occur in dogs and rodents and possibly in other animals in certain places. Some strains, however, appear to have become restricted to human beings.

The human malarias, some varieties of which have been experimentally transmitted to African anthropoid apes, are restricted to man and probably existed as such in prehumanoids.

Trematodes

The Opisthorchid trematodes of mammals and the Heterophyid trematodes of both mammals and birds appear to have little host specificity in their adult stages. Although normal to fish-eating hosts (including dogs and cats) in many parts of the world, they occur in human beings only when fish is eaten raw or considerably undercooked (Cameron 1945). In northern North

America for example *Metorchis conjunctus* is a common parasite of the liver of fish eating mammals and when it occurs in numbers it causes a serious morbidity and may cause death. It can and occasionally does occur in man but as its second intermediate host is the Common Sucker a fish rarely eaten by man it is not an important human parasite even though the fish is common and often heavily infected in the northeastern part of North America.

The blood flukes of man now collectively form one of the most important causes of human disease and it is estimated that they infect about one fifth of the total population of the world. The Far Eastern species has been known for many years to have numerous animal hosts and although I was able to show that *S. mansoni* in the West Indies had a sylvatic host in the naturalized African green monkeys there (Cameron 1928) it was only recently demonstrated that both African species occur in a wide variety of native animals (Kuntz 1955). Man however is still the main source of his own infection both in Africa and in South America. Blood flukes have existed as invaders of man for centuries but there is little doubt that irrigation has done much to increase their incidence in modern times.

Blood flukes capable of developing to sexual maturity in man are absent from North America but blood flukes of birds and rodents which have a developmental cycle similar to that of the common human pathogens have larvae capable of penetrating human skin that it reaches by way of infected water. These larvae die almost immediately but in dying they produce a sensitivity which when excited by subsequent exposure causes a sometimes violent and usually irritating dermatitis that is often called Swimmers Itch. Such schistosome dermatitis is a common condition throughout North America and it can be contracted even in salt water.

Tapeuorms

The two human species of *Taenia* occur as adults in man only but although the larvae normally are found in food animals the cysticercus of *Taenia solium* can develop aberrantly in man as well as in the pig. Growing as it does in an abnormal host it often develops in an abnormal situation. In the pig it is confined to the striated muscles in man it is often found in the central nervous system where it can cause serious and often fatal cerebral symptoms. This parasite is absent from the more northerly part of North America at present but it is still common in Central and South America as well as in parts of Europe and Asia.

Dipylidium although a normal parasite of urban dogs and cats is found not infrequently in human beings often in young children. As its normal intermediate host is a flea or a biting louse it is a little difficult to explain how humans become infected even when it is obvious that the original source of the infection was a pet dog or cat.

Hymenolepis is shared by man with domestic rodents and is not uncommon in the warmer parts of the continent. Its ability to dispense with an intermediate host facilitates infection of man by contamination of his food by mouse or rat feces.

In its adult stage the genus *Echinococcus* is confined to the Canidae and to

a lesser extent to the Felidae (Cameron 1927) Its larval stage is the hydatid cyst which may develop in many species of mammals Undoubtedly several different species exist but at least some of these have rather restricted ranges of intermediate hosts There is reason to believe however, that all of them can undergo larval development in human beings Man can be infected of course only by swallowing the tapeworm eggs, and these can be acquired only from contact of food or fingers with the feces of carnivores (It should be unnecessary to note that the cyst itself is *not* infective to man directly, and that it can be handled or even eaten with impunity The tapeworm will not develop in man and the sole source of human infection is carnivore feces) The usual carnivore involved is the dog and the usual intermediate hosts from which the dog becomes infected are domestic stock principally sheep and swine As a rule cattle are poor hosts and although they often become infected and develop cysts these cysts are very often sterile and abnormal In Great Britain and parts of continental Europe the fox is a common definitive host, and horses frequently are infected with the cyst In the northwestern part of North America there is a peculiar sylvatic cycle in which the wolf is the main carrier of the tapeworm and a variety of deer especially moose elk and caribou (including reindeer) are the carriers of the cyst Indications at present are that this species which also can infect dogs is not the common *F. granulosus* Indian hunters kill the deer and feed the lungs (containing the cyst) to their dogs The dogs in turn become infected and many villages have become grossly infected in consequence In some parts of the Canadian northwest the human incidence is very high and about one third of the local inhabitants are or have been infected

In the southern part of North America the cyst is reported in swine but only rarely in man In Canada the cyst is extremely rare in domestic stock and human cases diagnosed in the southern part of that country have in most cases contracted the parasite elsewhere The parasite is still common in many other parts of the world including the Mediterranean region Europe Asia Australia New Zealand South Africa and parts of South America It is one of the more important of the animal parasites transmissible to man and it is responsible for a considerable amount of disease

Fish tapeworms of the genus *Diphyllobothrium* are all of animal origin In North America the species are widely distributed throughout the northern part of the continent and occur in a number of fish eating mammals and possibly in birds *D. latum* which is the species most commonly reported from man occurs in bears as well as in dogs and there is some evidence suggesting that bears are the main reservoir from which man is ultimately infected When it is realized that the tapeworm can produce several million eggs daily for a decade it is easy to understand how a comparatively few reservoirs could keep the infections at a high level in fish (Cameron 1947) This is the commonest human tapeworm in North America the infection usually deriving from uncooked or undercooked fish Although it is known to produce some morbidity elsewhere and by its requirements of vitamin B₁₂ to cause a serious anemia virtually nothing is known of its pathogenicity in North America

The final larval stages of several species of this tapeworm are capable of

living in a somatic position in man although only after actual mechanical transfer from the final intermediate host to a wound or to the orbit. These pargana as they are called usually remain localized but they occasionally become malignant and metastasize as is also sometimes the case with old larval echinococci.

Nematodes

The human whipworm *Trichuris trichiura* is quite common in North America but it almost always occurs in such small numbers that it is difficult to understand how it can maintain itself without an alternate reservoir. It is common in swine however and it seems reasonable to believe that the pig acts as such a reservoir. The egg must be exposed to the atmosphere for some weeks before becoming infective and it has every opportunity of becoming a contaminant of vegetables during this period.

The genus *Capillaria* is closely related to *Trichuris* and like it is widespread in a great variety of mammals. Man however has no species proper to himself although he may be infected by the *Capillaria hepatica* of rodents. This parasite common in North America and elsewhere lives in the livers of rats and wild Muridae. The eggs are laid there and escape only after the death of the host either by natural decomposition or via the digestive tract of a carnivore. These eggs are not immediately infective and they must remain in the open for several weeks before an embryo develops. Infection is presumably by ingestion of the infective egg. A number of human cases are on record from North America but there is no ready means of diagnosis except by biopsy or autopsy and we have no knowledge of how common this parasite really is in man. In some areas however almost one third of the domestic rats are infected.

Related to these parasites is the much more serious *Trichinella spiralis*.

The trichina worm is really a parasite of carnivorous animals and its original home was probably the Arctic and sub-Arctic where carnivore feeds on carnivore. It is still absent from the tropics but it has become common in temperate lands where animal husbandry has converted the herbivorous pig into an omnivore and so by feeding it on pork scraps has made it a reservoir for the worm. The parasite has little host specificity and it can be induced to develop in almost any mammal. It is fairly common in rodents but there is no evidence that the rat enters into the pig-pig or pig-human cycle rather it is probable that it maintains a rat-dog or rat-cat cycle. In the Arctic trichinosis is a highly important human disease depending however not on swine but on wild carnivores especially bears and walrus. We have no evidence as yet that rodents play any part in the cycle although foxes both white and red as well as a great variety of terrestrial mammals are commonly infected. While white whales and seals are occasionally infected the sylvatic cycle seems to involve the polar bear and the walrus. Man becomes infected by eating either and in some parts of the Arctic about one third of the Eskimos are infected often seriously. Elsewhere in North America infection of man is usually from pork although it originates occasionally from bear flesh. While the index of infection is low in southern Canada it is high in the United States where it is estimated that about 20 per cent of the population have acquired

the parasite. However, infections are usually low and the clinical manifestations light. Nevertheless, this nematode can be considered as the most serious of all the helminth parasites of lower animals transmissible to man.

Ascaris lumbricoides has a problematical origin. Although several strains of this parasite may exist, there is no conclusive evidence that there is more than one species or variety of *A. lumbricoides*. The parasite occurs regularly in man, in swine and apparently in numerous other animals, including sheep. The available evidence suggests strongly that husbandry practices in temperate climates make it a predominantly porcine parasite while in warm climates hygienic conditions make it predominantly a human parasite. However, it is interchangeable between man and the pig and occasionally, between sheep and other animals. In northern lands where there is a fairly high incidence of very light infections in man, these infections are probably usually of porcine origin; in general, they are much too light to depend on man as their source. The reverse is probably true in the tropics where pigs are rarely infected and where the human incidence may exceed 60 per cent.

The relationship of this worm to human disease is still obscure. However, there is some evidence that both in its larval and adult form it may cause infections which, even when they are light, may cause toxic, nutritional and traumatic damage. Heavy infections, which are fortunately rare in the north, cause quite serious disturbances in health.

Most of the other helminth parasites of man are also accidental.

Toxocara canis is the common roundworm of puppies and infection of these animals is often prenatal. The female dog swallows the infective eggs and the larvae develop somatically and remain alive for a long period in a dormant condition until, during pregnancy, they migrate through the placenta to the fetus or to the intestine of the female. In either situation it can reach sexual maturity. Somatic migrations are less common in male dogs in which the worm appears to develop more normally. However, the relationship between this parasite and the female sex hormones of the host is still obscure. Beaver *et al.* (1952) have shown that somatic migrations of the larva can occur in children, although the worm does not appear to reach sexual maturity in them. However, this parasite can produce symptoms of liver disturbance and it can remain alive for months in the liver without further development.

The related *Toxocara felis* of cats also can develop in this way but, curiously, the very similar *Toxascaris leonina* of dogs and cats does not migrate from the intestinal tract and is not involved in the production of symptoms in man, although it sometimes develops to sexual maturity in his intestine.

Man's two common hookworms are specific to himself but the various hookworms of dogs and cats are sometimes able, in a manner analogous to that of the parasites of swimmers, to penetrate human skin and there to maintain themselves as larvae for some time. They ultimately die without further development. However, one species, *Uncinostoma ceylanicum*, is occasionally recorded as a sexually mature parasite in the human intestine but it may have entered the body by the mouth rather than through the integument.

Some of the other bursate nematodes that normally occur in other animals

occasionally succeed in parasitizing man. This is especially true of the various species of the genus *Trichostrongylus*.

However a number of reports of human infection by animal helminths are probably due to mistakes in identification. Eggs of helminths (for example the liver fluke) ingested in food may be passed unchanged in the feces. Similarly eggs of plant nematodes have been given specific names. Even more peculiar are species such as *Mecistocirrus* a parasite of swine and cattle that has been recorded as a human parasite on the sole authority of an abattoir attendant who found the worms lying on what he thought were human feces in the abattoir. Others have been recorded on the basis of the size of the egg. It would be unnecessary to mention these cases were it not for the fact that such instances have a way of getting into the textbooks and remaining there in subsequent editions.

Strongyloides stercoralis is a common parasite of man especially in warm countries where hygiene is poor. However the status of the various species of this genus is still doubtful. It is very widely distributed and many species are very similar morphologically. It is a comparatively recent parasite as shown by its peculiar life cycle in the free state and man may be infected by species normally occurring in domestic animals. Nevertheless it is highly probable that man is usually his own source of infection.

The history of the filarid worms which are carried by blood-sucking flies is so obscure that no valid deductions are possible but because of their rather exceptional mode of transmission wild animals may well be involved as the reservoir from which man originally acquired his infections. *Onchocerca* probably had a ruminant host and it may still have one in Africa. *Wuchereria bancrofti* now is probably an exclusively human parasite its intermediate hosts are almost always domestic insects and recent work on the related *W. malayi* suggests that an Asiatic monkey (*Macaca irus*) may have been the original host of this genus (Buckley and Edeson 1956).

Ectoparasites

Parasitic arthropods affecting man offer special problems and many of them are instrumental in the introduction of other organisms into him and can be of great importance even though they are not strictly speaking parasites of man. Thus the hard ticks are nonspecific parasites mainly of mammals and reptiles and none of them parasitizes man exclusively although several are important vectors of disease organisms. To a large extent the soft ticks are also nonspecific although a number of species of the genus *Ornithodoros* frequently attack man while one the African *O. moubata* has quite definite associations with man and may well be considered as adapted to his habits.

The Trombiculid mites which in their larval stage attack mammals and birds are equally nonspecific although quite a number of them will attach themselves to the human skin.

The role of *Sarcoptes scabiei* is more definite. There are a number of physiological varieties of this mange mite one of which is definitely adapted to living

within the human skin. Most if not all of the other varieties however, can live in the human skin for short periods but there is no evidence that any of these can become an habitual human parasite.

Demodex folliculorum is still one of the outstanding puzzles of medical entomology. The genus is widely distributed among animals, but there is no evidence that man can be infected from any source other than human. Until the life cycle of the parasite is understood both its pathogenicity and its mode of infection will remain doubtful.

Of the two species of lice infesting human beings there is little doubt that the body head species (*Pediculus humanus*) has evolved along with man from prehumanoid days. Nevertheless it has been found on South American monkeys but not on Old World monkeys and under rather exceptional laboratory conditions it has been reared on rabbits. The pubic louse is also exclusively human. No animal louse can attain more than a very transitory footing on the human skin.

This is not the case with fleas. In spite of the popular name of human flea applied to *Pulex irritans* no species is primarily a parasite of humans. *Pulex* itself is still basically a parasite of swine and originally was probably a parasite of badgers although in parts of the world such as North Africa, it has become acclimatized to live and to breed on man. It is extremely rare in North America and while not uncommon in pigsties in Europe is nearly as rare on man there as it is here. The closely related *Xenopsylla* is of course a rodent flea and like the common dog and cat flea it will readily attack man. *Xenopsylla* is apparently spreading in North America but the commonest fleas attacking human beings are those normally found on cats and dogs.

The jigger flea apparently originating in tropical America will also attack man but it also is basically a parasite of other animals. Bedbugs were probably originally parasites of bats and birds but one *Cimex lectularis* has become so domesticated that for all practical purposes it is a human parasite and is very widely distributed in all temperate and subtropical regions. Its tropical relative is less exclusively a human parasite. *C. lectularis* itself can be reared successfully on rabbits and other laboratory animals.

Larval parasitism by dipterous flies is particularly common in many parts of the world. In some cases it is a facultative type of parasitism with no particular host preferences and man is often infected with these nonspecific larvae. In other cases parasitism has become obligatory and host preferences have become more strict. This is especially true among the Oestridae but even here man is occasionally attacked. There are no strictly human species. *Dermatobia hominis* in spite of its name is not a specific human parasite and is common on monkeys and other animals including all domestic animals in South America.

Conclusions

Man was a late comer into the animal kingdom and he probably brought with him from his prehuman stage two or three malarial parasites, his pinworms and his lice. Being still quite generalized and primitive physiologically he offered a good habitat for the parasites proper to the animals with which he

surrounded himself and he easily became host to his taenias his hookworms his ascarids and his filariae. The more generalized his animals (in a physiological sense) and the more closely they associated with him the more likely they were to interchange parasites and swine and dogs on occasion still act as sources of human infection. Man's food habits exposed him in due course to other relatively nonspecific parasites such as most flukes and the fish tape worms but it is quite possible that this change caused him to lose some of his parasites. This applies for example to the tapeworm *Bertiella* which is still common in other primates and to which man is still susceptible. His bathing habits rendered him a suitable host for various skin penetrating blood flukes.

In this way man acquired his major parasites quite early in his history but the process did not cease in the early days of his development it is still going on and even now he is accidentally becoming infected with parasites from both domestic and wild animals.

As mentioned above it is necessary to distinguish between infection and disease. Except when accidents occur as when an ascarid worm occludes a bile duct adult helminths usually cause disease only when present in considerable numbers. As helminths cannot multiply continuously within a single host infection postulates exposure to a large number of infective forms usually over a considerable length of time (*Trichinella* infection is an obvious exception to the time factor). It follows therefore that disease seldom results from adult helminth parasites that are not normal human parasites. This is not true of larval stages that develop not only in abnormal situations but because of the long life span of man can often develop in abnormal ways, examples include such forms as cerebral cysticercosis larval toxocariasis and daughter hydatids. In these cases as in *Trichinella* a single exposure may produce clinical symptoms. This is true also of the protozoal infections because of their ability to multiply within the body.

As yet there are no preventive vaccines or sera for any of the parasitic infections. In many cases specific chemotherapy is available but in some of them (these include some of the more serious ones) no treatment has yet been discovered. The specificity of the life cycles of these parasites however makes practical preventive measures feasible often by such simple expedients as good hygiene or the adequate cooking of food.

References

- BEAVER P C C H SNYDER G M CARRERA J H DENT & J W LAFFERTY 1957 Chronic eosinophilia due to visceral larva migrans. *Pediatrics* 9(1) 7-19.
- BUCKLEY J J C & J F B FIDSON 1956 On the adult morphology of *Huchereria* sp. (mal. v.) from a monkey (*Macaca murina*) and from cats in Malaya and on *Huchereria* sp. from a mad gambia cat. *J. Helminthol.* 30(1) 1-20.
- CAMERON T W M 1977 The helminth parasites of animals and human disease. *Proc. Roy. Soc. Med.* 20 547-556.
- CAMERON T W M 1978 A new definitive host for *Strongylus edentatus*. *J. Helminthol.* 6(4) 219-222.
- CAMERON T W M 1945 Fish carried parasites in Canada. *Can. J. Comp. Med.* 9 245-254 283-286 307-311.
- KUNTZ R F 1955 Biology of the schistosome complex. *Am. J. Trop. Med. Hyg.* 4(3) 383-413.

within the human skin. Most if not all of the other varieties, however, can live in the human skin for short periods, but there is no evidence that any of these can become an habitual human parasite.

Demodex folliculorum is still one of the outstanding puzzles of medical entomology. The genus is widely distributed among animals, but there is no evidence that man can be infected from any source other than human. Until the life cycle of the parasite is understood, both its pathogenicity and its mode of infection will remain doubtful.

Of the two species of lice infesting human beings, there is little doubt that the body-head species (*Pediculus humanus*) has evolved along with man from prehumanoid days. Nevertheless, it has been found on South American monkeys, but not on Old World monkeys, and under rather exceptional laboratory conditions it has been reared on rabbits. The pubic louse is also exclusively human. No animal louse can attain more than a very transitory footing on the human skin.

This is not the case with fleas. In spite of the popular name of 'human flea' applied to *Pulex irritans*, no species is primarily a parasite of humans. *Pulex* itself is still basically a parasite of swine and originally was probably a parasite of badgers, although in parts of the world, such as North Africa, it has become acclimatized to live and to breed on man. It is extremely rare in North America and, while not uncommon in pigsties in Europe, is nearly as rare on man there as it is here. The closely related *Xenopsylla* is, of course, a rodent flea and, like the common dog and cat flea, it will readily attack man. *Xenopsylla* is apparently spreading in North America, but the commonest fleas attacking human beings are those normally found on cats and dogs.

The jigger flea, apparently originating in tropical America, will also attack man, but it also is basically a parasite of other animals. Bedbugs were probably originally parasites of bats and birds, but one *Cimex lectularis* has become so domesticated that, for all practical purposes, it is a human parasite and is very widely distributed in all temperate and subtropical regions. Its tropical relative is less exclusively a human parasite. *C. lectularis* itself can be reared successfully on rabbits and other laboratory animals.

Larval parasitism by dipterous flies is particularly common in many parts of the world. In some cases it is a facultative type of parasitism with no particular host preferences, and man is often infected with these nonspecific larvae. In other cases parasitism has become obligatory, and host preferences have become more strict. This is especially true among the Oestridae, but even here man is occasionally attacked. There are no strictly human species. *Dermatobia hominis*, in spite of its name, is not a specific human parasite and is common on monkeys and other animals, including all domestic animals in South America.

Conclusions

Man was a late comer into the animal kingdom, and he probably brought with him from his prehuman stage two or three malarial parasites, his pinworms, and his lice. Being still quite generalized and primitive physiologically, he offered a good habitat for the parasites proper to the animals with which he

surrounded himself and he easily became host to his taenias his hookworms his ascarids and his filariae. The more generalized his animals (in a physiological sense) and the more closely they associated with him the more likely they were to interchange parasites and swine and dogs on occasion still act as sources of human infection. Man's food habits exposed him in due course to other relatively non specific parasites such as most flukes and the fish tape worms but it is quite possible that this change caused him to lose some of his parasites. This applies for example to the tapeworm *Bertiella* which is still common in other primates and to which man is still susceptible. His bathing habits rendered him a suitable host for various skin penetrating blood flukes.

In this way man acquired his major parasites quite early in his history but the process did not cease in the early days of his development it is still going on and even now he is accidentally becoming infected with parasites from both domestic and wild animals.

As mentioned above it is necessary to distinguish between infection and disease. Except when accidents occur as when an ascarid worm occludes a bile duct adult helminths usually cause disease only when present in considerable numbers. As helminths cannot multiply continuously within a single host infection postulates exposure to a large number of infective forms usually over a considerable length of time (*Trichinella* infection is an obvious exception to the time factor). It follows therefore that disease seldom results from adult helminth parasites that are not normal human parasites. This is not true of larval stages that develop not only in abnormal situations but because of the long life span of man can often develop in abnormal ways examples include such forms as cerebral cysticercosis larval toxocarosis and daughter hydatids. In these cases as in *Trichinella* a single exposure may produce clinical symptoms. This is true also of the protozoal infections because of their ability to multiply within the body.

As yet there are no preventive vaccines or sera for any of the parasitic infections. In many cases specific chemotherapy is available but in some of them (these include some of the more serious ones) no treatment has yet been discovered. The specificity of the life cycles of these parasites however makes practical preventive measures feasible often by such simple expedients as good hygiene or the adequate cooking of food.

References

- BEAVER P C C H SNYDER G M CARRERA J H DENT & J W LAFFERTY 1952 Chronic eosinophilia due to visceral larva migrans. *Pediatrics* 9(1) 7-19.
- BICKLEY J J C & J F B EDESON 1956 On the adult morphology of *H. uclereria* sp. (mal y f) from a monkey (*Macaca* s) and from cats in Malaya and on *H. cherisia* sp. from a dog and a cat. *J. Helminthol.* 30(1) 1-20.
- CAMERON T W M 1971 The helminth parasites of animals and human disease. *Proc. Roy. Soc. Med.* 20 547-556.
- CAMERON T W M 1978 A new definitive host for *Sci. st. m. m. nson*. *J. Helminthol.* 6(4) 219-222.
- CAMERON T W M 1945 Fish carried parasite in Canada. *Can. J. Comp. Med.* 9 245-254 283-286 307-311.
- KUNTZ R F 1955 Biology of the schistosome complex. *Am. J. Trop. Med. Hyg.* 4(3) 383-413.

INDUSTRIAL ANTHRAX*

By Philip Sigmund Brachman†

Anthrax Investigations Unit Communicable Disease Center Atlanta Ga

and

Francis Robert Fekety‡

Epidemic Intelligence Service Communicable Disease Center Atlanta Ga

While anthrax remains a scourge in many parts of the world it has steadily declined in importance in the United States. Nevertheless anthrax has become increasingly interesting for there is still much to be learned about the biology of its causative agent its transmission and pathogenesis and the mechanisms of immunity to it. This presentation will summarize the major developments in the field of industrial anthrax since the appearance of the reviews of Smyth (1945) Wolff and Heimann (1951) and Steele and Helvig (1953). The term "industrial anthrax" refers to cases acquired in nonagricultural occupations that require contact with contaminated animal materials. In the United States about 90 per cent of industrial cases (about 25 to 35 cases per year) are found in workers exposed to carpet wools goat hair, or goat skins imported from countries where anthrax is prevalent. Occasional cases are traced to rarer sources such as elephant tusks and bone meal. There is a very slight but definite risk associated with residence in the neighborhood of factories processing these materials for a few cases of the disease have been reported in persons living or working near them.

FIGURE 1 shows the number of cases of human anthrax reported yearly to the National Office of Vital Statistics (NOVS) Washington D. C. since 1916. Both morbidity and mortality have shown definite declines and at the present time only 40 to 50 cases of anthrax are reported yearly. (These are minimum figures since under reporting of anthrax either advertent or inadvertent is not unexpected. For example from 1941 to 1956 NOVS recorded 85 cases of anthrax from one state. That state has made records available to us indicating that 122 cases occurred there during the same period.) As will be shown later however there is little justification for the optimistic view that this decline in morbidity will continue automatically. The cutaneous form of anthrax accounts for more than 95 per cent of anthrax in man reported in this country. An occasional case of inhalation anthrax or generalized (septicemic) anthrax is still reported but intestinal anthrax has never been documented in the United States. *Bacillus anthracis* is extremely sensitive *in vitro* to most of the common antibiotics (polymyxin is a notable exception) and deaths from anthrax are rare at present whereas a 10 to 20 per cent mortality was expected in the

The work reported in this paper is from the Communicable Disease Center Public Health Service Atlanta Ga. The Wistar Institute of Anatomy and Biology Philadelphia Pa. and the University of Pennsylvania School of Medicine and Veterinary Science Philadelphia Pa. It is Contribution No. 20 of Microbiology and Medicine (Wistar Institute) and was supported by a contract from the United States Army Chemical Corps Fort Detrick Md.

† Present address: The Wistar Institute University of Pennsylvania Philadelphia Pa.
‡ Present address: Department of Medicine Johns Hopkins Hospital Baltimore Md.




FIGURE 1 Anthrax in man in the United States 1916 to 1956 Data from the National Office of Vital Statistics Washington D C

preantibiotic era (FIGURE 2). A high fatality rate is still associated with inhalation or generalized anthrax. Cutaneous cases in which treatment is delayed, however, are frequently more severe than those cases in which therapy is initiated within 7 days after onset.

The nationwide distribution of cases over the past several decades is shown in FIGURE 3. It is readily apparent that most of the cases are reported from only a few states. It has been realized for some time that those cases reported from the northeastern states have been primarily industrial in origin. More than 50 per cent of the 3228 cases of anthrax reported since 1916 have been from five states (Pennsylvania, New York, New Jersey, Massachusetts, and New




FIGURE 2 Anthrax fatality rate in the United States 1916 to 1956 Data from the National Office of Vital Statistics Washington D C

INDUSTRIAL ANTHRAX*

By Philip Sigmund Brachman†

Anthrax Investigations Unit Communicable Disease Center Atlanta Ga

and

Francis Robert Fekety‡

Epidemic Intelligence Service Communicable Disease Center Atlanta Ga

While anthrax remains a scourge in many parts of the world it has steadily declined in importance in the United States. Nevertheless anthrax has become increasingly interesting for there is still much to be learned about the biology of its causative agent its transmission and pathogenesis and the mechanisms of immunity to it. This presentation will summarize the major developments in the field of industrial anthrax since the appearance of the reviews of Smyth (1945), Wolff and Heimann (1951) and Steele and Helvig (1953). The term industrial anthrax refers to cases acquired in nonagricultural occupations that require contact with contaminated animal materials. In the United States about 90 per cent of industrial cases (about 25 to 35 cases per year) are found in workers exposed to carpet wools goat hair or goat skins imported from countries where anthrax is prevalent. Occasional cases are traced to rarer sources such as elephant tusks and bone meal. There is a very slight but definite risk associated with residence in the neighborhood of factories processing these materials for a few cases of the disease have been reported in persons living or working near them.

FIGURE 1 shows the number of cases of human anthrax reported yearly to the National Office of Vital Statistics (NOVS) Washington D C since 1916. Both morbidity and mortality have shown definite declines and at the present time only 40 to 50 cases of anthrax are reported yearly. (These are minimum figures since under reporting of anthrax either advertent or inadvertent is not unexpected. For example from 1941 to 1956 NOVS recorded 85 cases of anthrax from one state. That state has made records available to us indicating that 122 cases occurred there during the same period). As will be shown later however there is little justification for the optimistic view that this decline in morbidity will continue automatically. The cutaneous form of anthrax accounts for more than 95 per cent of anthrax in man reported in this country. An occasional case of inhalation anthrax or generalized (septicemic) anthrax is still reported but intestinal anthrax has never been documented in the United States. *Bacillus anthracis* is extremely sensitive *in vitro* to most of the common antibiotics (polymyxin is a notable exception) and deaths from anthrax are rare at present whereas a 10 to 20 per cent mortality was expected in the

The work reported in this paper is from the Communicable Disease Center Public Health Service Atlanta Ga. The Wistar Institute of Anatomy and Biology Philadelphia Pa and the University of Pennsylvania School of Medicine and Veterinary Science Philadelphia Pa. It is Contribution No. 20 of Microbiology and Medicine (Wistar Institute) and was supported by a contract from the United States Army (Chemical Corps Fort Detrick Md).
† Present address: The Wistar Institute University of Pennsylvania Philadelphia Pa.
‡ Present address: Department of Medicine Johns Hopkins Hospital Baltimore Md.




FIGURE 1 Anthrax in man in the United States 1916 to 1956 Data from the National Office of Vital Statistics Washington D. C.

preantibiotic era (FIGURE 2). A high fatality rate is still associated with inhalation or generalized anthrax. Cutaneous cases in which treatment is delayed, however, are frequently more severe than those cases in which therapy is initiated within 7 days after onset.

The nationwide distribution of cases over the past several decades is shown in FIGURE 3. It is readily apparent that most of the cases are reported from only a few states. It has been realized for some time that those cases reported from the northeastern states have been primarily industrial in origin. More than 50 per cent of the 3228 cases of anthrax reported since 1916 have been from five states (Pennsylvania, New York, New Jersey, Massachusetts, and New




FIGURE 2 Anthrax fatality rate in the United States 1916 to 1956 Data from the National Office of Vital Statistics Washington D. C.




FIGURE 3 Distribution of cases of human anthrax by states 1970 to 1956 Data from the National Office of Vital Statistics Washington D C

Hampshire) which reflects the concentration of textile industries and tanneries in them. Cases in other sections of the country usually have been agricultural in origin. The latter patients were usually farmers, abattoir workers, or veterinarians who were infected as a result of handling carcasses of animals that had died of anthrax or of accidentally contaminating themselves while administering a veterinary spore vaccine. Texas, Mississippi, California, and Louisiana have accounted for more than 50 per cent of all agricultural cases.

For purposes of presentation in FIGURE 3, a state was classified as a source of industrial cases when the great majority of cases reported from it have been industrial in origin, even though it has occasionally reported agricultural cases. South Carolina and Nevada have not reported any human anthrax since 1916. Prior to 1953, the rare cases reported from North Carolina were agricultural. In 1953, a firm processing goat hair was established there, and since then 9 cases have been reported among its workers. Most of the anthrax reported from Virginia has been agricultural, but 4 cases in that state associated with a carpet manufacturing plant were reported in 1955.

Anthrax is primarily a disease of animals, which constitute its most important reservoir. Man is an incidental host only by virtue of contact with such animals directly or indirectly with their products. As a result, the incidence of human anthrax would be expected to reflect the incidence of the disease in




FIGURE 4 Industrial and agricultural human anthrax in the United States 1916 to 1956. Data from the National Office of Vital Statistics, Washington, D. C.

animals. That this is the case is shown by FIGURE 4 which demonstrates the number of cases of human anthrax diagnosed in the United States since 1916 by type of exposure. Anthrax in animals in the United States has declined since 1940 partly because of good animal husbandry and the widespread use of vaccine. The incidence of human agricultural anthrax has declined correspondingly. Industrial anthrax, however, is a reflection of the incidence of disease in foreign animals, since this has not changed appreciably; neither has the incidence of industrial anthrax. Notable variations in the number of yearly cases of industrial anthrax can be related to changes within the industry. The peaks of 1917 and 1941 probably represent the effect of wartime factors such as increased output and employment and the use of dirtier, less desirable and more highly contaminated raw materials. The trough in the 1930s parallels a decrease in employment and production. The agricultural peak in 1924 is a reflection of epizootics in Mississippi and Texas. The industrial peak in 1926 may reflect the tanning of hides from the domestic herds affected in 1924. All of these factors have become relatively stable, however, so that for the past 10 years about 80 per cent of the human anthrax in the United States (about 30 to 40 cases per year) can be called industrial. The industries involved have been far from stable, however.

In the last decade there has been a marked change in the distribution of cases throughout the industries involved, and some of the change can be attributed to the increased use of synthetic fibers and plastics. This has particularly affected tanneries and carpet wool concerns, which together contributed more than one half of the anthrax diagnosed in the United States for several decades (TABLE 1). In spite of the fact that anthrax has been declining, tanneries and woolen mills now account for only 11 per cent of the cases, or about 4 to 5 cases per year. Cases associated with the goat hair industry now predominate; their number has increased from an average of 3

TABLE 1
DISTRIBUTION OF ANTHRAX IN THE UNITED STATES BY SOURCE OF INFECTION
(Adapted from Wolff and Heiman 1951)

Source	Percentage reported cases traced to source				
	1919 to 1923	1924 to 1928	1929 to 1933	1934 to 1938	1939 to 1946
Hides and skins	53	38	30	35	2
Wool	3	25	27	15	9
Goat hair	17	9	4	5	65
Agriculture	13	14	26	34	21
Shaving brushes	18	4	2	2	0
Laboratory	—	2	3	2	0

cases per year during the 1930s to about 20 to 25 cases per year or 65 per cent of all anthrax in 1955 and 1956. The number of persons employed in woolen mills is at least 5 times the number employed in goat hair mills.

For several years therefore public health workers have concentrated their efforts on the control of anthrax in the goat hair industry. The major users of goat hair are a relatively small number of firms manufacturing haircloth interlinings used to lend shape to suits and coats. This highly competitive industry is one of the last of the textile industries to withstand the inroads of synthetic fibers. It should be understood that any preventive measure that would increase the cost of their product even a few cents per pound would probably be disastrous. Consequently protection against this disease has developed into a complicated and challenging public health problem.

The goat hair used in the United States is obtained primarily from middle eastern countries such as Pakistan, Iraq, Iran and India. It is impossible for domestic buyers to control the disease in herds in those countries or to predict or reduce the contamination of the hair at the time of export. Raw goat hair is shipped in jute or burlap-covered bales weighing several hundred pounds and stevedores and truckers handling these bales have been known to develop anthrax. When the bales are opened at the factory the material is fed into picking machines that shake the hair, separate the fibers and remove many of the coarse impurities such as dust, rocks, bones, dried blood and animal feces. Vacuum exhaust systems carry away much of this debris but it is an extremely dusty operation and one would expect the risk of contracting anthrax to be great at this stage.*

The hair fibers are then fed into carding and combing machines that further classify and separate the fibers. The loose strand of hair that is produced is then blended with various other fibers such as wool or rayon to produce a rope-like material called hair top. The top is drawn into a narrow strand and spun on frames. The resultant thread is then woven into fabric. The fabric

In one factory the hair is secured after it has been picked. This is done to improve the quality of the finished product rather than to prevent anthrax. The securing process consists of feeding bunches of the hair into a series of 3 vats of water and soap or soda at 40 to 50°C for about 30 minutes. The hair is then rinsed and conveyed to chamber in which it is dried in warm air at 80°C for 15 minutes.

TABLE 2

RELATION OF THE DEVELOPMENT OF ANTHRAX TO SURFACE CONTAMINATION AT VARIOUS STAGES IN THE PROCESSING OF GOAT HAIR 1945 TO 1946

Department	Total cases	Cases per 100 mill employees per year				Percent surface samples positive for B. anthracis 1954 to 1956
		Mill A	Mill B	Mill C	Combined	
Picking	22	4.4	4.4		4.4	66
Carding	40	4.4	4.4		4.4	62
Combing	25	4.4	2.2	4.4	4.0	59
Drawing	40	2.2	2.8	3.5	3.0	44
Spinning	100	1.4	2.3	1.8	2.2	39
Weaving	16	0.2	0.2	0.2	0.2	22
Finishing	16	0.9	0.9	0.8	0.8	19
Office	0	0.0	0.0		0.0	5
Entire mill	259	1.2	1.6	1.6	1.4	44

Excluded because of lack of reliable data.

interlining is finished by singeing (to remove extraneous hair) sizing washing Sanforizing and other processes

The danger of contracting anthrax has always seemed to be greatest in the early stages of this process. TABLE 2 summarizes information on this point collected from 3 goat hair factories. Two hundred and fifty cases of anthrax were diagnosed in these 3 factories in the past 13 years or 1.4 cases per 100 employees per year. The greatest number of cases came from the spinning rooms but the highest attack rates were found in the processes preceding spinning. Attack rates averaged more than 4 cases per 100 employees per year in the picking carding and combing departments or about twice that in the spinning department. Relatively low rates were found in the weaving and finishing stages. The trend toward a lower risk of contracting anthrax in successive stages was noted in all the factories studied. The striking similarity of departmental attack rates in each of the factories is noteworthy. It has been suggested that the risk of contracting the disease in one of these factories is related to the proximity of the employee to the picking and carding rooms and that most of the plant contamination results from the spread of the organism through the air from that source. The factories studied use similar methods of production but their physical plants are quite different. Consequently this striking correspondence in attack rates between factories suggests that the nature of the work is a more important determinant of the attack rate than is proximity to the hair storage and picking area. Mill C which scours its raw hair immediately after picking has as high an attack rate as those that do not. These attack rates may be compared to those in the wool industry furnished by Wolff and Heimann (1951) who recorded 0.078 cases per 100 employees per year in that industry or about one twentieth the rate in the goat hair industry. The rate for workers engaged in picking carding and scouring wool was about 0.21 cases per 100 employees per year or about one fifteenth the rate noted in the corresponding stages in the processing of goat hair. Our studies have




FIGURE 5 The relation of the risk of developing anthrax to the contamination of surfaces with *Bacillus anthracis* by department in three mills 1954 to 1956

shown that wool processing mills are less contaminated with the spores of *B. anthracis* than are goat hair mills.

TABLE 2 also summarizes the results of our studies on different occasions of the surface contamination of several goat hair mills with the spores of *B. anthracis*. Moist sterile cotton swabs were used to take cultures from floors and walls and from machinery and other equipment. The swabs were then used to inoculate blood agar plates that were incubated for 18 hours at 37°C. Colonies grossly and microscopically resembling *B. anthracis* were tested for mouse virulence and lysis by the gamma bacteriophage of Brown and Cherry (1955). Forty-four per cent of all samples in this series were positive for *B. anthracis*, indicating a relatively high contamination of the mills with this pathogen. The greatest numbers were isolated from the early dusty stages. FIGURE 5 shows that the departmental attack rate in these mills bears a direct relation to the level of surface contamination.

That contact by man with the organism does not necessarily lead to disease is demonstrated by the recent report of Carr and Rew (1957) who were able to recover *B. anthracis* from the nose or pharynx of 14 of 101 healthy workers in 2 of these mills. Again most of the recoveries were from persons working in the early stages of production. While the transmission of anthrax either from infected persons or workers such as these has never been documented, cases have been recorded in which asymptomatic workers might have been responsible. Transmission via the clothing of these workers is another likely explanation of these incidents.

It might be inferred from FIGURE 5 that if the surface contamination could be reduced to the 10 per cent level, a negligible attack rate might prevail. Our experience tends to confirm the importance of plant hygiene. However, it is probable that both the level of plant contamination and the attack rate are dependent upon the more important factor of hair contamination. Attempts to decontaminate the hair have met with many difficulties. In England, while the establishment of the Wool Disinfection Station at Liverpool and the passage of governmental regulations have satisfactorily controlled anthrax, such a plan does not seem applicable in the United States. The initial cost of the equipment would be enormous; it would be difficult to decide where it

should be located and the continuing costs (estimated at \$3 000 000 per year) could not be absorbed by the industry. A few firms here have recently begun to steam sterilize their hair in the top stage but this is not entirely satisfactory since the procedure alters the material and makes spinning difficult. In addition persons working in the early higher risk stages are unaffected by this measure. We have evaluated several other methods of disinfection (to be reported elsewhere) and have not found them to be effective or practicable. Considering the many facets of this problem it would appear that the use of an immunizing agent to protect these workers is the most logical and practical way to reduce the incidence of anthrax. Such a material has been developed by the United States Army Chemical Corps (Wright *et al* 1954) and is being field tested in some mills.

It has been hypothesized that many workers in these factories develop immunity to anthrax by virtue of some kind of subclinical infection in this highly contaminated environment. FIGURE 6 shows one of the reasons for this view. This graph shows the number of years 46 persons who developed anthrax had worked in the industry before developing the disease. The largest number of cases for a single year was observed in the first year of employment and more than one half the cases occurred in persons who had worked less than 4 years. However it has always been difficult to explain cases occurring after 15 to 20 years of contact with this environment. In an attempt to answer this question we obtained data on the length of employment of all the workers in these factories. From this information it has been possible to estimate the composition of the exposed population over this period and to determine the relation of the length of employment to the attack rate. The graphic distribution of the length of employment of all workers (FIGURE 6) was similar to that of the




FIGURE 6 The relation of length of employment in a goat hair mill to the development of anthrax




FIGURE 5 The relation of the risk of developing anthrax to the contamination of surfaces with *Bacillus anthracis* by department in three mills 1954 to 1956

shown that wool processing mills are less contaminated with the spores of *B. anthracis* than are goat hair mills

TABLE 2 also summarizes the results of our studies on different occasions of the surface contamination of several goat hair mills with the spores of *B. anthracis*. Moist sterile cotton swabs were used to take cultures from floors and walls and from machinery and other equipment. The swabs were then used to inoculate blood agar plates that were incubated for 18 hours at 31° C. Colonies grossly and microscopically resembling *B. anthracis* were tested for mouse virulence and lysis by the gamma bacteriophage of Brown and Cherry (1955). Forty four per cent of all samples in this series were positive for *B. anthracis* indicating a relatively high contamination of the mills with this pathogen. The greatest numbers were isolated from the early dusty stages. FIGURE 5 shows that the departmental attack rate in these mills bears a direct relation to the level of surface contamination.

That contact by man with the organism does not necessarily lead to disease is demonstrated by the recent report of Carr and Rew (1951) who were able to recover *B. anthracis* from the nose or pharynx of 14 of 101 healthy workers in 2 of these mills. Again most of the recoveries were from persons working in the early stages of production. While the transmission of anthrax either from infected persons or workers such as these has never been documented, cases have been recorded in which asymptomatic workers might have been responsible. Transmission via the clothing of these workers is another likely explanation of these incidents.

It might be inferred from FIGURE 5 that if the surface contamination could be reduced to the 10 per cent level a negligible attack rate might prevail. Our experience tends to confirm the importance of plant hygiene. However it is probable that both the level of plant contamination and the attack rate are dependent upon the more important factor of hair contamination. Attempts to decontaminate the hair have met with many difficulties. In England while the establishment of the Wool Disinfection Station at Liverpool and the passage of governmental regulations have satisfactorily controlled anthrax, such a plan does not seem applicable in the United States. The initial cost of the equipment would be enormous; it would be difficult to decide where it




FIGURE 7 Anthrax in two related goat hair mills

it was possibly more contaminated. Changes were made in the method of steam sterilization so as to ensure better penetration of steam to the center of the ball of hair top to reduce the possibility of error. Plant E was again washed with disinfectant. Since that time occasional cases have been diagnosed at Plant D (which uses unsterilized hair) but no further cases have been noted at its subsidiary. In November 1956 3 per cent of the surface samples taken at the subsidiary plant were positive for *B. anthracis*. If this low level of contamination can be maintained the number of cases of anthrax diagnosed there will probably be small.

Summary

The number of cases of anthrax in humans in the United States has steadily declined to the present level of 40 to 50 cases per year. Practically all diagnosed anthrax is now the cutaneous form and deaths from the disease have been rare since the introduction of antibiotics. The majority of cases reported are industrial in origin and they can most frequently be traced to contact with imported goat hair. The further reduction of the incidence of anthrax presents many problems but adequate ventilation and high standards of cleanliness in the mills seem to play an important role in the control of anthrax. The risk of contracting anthrax is greatest in the early steps in the processing of goat hair and is greater than the risk associated with wool processing. Prolonged exposure to the organism does not appear to lead to the development of subclinical infection or immunity to anthrax.

Acknowledgment

We are greatly indebted to Ida Sherman of the Communicable Disease Center for furnishing statistical advice and data on the incidence of anthrax. Much of the material discussed in this paper could not have been obtained

TABLE 3

RELATION OF THE LENGTH OF EMPLOYMENT TO THE DEVELOPMENT OF ANTHRAX IN FOUR COMPANIES PROCESSING GOAT HAIR 1945 TO 1956

Number of years employment	Cases of anthrax per year	Number of employees	Attack rate cases per 100 employees per year
0 to 4	9.8	546	1.8
5 to 8	3.4	262	1.3
9 to 12	1.9	209	0.9
13 to 16	1.1	119	0.9
17 to 20	1.1	90	1.2
Entire group	17.3	1226	1.4

anthrax cases for the majority of workers had been employed in this industry less than 5 years. TABLE 3 shows the attack rates for varying periods of employment. These rates ranged from 1.8 cases per 100 employees per year in the first 4 years to 0.9 for those who had worked for 9 to 16 years. The rate for 17 to 20 years' employment was 1.2 cases per 100 per year. None of these rates are significantly different. In addition they show much less variation than was observed between departments in these plants (TABLE 2). It was our conclusion, therefore, that these workers do not develop subclinical infection or immunity to anthrax by prolonged exposure to the organism. Actually, while a second attack of anthrax in the same individual is very rare, there are reports of its occurrence (Gold 1946) and immunity to anthrax in humans has never been demonstrated conclusively.

FIGURE 7 outlines a small epidemic of anthrax that illustrates some of the problems already described. It charts the occurrence of anthrax at one company (Plant D) and its subsidiary elsewhere (Plant I). Picking, carding and combing are performed at Plant D, and then part of the hair is sent to Plant I for further processing. Plant E was established early in 1953 and a few months later the first case of cutaneous anthrax was reported there. An investigation revealed extensive contamination of the plant environment with *B. anthracis*. The plant was closed and thoroughly cleaned with 5 per cent formaldehyde. A check of environmental contamination following this did not reveal any *B. anthracis*. At the same time Plant D began to steam sterilize the hair top before sending it to the subsidiary. However, a second case of anthrax was reported from Plant E in June 1955. At that time 43 per cent of the environmental cultures there were positive for *B. anthracis*. In addition, it was realized that the steam sterilization had not been completely effective, since *B. anthracis* was recovered from the inside of several balls of hair top shipped to Plant I. In December 1955 and in the early part of 1956 an unusual number of cases were reported from both mills: 7 cases were diagnosed at Plant I and 4 at Plant D. At that time 10 per cent of the surface samples at Plant D and 28 per cent of the surface samples at Plant I were positive for *B. anthracis*. This is probably related to the fact that the company began to use goat hair from Iran late in 1955. The management felt that the Iranian hair was dirtier than the hair normally used and that accordingly

NEWCASTLE DISEASE*

By R. P. Hanson

University of Wisconsin Madison Wis.

and

C. A. Brandly

University of Illinois Urbana Ill.

Newcastle disease is an infectious highly contagious disease that afflicts chiefly chickens and turkeys. Many species of birds as well as various mammals including man may contract the infection with serious consequences. In birds Newcastle disease may vary in its manifestations from an acute fulminating form to subacute mild or clinically inapparent forms manifesting various degrees of generalized and systemic involvement especially of the respiratory and nervous systems. It has been recognized in man chiefly as an acute transient conjunctivitis with or without malaise preauricular adenitis fever and chills. Generalization of the infection with presence of the virus in the blood urine and respiratory tissues and secretions has been encountered.

Properties of the Virus

In some of its characteristics the causative virus of Newcastle disease is similar to the viruses of influenza and mumps. This virus is admirably suited to survival in nature as it is resistant to the adversities of environment including temperature putrefaction drying moisture and light that it encounters in the habitat of its most common host domestic poultry. The virus of Newcastle disease survives long exposure to a wide range of pH values—extremes of 4 to 10 at 4 to 6 °C. for 1 week or more—that are inimical to most viruses (Moses Brandly and Jones 1947). In studies of heat stability involving 31 strains of Newcastle disease virus Hanson (1949) found that all strains remained infective after 15 min. at 56 °C. 3 were innocuous after 30 min. and of the remainder 3 retained their infectivity after such heating for 180 min.

Early work by Doyle (1924) and later that of others showed that drying of Newcastle disease virus infected secretions and tissues prolonged the survival time by months or years. H. E. Rhoades (personal communication 1937) found that strain KD NJ suspended and lyophilized in horse serum remained viable at room temperature (22 to 30 °C.) for 1 year in the refrigerator (6 to 7 °C.) for 10 years the latest period tested. Another strain 11 914 lived for 1 year but not for 2 under each condition. Seventeen field strains were also viable after storage in the refrigerator (6 to 4 °C.) for 10 years.

Definitive data on the effect of light are limited. Exposure of infected organ suspensions to direct sunlight for 1 hour had no effect on the virus (Iyer 1943). Exposure of virus bearing fluid to short ultraviolet rays (1600 to 1800 Å) inactivated the strains studied in 0.8 to 1.08 seconds (Brandly *et al.* 1946a).

The literature on Newcastle disease has become so voluminous that references in this paper have been limited largely to reviews and to the selected reports that are available to the authors.

without the generous cooperation of plant managers and public health officials especially Cleon J. Gentzkow (Pennsylvania State Department of Public Health), Forrest Bumford (New Hampshire Department of Health), Milton Werrin and A. LaBocetta (Philadelphia Department of Health), Alice Broadhurst (Massachusetts Department of Labor), Herman Gold Chester, Ia. and Martin Hines (North Carolina Department of Health). The assistance of Alexander D. Langmuir is gratefully acknowledged. Leighton Cluff, Ivan L. Bennett, and C. Brooke Worth assisted in the preparation of this manuscript.

References

- BROWN, E. R. & W. B. CHERRY. 1935. Specific identification of *Bacillus anthracis* by means of a variant bacteriophage. *J. Infectious Diseases* 96: 34-39.
- CARR, E. A. & R. R. REW. 1937. Recovery of *Bacillus anthracis* from the nose and throat of apparently healthy workers. *J. Infectious Diseases* 100: 169-171.
- GOLD, H. 1946. Anthrax. In Nelson Loose Leaf Medicine 1: 641-652. Nelson & Sons, New York, N. Y.
- SMYTH, H. F. 1945. Report of the committee on industrial anthrax (anthrax in the United States 1939-1943). *Am. J. Public Health* 35: 850-858.
- STEELE, J. D. & R. J. HFLVIG. 1953. Anthrax in the United States. *Public Health Repts U. S.* 68: 616-623.
- WOLFF, A. H. & H. HFIMANN. 1951. Industrial anthrax in the United States: an epidemiologic study. *Am. J. Hyg.* 53: 80-109.
- WRIGHT, G. G. T. W. GREEN & R. G. KANODE, JR. 1954. Studies on immunity in anthrax. V. Immunizing activity of alum precipitated protective antigen. *J. Immunol.* 73: 387-391.

Discussion of the Paper

OSCAR SUSSMAN (*Bureau of Veterinary Public Health, State of New Jersey Department of Health, Trenton, N. J.*) There was a known case of anthrax originating from bones imported from abroad; these bones were utilized in the production of gelatin in a plant located in New Jersey. Apparently the procedure used for the production of this gelatin did not kill all of the anthrax spores. This is a case in which known disease organisms are present in a product that will be used as human food. My question is whether or not there is considerable danger of air contamination of the finished product which is made in the same plant in which the original infected bone material was processed?

PHILIP SIGMUND BRACHMAN. The question as to whether there is considerable hazard in air contamination of the finished gelatin product is related directly to the degree of contamination of the processed dried bones to the types of equipment used and to the ventilation of the plant as well as to the stages in the manufacturing process at which the actual grinding and mixing take place. These factors will be all important in determining the type of aerosol that may be established in such a plant. Certainly if conditions are favorable it would be possible for the finished gelatin to be contaminated with bacteriological organisms originating from the raw material via the air. However, one cannot hazard a guess as to the probability of such an occurrence without more information about the plant, about the raw material and about other factors already cited.

NEWCASTLE DISEASE*

Py R I Hanson

University of Wisconsin-Madison, Wis.

and

C A Brandly

University of Illinois-Urbana, Ill.

Newcastle disease is an infectious highly contagious disease that afflicts chiefly chickens and turkeys. Many species of birds as well as various mammals including man may contract the infection with serious consequences. In birds Newcastle disease may vary in its manifestations from an acute fulminating form to subacute mild or clinically inapparent forms manifesting various degrees of generalized and systemic involvement especially of the respiratory and nervous systems. It has been recognized in man chiefly as an acute transient conjunctivitis with or without malaise, preauricular adenitis, fever and chills. Generalization of the infection with presence of the virus in the blood, urine and respiratory tissues and secretions has been encountered.

Properties of the Virus

In some of its characteristics the causative virus of Newcastle disease is similar to the viruses of influenza and mumps. This virus is admirably suited to survival in nature as it is resistant to the adversities of environment including temperature, putrefaction, drying, moisture and light that it encounters in the habitat of its most common host, domestic poultry. The virus of Newcastle disease survives long exposure to a wide range of pH values—extremes of 4 to 10 at 4 to 6 °C for 1 week or more—that are inimical to most viruses (Moses, Brandly and Jones 1941). In studies of heat stability involving 31 strains of Newcastle disease virus, Hanson (1949) found that all strains remained infective after 15 min at 56 °C; 3 were innocuous after 30 min and of the remainder 3 retained their infectivity after such heating for 180 min.

Early work by Doyle (1927) and later that of others showed that drying of Newcastle disease virus infected secretions and tissues prolonged the survival time by months or years. H. E. Rhoades (personal communication 1937) found that strain KD N₁ suspended and lyophilized in horse serum remained viable at room temperature (22 to 30 °C) for 1 year, in the refrigerator (6 to 7 °C) for 10 years, the latest period tested. Another strain 11 914 lived for 1 year but not for 2 under each condition. Seventeen field strains were also viable after storage in the refrigerator (6 to 7 °C) for 10 years.

Definitive data on the effect of light are limited. Exposure of infected organ suspensions to direct sunlight for 1 hour had no effect on the virus (Iyer 1943). Exposure of virus bearing fluid to short ultraviolet rays (1600 to 1800 Å) inactivated the strains studied in 0.8 to 1.08 seconds (Brandly *et al.* 1946a).

The literature on Newcastle disease has become so voluminous that references in this paper have been limited largely to reviews and to the selected reports that are available to the authors.

The tenacity of Newcastle disease virus under poultry house environments is well demonstrated by Dobson's experience (1939). One-day-old chicks became infected after being placed in 'dirty' battery brooders 7 weeks after removal of Newcastle disease infected chickens. Chickens placed in an uncleaned pen 2 weeks after the removal of fowls sick with Newcastle disease developed specific viral antibodies without prior overt signs (Jungherr 1948). Newcastle disease virus remained viable for at least 6 months in the bone marrow and muscle tissue of fowl carcasses held under trade chilling conditions (Doyle 1933). The skins of both dressed and plucked infected carcasses of market poultry stored at 24 to 36° F (-5 to 2° C) retained their infectivity for at least 60 days (Gordon Reid and Asplin 1948). Similar studies with carcasses and tissues of artificially infected fowl showed that Newcastle disease virus survived in them for 98 to 160 days (Asplin, 1949). According to Olesiuk (1951) strain 9251 survived in fresh eggs under certain storage conditions for the following periods: incubator 126 days, normal room 235 days, hen house 255 days, refrigerator 538 days (last test). The virus remained viable on chick down for the following numbers of days: incubator 87, normal room 192, hen house 255 (last test), refrigerator 538, freezer chest 538 (last test).

Substantial resistance of Newcastle disease virus to destruction by various chemical agents commonly employed for disinfection including the phenolic compounds and formalin as suggested by the work of Doyle (1921) and Farnas (1930) has been verified by more critical later studies (Tilley and Anderson 1947, Cunningham 1948, Moses 1948, Beamer and Prier 1950). Triethylene glycol vapors in saturated concentrations at 23° C and 50 per cent relative humidity were found to be quickly lethal to Newcastle disease virus aerosols (Krugman and Sverdlov 1949) yet in trials in a large broiler plant (Ellis Brandly and Hanson 1952) and with artificial infection among chicks (University of Illinois unpublished data) triethylene glycol vapors failed to prevent the spread of even benign Newcastle disease infection.

The properties of virulence and pathogenicity which are inseparable in the complex interaction between the host and its viral parasite are paramount in assessing the significance and potential of Newcastle disease. Certain strains of Newcastle disease virus are lethal for chickens, embryonating avian eggs and sometimes for various other species whereas other strains which may be variants of the lethal ones are of low or indifferent pathogenicity. Prior or concurrent devitalizing factors including environment and other diseases may increase the susceptibility of the host and later the severity of Newcastle disease (Francis and Kish 1952, Sinha, Hanson and Brandly 1957). This may be seen in the markedly adverse reactions resulting from complications of vaccinal infection (Brandly and Winslow 1947, McDougale 1948, Binns, Nelsen and Miner 1949). Age as a susceptibility factor is well recognized in mammals (Upton *et al.* 1955) as well as in poultry (Brandly 1950 and elsewhere). Nevertheless deviations from the highly fatal fulminating disease that identified the initial and early outbreaks of Newcastle disease (Doyle 1927, Kraneveld 1926) were not recognized until several years after the first appearance of Newcastle disease in the United States (Brandly 1953, Beach

1944 Beaudette and Hudson 1956) Wild strains of low pathogenicity identified later include the low form from England (Anon 1948) the English F strain (Asplin 1947) the B₁ or Blacksburg strain (Hitchner Reising and Van Roekel 1951) one associated with an outbreak in Germany (Hilbrich 1952) and the Roakin strain (Beaudette Bivins and Miller 1949)

Markedly emphasizing the great variability in pathogenicity and other properties of Newcastle disease virus were the differences found by Hanson (1949) among various strains of Newcastle disease virus. Among 18 strains studied pathogenicity varied from 0 to 10⁶ LD₅₀ doses per 0.5 ml. of infective allanto-innocuous fluid and the tropisms varied from low to high degree of pneumal visceral and neural. Aside from its capacity to invade and involve the central nervous system of the postnatal fowl and mammal Newcastle disease virus was shown to produce developmental defects in chicken embryos following inoculation of the embryonating egg (Williamson Blattner and Robertson 1953)

Differences or changes in tissue tropisms among strains of Newcastle disease virus can account for variations in such properties as invasiveness of the virus and in the signs, course and pathological changes resulting from Newcastle disease infection (Jungherr and Terrell 1946). Furthermore evidence was advanced to indicate that aside from individual and host factors differences in pathogenicity and infective dosage, route of infection and particularly degrees of partial immunity prior to infection act as principal modifiers of pathologic expressions of the disease.

The established tenacity of Newcastle disease virus obviously contributes significantly to its perpetuation and spread. In addition its pneumotropic affinities favor air borne liberation especially via infective droplets and particles from the respiratory system. Infection by way of the alimentary tract with eggs and through coitus together with elimination via all body orifices and from the integument during life of fowl afflicted with Newcastle disease further assures extensive dissemination of the virus. With ready spread of often concentrated infection there is provided opportunity which history indicates Newcastle disease virus has utilized efficiently for variation and mutation. The latter characteristic reinforced by the multiple tissue tropisms of Newcastle disease virus can account in part at least for its recognized adaptability to host species other than fowl.

Continued frequent contact via respiratory conjunctival and other routes may facilitate development of Newcastle disease to person to person contagions and among individuals of other mammalian species (Brandly 1950 1952 Quinn *et al.* 1952 Kyle MacFadden and Gunderson 1952). Evans (1955) pointed out that emergence of variants of Newcastle disease virus caused the respiratory form of the disease in man and may result in man to man transmission of the infection in the future.

World Wide Dissemination of the Virus

The recorded history of Newcastle disease emphasizes the consequences of lack of sanitary regulation and control of the disease upon its further dissemina-

tion from its point of origin in Indonesia, where it was described in 1926 as a previously unknown fulminating malady. In England Newcastle disease was summarily eradicated by slaughter and terminal disinfection following its first outbreak there in the same year (Doyle, 1927). During the decade following recognition of what is now generally called Newcastle disease the malady was reported from India, the Philippines, Japan, Korea, Australia, Ceylon, and Kenya. Subsequently it was recognized in Syria, Palestine, and the Middle Congo and during World War II it spread to Europe by way of Sicily and continental Italy. It was first identified in the United States in 1944 (Brandly *et al.*, 1944, 1946; Beach, 1944) but it may have found its way there as early as 1935 (Beach, 1944; Beaudette and Hudson, 1956). The present world wide diffusion of Newcastle disease was inevitable in the absence of land and sanitary barriers between adjacent areas or across water and air routes of transport. Inevitable also is the continued opportunity for Newcastle disease virus to adapt to new animal hosts including man.

The identity of Newcastle disease in its initial appearance in these and later foci was seldom recognized with the result that many synonyms including pseudofowl pest, *atypische geflügelpest*, pseudopoultry plague, avian pest, *Ranikhet*, avian distemper, and pneumoencephalitis have been recorded (Brandly *et al.*, 1946). Handicapped by lack of definitive laboratory diagnostic procedures and by the differences in clinicopathological features of the disease as it appeared in new areas, recognition of the malady and application of control measures were frequently delayed. The delay in suspecting and identifying Newcastle disease after it had invaded the United States can be ascribed largely to the previously undescribed features of the disease, namely, low mortality and predominance of respiratory and nervous signs.

The low mortality rate was also responsible in large part for the attitude of the poultry industry of the United States in opposing the accepted principle of eradicating newly introduced exotic maladies as was done with foot and mouth disease. Nevertheless the development of mild or subclinical expressions of the disease, whether through change in the virus, the host, or both, make it more difficult to detect and, especially, to eradicate Newcastle disease being a contagion that is highly diffusible.

Source of Infection

A primary source of infection for chickens and one source of infection for man is the chicken in the initial stage of disease. Beginning as early as 24 hours before the development of respiratory signs and continuing for several days thereafter, diseased birds discharge virus into the air that they exhale (Sinha, Hanson, and Brandly, 1952). The virus has been recovered from small samples of air taken within a few inches of the heads of such birds and it also has been obtained from the air in poultry houses in which a number of diseased birds were present (DeLay *et al.*, 1948; DeOme *et al.*, 1948). The respiratory discharges contribute almost all of the virus that exists in the form of the natural aerosol to which poultrymen are exposed. Fecal discharges, except in the instance of infection with one of a few exotic strains, are relatively free of virus.

Artificial aerosols are also a considerable hazard to man. Fluid virus is sprayed and nebulized as a means of vaccinating chickens (Johnson and Cross 1951 Hitchner and Keising 1952). Virus dust preparations are used for the same purpose. Those who provide vaccinating service to poultrymen and poultrymen who do their own vaccinating are subject to exposure to these aerosols for periods varying from a few minutes to several hours and at intervals as frequent as 3 times weekly and as infrequent as once or twice a year. Strains of low pathogenicity for chickens are used for vaccines but it has not been shown that these strains are less hazardous for man (Quinn Hanson and Brown 1952).

The laboratories in which the vaccines are produced and in which the virus is studied provide other and important opportunities for exposure of man to viral aerosols. The virus is isolated in embryonating chicken eggs for diagnosis and is propagated in them for the production of vaccine. The shell is broken in harvesting, the infective fluids and tissues and the desired materials are removed with a suction device and transferred to mixing containers. The virus material is ground, centrifuged and lyophilized before packaging. Aerosols may be created in transferring, grinding or centrifuging operations (Stein and Rogers 1950). When lyophilization is done in a bulk drum the dried virus is ground to a fine powder and distributed in measured quantities to the final vial. This process exposes the operator to a high and continuous virus aerosol (Evans 1954). Such individuals usually have good serologic titers.

Even in the incubative stages of the disease the carcass of the chicken contains some virus (Hofstad 1951). The concentration is the highest in the viscera particularly in the lungs and spleen although detectable quantities may be found in the blood and even in the muscle. Individuals working in eviscerating plants and in diagnostic laboratories are exposed when handling such materials. Virus may be carried to the eyes or nose by the spattering of blood and tissue substances or it may be transferred to the eye by contaminated hands. The edible carcass should under usual conditions of marketing be quite free of virus and an instance of human infection by the ingestion or handling of dressed poultry is unknown.

Diseased carcasses and particularly viscera may be a source of infection to chickens (Hess 1951 Lloyd 1952). It is possible that scavengers among rodents and carnivores may distribute the disease by eating dead chickens and then contaminating food supplies with their excreta (Perci and Silvagni 1955 Bolin 1948 Walker and McKercher 1954).

Since a viremia occurs in Newcastle disease of chickens research workers have suspected that biting arthropods may play some role in the epizootiology of the disease. Little support has been found for this idea. Lice and mites do become engorged with the virus and it has been detected in them by grinding and injecting material into eggs but no virus has been transferred by the bites of arthropods (Bolin 1948 Hofstad 1949). Furthermore instances in which Diptera were reported to transmit the disease have not been confirmed (Lucard 1934). It is doubtful that man is exposed to the virus by the agency of arthropods.

Disease in Man

Burnet (1943) was the first to report the pathogenicity of Newcastle disease for man. An individual harvesting virus from an embryonated egg in his laboratory was infected by material that splashed into an eye. Conjunctivitis developed within 24 hours and the virus was isolated from washings obtained from the affected eye. Most of the subsequent cases in man have followed a similar course with a mild to severe conjunctivitis and spontaneous recovery.

More than 100 cases have been observed since this initial one and numerous reports have appeared in the literature (TABLE 1). The incubation period is usually 1 or 2 days but it may occasionally be 3 or 4 days. The initial signs of infection of the eye are irritation and lacrimation. The eye appears red from the injection and swelling of the subconjunctival tissues. The inflammation may become more severe in the next day or two. The lid may also become edematous and the preauricular lymph node may become palpable. Nelson *et al.* (1952) who observed 40 cases in man reported that the preauricular adenitis developed on the infected side of 70 per cent of the patients. They found only one eye infected in almost all of the cases and observed that constitutional symptoms were rare. The conjunctivitis lasts for 3 or 4 days in most patients but courses of 7 to 8 days and even of 21 days have been reported.

A description of cytoplasmic inclusions presumed to be pathognomonic in cells obtained from conjunctival scrapings on the third, fourth and fifth days (Keeney and Hunter 1950) has not been confirmed by other observers (Orlandella 1955). Blood changes do not appear to be consistent. The cornea, the extraocular movements and the visual field remain normal and the pupil reacts normally. Recovery is usually uneventful and complete within a week or two. Occasionally subsequent impairment of visual accommodations has been reported.

Virus has been isolated from conjunctival washings obtained on the second, third, fourth and fifth days after the onset of symptoms. In some instances virus has been isolated from other secretions and tissues. Antibodies have been detected as early as 1 day after onset but more frequently 14 to 21 days later. The immunological response is variable. Some individuals from whom the virus has been isolated do not develop hemagglutination inhibition or serum neutralizing antibodies. In others hemagglutination titers of 1:16 to 1:160 and serum neutralizing titers of 1:100 to 1:100,000 have been reported. Among employees of a chicken processing plant Nelson *et al.* (1952) found that 64 per cent of the exposed individuals not reporting conjunctivitis had high neutralizing titers and only 25 per cent of those reporting conjunctivitis had high neutralizing titers. None of the unexposed controls had significant titers.

A generalized infection of man has been reported by numerous observers (TABLE 1). In some instances it has been a mild influenza like infection characterized by rise in body temperature of 2 or 3 °F with chills and pharyngitis (Mitchell and Walker 1951). Conjunctivitis may or may not be associated with this syndrome (Quinn *et al.* 1952). The infection lasts 3 to 4 days. The exposure of most individuals in whom the generalized infection has devel-

TABLE I
NEWCASTLE DISEASE OF MAN

		Year	Notes
BURNET F M	Med J Australia 2 313	1943	first isolation
ANDERSON I I	Med J Australia 2 371	1946	
SHIMKIN A I	Brit J Ophthalmol 30 260	1946	
LATOW J	Harefuah 30 47	1946	17 clinical cases
NADRAY M	Ophthalmol 113 106	1947	
HOWITT B I J & BISHOP & R F KISSLING	Am J Public Health 38 1263	1948	(see Howitt 1950)
HURT L M	Los Angeles I D Rept	1948	
KUSJMGIEA J	Zoogroßkass 3 76	1948	
POMEROY B S & R FENSTER MACHER	U S Egg Poultry 64 19	1948	
BARWELL M B M LE GIER T MURRAY W SCHOFIELD & G O BROWN	J Lab Clin Med 34 1591	1949	isolation from lung
BERKE Z & S B GOLFEN	Türk İylen Tecrübi Biol Dergisi 9 1	1949	
FREYMANN M W & I B BANG	Bull Johns Hopkins Hosp 84 409	1949	
JACALLS W L & A MAHONEY	Am J Public Health 39 737	1949	
MCGOLGH I F	Ohio State Med J 45 25	1949	later invalidated
SINKOVITS J	Kisérletes O v ostul mány 1 134	1949	
HOWITT B I	J Immunol 64 73	1950	(see Howitt 1948)
JACOBY H A VALLER & A LE PRIOL	Bull acad natl m'l Paris 134 106	1950	
KEENEY A H & M C HUNTER	A M A Arch Ophthalmol 44 573	1950	
LÉPINE P P ATANASU & H G GABREAU	Rev m d vet y parasitol 6 1054	1950	
SHEFRAN G A	J Am Vet Med Assoc 66 219	1950	
IVANS A S	J Immunol 67 529	1951	serologic survey
CLYDESON D P & H T MOSES	J Am Vet Med Assoc 118 1	1951	
HUNTER M C A H KEENEY & M M SICE	J Infectious Diseases 88 272	1951	(see Keeneey 1950)
LATTE B	Bull Ocul 30 553	1951	
MELNICK J L & V J EDINBO	Am J Hyg 64 351	1951	serologic survey
MITCHELL C A & R V L WALKER	Can J Comp Med Vet Sci 15 226	1951	influenza like disease
DIV A & A FI	Bol inst invest vet 4 644	1952	case of parotiditis
LEJUEM A & J DAI 4ET	Ann inst Pasteur 63 407	1952	clinical survey
LIPPMAN O	Am J Ophthalmol 36 104	1952	
MOLTER S E & J CLARK	A M A Arch Internal Med 89 20	1952	case of anemina
NELLS A C B B S POMEROY B SCHRALL W F PARKS & R J JENDENAN	Am J Public Health 42 672	1952	outbreak in a processing plant

Disease in Man

Burnet (1943) was the first to report the pathogenicity of Newcastle disease for man. An individual harvesting virus from an embryonated egg in his laboratory was infected by material that splashed into an eye. Conjunctivitis developed within 24 hours and the virus was isolated from washings obtained from the affected eye. Most of the subsequent cases in man have followed a similar course with a mild to severe conjunctivitis and spontaneous recovery.

More than 100 cases have been observed since this initial one and numerous reports have appeared in the literature (TABLE 1). The incubation period is usually 1 or 2 days but it may occasionally be 3 or 4 days. The initial signs of infection of the eye are irritation and lacrimation. The eye appears red from the injection and swelling of the subconjunctival tissues. The inflammation may become more severe in the next day or two. The lid may also become edematous and the preauricular lymph node may become palpable. Nelson *et al* (1952), who observed 40 cases in man, reported that the preauricular adenitis developed on the infected side of 50 per cent of the patients. They found only one eye infected in almost all of the cases, and observed that constitutional symptoms were rare. The conjunctivitis lasts for 3 or 4 days in most patients but courses of 7 to 8 days and even of 21 days have been reported.

A description of cytoplasmic inclusions presumed to be pathognomonic in cells obtained from conjunctival scrapings on the third, fourth, and fifth days (Keeney and Hunter 1950) has not been confirmed by other observers (Orlandella 1955). Blood changes do not appear to be consistent. The cornea, the extraocular movements, and the visual field remain normal and the pupil reacts normally. Recovery is usually uneventful and complete within a week or two. Occasionally subsequent impairment of visual accommodations has been reported.

Virus has been isolated from conjunctival washings obtained on the second, third, fourth, and fifth days after the onset of symptoms. In some instances virus has been isolated from other secretions and tissues. Antibodies have been detected as early as 7 days after onset but more frequently 14 to 21 days later. The immunological response is variable. Some individuals from whom the virus has been isolated do not develop hemagglutination inhibition or serum neutralizing antibodies. In others hemagglutination titers of 1:16 to 1:160 and serum neutralizing titers of 1:100 to 1:100,000 have been reported. Among employees of a chicken processing plant, Nelson *et al* (1952) found that 64 per cent of the exposed individuals not reporting conjunctivitis had high neutralizing titers and only 25 per cent of those reporting conjunctivitis had high neutralizing titers. None of the unexposed controls had significant titers.

A generalized infection of man has been reported by numerous observers (TABLE 1). In some instances it has been a mild influenza-like infection characterized by rise in body temperature of 2 or 3 °F with chills and pharyngitis (Mitchell and Walker 1951). Conjunctivitis may or may not be associated with this syndrome (Quinn *et al* 1952). The infection lasts 3 to 4 days. The exposure of most individuals in whom the generalized infection has devel-

TABLE I
NEWCASTLE DISEASE OF MAN

		Year	Notes
BURNET F M	Med J Australia 2 313	1943	first isolation
ANDERSON E E	Med J Australia 2 31	1946	
SHIMKIN N I	Brit J Ophthalmol 30 760	1947	
YATOM J	Harefuah 30 4	1946	1 clinical cases
RADRAY M	Ophthalmol 113 106	194	
HOWITT B F L K BISHOP & R E KLING	Am J Public Health 35 1263	1948	(see Howitt 1950)
HURT L M	Los Angeles L D Rept	1948	
KUJUMGIEN J	Zoogonomas 3 6	1948	
POMEROY B S & R FENNER	L S Egg Poultry 54 19	1948	
WACHS			
BARTWELL M B M LUGHER F	J Lab Clin Med 34 1381	1949	isolation from lung
MCKENNA W SCHOTFIELD & G O BROWN			
BERKE Z & S B GOLEN	Turk J en Tecribi B ol Dergisi 9 1	1949	
FREYMAN M W & F B BANG	Bull Johns Hopkins Hosp 84 409	1949	
INGALLS W L & A MAHONEY	Am J Public Health 39 3	1949	
MCGOUGH I F	Ohio State Med J 45 25	1949	later invalidated
SINKOVITS J	Kisrletes Orvostudomány 1 134	1949	
HOWITT B F	J Immunol 64 3	1950	(see Howitt 1948)
JACOTOT H, A VALLÉE & A LE PRIOL	Bull acad natl méd Paris 134 107	1950	
KEENEY A H & M C HUNTER	A M A Arch. Ophthalmol 44 53	1950	
LÉPINE P, P ATANASIC & H G GARREAU	Rev med vet v parasitol 5 1054	1950	
SHEERAN G A	J Am Vet. Med Assoc 66 719	1950	
EVANS A S	J Immunol 67 579	1951	serologic survey
GUSTAFSON D P & H E MOSES	J Am. Vet. Med Assoc. 118 1	1951	
HUNTER M C A H KEENEY & M M SICE	J Infectious Diseases 83 27	1951	(see Keenev 1950)
LATTE B	Bull. Ocul 30 553	1951	
MELNICK J L & N LEDINCO	Am J Hyg 54 354	1951	serologic survey
MITCHELL C A & R V L WALKER	Can J Comp Med Vet Sci 15 226	1951	influenza like disease
DINO A & A FEGO	Bol inst. invest. vet 4 644	1952	case of paratuberculosis
ENQUEM A & J DAUS ET	Ann inst. Pasteur 83 407	1952	clinical survey
LIPPMAN O	Am J Ophthalmol 35 104	1952	
MCKINTY S E & E CLARK	A M A Arch. Internal Med 89 20	1952	case of anemia
NELSON C B B S POMEROY K SCHRAILL W E PARKS & R J LINDEMANN	Am J Public Health 42 62	1952	outbreak in a processing plant

TABLE 1—Continued

		Year	Notes
NEGRI R A D'AMORE & L RAVAIOLI	Attisoc ital sci vet 6 567	1957	
QUINN R W R P HANSON J W BROWN & C A BRANDLY	J Lab Clin Med 40 736	1952	isolation from blood and urine
SLONIM D & V STRANUKOVA	Bull Hyg 27 708	1952	
TOPOLNIA F & A H BEGANOVIC	Vet Arch 21 572	1952	
WENNER H A M H JENSON & A MONLEY	J Immunol 68 343	1952	
BIELING R & O SIEGMANN	Deut med Wochschr 78 1037	1953	
MITCHELL C A	Proc Book Am Vet Med Assoc 432	1953	encephalitis like disease (see Moolten 1952)
MOOLTEN S E F CLARK B F GLASSER E KATZ & B S MILLER	Am J Med 14 294	1953	
NEGRI R A D'AMORE & L RAVAIOLI	Rend ist super sanità 16 183	1953	
PASCUAL	J Am Med Assoc 151 567	1953	case of jejunitis
RODOT M	Thesis Paris (Alfort)	1953	
DINTER Z & K BAKOS	Nord Med 51 853	1954	
EVANS A S	Am J Hyg 60 204	1954	
SCHOOP G	Deut tierärztl Wochschr 61 167	1954	
SIEGERT R H G HAUSMANN & E MANNWEILER	Klin Wochschr 32 8	1954	
AMADEO G	Caz Vet 3 17	1955	
ANON	Zootria Chile 4 75	1955	
EVANS A S	Am J Public Health 45 742	1955	
FRANCO A S	Bol inform S y V 1 7 3	1955	
JACOTOT H A VALLÉE & A LE PRIOL	Ann inst Pasteur 88 111	1955	
ORLANDO L A	Nuova vet 31 206	1955	
VETTERLEIN W	Deut Gesundheitsw 10 1327	1955	

oped has been through an aerosol of the virus. Isolation of virus has been made from nasopharyngeal washings saliva blood and urine. Bawell *et al* (1949) also isolated the virus from lung tissue obtained by biopsy.

Several other forms of this disease have been suggested. Moolten and Clark (1952) reported a hemolytic anemia. The virus was isolated from the blood of three patients over an extended period. Recovery took several months. A Spanish investigator (Pascual 1953) has described a jejunitis associated with Newcastle disease infection of man. Invasion of the central nervous system with consequent production of encephalitis was initially suspected by Howitt, Bishop, and Kissling (1948) upon serologic grounds that were subsequently found to be untenable (Howitt 1950). Canadian workers have described a mild encephalitis in a man from whom Newcastle disease virus was isolated (Mitchell 1953). Wenner, Montley, and Todd (1950) have shown that the virus is capable of producing fatal encephalitis in rhesus monkeys if inoculated into the central nervous system but that the virus is unable

to invade the central nervous system through the blood system or even along the peripheral nerves. While there is considerable evidence that systemic infection of man sometimes occurs some of the reports are open to criticism (Morgan 1952).

The immunology of Newcastle disease in man is perhaps the most controversial aspect of the disease. In chickens the virus induces the production of antibodies that inhibit the hemagglutinin of the virus and neutralize its infectivity for a susceptible host. Hemagglutination inhibition titers of 1:1280 and serum neutralizing titers of 1:1 000 000 are not uncommon and they develop within 14 to 21 days after the initiation of infection. Birds with circulating antibodies are immune to challenge. Specific antibodies detectable by hemagglutination inhibition, serum neutralization and complement fixation tests may be produced in rabbits, cattle and horses. While some serologic differences exist among strains of the virus all strains cross react to a considerable degree with antibodies produced to heterologous strains.

In man infection may or may not result in the production of antibody detectable by hemagglutination inhibition and serum neutralization tests. Man may or may not be refractory to a second exposure to the virus (Nelson *et al* 1952). Reinfection has been reported within 3 or 4 months of the initial infection. Further complicating the problem are the nonspecific neutralizing substances that may be present in the serum of man. Howitt (1950) has described a heat labile neutralizing substance found in many human sera. This substance is destroyed at 56 C in 30 minutes and it is also irreversibly fixed to filter paper upon which the serum has been dried. A second heat stable substance that neutralizes Newcastle disease virus and inhibits the hemagglutinin is found in some sera. It is associated with the presence of neutralizing antibodies for mumps virus. Wenner, Jensen and Monley (1952) contend that the substance may be partially inactivated by heating at 56 C for 30 minutes and that accordingly it is not specific and is not an antibody shared by mumps and Newcastle disease. Evans (1955) utilized cross adsorption procedures and found that in human sera that neutralized both mumps and Newcastle disease virus there were two possibilities: (1) Newcastle disease virus alone could adsorb what was Newcastle disease virus antibody and (2) either the Newcastle disease virus or mumps virus could adsorb a Newcastle disease neutralizing substance. It has been shown repeatedly that Newcastle disease antibody will not neutralize mumps virus and that man or animals immunized against a single strain of mumps virus fail to produce substances capable of neutralizing Newcastle disease virus. Nevertheless such cross reactions are common during convalescence from naturally acquired mumps infection. Evans (1955) interprets this to mean that more than one strain of mumps virus exists and that at least one of these has an antigen that is common or related to those present in some strains of Newcastle disease virus. One factor that had contributed to the controversy is the lack of uniformity in the procedure by which the neutralization test is conducted. Investigators of the disease in birds have found that serial dilutions of virus and undiluted serum gave a sensitive and reproducible result while the reverse

TABLE 1—Continued

		Year	Notes
NEGRI R A D'AMORE & L RAVAIOLI	Attisoc ital sci vet 6 562	1952	
QUINN R W R I HANSON J W BROWN & C A BRANDLY	J Lab Clin Med 40 736	1952	isolation from blood and urine
SLONIM D & V STRANUKOVA	Bull Hyg 27 108	1952	
TOPOLNIK I & A H BEGANOVIC	Vet Arch 21 572	1952	
WENNER H A M H JENSON & A MONTLEY	J Immunol 68 343	1952	
BIELING R & O SIEGMANN	Deut med Wochschr 78 1037	1953	
MITCHELL C A	Proc Book Am Vet Med Assoc 432	1953	encephalitis like dis- ease (see Moolten 1952)
MOOLTEN S E F CLARK B F GLASSER I KATZ & B S MIL- LER	Am J Med 14 294	1953	
NEGRI R A D'AMORE & L RAVAIOLI	Rend ist super sanit 16 183	1953	
PASCUAL	J Am Med Assoc 151 567	1953	case of je- junitis
RODOT M	Thesis Paris (Alfort)	1953	
DINTER Z & K BAKOS	Nord Med 51 853	1954	
EVANS A S	Am J Hyg 60 204	1954	
SCHOOP G	Deut tierarztl Wochschr 61 167	1954	
SIEGERT R H G HALSMANN & I MANNWEILER	Klin Wochschr 32 8	1954	
AMADEO G	Gaz Vet 3 17	1955	
ANON	Zootria Chile 4 25	1955	
EVANS A S	Am J Public Health 45 742	1955	
FRANCO A S	Bol inform S y V A 7 3	1955	
JACOTOT H A VALLÉE & A LE PRIOL	Ann inst Pasteur 88 111	1955	
ORLANDELLA V	Nuova vet 31 206	1955	
VETTERLEIN W	Deut Gesundheitsw 10 1327	1955	

oped has been through an aerosol of the virus. Isolation of virus has been made from nasopharyngeal washings saliva blood and urine. Bawell *et al* (1949) also isolated the virus from lung tissue obtained by biopsy.

Several other forms of this disease have been suggested. Moolten and Clark (1952) reported a hemolytic anemia. The virus was isolated from the blood of three patients over an extended period. Recovery took several months. A Spanish investigator (Pascual 1953) has described a jejunitis associated with Newcastle disease infection of man. Invasion of the central nervous system with consequent production of encephalitis was initially suspected by Howitt, Bishop and Kissling (1948) upon serologic grounds that were subsequently found to be untenable (Howitt 1950). Canadian workers have described a mild encephalitis in a man from whom Newcastle disease virus was isolated (Mitchell 1953). Wenner, Montley and Todd (1950) have shown that the virus is capable of producing fatal encephalitis in rhesus monkeys if inoculated into the central nervous system but that the virus is unable

lishments on which several million chickens are raised each year to small flocks of less than 100 that are kept to supply the farmer's wife with egg money. To this population of 7 or 8 million people who feed and water chickens and clean out chicken houses must be added the several hundred thousand people engaged in the processing of poultry and a few thousand more employed in occupations in which the virus is handled. It should not be inferred that all of these people are at equal risk; almost all of the infections that have been reported have come from the two smaller groups which however are under much closer medical observation than the larger group. Every recorded instance of human infection has been an initial case, the transmission being from bird to man and not from man to man.

The greatest importance of Newcastle disease is its cost to the consumer as reflected in higher prices for poultry products and in poorer quality of the product. The disease may or may not produce a high mortality in chickens depending on the strain of the virus, the age and condition of the individual bird and certain environmental factors. A strain of virus isolated in Texas a few years ago killed 9 out of 10 chickens irrespective of age (Boney 1951). Other strains have killed only 1 of 100 chicks a day old.

A fallacy believed by many consumers is that the death of the chicken is the important loss to the producer and that if the chicken survives the disease the producer does not suffer. Actually when the disease does not kill it usually affects the chicken by interrupting its growth and egg production and sometimes by damaging the reproductive tract so that the hen lays eggs with poor shells and an inferior albumen (Biswall 1954; Parnell 1950; Knox 1950).

Interruption of growth for one week would mean added feed cost of 10 cents per bird. Such an increase could wipe out the margin of profit in broiler production and mean a net loss to the producer. Another factor adding to his loss is the increase in the number of culls that must be rejected from the dressing line. Interruption of egg production for 2 to 4 weeks is equally disastrous. To this is added the production of eggs that are downgraded and consequently sell for a lower price. All of these things are reflected in the amount of meat and eggs that are available to the consumer in the United States at a reasonable cost.

The control of animal disease in the United States has progressed successfully to complete eradication of but one poultry disease, namely fowl plague. To date there has been no serious discussion of an eradication program for Newcastle disease or even of an organized control program. When the disease was first recognized in the United States an attempt was made to stay its spread by quarantine but this approach was abandoned as vaccines came into use. The disease is now found in all sections of the country in which poultry is raised. *The incidence varies from year to year but there is no evidence of retrenchment.* Individual poultrymen attempt to protect their flocks against the infection by vaccination if the risk warrants it. This decision is based on the experience of the previous seasons and the current reports of the disease in the neighborhood. Vaccination adds about one or one and one half cents to the cost of producing a broiler. On a national basis this would amount to more

procedure, serial dilution of the serum with a constant amount of virus was unsatisfactory. Evans (1955) demonstrated that the same situation holds for human sera in a neutralization test for Newcastle disease virus. While more sera are found to contain antibody and while the titers are higher the use of heat inactivated undiluted sera in neutralization tests for Newcastle disease virus does not in all instances result in the detection of antibody following exposure.

There appears to be little question that where an opportunity for contact with Newcastle disease virus exists a considerable proportion of the humans so exposed will develop a transient conjunctivitis. Systemic infection of man may sometimes occur. The history of Newcastle disease in man with its conflicting reports gives ample reason for a questioning attitude toward systemic infection of man. Serology while sometimes helpful is undependable.

Isolation of the virus is a difficult procedure for routine diagnosis but it is much more meaningful than are serologic findings. Unfortunately isolation also can be misleading for two reasons.

First spurious isolation may occur by contamination of the tissue under test with virus already present in the laboratory in which the isolation is being made. It is possible but very unlikely that the virus might be present in the eggs used for isolation. These problems can be avoided by taking adequate precautions in the selection of procedures and of materials and by demonstrating that the virus can be reisolated from another aliquot of the original material.

The second reason for doubt is the frequently inadequate attention given to the possibility of other concurrent infections either bacterial or viral. For example the common cold may be acting simultaneously with a specific Newcastle disease infection of the eye to produce what appears to be Newcastle disease conjunctivitis associated with a generalized Newcastle disease infection. This objection can be answered only after study of many case histories reveals one or more consistent disease patterns associated with Newcastle disease infection. It is necessary that consideration always be given to the possibility that more than one clinical pattern of the disease exists since there are many strains of Newcastle disease virus and since they may be introduced by more than one route. Conjunctivitis only may occur when some strains of virus are introduced by splattering while a generalized infection may occur when other strains are introduced as aerosols.

A diagnosis of Newcastle disease of man should be dependent upon a history of contact with the virus, a short incubation period, the development of conjunctivitis or possibly influenzalike symptoms and the isolation of the virus under satisfactory conditions.

Newcastle disease of man has been a minor occupational disease limited to the people employed in the production of chickens and eggs and in processing them for the market and to those concerned with the diagnosis of the disease or the production of vaccine. The population at risk is rather sizable as there are approximately 3,418,000 farms in the United States on which poultry are kept (1954 United States Census Report). This includes huge poultry estab-

- D'OME K B R A HANSKOWSKI & P D DELAY 1948 Poultry Sci 27 (5)
 DOBSON V 1939 Proc 7th World's Poultry Congr Exposition 250-253
 DONALD T M 1917 J Comp Pathol Therap 40 144
 DONALD T M 1933 J Comp Pathol Therap 46 81
 DONALD T M 1935 J Comp Pathol Therap 48 1
 EVANS A S 1954 Am J Hyg 60 204-213
 EVANS A S 1955 Am J Public Health 45 749
 FLEMING C C L A BRANDLY & K I HANSON 1952 Poultry Sci 31 394
 FARRAS I L 1930 Philippine J Agr 2 312
 FRANCIS D W & A F KILPATRICK 1922 Poultry Sci 31 916
 GILLESPIE J H B KESSEL & J FARRICANT 1950 Cornell Vet 40 93-94
 GORDON R F J REID & F D ASKIN 1948 Proc World's Poultry Congr Exposition (Copenhagen) 1 642-650
 GUTTAPOON D I & H F MOORE 1953 Am J Vet Research 14 592-95
 HANSON K P 1949 Doctoral thesis Univ. of Malaya
 HE P 1951 Bull Off Intern Epiz 35 26-29
 HILBRICH J 1952 Deut Wirt 4 751
 HITCHNER S B G KEISING & H VAN KOPPEL 1951 Am J Vet Research 12 746
 HITCHNER S B & C REISING 1951 Proc Am Vet Med Assoc 258-264
 HOFFMANN M S 1949 Am J Vet Research 10 370-371
 HOFFMANN M S 1951 Am J Vet Research 12 334-339
 HOWITT B F I K HANSON & R F KILPATRICK 1948 Am J Public Health 38 1263-1271
 HOWITT B F 1950 J Immunol 64 73-84
 IVER S G 1943 Indian J Vet Sci 13 1
 JOHNSON T I & W B CRUICKSHANK 1951 Vet Med 46 55-57
 JUNGHER E L & A TERFILL 1946 Proc 50th Ann Meeting U S Livestock Sanitary Assoc 168
 JUNGHER E L 1948 J Am Vet Med Assoc 112 124-125
 KEEFEY A H & M C HUNTER 1950 N M J Arch Ophthalmol 44 574-580
 KNOX C W 1950 Poultry Sci 29 907-911
 KRAVITZ F C 1976 Ned Ind Med J 48 448
 KREIGMAN S & B SWERDLOW 1949 J Soc Exptl Biol Med 71 689
 KYLE T S H W MACFADDEN JR & M F GENDER JR 1957 Cited by M Galtin 1953 Communicable Disease Center Public Health Service F S A Atlanta Ga
 LLOYD I C 1952 Newcastle disease World's Poultry Sci 8 99-107
 McDONALD H L 1948 Personal communication
 MORGAN H K 1952 J Lab Clin Med 40 924
 MOSE H F 1948 J Am Vet Med Assoc 112 126
 MOSE H J C A BRANDLY & F J JONES 1947 Science 105 477
 MITCHELL C A & R V L WALKER 1951 Can J Comp Med Vet Sci 15 226-227
 MITCHELL C A 1953 Proc 80th An Vet Med Assoc 432-434
 NELSON C B B S POWEROY & SCHERLE W F PARAS & R J INDEMAN 1952 Am J Public Health 42 672-678
 OLEICK O M 1951 Am J Vet Research 12 152
 ORLANDO A 1950 Nuo x vet 31 206-208
 ORWELL E D 1950 Poultry Sci 29 153-155
 PAULAL-FOREIGN LETTERS-SPAIN 1953 J Am Med Assoc 151 (7) 516
 PICARD W K 1934 Zentr Bakteriol Parasitenkult u. Inf Dis Orig 132 440-447
 PILLAI N & T SUGANATHAN 1953 J Lab Clin Med 42 637
 QUINN R W & I K HANSON J W BROWN & C A BRANDLY 1952 J Lab Clin Med 40 736-743
 SINHA S K R P HANSON & C A BRANDLY 1952 J Infectious Diseases 91 276-282
 STEIN C D & H KOGERS 1950 Am J Vet Research 11 339-344
 TYLE P N & W A ANDERSON 1947 Vet Med 42 2-9
 TYLE P N & R F HANSON D DOW & C A BRANDLY 1955 J Infectious Diseases 96 24
 U S GOVT REPORT 1954
 WALKER R V I & I D M HERCHER 1954 Can J Comp Med Vet Sci 18 451
 WANNER H A A MONTGOMERY & R N TODD 1950 J Immunol 64 303-321
 WANNER H A M H JENSEN & A MONLEY 1952 J Immunol 68 343-356
 WILLIAMSON A I R J BLATTNER & G C ROBERTSON 1953 J Immunol 71 207

than 4 million dollars exacted as a tribute to Newcastle disease in 1956 a year in which 474 million broilers were marketed. Newcastle disease will probably remain a serious problem for some time to come. The trend to larger and larger plants for the production of eggs and poultry and the increased concentration of these plants in selected areas increase the problem of air borne infection. This increases the need for periodic vaccination a requirement that is being met by the use of aerosol products requiring less labor in their administration than did the older methods. The direction is toward fewer men in the production of poultry products but these individuals are exposed to aerosols of Newcastle disease virus more frequently and over prolonged periods.

Newcastle disease has been reported in many species of wild and domestic birds (Blayland 1951 Gillespie Kassell and Fabricant 1950 Gustafson and Moses 1953). The virus may become better adapted to sparrows or pigeons, and these birds could then serve as sources of infection to man. Whether such a virus would be more or less pathogenic for man is pure speculation. Burnet suggested in 1942 that Newcastle disease virus might be a potential human pathogen still in its avian form, since eventually it might, through selection afforded by frequent exposure develop a mammalian form capable of perpetuation in man. Burnet has postulated that such a development from an avian form to a mammalian form was the origin of influenza.

References

- ANON 1948 *Vet J* 104 273-274
 ASPLIN F D 1947 *Vet Record* 59 621-623
 ASPLIN F D 1949 *Vet Record* 61 159-160
 BAWELL M B M LE GIER F MURREY W SCHOFIELD & G O BROUN 1949 *J Lab Clin Med* 34 1581
 BEACH J K 1944 *Science* 10 361
 BEAMER P D & J E IRIER 1950 *Am J Vet Research* 40 56
 BEAUDETTE F R 1951 *Proc 55th Ann Meeting U S Livestock Sanitary Assoc* 108
 BEAUDETTE F R BIVINS & MILLER 1949 *Cornell Vet* 39 3
 BEAUDETTE F R & C B HUDSON 1956 *Cornell Vet* 46 227
 BINS W H W NELSON & M L MINER 1949 *Farm & Home Sci Utah Agr Expt Sta* 10 17
 BISWAL G 1954 *Poultry Sci* 33(4) 843-851
 BLAYLAND J D 1951 *Vet Record* 63 731-733
 BOLIN F M 1948 *Proc Soc Am Bacteriol* 1 43
 BONEY W A 1951 *Southwestern Vet* 5 19-21
 BRANDLY C A 1950 *J Am Vet Med Assoc* 116 139
 BRANDLY C A 1951 *Public Health Repts U S* 66 668
 BRANDLY C A 1952 *In Diseases of Poultry Chap 21* 531 Biester & Schwarte Eds Iowa State College Press Ames Iowa
 BRANDLY C A 1953 *Proc 15th Intern Vet Congr*
 BRANDLY C A H E MOSES & E F JONES 1944 *Spec Rept Huntington Lab to War Department* Washington D C
 BRANDLY C A H E MOSES F L JONE & E I JUNGHER 1946 *Am J Vet Research* 7 243
 BRANDLY C A H F MOSES E I JUNGHER & F I JONES 1946a *Am J Vet Research* 7 239
 BRANDLY C A & N WINSLOW 1947 *Unpublished data*
 BURNET F M 1942 *Australian J Exptl Biol Med Sci* 20 81 89
 BURNET F M 1943 *Med J Australia* 2 313 314
 CLUNNINGHAM C H 1948 *Am J Vet Resea ch* 9 195
 DELAY P D K B DEOME & R A BANKOWSKI 1949 *Science* 107 474-475

TABLE 1
TOTAL REPORTED CASES IN THE UNITED STATES 1946 TO 1955

	Typhoid	Paratyphoid and salmonella
1946	3768	723
1947	3075	951
1948	2840	882
1949	2495	1243
1950	2481	1233
1951	2128	1733
1952	2341	2596
1953	2252	3946
1954	2169	5375
1955	1704	5447
Total	23352	24129

TABLE 2
FOOD POISONING IN THE UNITED STATES BY OUTBREAKS

	Salmonella	Staphylococcus	Others	Total
1953	21	81	84	186
1954	27	100	126	252
1955	16	102	77	195
Total	63	283	287	633

Including 236 outbreaks of gastroenteritis of undetermined etiology

TABLE 3
FOOD POISONING IN ENGLAND AND WALES 1950 TO 1955 BY INCIDENTS

	Salmonella	Staphylococcus	Others	Total
1950	2071	87	1816	3979
1951	1668	65	1614	3347
1952	2098	8	1339	3519
1953	3114	137	2031	5277
1954	3508	127	2381	6016
1955	5763	138	3554	8961
Total	17618	656	12195	31099

Including 12464 incidents of undetermined etiology

the Ministry of Health of Great Britain^{6,10} for the years 1950 to 1955 inclusive (TABLE 3). These figures represent incidents or episodes which are made up of outbreaks involving more than one family, family outbreaks and sporadic cases. Food poisoning in all categories increased during the period illustrated and in each year salmonellosis was responsible for approximately 50 per cent or more of the incidents. In contrast staphylococcal poisoning comprised only 2 per cent of the incidents. If one considers general outbreaks, family out

SALMONELLOSIS OBSERVATIONS ON INCIDENCE AND CONTROL

By P. R. Edwards

Communicable Disease Center, Public Health Service, Atlanta, Ga.

In any discussion of the prevention of the transmission of a disease from animals to man it is desirable first to establish the incidence of the condition in man and in the animal hosts from which it was transmitted. In the case of salmonellosis it is impossible to determine accurately the incidence of the condition either in man or animals. No systematic method of reporting salmonellosis of animals has been established and one must rely only on summaries of isolation and identification of causative agents that are dispersed throughout the medical and veterinary literature to estimate the frequency with which the bacteria occur among the lower animals.

The difficulty of establishing the incidence of salmonellosis in man has been dwelt upon at length by Meyer¹ in his excellent monograph on food poisoning. Here it was pointed out that the summaries published annually by the National Office of Vital Statistics, Public Health Service, Bethesda, Md. were based upon woefully inadequate reporting from a minority of the states and that in most instances the etiological agents of food borne infections were not determined accurately. In spite of these deficiencies one must rely upon these summaries for any estimate of the prevalence of *Salmonella* infections in the United States. In TABLE 1 are listed the reported cases of typhoid fever and other salmonellosis for the ten year period 1946 to 1955. It is apparent that while typhoid fever gradually declined during this period paratyphoid fever and other salmonellosis increased evenfold. MacCready, Reardon and Saphra² observed a fourfold increase in incidence in Massachusetts from 1940 to 1955. It is not possible to say what part of this rise represents an actual increase in the number of cases and what portion is due to more accurate diagnosis and better reporting. Probably each factor has some influence on the figures which, inadequate though they are, demonstrate a growing awareness of the problem on the part of state health agencies.

If one turns to outbreaks of salmonellosis and other food poisonings the figures included in TABLE 2 may be extracted from the data compiled by Dauer and Sylvester.³⁻⁵ Outbreaks of bacillary dysentery have been excluded for obvious reasons. Only 10 per cent of the recorded outbreaks were attributed to salmonellosis and these included but 18 per cent of the cases of *Salmonella* infection reported for the corresponding years. On the contrary outbreaks of staphylococcal poisoning reported were 4.5 times as numerous as outbreaks of salmonellosis. It is noteworthy that insofar as can be ascertained from the publications salmonellae were isolated in connection with each outbreak of salmonellosis reported but in the 202 staphylococcal outbreaks reported for 1954 and 1955 staphylococci were isolated from food, food handlers or patients in only 20 per cent of the episodes.

The data described above may be contrasted with the figures compiled by

TABLE 1
TOTAL REPORTED CASES IN THE UNITED STATES 1946 TO 1955

	Total	Percentage of total cases
1946	3268	723
1947	3075	951
1948	2840	882
1949	2795	1243
1950	2481	1233
1951	2128	1733
1952	2341	2506
1953	2252	3916
1954	2169	5375
1955	1704	5447
Total	23 352	24 129

TABLE 2
FOOD POISONING IN THE UNITED STATES BY OUTBREAKS

	Salmonella	Staphylococcus	Other	Total
1953	21	91	84	186
1954	26	100	126	252
1955	16	107	77	195
Total	63	293	287	633

Including 236 outbreaks of gastroenteritis of undetermined etiology

TABLE 3
FOOD POISONING IN ENGLAND AND WALES 1950 TO 1955 BY INCIDENTS

	Salmonella	Staphylococcus	Other	Total
1950	2071	82	1876	3949
1951	1668	65	1614	3347
1952	2098	82	1339	3519
1953	3114	132	2031	5277
1954	3508	127	2381	6016
1955	5269	138	3554	8961
Total	17 678	626	12 955	31 099

Including 12 464 incidents of undetermined etiology

the Ministry of Health of Great Britain^{8,10} for the years 1950 to 1955 inclusive (TABLE 3). These figures represent incidents or episodes which are made up of outbreaks involving more than one family, family outbreaks and sporadic cases. Food poisoning in all categories increased during the period illustrated and in each year salmonellosis was responsible for approximately 50 per cent or more of the incidents. In contrast staphylococcal poisoning comprised only 2 per cent of the incidents. If one considers general outbreaks, family out-

TABLE 4
FOOD POISONING IN ENGLAND AND WALES 1950 TO 1955 BY INCIDENTS

	General outbreaks	Family outbreaks	Sporadic cases	Total incidents
<i>Salmonella</i>	920	1986	14 772	17 678
<i>Staphylococcus</i>	303	182	141	626
Other	1641	687	10 467	12 795
Total	2864	2855	25 380	31 099

TABLE 5
OUTBREAKS PER 100 000 POPULATION 1953-1955

	England and Wales	United States
<i>Salmonella</i>	1.17	0.042
<i>Staphylococcus</i>	0.4	0.18

breaks and sporadic cases separately for the same period the data in TABLE 4 are obtained. The proportion of staphylococcal incidents is greater among the general outbreaks than among family outbreaks and sporadic cases but even when distinct outbreaks only are considered *Salmonella* infections outnumber staphylococcal poisonings by 3 to 1. Furthermore it is evident that the great majority of the former occurred as sporadic cases—a fact borne out by our own experience.

Objection has been taken to the British data on the ground that workers in England were prone to ignore staphylococcal poisonings and that only in recent years have British bacteriologists taken full cognizance of the disease.¹ It is felt that these objections are based only upon the preponderance of *Salmonella* infections in the British reports. The data in TABLE 5 are presented in confirmation of this view. From these data it is evident that more than twice as many outbreaks of staphylococcal poisoning per capita were reported in 1953, 1954, and 1955 in England and Wales as in the United States. For the same period the per capita reports of outbreaks of salmonellosis were 28 times as numerous as those in the United States. It seems very unlikely to me that the incidence of the two diseases in the two countries actually could be so different particularly in view of the report of a committee of the Medical Research Council, London, England,¹¹ that importation of egg powder from the United States added greatly to the problem of salmonellosis in Great Britain. Rather it would seem that reporting of salmonellosis in England and Wales is much more complete than in the United States and that only a small fraction of the outbreaks in the latter country are diagnosed and reported. Furthermore it must be remembered that sporadic cases and family outbreaks greatly outnumber the dramatic occurrences that draw the attention of the press and the public as well as of the agencies that deal with public health. It can be concluded only that endemic salmonellosis constitutes a considerable problem in the United States as it does in other countries.

That salmonellae are widely dispersed and occur frequently among the various species of lower animals is evident from numerous reports in the literature. This statement applies not only to the host adapted types (the so-called primary salmonellosis of many species) but to the nonhost adapted types as well. Improved methods of isolation and identification have made the prevalence of the bacteria increasingly apparent within recent years. Not only do some types occur both in animals and man¹⁻¹⁴ but there is a distinct correlation between the presence of the organisms in lower animals and in the human population. For instance Ranta and Dolman¹⁵ noted the high incidence of *S. thompson* in man in western Canada and Gibbons and Moore¹⁶ isolated many cultures of this type from egg powder produced in the same locality. Atkinson *et al.*¹⁷⁻²⁰ called attention to the frequent occurrence of *S. botus morbillicans* in animals and man in Australia. Neither of these types was found frequently in material from similar sources in the United States at the time the observations were made. Likewise Kauffmann²¹ and Van Oye²² have noted the presence of a number of rare and exotic serotypes in man and various animal species in the Belgian Congo and Stewart and DeCaputo³ noted the similarity of types isolated from man and animals in the southeastern United States. Furthermore dramatic increases in the incidence in both man and animals of serotypes previously seen very rarely or not at all in a given region often are observed. In the United States *S. heidelberg* and *S. blockley* are excellent examples. Such a list of occurrences could be extended greatly and the observations imply that there is a direct correlation between salmonellosis of animals and man in any given locality although the methods whereby the organisms are transmitted from animals to man and vice versa may vary from one region to another depending upon sanitary facilities and habits of the populations. The literature dealing with the presence of salmonellae in animals is so voluminous that it is possible only to mention representative observations that deal with the presence of the bacteria in animals used widely as food and in animals which in one way or another come into close contact with man.

Ruminants. In many parts of the world *Salmonella dublin* is endemic in cattle and often has been transmitted to man through infected carcasses and milk. Bartel²⁴ Henning²⁵, Savage²⁷, Field²⁸ and McCall²⁹ have reviewed the literature pertaining to the transmission of *S. dublin* infection from cattle to man. Murdock and Gordon³⁰ noted the presence of *S. dublin* in the feces of 24% per cent of apparently healthy slaughter cattle from Fire indicating the extent of the problem in certain areas. It is known that many healthy cattle are either temporary or permanent carriers of the bacteria that large numbers of the bacteria are excreted in the feces and that mastitis occurs occasionally in infected cows resulting in the excretion of very large numbers of the organisms in the milk. Not only is *S. dublin* found in cattle but other *Salmonella* types also occur among them. Edwards, Bruner and Moran¹, Collard and Sen² and Seeliger¹⁵ noted the occurrence of a variety of salmonellae in cattle and Norton and Armstrong³¹ described a milk borne outbreak of *S. typhimurium* infection due to the presence of an infected cow in a dairy herd.

Bartel⁴ and Lutje³³ noted the frequent occurrence of *S. typhimurium* among cultures from adult slaughter cattle. As regards other ruminants it is known that sheep frequently are infected with *S. typhimurium* and a variety of other types.^{12, 34-37}

Swine For some time swine have been known to carry salmonellae and the older European literature abounds in descriptions of mass outbreaks of *S. choleraesuis* infections in man due to the consumption of infected pork. However it was not until Hormaeche and Salsamendi^{38, 39} systematically examined the feces and lymph glands of apparently normal hogs that the number and variety of the bacteria in swine were realized fully. These observations were amply confirmed by many workers.⁴⁰⁻⁴³ Galton *et al.*⁴¹ noted that isolations from swine in abattoirs were much more frequent than from animals on the farm and mentioned the importance of the sales yard, the holding lot, and the abattoir in the spread of the organisms.

Fowl When one considers the extremely large population of domestic fowl and the frequency with which salmonellae are isolated from them,^{44, 45} birds probably constitute the largest single reservoir of these bacteria among animals. The isolation of salmonellae by Perelli Minetti *et al.*⁴⁶ from 41.4 per cent of all turkeys presented for autopsy in California well illustrates the extent of the problem of salmonellosis in domestic fowl. From the results of a number of investigators^{41, 48, 47, 52} it is evident that the bacteria frequently are present in egg products. Salmonellae may gain entrance to eggs either as a result of ovarian infection or by penetration of the eggshell. The conditions under which penetration of the eggshell takes place have been studied by Stokes Osborne and Bayne⁴³ who found that penetration did not occur in eggs stored at 10° C. or below. The frequent occurrence of the bacteria in processed eggs was attributed by Solowey *et al.*⁴⁴ largely to contamination of eggshells. These investigators emphasized the desirability of using only clean eggs in processing plants. As long as cracked and soiled eggs that cannot be marketed as shell eggs are incorporated into dried and frozen egg products one can expect a high incidence of salmonellae in those products.

Other animals Dogs and cats have been incriminated as carriers of salmonellae.^{45, 48} In some instances direct transmission of the infections from dogs to man has been observed. Rodents long have been known to carry *Salmonella enteritidis* and *S. typhimurium* and for many years it was thought that rats and mice carried these types exclusively. It now is known that wild rats may carry a variety of serotypes⁴⁵ and that the types that are carried are largely the same as those found in the environment of the animals.

Salmonellae occur also in insects and other cold blooded animals but it is not clear whether their occurrence in these animals constitutes a true carrier state or whether it is merely a response to their environment. From the data available on insects it would seem that flies long suspected as vectors of enteric infections carry the organisms both on the exterior of their bodies and in their intestinal tracts but only for a short period after exposure.^{45, 49} From the results obtained by Watt and Lindsay⁷⁰ in fly control studies flies are of less importance in the transmission of salmonellae than of shigellae. Cold

blooded animals such as snakes⁶⁸ tortoises^{71, 72} and ticks⁷³ have been found to carry salmonellae and Thomas⁷⁴ observed transmission from a tortoise to a child. Salmonellae also have been isolated from fish⁶⁰ and from shellfish.^{81, 82} Evidence indicates that waters contaminated by man or animals were responsible for the presence of the organisms in these aquatic animals.

Within recent years a number of studies of the presence of salmonellae in sewage polluted waters of rivers and harbors and in wild fowl that inhabit these waters were undertaken in Europe.^{83, 84} It was demonstrated that many river and harbor waters contained numerous salmonellae of a variety of types and that many of these types also were present in gulls, terns, ducks and other birds that inhabit these waterways. Flooding of meadows with these polluted waters led to contamination of the soil and the infection of domestic animals.⁸⁴ The presence of salmonellae in the irrigation waters of the western United States also has been demonstrated repeatedly.^{87, 88}

Salmonellae in foods It is evident that numerous *Salmonella* types are widely distributed throughout the animal kingdom. It is small wonder therefore that the organisms frequently are present in food products of animal origin. While it has been known for some time that salmonellae occasionally are present in meats, recent investigations have revealed their presence in an unsuspectedly high percentage of them. Holtz⁸⁹ recovered the bacteria from 3 per cent of 24,442 meat samples and Walzburg⁹⁰ did so in 4 per cent of the samples of beef examined. Diageler and Kotter⁹¹ found salmonellae in 951 samples of meats and other foods. Seeliger¹⁴ recorded the isolation of a number of *Salmonella* types from a variety of food products in Germany. From the publications of Cherry, Scherago and Weaver⁹² and Ielsenfeld, Young and Yoshimura⁹³ it was evident that salmonellae found their way into commercial meat products in the United States, but it was not until Calton, Lowery and Hardy⁹⁴ isolated the organisms from 23 per cent of the samples of fresh sausage and from 12.5 per cent of the samples of smoked sausage examined that the extent of the problem was fully apparent. The presence of salmonellae in eggs and egg products has been mentioned. That these organisms also can be found in dressed poultry is evident from the reports of Cherry, Barnes and Edwards⁹⁴, Schneider and Gunderson⁹⁵, Prowne⁹⁶, Galton *et al.*⁹⁷ and Mackel and Payne (personal communication).

The transmission of salmonellosis from animals to man through contaminated animal food products has been recognized since the days of Gartner⁹⁸ and Basebau.⁹⁹ The literature regarding the transmission of the infections from animals to man is so voluminous that it is possible to mention only a few examples. The part played by beef in the production of the disease was reviewed recently by Clarenburg.¹⁰⁰ Trub and Schneider¹⁰¹ noted 418 outbreaks of salmonellosis in Germany in 1951 and 1952, in many of which a variety of animal food products were proved to be vehicles of infection. Hauser, Treutling and Breiffel¹⁰², Jones and Symons⁹⁸ and Premavesi¹⁰³ described outbreaks due to infected sausage and the last named author was able to trace the source of the infection to the swine from which the meat was derived.

The role of duck eggs in salmonellosis of man was reviewed by Clarenburg.¹⁰⁴

and by Lerche¹⁰⁸ Clarenburg Vink and Noordhoff¹⁰⁷ and Miller¹⁰⁸ observed cases of salmonellosis due to duck eggs and Yamamoto *et al*¹⁰⁹ incriminated goose eggs in the production of the disease. Hen eggs have been found to give rise to human infections¹¹⁰⁻¹¹ while frozen whole eggs were incriminated as a vehicle of infection by Newell¹¹² and Newell Hobbs and Wallace¹¹⁴. The widespread outbreak of *S. montevideo* infection in babies in the United States due to infected dried egg yolk was described by Abramson *et al*¹¹⁵. Direct transmission from fowl to man by contact infection frequently has been noted¹²⁻⁴³⁻¹¹⁶⁻¹¹⁸.

The Role of Food Processing and the Human Carrier in Salmonellosis

While salmonellae obviously are indigenous to many animal food products the role of the human carrier and the processing plant in the contamination of the final product must not be underestimated. Clarenburg¹⁰⁰ called attention to the necessity of differentiating between intravital and post mortem contamination of food products. Galton *et al*⁴¹ noted that salmonellae were found in carcasses in abattoirs with greater frequency than in swine on farms and they emphasized the role of the abattoir environment and processing procedures in the spread of the organisms. Hardy and Galton¹¹⁹ also emphasized the role of the processing plant in the spread of the bacteria. Schonberg¹⁰ cited the inadequate facilities extant in many abattoirs and suggested control measures to lessen the contamination of carcass meats. Browne⁵⁶ and Galton *et al*⁹⁷ found salmonellae widely distributed in poultry processing plants. The bacteria were found on tables trays mechanical equipment the hands of workers animal carcasses and edible viscera. The recovery of *S. typhi*, *S. paratyphi* 4 and shigellae from market meats by Floyd Baranski and El Cannani¹¹ obviously represents contamination from human sources. Outbreaks in which human carriers were responsible for the contamination of meat products have been described¹⁻¹²⁵ at length. Felsenfeld and Young¹⁴ reported that 26 of 56 outbreaks due to nonhost adapted salmonellae the sources of which were established were caused by human carriers. The frequent occurrence of the normal human carrier has often been stressed¹⁻¹⁵⁻¹⁶⁻¹⁷⁻¹⁸. In a large series of *Salmonella* isolations Edwards Bruner and Moran¹ noted that 29.6 per cent of the cultures from man were isolated from asymptomatic persons while Saphra and Winter¹⁶ reported 15.5 per cent and MacCready Reardon and Saphra² 21.9 per cent of cultures from carriers. Saphra and Winter¹⁶ estimate the carrier rate in the general population at 0.2 per cent while Savage¹⁷⁹ cites the results obtained by the Public Health Laboratory Service London England in the examination of 9829 persons of whom 0.24 per cent were carriers. In the series of cultures described by Galton and Hardy¹¹⁹ no less than 63 per cent of the cultures of known origin were isolated from asymptomatic carriers. Inasmuch as many of these cultures were isolated from food handlers (Galton personal communication) and since Galton and Quan¹²² and Galton and Hardy¹²¹ comment upon the frequency with which the organisms are isolated from food handlers it becomes apparent that the carrier state in food handlers occurs much more frequently than in the general

population. It might be said that the carrier state is an occupational hazard of those persons who continually handle uncooked meats and carcasses. Carriers occur among infants as well as among adults.^{70 11 122} These nonhost adapted salmonellae may persist in man for only a short time or they may persist for very long periods.^{2 16 120 125} Thomson¹²² found that the numbers of salmonellae excreted by asymptomatic carriers varied widely but that the numbers found per gram of feces sometimes exceeded those found in clinical cases. Direct transmission from man to man may occur without the intervention of food. This phenomenon has been observed frequently in infants in hospital nurseries who may contract infections from other infants or from asymptomatic staff members.^{122 125} Infection of babies at parturition by mothers has been observed repeatedly.^{127 121} It is well known that both clinical cases and the carrier state in adults may result from contact with clinical cases.

Control. In any consideration of the control of infections due to nonhost adapted types of salmonellae it is necessary to consider animal reservoirs of infection, excretion of the bacteria by human cases, convalescents and carriers, the presence of the organisms in the environment, and the methods whereby they are transmitted among and between man and the lower animals.

While there is a large reservoir of salmonellae in the lower animals, this situation can be expected to improve gradually. Tests for detection of the infections and methods of eradication are being studied in fowl and cattle. Progress has been made in the eradication of salmonellae from poultry flocks.¹² Lukas and Bradford¹⁴⁵ noted a 50 per cent decrease in the number of turkeys presented for autopsy and a decline in the rate of salmonellosis from 41 per cent to 21 per cent in the cases examined. This decline was attributed to an agglutination testing program for *S. typhimurium* infection. Field¹⁴ and Henning⁶ have studied the possibilities of the detection of *S. dublin* infection by agglutination tests. While the elimination of salmonellosis from flocks and herds cannot be accomplished in the immediate future, it is encouraging that animal pathologists, particularly poultry pathologists, are thinking in terms of eradication of the infections rather than of their control. It must be admitted, however, that this attitude is dictated largely by economic considerations more than by regard for public health. In any effort to eradicate salmonellae from domestic animals it is necessary to take into consideration the continuous seeding of the population through infected feedstuffs. Galton, Harless and Hardy¹⁴⁶ and Griffin¹⁴⁷ have isolated salmonellae from animal feeds. Recently fish meal has been incorporated into many feeds and this supplement is known to be a fertile source of salmonellae.¹⁵ Rohde¹⁴⁸ isolated ten different *Salmonella* types from one sample of Angola fish meal and Boring⁴⁹ repeatedly isolated the bacteria from fish meal produced in the United States. Unfortunate incidents have resulted from the addition of egg powder to feeds. It will be necessary to eliminate such sources of contamination if *Salmonella* infections are to be eliminated from flocks and herds.

As regards dissemination from person to person, every effort should be made to prevent the spread of the organisms among the population through the excreta of clinical cases, convalescents and carriers. As emphasized by

and by Lerche¹⁰⁸ Clarenburg Vink and Noordhoff¹⁰⁷ and Miller¹⁰⁶ observed cases of salmonellosis due to duck eggs and Yamamoto *et al*¹⁰⁹ incriminated goose eggs in the production of the disease. Hen eggs have been found to give rise to human infections¹¹⁰⁻¹¹ while frozen whole eggs were incriminated as a vehicle of infection by Newell¹¹² and Newell Hobbs and Wallace¹¹¹. The widespread outbreak of *S. montevideo* infection in babies in the United States due to infected dried egg yolk was described by Abramson *et al*¹¹³. Direct transmission from fowl to man by contact infection frequently has been noted.^{1 45 114 115}

The Role of Food Processing and the Human Carrier in Salmonellosis

While salmonellae obviously are indigenous to many animal food products the role of the human carrier and the processing plant in the contamination of the final product must not be underestimated. Clarenburg¹⁰⁰ called attention to the necessity of differentiating between intravital and post mortem contamination of food products. Galton *et al*⁴¹ noted that salmonellae were found in carcasses in abattoirs with greater frequency than in swine on farms and they emphasized the role of the abattoir environment and processing procedures in the spread of the organisms. Hardy and Galton¹¹⁹ also emphasized the role of the processing plant in the spread of the bacteria. Schonberg¹⁰ cited the inadequate facilities extant in many abattoirs and suggested control measures to lessen the contamination of carcass meats. Browne⁹⁶ and Calton *et al*⁹⁷ found salmonellae widely distributed in poultry processing plants. The bacteria were found on tables, trays, mechanical equipment, the hands of workers, animal carcasses and edible viscera. The recovery of *S. typhi*, *S. paratyphi A* and *higellae* from market meats by Floyd Baranski and El Gannani¹¹ obviously represents contamination from human sources. Outbreaks in which human carriers were responsible for the contamination of meat products have been described^{1 15} at length. Felsenfeld and Young⁸ reported that 26 of 56 outbreaks due to nonhost adapted salmonellae the sources of which were established were caused by human carriers. The frequent occurrence of the normal human carrier has often been stressed.^{1 15 16 17 18} In a large series of *Salmonella* isolations Edwards, Bruner and Moran¹ noted that 29.6 per cent of the cultures from man were isolated from asymptomatic persons while Saphra and Winter¹⁶ reported 15.5 per cent and MacCreedy, Keardon and Saphra 21.9 per cent of cultures from carriers. Saphra and Winter¹⁶ estimate the carrier rate in the general population at 0.2 per cent while Savage¹⁹ cites the results obtained by the Public Health Laboratory Service, London, England, in the examination of 9829 persons of whom 0.24 per cent were carriers. In the series of cultures described by Galton and Hardy¹¹⁹ no less than 63 per cent of the cultures of known origin were isolated from asymptomatic carriers. Inasmuch as many of these cultures were isolated from food handlers (Galton personal communication) and since Calton and Quan¹²⁰ and Calton and Hardy¹²¹ comment upon the frequency with which the organisms are isolated from food handlers, it becomes apparent that the carrier state in food handlers occurs much more frequently than in the general

population. It might be said that the carrier state is an occupational hazard of those persons who continually handle uncooked meats and carcasses. Carriers occur among infants as well as among adults.^{70 127 133} These nonhost adapted salmonellae may persist in man for only a short time or they may persist for very long periods.^{2 14 136 137} Thomson¹²⁸ found that the numbers of salmonellae excreted by asymptomatic carriers varied widely but that the numbers found per gram of feces sometimes exceeded those found in clinical cases. Direct transmission from man to man may occur without the intervention of food. This phenomenon has been observed frequently in infants in hospital nurseries who may contract infections from other infants or from asymptomatic staff members.^{132 135} Infection of babies at parturition by mothers has been observed repeatedly.^{137 141} It is well known that both clinical cases and the carrier state in adults may result from contact with clinical cases.

Control. In any consideration of the control of infections due to nonhost adapted types of salmonellae it is necessary to consider animal reservoirs of infection, excretion of the bacteria by human cases, convalescents, and carriers, the presence of the organisms in the environment, and the methods whereby they are transmitted among and between man and the lower animals.

While there is a large reservoir of salmonellae in the lower animals, this situation can be expected to improve gradually. Tests for detection of the infections and methods of eradication are being studied in fowl and cattle. Progress has been made in the eradication of salmonellae from poultry flocks.¹⁴² Lukas and Bradford¹⁴³ noted a 50 per cent decrease in the number of turkeys presented for autopsy, and a decline in the rate of salmonellosis from 41 per cent to 21 per cent in the cases examined. This decline was attributed to an agglutination testing program for *S. typhimurium* infection. Field¹⁴⁴ and Henning⁶ have studied the possibilities of the detection of *S. dublin* infection by agglutination tests. While the elimination of salmonellosis from flocks and herds cannot be accomplished in the immediate future, it is encouraging that animal pathologists, particularly poultry pathologists, are thinking in terms of eradication of the infections rather than of their control. It must be admitted, however, that this attitude is dictated largely by economic considerations more than by regard for public health. In any effort to eradicate salmonellae from domestic animals it is necessary to take into consideration the continuous seeding of the population through infected feedstuffs. Galton, Harless, and Hardy¹⁴⁵ and Ciffin¹⁴⁶ have isolated salmonellae from animal feeds. Recently fish meal has been incorporated into many feeds, and this supplement is known to be a fertile source of salmonellae.¹⁵ Rohde¹⁴⁷ isolated ten different *Salmonella* types from one sample of Angola fish meal, and Boring¹⁴⁸ repeatedly isolated the bacteria from fish meal produced in the United States. Unfortunate incidents have resulted from the addition of egg powder to feeds. It will be necessary to eliminate such sources of contamination if *Salmonella* infections are to be eliminated from flocks and herds.

As regards dissemination from person to person, every effort should be made to prevent the spread of the organisms among the population through the excreta of clinical cases, convalescents, and carriers. As emphasized by

Grant¹³⁵ ¹⁵⁰ this aspect of the problem is a joint responsibility of the practicing physician the health officer and the laboratory Prompt isolation of patients with symptoms of intestinal infections and proper disinfection of their dejecta must be carried out until such time as a definite diagnosis is established Contacts should be examined bacteriologically and the strictest personal hygiene should be practiced within the household Convalescents should be examined systematically until the bacteria no longer can be isolated from their excreta Cheever¹⁵¹ in particular has called upon the practicing physician to take a more active part in the control of the infections and the education of the general public

In the United States there is an unfortunate tendency no longer to require stool cultures in the examination of food handlers When one considers the increasing incidence of reported salmonellosis one must question the wisdom of such an omission Without exception all of the writers who have dealt with the problem of the prevention of salmonellosis have emphasized the necessity of stool examinations and most of them have insisted that these be frequent When one considers that salmonellosis has been transmitted through the medium of such foods as coconut ¹⁵ dried yeast ¹⁵² soya milk ¹⁵⁴ and smoked fish¹⁵⁵ in which salmonellae are not indigenous the role of the human carrier in the processing plant becomes increasingly apparent The systematic and frequent examination of food handlers should extend not only to workers in food dispensing establishments but to those in food processing plants as well Any individual who exhibits any symptoms of intestinal infection should be prevented from handling food until it is established that pathogenic bacteria cannot be demonstrated in his stools Considering the possibility of transmission of the infections through food one cannot refrain from noting that it seems rather futile to require a blood test of food handlers for venereal disease but to make no effort to determine the presence of pathogenic microorganisms in their stools

The control of hygienic conditions in food processing plants as well as food-dispensing establishments must be improved Galton *et al*¹¹ demonstrated that thorough and consistent cleansing of abattoirs and equipment resulted in a marked reduction in the recovery of salmonellae from carcasses and from the environment It is probable that enforcement of the newly adopted poultry ordinance will result in an improvement of conditions in poultry processing plants Savage¹²² has suggested that the adoption of bacteriological standards for the control of abattoir and food processing establishments would be very helpful Hardy and Galton¹¹⁹ cite the successful adoption of coliform organisms as an index of pollution in dairy products and shellfish and while calling attention to the need for proper concern for the bacteriological environment of meat processing plants these authors feel that such an index would be overly sensitive at present Macready Keardon and Saphra² feel that an approach similar to that employed in dairy products should be applied to other products of animal origin including fish and poultry It must be remembered that food habits are changing rapidly in many countries particularly in the United States Food technology has made giant strides

within recent years and Savage¹²⁹ in particular has called attention to the removal of food preparation from the family kitchen to the large establishment. In many instances it seems that methods of control and measures for their enforcement have not kept pace with advances in food processing.

More rigid control of products becomes even more important and practical when it is remembered that salmonellae may survive temperatures to which the food products that contain these organisms sometimes are subjected in cooking.^{11, 134} Since it is possible to pasteurize certain food products effectively^{137, 141} or to destroy salmonellae in them through a combination of acidification and mild heating¹³⁹ it would not seem unreasonable to adopt and to enforce bacteriological standards for such products. Furthermore it is not uncommon for untoward incidents to occur in connection with the newer method of food preservation and preparation; these incidents may lead to the multiplication of organisms present in the food and in some instances to the contamination of previously cooked products. Large numbers of salmonellae have been found in turkey carcasses that had been frozen, thawed and refrozen and which, due to improper method of preparation, gave rise to a severe outbreak of salmonellosis (Mackel and Payne, personal communication). Chicken carcasses that had been thoroughly cooked were contaminated during boning by persons who had prepared the raw carcasses.¹⁴⁰ These examples are cited only to call attention to the need of improved procedures and practices in the mass preservation and preparation of foods.

All of the writers who have given serious thought to the control of salmonellosis have arrived at similar conclusions. Grant¹⁴² Hinshaw and McNeil¹⁴³ and Savage¹²⁹ all have published excellent reviews dealing with the epidemiology and control of salmonellosis. All are agreed that effective control can result only through the cooperation of governmental agencies dealing with health, agriculture and food. Further, it is agreed that salmonellosis should be a reportable disease of both man and animals and that effective reporting and control measures should be enforced. The responsibilities of the physician and the health officer, as well as of the practicing veterinarian, the veterinary officer and the sanitarian are indicated. It is only through close cooperation of these individuals and the aforementioned governmental agencies and through education of the public in the fundamental sanitary aspects of the preparation and preservation of foods that control of the infections may be expected.

References

1. MEYER, K. F. 1953. Food poisoning. *New Engl J Med* 249: 765-773, 804-812, 843-852.
2. MACCREADY, R. A., J. I. REARDON & I. SAPIRA. 1953. Salmonellosis in Massachusetts. *New Engl J Med* 256: 1121-1123.
3. DAUER, C. C. & G. SYLVESTER. 1954. 1953 summary of disease outbreaks. *Public Health Repts U S* 69: 538-546.
4. DAUER, C. C. & G. SYLVESTER. 1955. 1954 Summary of disease outbreaks. *Public Health Repts U S* 70: 536-544.
5. DAUER, C. C. & G. SYLVESTER. 1956. 1955 Summary of disease outbreaks. *Public Health Repts U S* 71: 797-803.

- 6 COCKBURN W C & F SIMPSON 1951 Food poisoning in England and Wales 1950 Med Research Council Brit Monthly Bull Min Health 10 278-239
- 7 COCKBURN W C & E SIMPSON 1954 Food poisoning in England and Wales 1951 and 1952 Med Research Council Brit Monthly Bull Min Health 13 12-24
- 8 COCKBURN W C & E VERNON 1955 Food poisoning in England and Wales 1953 Med Research Council Brit Monthly Bull Min Health 14 34-42
- 9 COCKBURN W C & E VERNON 1955 Food poisoning in England and Wales 1954 Med Research Council Brit Monthly Bull Min Health 14 203-216
- 10 COCKBURN W C & E VERNON 1956 Food poisoning in England and Wales 1955 Med Research Council Brit Monthly Bull Min Health 15 263-273
- 11 MEDICAL RESEARCH COUNCIL OF BRITAIN 1947 Bacteriology of spray dried egg powder with particular reference to food poisoning Med Research Council Brit Mem Spec Rept Ser 260
- 12 EDWARDS P R D W BRUNER & A B MORAN 1948 The genus *Salmonella* its occurrence and distribution in the United States Kentucky Agr Expt Sta Univ Kentucky Ann Rept Bull No 525
- 13 TAYLOR J 1950 Salmonellosis in man and animals Public Health Pepts London 63 168-169
- 14 EDWARDS P R 1956 *Salmonella* and salmonellosis Ann N Y Acad Sci 66(1) 44-53
- 15 SEELIGER H P R 1956 Salmonellosen in Deutschland (1953-1955) Arch Hyg u Bakteriol 140 499-528
- 16 SAPHRA I & J W WINTER 1957 Clinical manifestation of salmonellosis in man New Engl J Med 256 1128-1134
- 17 RANTA L E & C E DOLMAN 1947 Experience with *Salmonella* typing in Canada Can J Public Health 38 286-294
- 18 GIBBONS N F & R L MOORE 1944 Dried whole egg powder VI Occurrence and distribution of *Salmonella* organisms in Canadian powder Can J Research 22 48-63
- 19 ATKINSON N G M WOODROOFE & D E CULVER 1952 The occurrence of *Salmonella* types in Australia VIII Australian J Exptl Biol Med Sci 30 73-80
- 20 ATKINSON N M C CARTER J M WOLLASTON & M WALL 1953 The occurrence of *Salmonella* types in Australia X Australian J Exptl Biol Med Sci 31 465-471
- 21 KAUFFMANN F 1953 On the occurrence of *Salmonella* types in the Belgian Congo Acta Pathol Microbiol Scand 32 549-553
- 22 VAN OYE E 1955 Les salmonellae du Congo belge Ann soc belge med trop 35 229-244
- 23 STEWART W H & T DeCAPITO 1953 Salmonellosis in man and animals in south west Georgia Am J Trop Med Hyg 2 273-278
- 24 BARTEL H 1938 Die geographische Verteilung der von Schlachttieren isolierten Enteritisbakterienstämme in Deutschland Tierarztl Rundschau 44 601-606
- 25 HENNING M W 1939 The antigenic structure of *Salmonella* obtained from domestic animals and birds in South Africa Onderstepoort J Vet Research 13 179-189
- 26 HENNING M W 1953 Calf paratyphoid I A general discussion of the disease in relation to animals and man Onderstepoort J Vet Research 26 3-73
- 27 SAVAGE W G 1940 Discussion of *Salmonella* infections Proc Roy Soc Med 33 357-362
- 28 FIELD H I 1949 *Salmonella* infections in cattle Vet Record 61 109-112
- 29 MCCALL A M 1953 An explosive outbreak of food poisoning caused by *Salmonella dublin* Lancet 1953 (1) 1307-1304
- 30 MURDOCK C R & W A M GORDON 1953 The incidence of *Salmonella* in healthy cattle in Northern Ireland Med Research Council Brit Monthly Bull Min Health 12 72-76
- 31 COLLARD P & R SEN 1956 Isolation of salmonellae from cattle in Ibadan W African Med J 5 118-120
- 32 NORTON R & F C ARMSTRONG 1954 A milk borne outbreak due to *Salmonella typhi muenchen* Med Research Council Brit Monthly Bull Min Health 13 90-95
- 33 LUTJE F 1939 Über das Vorkommen von Bakterien der Paratyphus Enteritidis Gruppe beim Kinde Deut tierarztl Wochsch 48 227-234 246-250 257-261
- 34 NEWSOME I E & F CROSS 1935 Third outbreak of paratyphoid dysentery in lambs J Vet Med Assoc 86 534 536
- 35 STEWART D F 1940 Two mortalities among sheep due to infection with species of the *Salmonella* group of bacteria Australian Vet J 16 169-172

- 36 WATTS P S & M WALL 1952 The 1951 *Salmonella typhimurium* epidemic in sheep in South Australia Australian Vet J 28 165-168
- 37 LEDA S 1953 On an occurrence of *Salmonella pullorum* infection in a ewe Kitasato Arch Fptl Med 18 275-277
- 38 HORMAECHIE I & K SALSAMENDI 1936 Sobre la presencia de salmonelas en los gauchos mesentericos de cer los normales Arch uruguay med cir y especial Mon tevideo 9 66-672
- 39 HORMAECHIE F & R SALSAMENDI 1939 El cerdo normal como portador de salmonelas Arch uruguay med cir y especial Montevideo 14 375-387
- 40 RUBIN H I 1940 The incidence of *Salmonella* in normal hogs J Bacteriol 40 463
- 41 CALTON M M W & SMITH H B McLEATH & A V HARDY 1954 *Salmonella* in swine cattle and the environment of abattoirs J Infectious Diseases 95 236-245
- 42 KRANEVELD I C M FRER & M MANSJOER 1951 *Salmonella* Draggers onder klinisch gersonde varkens op Java Hemera zoa 68 48-73
- 43 BUTIAUX I R CALMONT & I MORFL 1951 *Salmonella* dans les ganglions mesenteriques de porcs Ann inst Pasteur 81 236-238
- 44 DARBY C W & H J STAPFETH 1942 *Salmonella* infections common to man animals and birds Proc 46th Meeting U S Livestock San Assoc 189-207
- 45 HINSHAW W P E McNEIL & T J TAYLOR 1944 Avian salmonellosis Types of *Salmonella* isolated and their relation to public health Am J Hyg 40 264-278
- 46 PERELLI MINETTI J I J C RUCKER & F K ROSS 1948 Paratyphosis Calif Dept Agr Bull 37 151
- 47 SOLOWEY M F E H McFARLANE F H SPALDING & C CHMERDA 1947 Microbiology of spray-dried whole egg II Incidence and types of *Salmonella* Am J Public Health 37 971-982
- 48 SCHNEIDER M D 1945 The isolation of *Salmonella* from frozen whole and powdered egg Army Med Bull 4 47
- 49 SCHNEIDER M D 1946 Investigation of *Salmonella* content of powdered whole egg with not more than 2% moisture content II General survey on the occurrence of species of *Salmonella* in high quality egg powder Food Research 11 313
- 50 MEDICAL RESEARCH COUNCIL BRITISH PUBLIC HEALTH LABORATORY SERVICE 1954 *Salmonella* in duck eggs Monthly Bull Min Health 13 38-43
- 51 MEDICAL RESEARCH COUNCIL BRITISH PUBLIC HEALTH LABORATORY SERVICE 1955 *Salmonella* in liquid egg including frozen liquid egg Monthly Bull Min Health 14 65-69
- 52 MURDOCK C R 1954 The occurrence of *Salmonella* in liquid egg Med Research Council Brit Monthly Bull Min Health 13 43-44
- 53 STOKES J L W W OSBORNE & H G BAYNE 1956 Penetration and growth of *Salmonella* in shell eggs Food Research 21 510-518
- 54 SOLOWEY M F E H SPALDING & H F GORESLINE 1946 An investigation of a source and mode of entry of *Salmonella* organisms in spray dried whole egg powder Food Research 11 380-390
- 55 MAGNUSSEN K E 1938 Ein Hund als Ansteckungsquelle von Paratyphusinfektionen Z Hyg Infektionskrankh 121 136-138
- 56 KALFFMANN F & F J HENNINGSEN 1939 A new *Salmonella* type from man and dog Acta Pathol Microbiol Scand 16 99-102
- 57 BALL M R 1951 *Salmonella* in dogs and cats of the Los Angeles Honolulu and Bermuda areas J Am Vet Med Assoc 118 164-166
- 58 CARTER H S & B L WEIR 1952 A dog as a probable source of human infection with *Salmonella paratyphi B* J Pathol Bacteriol 64 230
- 59 GALTON M M J I SCATTERDAY & A V HARDY 1952 Salmonellosis in dogs I Bacteriological epidemiological and clinical considerations J Infectious Diseases 91 1-5
- 60 BLAZEK K K HRUSKA & J KRAL 1953 Salmonellosis u psu Veterinarství Brno 3 273-274
- 61 JUDEVID T & P R EDWARD 1953 Unpublished data
- 62 MACKERRAS I M & M J MACKERRAS 1949 An epidemic of infantile gastroenteritis in Queensland caused by *Salmonella typhimurium* (Basenau) J Hyg 47 166-181
- 63 IYVINE M J R ENRIGHT G CHING & I TANIMOTO 1950 Observations on salmonellosis in the Hawaiian Islands Hawaii Med J 9 149-155

- 6 COCKBURN W C & E SIMPSON 1951 Food poisoning in England and Wales 1950 Med Research Council Brit Monthly Bull Min Health 10 228-239
- 7 COCKBURN W C & E SIMPSON 1954 Food poisoning in England and Wales 1951 and 1952 Med Research Council Brit Monthly Bull Min Health 13 12-24
- 8 COCKBURN W C & E VERNON 1955 Food poisoning in England and Wales 1953 Med Research Council Brit Monthly Bull Min Health 14 34-42
- 9 COCKBURN W C & E VERNON 1955 Food poisoning in England and Wales 1954 Med Research Council Brit Monthly Bull Min Health 14 203-216
- 10 COCKBURN W C & E VERNON 1956 Food poisoning in England and Wales 1955 Med Research Council Brit Monthly Bull Min Health 15 263-273
- 11 MEDICAL RESEARCH COUNCIL OF BRITAIN 1947 Bacteriology of spray dried egg powder with particular reference to food poisoning Med Research Council Brit Mem Spec Rept Ser 250
- 12 EDWARDS P R D W BRUNER & A B MORAN 1948 The genus *Salmonella* its occurrence and distribution in the United States Kentucky Agr Expt Sta Univ Kentucky Ann Rept Bull No 525
- 13 TAYLOR J 1950 Salmonellosis in man and animals Public Health Repts London 63 168-169
- 14 EDWARDS P R 1956 *Salmonella* and salmonellosis Ann N Y Acad Sci 66(1) 44-53
- 15 SEELIGER H P P 1956 Salmonellosen in Deutschland (1953-1955) Arch Hyg u Bakteriol 140 499-528
- 16 SAPHIRA I & J W WINTER 1957 Clinical manifestations of salmonellosis in man New Engl J Med 256 1128-1134
- 17 JANTA L E & C E DOLMAN 1947 Experience with *Salmonella* typing in Canada Can J Public Health 38 286-294
- 18 GIBBONS V F & R L MOORE 1944 Dried whole egg powder VI Occurrence and distribution of *Salmonella* organisms in Canadian powder Can J Research 22 48-63
- 19 ATKINSON N G M WOODROOFE & D E CULVER 1952 The occurrence of *Salmonella* types in Australia VIII Australian J Exptl Biol Med Sci 30 73-80
- 20 ATKINSON N M C CARTER J M WOLLASTON & M WALL 1953 The occurrence of *Salmonella* types in Australia X Australian J Exptl Biol Med Sci 31 463-471
- 21 KAUFFMANN F 1953 On the occurrence of *Salmonella* types in the Belgian Congo Acta Pathol Microbiol Scand 32 549-553
- 22 VAN OYE E 1955 Les salmonellae du Congo belge Ann soc belge med trop 35 229-244
- 23 STEWART W H & T DECAPITO 1953 Salmonellosis in man and animals in south west Georgia Am J Trop Med Hyg 2 273-278
- 24 BARTEL H 1938 Die geographische Verteilung der von Schlachttieren isolierten Enterisbakterienstämme in Deutschland Tierarztl Rundschau 44 601-606
- 25 HENNING M W 1939 The antigenic structure of *Salmonella* obtained from domestic animals and birds in South Africa Onderstepoort J Vet Research 13 19-189
- 26 HENNING M W 1953 Calic paratyphoid I A general discussion of the disease in relation to animals and man Onderstepoort J Vet Research 26 3-23
- 27 SAVAGE W G 1940 Discussion of *Salmonella* infections Proc Roy Soc Med 33 351-362
- 28 FIELD H I 1949 *Salmonella* infections in cattle Vet Record 61 109-112
- 29 MCCALL A M 1953 An explosive outbreak of food poisoning caused by *Salmonella dublin* Lancet 1953 (i) 1302-1304
- 30 MURDOCK C R & W A M GORDON 1953 The incidence of *Salmonella* in healthy cattle in Northern Ireland Med Research Council Brit Monthly Bull Min Health 12 72-76
- 31 COLLARD P & R SEN 1956 Isolation of salmonellae from cattle in Ireland W African Med J 5 118-120
- 32 NORTON R & I C ARMSTRONG 1954 A milk borne outbreak due to *Salmonella typhi murium* Med Research Council Brit Monthly Bull Min Health 13 30-9
- 33 LUTJE F 1939 Über das Vorkommen von Bakterien der Paratyphus Enterit Gruppe beim Rinde Deut tierarztl Wochschr 48 221-234 246-250 257 261
- 34 NEWSOME I E & F CROSS 1955 Third outbreak of paratyphoid enteritis in lambs J Vet Med Assoc 86 534-536
- 35 STEWART D F 1940 Two mortalities among sheep due to infection with species of the *Salmonella* group of bacteria Australian Vet J 16 169-172

- 93 CALTON M M W V LOWERY & V V HARDY 1954 *Salmonella* in fresh and smoked pork sausage. J Infectious Diseases 94 237-238
- 94 CHERRY W B L A BARNES & I R EDWARDS 1946 Observations on a monophasic *Salmonella* variant. J Bacteriol 51 235-243
- 95 SCHNEIDER M D & M F CUNTERON 1949 Investigators shed more light on *Salmonella typhimurium*. U S Egg and Poultry Mag 55 10-11-27
- 96 BROWN A S 1947 The public health significance of *Salmonella* on poultry and poultry products. Thesis Univ Calif Berkley Calif
- 97 CALTON M M D C MACKENZIE A I LEWIS W C HAIR & V V HARDY 1955 *Salmonella* in poultry and poultry processing plants in Florida. Am J Vet Research 16 132-137
- 98 GARTNER A 1898 Über die Fleischvergiftung in Frankenhäusern und den Erreger derselben. Correspondenzblatt allg ärztl Ver Thüringen 17 573
- 99 BAENAT I 1894 Über eine im Fleisch gefundene infectiöse Bacterie. Arch Hyg u Bakteriol 20 242-294
- 100 CLARENBERG A 1953 Sources of contamination in outbreaks of food poisoning caused by meat in the Netherlands. Proc Soc Appl Bacteriol 16 169-172
- 101 TRUB C I I & I SCHNEIDER 1952 Auftreten gehäufter Einzelerkrankungen und Gruppenerkrankungen durch *Salmonella panama* und *Salmonella typhimurium* in Paratyphus C Gruppe im Lande Nordrhein Westfalen 1950-51. Z Hyg Infektionskrankh 135 121-145
- 102 HAUSER G H W L TREUTING & I A BRUFFELH 1945 An outbreak of food poisoning due to a new etiological agent—*Salmonella berli*. Public Health Repts U S 60 1138-1142
- 103 JONES A C & A D SYMONS 1948 An outbreak of food poisoning due to *Salmonella typhimurium* conveyed by sausages and sausage meat. Med Research Council Brit Monthly Bull Min Health 7 207-208
- 104 PREMAYE I K A 1946 Über eine durch *Salmonella heidelberg* hervorgerufene Krankheitsform. Med Klin 51 1103-1104
- 105 CLARENBERG A 1939 Paratyphus in ducks in relation to public health. Proc 7th World's Poultry Congr 253-256
- 106 LERCHE M 1937 *Salmonella* Infektion beim Geflügel und ihre Bedeutung für die Epidemiologie der Salmonellabakterien. Proc 7th World's Poultry Congr 274-278
- 107 CLARENBERG A H H VINK & B C NOORDHOFF 1940 Infectie met *Salmonella saint paul* in verband met Salmonellosis bij eenden. Ned Tijdschr Geneesk 94 1615-1618
- 108 MILLER A A 1952 Human *Salmonella typhimurium* infection due to duck eggs with special reference to flocks of ducks. Brit Med J 1952 (ii) 125-128
- 109 YAMAMOTO A K HARADA H FUKUMI & R NAKAYA 1953 On food poisoning due to *Salmonella typhimurium* from goose eggs. Gunma J Med Sci Japan 1 87-95
- 110 WATT J 1945 An outbreak of *Salmonella typhimurium* infection in man from infected chicken eggs. Public Health Repts U S 60 835-837
- 111 GERNEZ RIUEUX C R BUTTIAUX A KESTELOOT & R SIFFRAN 1949 Intoxications alimentaires par *Salmonella pullorum*. Presse med 57 419-420
- 112 MOLLER A 1935 Breslau Gruppenerkrankung durch Hühnereier. Zentr Bakteriol Parasitenk I Orig 164 535-539
- 113 NEWELL K 1955 Paratyphoid fever possibly associated with Chinese frozen egg. Med Research Council Brit Monthly Bull Min Health 14 146-160
- 114 NEWELL K B C HOBBS & E J G WALLACE 1955 Paratyphoid fever associated with Chinese frozen whole egg. Brit Med J 1955 (i) 1246-1297
- 115 ABRAMSON H H GREYENBERG S PLOTKIN & C OLDENBURCH 1954 Food poisoning in infants caused by egg yolk powder. Am J Diseases Children 87 1-6
- 116 ABELSMITH M K & H E ROBERTSON 1953 *Salmonella typhimurium* in 1952 turkey flocks—A public health hazard. Can J Public Health 44 263-265
- 117 PERIERA M S & J D BLANLAND 1955 *Salmonella typhimurium* infection in man contracted from turkeys. Med Research Council Brit Monthly Bull Min Health 14 52-55
- 118 CLARENBERG A G D HEMMES & W WAGENVOORT 1952 On the occurrence of *Salmonella typhimurium* in man and animal and the mode of transmission of the salmonellosis caused by this species. Antonie van Leeuwenhoek J Microbiol Serol 18 171-181
- 119 HARDY A V & M M GALTON 1955 The role of food processing plants in the dissemination of *Salmonella*. Am J Trop Med Hyg 4 725-730

- 64 IUDLAM G B 1954 *Salmonella* in rats with special reference to findings in a butcher's by products factory Med Research Council Brit Monthly Bull Min Health 13 196-207
- 65 LEE P E 1955 *Salmonella* infections of urban rats in Brisbane Queensland Australia J Exptl Biol Med Sci 33 113-116
- 66 HORMAECHÉ E U L SERRACO C A PELUFFO C P I ALEPPO 1943 Causes of infantile summer diarrhea Am J Diseases Children 66 539-551
- 67 OSTROLEK M & H WELCH 1942 The house fly as a vector of food poisoning organisms in food producing establishments Am J Public Health 32 487-494
- 68 MCNEIL E & W J HINSHAW 1944 Snakes cats and flies as carriers of *Salmonella typhimurium* Poultry Sci 23 456-457
- 69 GROSZ H & U PRELSZ 1951 Infektionsversuche an Fliegen mit Darm pathogenen Keimen I Versuche mit Typhus und Paratyphusbakterien Zentr Bakteriell Parasitenk I Orig 156 371-377
- 70 WATT J & D I LINDSAY 1948 Diarrheal disease control studies Effect of fly control in a high mortality area Public Health Pepts U S 63 1319-1334
- 71 BOYCOTT J A J TAYLOR & S H DOUGLAS 1953 *Salmonella* in tortoises J Pathol Bacteriol 65 401-411
- 72 HIRSCH W & R SHAPIRO HIRSCH 1954 *Salmonella* in tortoises J Med Assoc Israel 46 237-238
- 73 ALIV K 1956 *Salmonella* in tortoises Acta Pathol Microbiol Scand 38 71-72
- 74 THOMAS M E M 1957 Enteritis from tortoises Med Research Council Brit Monthly Bull Min Health 16 29-31
- 75 MACKEY J P 1955 Salmonellosis in Dar es Salaam E African Med J 32 1-6
- 76 IEF P E & I M MACLERRAS 1955 *Salmonella* infections in Australian native animals Australian J Exptl Biol Med Sci 33 117-126
- 77 COLLARD P 1956 *Igama igama* as a reservoir of *Salmonella* infection in Ibadan W African Med J 5 154-156
- 78 COLLARD P & D MONTEFIORE 1957 The Agama lizard as a reservoir of *Salmonella* infection in Nigeria Nature 179 164
- 79 FLOYD T M & H HOOCHSTRAAL 1956 Isolation of *Salmonella* from ticks in Egypt J Egypt Public Health Assoc 31 119-128
- 80 GLASEHARAM J T VELALDAPILLAI & G R NILES 1956 The isolation of *Salmonella* organisms from fresh fish sold in a Colombo fish market J Hyg 54 581-584
- 81 BIDWELL M H & C B KELLY JR 1950 Ducks and shellfish sanitation Am J Public Health 40 923-928
- 82 STEINIGER F 1956 Zur Freilandbiologie der Salmonellen im Bereich des westlichen Mittelmeeres Zentr Bakteriell Parasitenk I Orig 166 245-265
- 83 STEINIGER F 1953 Aus Freilandbiologie der Salmonellen Proc 6th Intern Congr Microbiol 4 193-199
- 84 POPP L 1956 Vorfluter—ein reichhaltiges *Salmonellen* Reservoir Zentr Bakteriell Parasitenk I Orig 166 90-94
- 85 STRALCH D & W MUCKER 1956 Bakteriologische Wasseruntersuchungen in einem oberhessischen Fluss in Zusammenhang mit in diesem Flusstal aufgetretenen *Salmonellen* Infektionen bei Haustieren Berlin u Münch tierarztl Wochschr 69 200-208
- 86 HOLTZ K 1956 Über die Bedeutung der durch bakteriologische Fleischuntersuchungen ermittelten *Salmonellen* im Gebiet des Niederrheins Arch Lebensmittel Hyg 7 121-123
- 87 DUNLOP S C R M TWEDT & W L WANG 1952 *Salmonella* in irrigation water Sewage and Ind Wastes 24 1015-1020
- 88 GREENFIELD M & F FISH 1952 Personal communication
- 89 WALZBURG U 1955 Zur Verbreitung der Salmonellose des Rindes im Nordwest deutschen Küstengebiet unter besonderer Berücksichtigung des Raumes Weser Ems Arch Lebensmittel Hyg 6 223-226
- 90 DIAGELER A & J KOTTER 1956 Über das Vorkommen von *Salmonellen* in Bayern während der Jahre 1946 bis 1954 Berlin u Münch tierarztl Wochschr 69 281-285
- 91 CHERRY W B M SCHERAGO & R H WEAVER 1943 The occurrence of *Salmonella* in retail meat products Am J Hyg 37 211-215
- 92 FELSSENFIELD O A M YOUNG & T YOSHIMURA 1950 A survey of *Salmonella* organisms in market meat eggs and milk J Am Vet Med Assoc 116 17-21

- 93 CALTON M M W & LOWERY & V V HARDY 1954 *Salmonella* in fresh animal meat
pork sausage J Infectious Diseases 95 232-235
- 94 CHERRY W B I & BARNES & I R EDWARDS 1947 Observation on a mono-
phasic *Salmonella* variant J Bacteriol 61 235-243
- 95 SCHIFFER M D & M I CUNIFFON 1949 Investigations shall more light on
Salmonella typhimurium U S J of Poultry Mag 55 10-11 27
- 96 BROWN A S 1949 The public health significance of *Salmonella* on poultry and
poultry products Thesis Univ Calif Berkeley Calif
- 97 CALTON M M D C MACFARLANE W C HARRIS & V V HARDY 1955
Salmonellosis in poultry and poultry processing plants in Florida Am J Vet Re-
search 16 132-137
- 98 CARTNER A 1888 Über die Fleischvergiftung in Frankenhausen und den Erreger
derselben Correspondenzblatt allg ärztl Ver Thüringen 17 573
- 99 BASENAT F 1894 Über eine im Fleisch gefundene infectiöse Bacterie Arch
Hyg u Bakteriell 20 242-294
- 100 CLARENBURG A 1953 Sources of contamination in outbreaks of food poisoning
caused by meat in the Netherlands Proc Soc Appl Bacteriol 16 169-172
- 101 TRUB C I P & I SCHNEIDER 1952 Auftreten gehäufter Einzelerkrankungen
und Gruppenkrankungen durch *Salmonella panama* und *Salmonella* Typen der
Paratyphus C Gruppe im Lande Nordrhein Westfalen 1950-51 Z Hyg Infect
ionskrankh 135 121-145
- 102 HAUSER G H W L TREUTING & I A BREITFELDER 1945 An outbreak of food
poisoning due to a new etiological agent—*Salmonella berta* Public Health Repts
U S 60 1138-1142
- 103 JONES A C & A D SYMONS 1948 An outbreak of food poisoning due to *S. dublin*
conveyed by sausages and sausage meat Med Research Council Brit Monthly
Bull Min Health 7 207-206
- 104 IREMAVESTI K A 1956 Über eine durch *Salmonella heidelberg* hervorgerufene
Krankheitsform Med Klin 51 1103-1104
- 105 CLARENBURG A 1939 Paratyphoid in ducks in relation to public health Proc
7th World S Poultry Congr 233-236
- 106 LERCHE M 1939 *Salmonella* Infektionen beim Geflügel und ihre Bedeutung für die
Epidemiologie der *Salmonella* Infektion Proc 7th World S Poultry Congr 274-278
- 107 CLARENBURG A H H VINE & B C NOORDHOFF 1950 Infectie met *Salmonella*
saint pauli in verband met *Salmonellosis* bij eenden Ned Tijdschr Geneesk 94
1615-1618
- 108 MILLER A A 1952 Human *Salmonella typhimurium* infection due to duck eggs
with special reference to flocks of ducks Brit Med J 1952 (ii) 125-129
- 109 YAMAMOTO A K HARADA H FUKAMI & R NAKAYA 1952 On food poisoning due
to *S. typhimurium* from goose eggs Gann J Med Sci Japa 1 87-95
- 110 WATT J 1945 An outbreak of *Salmonella* infection in man from infected chicken
eggs Public Health Repts U S 60 835-839
- 111 GERNETZ C R BUTTACUCCI A KESTELOOT & R SUFFRAN 1949 Intoxica-
tions alimentaires par *Salmonella paratyphi* Presse med 57 479-480
- 112 MÖLLER A 1955 Breslau Cruppenkrankung durch Hühnererzeuger Zentr Bakteriell
Paastenk 1 Orig 164 535-539
- 113 NEWELL K 1955 Paratyphoid B fever possibly associated with Chinese frozen egg
Med Resea ch Council Brit Monthly Bull Min Health 14 146-160
- 114 NEWELL K B C HOBBS & J C WALLACE 1955 Paratyphoid fever associated
with Chinese frozen whole egg Brit Med J 1955 (ii) 1296-1297
- 115 ABRAMSON H H GREENBERG S FLOTJIN & C OLDENBUSCH 1954 Food poisoning
in infants caused by egg yolk powder Am J Diseases Children 87 1-6
- 116 ABELSETH M K & H F ROBERTSON 1953 *Salmonella typhimurium* in 1952 tur-
key flock—A public health hazard Can J Public Health 44 263-265
- 117 PERIERA M S & J D BLAXLAND 1955 *Salmonella typhimurium* infection in man
contracted from turkeys Med Research Council Brit Monthly Bull Min Health
14 52-55
- 118 CLARENBURG A G D HEMMES & W WAGENVOORT 1952 On the occurrence of
Salmonella dublin in man and animals and the mode of transmission of the salmonel-
losis caused by this species Antonie van Leeuwenhoek J Microbiol Serol 18 171-
181
- 119 HARDY A V & M M GALTON 1955 The role of food processing plants in the dis-
semination of *Salmonella* Am J Trop Med Hyg 4 725-730

- 64 LUDLAM G B 1954 *Salmonella* in rats with special reference to findings in a butcher's by products factory Med Research Council Brit Monthly Bull Min Health 13 196-207
- 65 LEE P E 1955 *Salmonella* infections of urban rats in Brisbane Queensland Australian J Exptl Biol Med Sci 33 113-116
- 66 HORMAECHÉ E U I SURRACO C A PELUFFO C P L ALEFFO 1943 Causes of infantile summer diarrhea Am J Diseases Children 66 539-551
- 67 OSTROLFNA M & H WELCH 1942 The house fly as a vector of food poisoning organisms in food producing establishments Am J Public Health 32 487-494
- 68 MCNEIL E & W R HINSHAW 1944 Snakes cats and flies as carriers of *Salmonella typhimurium* Poultry Sci 23 456-457
- 69 GROSZ H & U IREUSZ 1951 Infektionsversuche an Fliegen mit Darm pathogenen Keimen I Versuche mit Typhus und Paratyphusbakterien Zentr Bakteriell Parasitenk I Orig 166 371-377
- 70 WATT J & D R LINDSAY 1948 Diarrheal disease control studies Effect of fly control in a high morbidity area Public Health Repts U S 63 1319-1334
- 71 BOYCOTT J A J TAYLOR & S H DOUGLAS 1953 *Salmonella* in tortoises J Pathol Bacteriol 65 401-411
- 72 HIRSCH W & R SHAPIRO HIRSCH 1954 *Salmonella* in tortoises J Med Assoc Israel 46 237-238
- 73 ALIN K 1956 *Salmonella* in tortoises Acta Pathol Microbiol Scand 38 71-72
- 74 THOMAS M E M 1957 Enteritis from tortoises Med Research Council Brit Monthly Bull Min Health 16 29-31
- 75 MACKEY J P 1955 Salmonellosis in Dar es Salaam E African Med J 32 1-6
- 76 LEE P E & I M MACKERRAS 1955 *Salmonella* infections in Australian native animals Australian J Exptl Biol Med Sci 33 117-126
- 77 COLLARD P 1956 *Agama agama* as a reservoir of *Salmonella* infection in Ibadan W African Med J 5 154-156
- 78 COLLARD P & D MONTEFIORE 1957 The Agama lizard as a reservoir of *Salmonella* infection in Nigeria Nature 179 164
- 79 FLOYD T M & H HOOGSTRAAL 1956 Isolation of *Salmonella* from ticks in Egypt J Egypt Public Health Assoc 31 119-128
- 80 GULASEKHARAM J T VELAUDAPILLAI & G R NILFS 1956 The isolation of *Salmonella* organisms from fresh fish sold in a Colombo fish market J Hyg 54 581-584
- 81 BIDWELL M H & C B KELLY JR 1950 Ducks and shellfish sanitation Am J Public Health 40 923-928
- 82 STEINIGER F 1956 Zur Freilandbiologie der Salmonellen im Bereich des westlichen Mittelmeeres Zentr Bakteriell Parasitenk I Orig 166 245-265
- 83 STEINIGER F 1953 Aus Freilandbiologie der Salmonellen Proc 6th Intern Congr Microbiol 4 193-199
- 84 POPI L 1956 Vorfluter—ein reichhaltiges Salmonellen Reservoir Zentr Bakteriell Parasitenk I Orig 166 90-94
- 85 STRALCH D & W MUNKER 1956 Bakteriologische Wasseruntersuchungen in einem oberhessischen Fluss in Zusammenhang mit in diesem Flusstal aufgetretenen Salmonellen Infektionen bei Haustieren Berlin u Munch tierarztl Wochschr 69 200-208
- 86 HOLTZ K 1956 Über die Bedeutung der durch bakteriologische Fleischuntersuchungen ermittelten Salmonellen im Gebiet des Niederrheins Arch Lebensmittelhyg 7 171-173
- 87 DUNLOP S C R M TWIFT & W L WANG 1952 *Salmonella* in irrigation water Sewage and Ind Wastes 24 1015-1070
- 88 GREENFIELD M & F FISH 1952 Personal communication
- 89 WALZBURG U 1955 Zur Verbreitung der Salmonellose des Rindes im Nordwest deutschen Küstengebiet unter besonderer Berücksichtigung des Raumes Weerfms Arch Lebensmittelhyg 6 273-226
- 90 DIAGELEA A & J KOTTER 1956 Über das Vorkommen von Salmonellen in Bayern während der Jahre 1946/1 1954 Berlin u Munch tierarztl Wochschr 69 231-285
- 91 CHERRY W B M SCHFRAGO & R H WEAVER 1943 The occurrence of *Salmonella* in retail meat products Am J Hyg 37 211-215
- 92 FELSENFELD O A M YOUNG & T YOSHIMURA 1950 A survey of *Salmonella* organisms in market meat eggs and milk J Am Vet Med Assoc 116 17-21

- 147 CRIFFIN C A 1952 A study of prepared feeds in relation to *Salmonella* infection in laboratory animal J Am Vet Med 121 197-200
- 148 KOHDE I 1956 Personal communication
- 149 BORING J 1957 Personal communication
- 150 CRANT J 1950 Salmonellosis. Med Officer 84 59-19-22
- 151 CHEEVER F S 1955 The acute diarrheal diseases of bacterial origin. Bull N Y Acad Med Sci 2 31 611-77
- 152 MACKENZIE F F 1953 Typhoid and coconut. An unexpected and dangerous combination. Health Bull Melbourne 110 19-23
- 153 KINZ L J & O T C OCCHIERONI 1955 Salmonellosis originating in a hospital. A newly recognized source of infection. New Engl J Med 253 161-163
- 154 BROWNE A S 1955 Personal communication
- 155 OLITZKY I A M IERRE M A SHEFFMAN & M WERRIN 1956 Smoked fish as a vehicle of salmonellosis. Public Health Reps U S 71 773-779
- 156 SOLOWAY M & F J CALESSNICK 1948 Survival of *Salmonella* in reconstituted egg powder subjected to blinding and scrambling. Food Research 13 216-226
- 157 WINTER A I G F STEWART A H McFARLANE & M SOLOWAY 1946 Pasteurization of liquid egg products. III Destruction of *Salmonella* in liquid whole egg. Am J Public Health 36 451
- 158 CRESLINE H E K M HAYES R E MOSER JR M A HOWE JR & E F DREWNAK 1951 Pasteurization of liquid whole egg under commercial conditions to eliminate *Salmonella*. U S Dept Agr Circ 897
- 159 ILLMANN C 1956 Uskadelig pårelse af salmonellabakterier i ikke varmebehandlede levned midler tilberedt af rå hønseæg. Nord Vete 8 859-864
- 160 HINSHAW W R & F McNEIL 1951 *Salmonella* infection as a food industry problem. Advances in Food Research 3 209-240

- 120 SCHONBERG F 1953 The place of the veterinary surgeon in food sanitation with particular attention to food of animal origin Proc 15th Intern Vet Congr 1-8
- 121 FLOYD T M J R BARANSKI & M EL GANNANI 1953 Recovery of human enteric pathogens on meat from butcher shops in Cairo Egypt J Infectious Diseases 92 224-225
- 122 BUTTIAUX R V LESAFFRE & J MORTAMEZ 1953 Les tox. infections alimentaires par *Salmonella* Necessite d'une surveillance des manipulateurs de denrees alimentaires Presse med 61 47-748
- 123 NAKAYA R G SAKAGUCHI & H FUKUMI 1952 *S. new b unsr* sick from a food poisoning outbreak and from a healthy carrier Japan J Med Sci & Biol 6 125-128
- 124 POIDE R 1955 Über das Auftreten verschiedener Salmonellosen in Hamburg während 1954 und die sich daraus ergebenden epidemiologischen Konsequenzen Arch Hyg u Bakteriol 139 148-157
- 125 SILVERSTOPE L & M WRANNE 1955 *Salmonella*epidemierna i Östergötlands län år 1953 Nord Hyg Tidskr 36 213-229
- 126 FELSENFELD O & V M YOLAC 1949 A study of salmonellosis in North and South America Am J Trop Med 29 483-491
- 127 GALTON M M & A V HARDY 1953 The distribution of *Salmonella* infections in man in Florida during the past decade Public Health Lab 11 88-93
- 128 EISENBERG G M A J PALAZZOLO & H F FLIPPEY 1955 Clinical and microbiologic aspects of salmonellosis New Engl J Med 253 90-94
- 129 SAVAGE SIR W 1956 Problems of *Salmonella* food poisoning Brit Med J 1956 (ii) 317-321
- 130 GALTON M M & M S QUAN 1944 *Salmonella* isolated in Florida during 1943 with the combined enrichment method of Kauffmann Am J Public Health 34 1071-1075
- 131 GALTON M M & A V HARDY 1948 Studies of the acute diarrheal diseases. XVI Salmonellosis in Florida Public Health Repts U S 63 84, 851
- 132 MACKERRAS I M & A M PASK 1949 Infant *Salmonella* carriers Lancet 1949 (ii) 940
- 133 LEEDER F S 1956 An epidemic of *Salmonella panama* infections in infants Ann N Y Acad Sci 66(1) 54-60
- 134 D'ALBORA J B A P INGENO & J M EDSON 1945 Outbreak of food poisoning due to *Salmonella montevideo* in an army general hospital J Am Med Assoc 129 10-12
- 135 GRANT J 1951 Human salmonellosis Some further observations Med Officer 85 117-119
- 136 THOMSON S 1955 The numbers of pathogenic bacilli in faeces in intestinal diseases J Hyg 63 211-224
- 137 WATT J & E CARLTON 1945 Studies on the acute diarrheal diseases. XVI An outbreak of *Salmonella typhimurium* infection among newborn premature infants Public Health Repts U S 60 734-740
- 138 ABRAMSON H 1947 Infection with *Salmonella typhimurium* in the newborn Am J Diseases Children 74 516-586
- 139 SELIGMANN E 1949 Mass invasion of salmonellae in a babies ward Ann Paediat 172 406-408
- 140 NETER E 1950 Observations on the transmission of salmonellosis in man Am J Public Health 40 929-933
- 141 EPSTEIN H C A HOCHWALD & K ASLEE 1951 *Salmonella* infections of the newborn infant J Pediatr 38 23, 31
- 142 HINSHAW W R & E MCNEIL 1943 Use of the agglutination test in detecting *Salmonella typhimurium* carriers in turkey flocks Proc 44th Ann Meeting U S Livestock San Assoc 106-121
- 143 GORDON K F & J S CARIDE 1944 *Salmonella* infection in ducks Observations on the value of the agglutination test in the eradication of the infection and in estimations on the cycle of infection via the egg J Comp Pathol Therap 54 61-6
- 144 DELAY P D T W JACKSON E F JONE & D F STICHER 1944 A testing series for the control of *Salmonella typhimurium* infection in turkeys Am J Vet Research 15 122-129
- 145 LUKAS G N & D R BRADFORD 1954 Salmonellosis in turkey poult as observed at routine necropsy of 1144 cases J Am Vet Med Assoc 125 215-218
- 146 GALTON M M M HARLES & A V HARDY 1955 *Salmonella* isolations from dehydrated dog meals J Am Vet Med Assoc 126 57-58

- 147 GRIFFIN C A 1952 A study of prepared feeds in relation to *Salmonella* infection in laboratory animals J Am Vet Med 121 197-200
- 148 ROSENFELD I 1961 Personal communication
- 149 BORING J 1967 Personal communication
- 150 CRANT J 1950 Salmonellosis Med Officer 84 5-9 19 22
- 151 CHIFFER F S 1955 The acute diarrheal diseases of bacterial origin Bull N Y Acad Med Ser 2 31 611-676
- 152 MACKENZIE F F 1953 Typhoid and cocnut An unexpected and dangerous combination Health Bull Melbourne 110 19-23
- 153 KENZIE J & O T C OUCHTERLONY 1955 Salmonella is originating in a hospital A newly recognized source of infection New Engl J Med 253 761 763
- 154 BROWNE A S 1955 Personal communication
- 155 OLITSKY I A M FERRI M A SHEPPMAN & M WERRIN 1956 Smoked fish as a vehicle of salmonellosis Public Health Repts U S 71 713 719
- 156 SOLOWAY M & F J CALE NICK 1948 Survival of *Salmonella* in reconstituted egg powder subjected to holding and scrambling Food Research 13 216-226
- 157 WINTER A R G F STEWART A H MCFARLANE & M SOLOWAY 1946 Pasteurization of liquid egg products III Destruction of *Salmonella* in liquid whole egg Am J Public Health 36 451
- 158 CORFSLINE H F K M HAYES R F MOSER JR M A HOWE JR & E L DREW NAK 1951 Pasteurization of liquid whole egg under commercial conditions to eliminate *Salmonella* U S Dept Agr Circ 897
- 159 ILLFMAN G 1966 Uskadeliggdørelse af salmonellaaktener i ikke varmebehandlede levnedsmidler tillæret af rå hønseæg Nord Veterinarmed 8 859-874
- 160 HINSHAW W R & F McNEIL 1951 *Salmonella* infection as a food industry problem Advances in Food Research 3 209-240

THE EXOTIC ZONOSIS

By M. S. Shahan and J. Traum*

*Plum Island Animal Disease Laboratory Animal Disease and Parasite Research Division
Agricultural Research Service United States Department of Agriculture Greenport N. Y.*

According to Webster's International Dictionary (second edition 1933) zoonosis is defined either as a disease communicable from animals to each other or to man or as a disease due to animal parasites. In a broad sense this inclusive definition may be applied to almost any communicable disease of animals regardless of whether it is transmissible to man.

Despite this definition the term zoonosis appears only once in the titles of other papers presented in this monograph. In some circles there seems to be a feeling of dissatisfaction with the various applications of this term. This attitude has been voiced in conversations, published papers¹ and discussions. In this paper the word zoonosis when not otherwise qualified is used to refer to disease of animals.

It is not intended to develop a discussion on the use of the term itself but it is believed desirable to call attention to a recent article by Wagener in which he succinctly reviews the history, introduction, various applications and modifications of and substitutions for the word zoonosis. Wagener's own summary translated is as follows: 'The present significance of zoonosis is investigated. The new words anthroponosis and zooanthroponosis are discussed with regard to the epidemiological value and the linguistic significance. In order to distinguish the pathogenic effect on men (man) and animals diseases of animals infectious for men (man) are zooanthroponosis (zooanthroponoses), anthroponosis (anthroponoses) are those human diseases which are infectious for animals.'

In such a comprehensive monograph as this it is to be expected that there will be some degree of overlapping and duplication of the subject matter. Insofar as practicable references to diseases covered by other contributors have been omitted in this paper.

The importance of animals to human health has been expressed aptly by Middleton² as follows: 'When man domesticated certain lower animals for his personal comfort and gain he assumed the obvious hazard of sharing their diseases. So far as is known however animals are reservoirs of many but far from all of the diseases that affect man. On the other hand man may be the primary source of some diseases transmissible to the lower animals.'

As pointed out by Shope³ there is good indirect evidence to show that swine acquired influenza from man in 1918. An outbreak of influenza virus infection in a ferret colony has been attributed to human sources. It appears therefore that man and his influenza viruses constitute a health hazard for at least these two species of lower animals. Another example is tuberculosis of the human type which on occasion has complicated efforts to eradicate the disease in animals by means of the tuberculin test.⁴ Except for occasional

* Present address: Department of Bacteriology, University of California, Berkeley, Calif.

transitory sensitization to tuberculin infection with the human type of tuberculosis is of little importance in cattle although it may be so in some other species

As has long been known and as again reported recently ⁸ infection with the human type of *Mycobacterium tuberculosis* may on rare occasions persist for a long time in cattle whether acquired through exposure to human cases or by experimental inoculation Attempts to immunize cattle against bovine tuberculosis ⁷ ⁸ by injecting them with the human type of tubercle bacilli usually result in establishment of nonprogressive lesions in one or more organs that are difficult to detect by the most careful autopsy In a few instances it was found that the organism was shed in the milk for various lengths of time without showing detectable lesions in the udder Under such conditions bovine infection with human tuberculosis would be dangerous to consumers of raw milk and could produce slight lesions in suckling calves or render them tuberculin sensitive

In contrast with the e examples is rabies primarily a disease of lower mammals among which an increasingly greater number is being discovered as infected in nature Between these two extremes lies a broad field of interest essentially common both to veterinarians and others concerned primarily with animal diseases and to physicians and public health officials fundamentally responsible for human health Overemphasis of either of these basic interests can only be detrimental to both Consequently increasingly close cooperation among physicians veterinarians epidemiologists and epizootiologists is indicated

Concerning possibilities of disease in man attributable to infection from animals it may be stated that lack of proof of human infection with an agent known to be infectious for animals does not preclude the possibility of such a development to a greater or less extent at some future time Vesicular stomatitis a disease that has occurred sporadically in North America and in portions of Central and South America for decades affecting cattle horses and swine has been found capable of producing an influenza like syndrome in man A few years ago this virus would have been classed generally as noninfectious for man

The specific character of infectious processes in man may be undetermined or even ignored as in the case of animals Also to be considered is the interesting hypothesis that any animal may be attacked by any infectious agent under appropriate conditions

The growing speed and extent of world wide commerce and travel accentuate the danger of the introduction of new diseases whether they be localized in certain regions or be widespread From the broad international viewpoint infection in any quarter of the globe potentially may be carried to any other portion Diseases that are exotic or foreign to one area today may become epizootic or epidemic there in a very short time A notable example is the 1918 pandemic of human influenza

Meyer and Hjarre (cited by Wagener) report at least eighty diseases of animals that are transmissible to man Of these only a very small number are of importance from the standpoint of public health

THE EXOTIC ZOONOSSES

By M S Shahan and J Traum*

*Plum Island Animal Disease Laboratory Animal Disease and Parasite Research Division
Agricultural Research Service United States Department of Agriculture Greenport N Y*

According to Webster's International Dictionary (second edition 1933) 'zoonosis' is defined either as a disease communicable from animals to each other or to man or as a disease due to animal parasites. In a broad sense this inclusive definition may be applied to almost any communicable disease of animals regardless of whether it is transmissible to man.

Despite this definition the term zoonosis appears only once in the titles of other papers presented in this monograph. In some circles there seems to be a feeling of dissatisfaction with the various applications of this term. This attitude has been voiced in conversations, published papers,¹ and discussions. In this paper the word zoonosis when not otherwise qualified is used to refer to disease of animals.

It is not intended to develop a discussion on the use of the term itself but it is believed desirable to call attention to a recent article by Wagener² in which he succinctly reviews the history, introduction, various applications and modifications of and substitutions for the word zoonosis. Wagener's own summary translated is as follows: 'The present significance of zoonosis is investigated. The new words anthroozoonosis and zooanthroponosis are discussed with regard to the epidemiological value and the linguistic significance. In order to distinguish the pathogenic effect on men (man) and animals diseases of animals infectious for men (man) are zooanthroponosis (zooanthroponoses), anthroozoonosis (anthroozoonoses) are those human diseases which are infectious for animals.'

In such a comprehensive monograph as this it is to be expected that there will be some degree of overlapping and duplication of the subject matter. Insofar as practicable references to diseases covered by other contributors have been omitted in this paper.

The importance of animals to human health has been expressed aptly by Middleton³ as follows: 'When man domesticated certain lower animals for his personal comfort and gain he assumed the obvious hazard of sharing their diseases. So far as is known however animals are reservoirs of many but far from all of the diseases that affect man. On the other hand man may be the primary source of some diseases transmissible to the lower animals.'

As pointed out by Shope⁴ there is good indirect evidence to show that swine acquired influenza from man in 1918. An outbreak of influenza virus infection in a ferret colony has been attributed to human sources. It appears therefore that man and his influenza viruses constitute a health hazard for at least these two species of lower animals. Another example is tuberculosis of the human type which on occasion has complicated efforts to eradicate the disease in animals by means of the tuberculin test.⁵ Except for occasional

part of the nineteenth Century. Its prevalence here led to the establishment in 1884 in the United States Department of Agriculture of the Bureau of Animal Industry which was given responsibility for control of this and other animal diseases. This disease was subsequently eradicated completely from the United States through slaughter of affected and exposed animals. The disease has been eliminated elsewhere from North and South America and Western Europe but it persists in other parts of the world especially in Africa and Australia. Although the causative organism variously known as *Asterococcus mycoides*, *Bovimycetes pleuropneumoniae* and *Mycoplasma mycoides* is not known to affect man many so called pleuropneumonia like organisms (PPLO) have been discovered in cattle sheep goats dogs rodents and birds as well as in man in various parts of the world^{11, 12}. The classification and nomenclature of this large group of organisms are not firmly established. The latest classification based on recommendations of both F. A. Irendt and D. G. Edward is used by Merchant and Lacker¹³.

Glanders primarily a disease of the equidae was once rampant in the United States. This disease may now be classed as exotic to that country and to Canada although it persists in many other parts of the world. *Malleomyces mallei* causes a serious granulomatous tuberculosis like or ulcerative disease in man that is contracted directly from affected animals. Sheep goats dogs and cats occasionally contract the disease naturally. A disease etiologically and pathologically resembling glanders namely melioidosis recently has been identified in sheep goats and pigs in the Netherlands West Indies¹⁴. Infection due to the causative organism *M. pseudomallei* is said to produce infections in man chiefly in Southeast Asia. The presence of the disease which causes polyarthritis emaciation and death should be kept in mind in relation to suggestive unidentified infections of either man or animals.

Bluetongue a viral disease that affects sheep primarily and cattle occasionally was believed originally to be confined to the African continent. Lately it has spread to other countries in the Middle East and has become a major problem in the United States. Insofar as is known man is not susceptible to this infection. The disease is an example of what a few years ago would have been classed as exotic.

Another example of a formerly exotic animal disease that has become established in the United States presumably through importations of sheep is scrapie. As compared with other viral diseases there is an extremely extended incubation period in scrapie—18 months or longer in natural cases. This and its insidious nature result in major problems in relation to control and eradication. The disease is not known to affect man.

In Australia and the United States at about the same time in the late 1930s an acute febrile human illness was given the name Q fever. The causative organism *Coxiella burnetii* (*R. burnetii*) has been recovered from *Dermacentor andersoni*, *Imblyomma americanum*, *Haemaphysalis humerosa* and *H. bispinosa* and also from cattle sheep goats and various species of wild animals. The organism which is shed in milk has been found unusually resistant to common methods of pasteurization. Once considered exotic to the United States Q fever must now be classed as an indigenous disease of both man and animal¹⁵.

In a 265 page booklet entitled *Foreign Animal Diseases* ⁴ compiled by a committee of the United States Livestock Sanitary Association Trenton N J twenty six more or less important exotic animal diseases are listed as follows African swine fever Teschen disease bovine infectious petechial fever contagious bovine pleuropneumonia East Coast fever ephemeral fever infectious infertility of cattle, lumpy skin disease malignant catarrh rinderpest blue tongue contagious agalactia enzootic abortion louping ill Nairobi sheep disease sheep pox, scrapie heartwater nagana Rift Valley fever streptothricosis anthrax foot and mouth disease Venezuelan equine encephalomyelitis fowl plague and Newcastle disease Of these only seven (louping ill Rift Valley fever, Venezuelan equine encephalomyelitis foot and mouth disease sheep pox anthrax and Newcastle disease) are listed as communicable to man

Anthrax and Newcastle disease are included in the list of foreign diseases referred to above despite the fact that both are enzootic to the United States Anthrax is probably included because of the incrimination of imported bone meal as the source of outbreaks in animals and also because imported hides hair and wool are known to be responsible for anthrax in man In 1952 anthrax occurred extensively in animals in some areas of the United States particularly in the Middle Western states It is believed that at least one outbreak was caused by the feeding of imported bone meal containing anthrax spores Since then the United States Department of Agriculture regulations have been amended to provide that only steamed or degelatinized bone meal is eligible for unrestricted entry Such bone meal is commonly referred to as feeding bone meal Other kinds of bone meal are required to go to designated approved industrial establishments for further processing before they are released Bone meals that have been permitted unrestricted entry have been shown on analysis to contain not more than 2 per cent nitrogen and not less than 25 per cent phosphoric acid Such analyses indicate that the bone meal has been processed by cooking at 250 °F for at least 1 hour under 20 lb pressure

Newcastle disease of poultry so called because of its appearance in the vicinity of Newcastle upon Tyne in England was not recognized as such in the United States until about 1944 The disease had existed previously in California poultry for a number of years having been identified by the pathologically descriptive name pneumoencephalitis This malady recognized eventually as Newcastle disease spread throughout the United States in a generally mild form describing it from the standpoint of mortality A wide range of species of birds is susceptible and infection has been observed sporadically in man generally as the result of laboratory exposure¹¹ Although immunologically indistinguishable most strains of this virus in the United States may be differentiated from Asian and European strains on the basis of clinical manifestations pathology and mortality in inoculated birds Outbreaks due to foreign Newcastle disease viruses have been eradicated by destruction of affected and exposed birds

Among the animal diseases there are numerous examples of infections that have been spread through normal traffic channels Bovine pleuropneumonia long a scourge in Europe was introduced into the United States in the early

part of the nineteenth Century. Its prevalence here led to the establishment in 1884 in the United States Department of Agriculture of the Bureau of Animal Industry which was given responsibility for control of this and other animal diseases. This disease was subsequently eradicated completely from the United States through slaughter of affected and exposed animals. The disease has been eliminated elsewhere from North and South America and Western Europe but it persists in other parts of the world especially in Africa and Australia. Although the causative organism variously known as *Istero coccus mycoides*, *Botrymyces pleuropneumoniae* and *Mycoplasma mycoides* is not known to affect man many so called pleuropneumonia like organisms (PILO) have been discovered in cattle sheep goats dogs rodents and birds as well as in man in various parts of the world¹¹⁻¹⁸. The classification and nomenclature of this large group of organisms are not firmly established. The latest classification based on recommendations of both F. A. Irendt and D. C. Edward is used by Merchant and Packer¹⁹.

Glanders primarily a disease of the equidae was once rampant in the United States. This disease may now be classed as exotic to that country and to Canada although it persists in many other parts of the world. *Malleomyces mallei* causes a serious granulomatous tuberculosis like or ulcerative disease in man that is contracted directly from affected animals. Sheep goats dogs and cats occasionally contract the disease naturally. A disease etiologically and pathologically resembling glanders namely melioidosis recently has been identified in sheep goats and pigs in the Netherlands West Indies²⁰. Infection due to the causative organism *M. pseudomallei* is said to produce infections in man chiefly in Southeast Asia. The presence of the disease which causes polyarthritis emaciation and death should be kept in mind in relation to suggestive unidentified infections of either man or animals.

Bluetongue a viral disease that affects sheep primarily and cattle occasionally was believed originally to be confined to the African continent. Lately it has spread to other countries in the Middle East and has become a major problem in the United States. Insofar as is known man is not susceptible to this infection. The disease is an example of what a few years ago would have been classed as exotic.

Another example of a formerly exotic animal disease that has become established in the United States presumably through importations of sheep is scrapie. As compared with other viral diseases there is an extremely extended incubation period in scrapie—18 months or longer in natural cases. This and its insidious nature result in major problems in relation to control and eradication. The disease is not known to affect man.

In Australia and the United States at about the same time in the late 1930s an acute febrile human illness was given the name Q fever. The causative organism *Coxiella burnetii* (*R. burnetii*) has been recovered from *Dermacentor andersoni*, *Amblyomma americanum*, *Ixodes humerosus* and *I. bispinosus* and also from cattle sheep goats and various species of wild animals. The organism which is shed in milk has been found unusually resistant to common methods of pasteurization. Once considered exotic to the United States Q fever must now be classed as an indigenous disease of both man and animal²¹.

Sheep pox (*variola ovina*) is an acute highly lethal disease accompanied by formation of elementary bodies. It is a serious malady producing high mortality in Asia Southwest Europe and the Middle East. While it is a true pox in character it also appears to be peculiar to sheep and has been reported to be distinct from goat pox. Evidence that man may be affected is scanty.

The arthropod borne encephalitides have acquired major interest among epidemiologists and epizootiologists particularly since identification of equine encephalomyelitis virus in California by Meyer and his associates. Since this important finding two immunologically distinct types of the virus eastern and western have been identified in the United States and elsewhere particularly in the Western Hemisphere. A third immunologically different type the Venezuelan was identified in 1938²². Fortunately this disease in the strict sense an exotic zoonosis has not yet been diagnosed in the United States. Man is susceptible to all three of these viruses.

An endemic encephalomyelitis of horses and possibly of other animals commonly called Borna disease from a locality in Saxony where a severe epizootic occurred from 1894 to 1896 persists in Germany. It has a considerably longer incubation period than the encephalomyelitides known in the United States and is apparently distinct from them etiologically. It has not been reported to affect man.

Rift Valley fever sometimes called enzootic hepatitis an acute mosquito borne viral disease of sheep and cattle has not been identified outside of Africa where it exists particularly in and near the Great Rift Valley. It causes abortions in animals and is highly fatal especially to lambs. Shepherds farmers butchers and veterinarians often contract the disease in the course of handling infected animals or carcasses. The disease in man in whom it has an incubation period of 4 to 6 days resembles influenza. The temperature curve resembles that of dengue or yellow fever. It is rarely fatal but serious sequelae may occur²¹. The virus has been isolated from 8 or more native species of mosquitoes of 2 genera and experimental transmission has been accomplished with some species²³.

Laboratory workers have repeatedly become infected with the virus resulting in illness that cannot be considered inconsequential. While the disease has never occurred in animals in the United States or elsewhere outside of Africa there have been a number of laboratory infections in the United States and other countries to which the virus was imported for study. Sabin and Blumberg⁶ have reviewed cases that occurred in laboratories in various parts of the world including at least 8 cases in the United States. In one instance neutralizing antibodies were demonstrable 12 years after the attack.

A previously undescribed virus²⁷ was isolated in 1954 and 1955 from cases of a disease resembling Rift Valley fever that produced abortions in ewes and high mortality in lambs in Wesselsbron Orange Free State Union of South Africa. Serum neutralization and complement fixation tests proved this virus to be antigenically different from representative strains of Rift Valley fever virus. In fact the disease was identified in ewes previously vaccinated against Rift Valley fever. Neutralization tests of cattle sera collected rou

tinely for testing for brucellosis in many parts of the Orange Free State demonstrated antibodies against the W strain of this virus. On the basis of unpublished reports it seems that the virus is mosquito borne. It is important to note that during the course of study of the virus at the laboratory at Onderstepoort Transvaal Union of South Africa five of the six people engaged in the work became sick but not critically so. However muscular pain was experienced for a period of weeks.

Louping ill²⁴ a viral encephalomyelitis particularly of sheep and occasionally of cattle has been known for more than a century in Scotland and northern England more recently the disease has been diagnosed in the Union of Soviet Socialist Republics and Czechoslovakia. At times the disease produces heavy losses. Serologically the virus has been found to be related closely to that of Russian spring summer encephalitis. Similarly louping ill is tick borne chiefly by *Ixodes ricinus*. While the sheep disease has never been diagnosed in the United States several human infections have occurred in laboratories in the United States²⁵. As in sheep the disease in man is diphasic beginning with viremia and concluding after various secondary nervous manifestations. Natural infections among humans have also been reported^{29, 30}.

Foot and mouth disease (FMD) is an acute highly communicable disease affecting almost exclusively cloven footed animals both domesticated and wild. It is characterized by the formation of vesicles followed by erosions in the mucosa of the mouth (including the tongue lips cheeks gums dental pad and palate) and sometimes of the gastrointestinal tract. It affects particularly the rumen the skin (especially between and above the hoofs) the dew claws the teats and the udder.

Very few cases of FMD in man have been authenticated^{31, 32} and it has never become a public health problem except as the disease in animals may affect food supplies (milk and meat) it is mentioned here however because of its wide distribution and persistence in animals in Africa Asia Europe and South America. From time to time it has occurred in practically every country. It has invaded the United States nine times since 1870 the last occasion being in 1929. It was last eradicated from Canada in 1953 and from Mexico in 1954. The only major livestock producing areas presently free of the disease are North America Central America Australia and New Zealand.

England has succeeded repeatedly in eradicating FMD only to be reinfected through meat imported from infected countries or in other ways from nearby Europe³³. In countries where the disease becomes epizootic there is an estimated loss of 25 per cent in productivity of cattle swine sheep and goats not considering deaths which are usually 5 per cent or less but may be much greater.

Although there has been a total of nine outbreaks of FMD in the United States some very widespread there is not one recorded proved instance of the disease in man in this country. Vetterlein³² has recently reviewed critically the cases reported from 1926 to 1954. He accepts a total of 21 cases as having been appropriately proved. Recently Ratner *et al*³⁴ have described a very unusual lingering case in the Soviet Union from which type A virus was identified. The virus is reported to have been obtained on the 25th and 159th days

from lesions on the foot on the 172nd day from the saliva and finally on the 178th day from saliva from so called dry lesions on the palms of the hand and from the feces. The character of the lesions and the long duration of the illness are distinctly different from any previously described case.

In addition to the diseases and agents previously referred to mention should be made of others.

Discovery of yellow fever antibodies in man in regions in South America where *Aedes aegypti* were not universally present led to identification of the so called jungle yellow fever as distinguished from the classic urban type transmitted by that vector. Yellow fever antibodies have been found in various species of monkeys other primates and marsupials in these same jungle areas and in the absence of *A. aegypti* this type of the disease may thus be classed as a zoonosis that is exotic to North America.

Because they are exotic to the United States some other animal diseases not known to affect man but recognized as serious livestock problems should be mentioned. Perhaps the most important of these is rinderpest or cattle plague one of the most destructive of all animal plagues which persists in Africa Asia and Eastern Europe. Except for a single occurrence in Brazil the disease has never appeared in the Western Hemisphere. Although it is essentially a bovine disease it sometimes affects other ruminants and swine. In many outbreaks mortality exceeds 90 per cent. There is considerable variation in susceptibility among various breeds and types of cattle. Accordingly for immunization purposes varying virulent virus is used after being adapted to chicken embryos rabbits or goats. Some cattle may be so highly susceptible as to require simultaneous administration of antiserum with even the most attenuated virus in order to eliminate losses due to attempted immunization. There is no evidence of human infections with this virus.

So called Teschen disease⁴ a viral encephalomyelitis of swine is reported to be quite widespread in Czechoslovakia where it was first discovered about 1929 near the town of Teschen whence it has spread to other countries in Central and Western Europe. Some of the similarities between this disease and poliomyelitis have been explored and there is apparently no relationship between the two viruses.³⁵

African swine fever⁴ is an acute highly fatal septicemic viral disease of swine that symptomatically and pathologically closely resembles hog cholera or as it is designated in many countries other than the United States swine fever. The African disease is not known to exist beyond that continent. Present information indicates that like hog cholera the disease is not transmissible to man.

Wild bush pigs or wart hogs have been found to be symptomless carriers of the virus of African swine fever. Domestic swine frequently contract the disease through contact with such feral animals. Biologicals that protect against hog cholera have been found ineffective as preventives of African swine fever. Indeed no effective means of prophylaxis other than avoiding contact with wild pigs or wart hogs and the eradication of affected domestic herds has yet been developed.

The broad scope of the title of this paper would permit continuation beyond

the space allotted however the diseases omitted here or skipped over in a manner that may appear cursory are covered elsewhere in these pages. As has been pointed out by Hagan²² in addition to the human health hazard animal diseases also affect the welfare of man through their serious economic effects. It has been estimated²³ that in the United States death losses among livestock from all causes in 1934 amounted to 1.5 million cattle 2.5 million calves 4 million sheep 10.5 million swine 23.5 million chickens and 7.2 million turkeys. These figures do not reflect the whole toll since they do not include losses in other species feed labor and capital investment nor in untold numbers of newborn animals. It is estimated that 20 to 35 per cent of the pigs born never reach weaning age. Internal parasites have been estimated to exact 400 million dollars annually from incomes from livestock. Some of these total losses involve diseases dangerous to man but many do not.

Before concluding it is considered appropriate to mention an important aspect of the animal disease problem. On October 8 1932 supplemented May 12 1933 notices were sent to laboratories research institutions and individuals studying animal diseases to advise that under regulations of the Department of Agriculture it is necessary to obtain authorization by the Secretary of Agriculture for the importation and interstate transportation of certain animal disease organisms and vectors.²⁴ This notice which was widely published in many scientific journals and other publications directed attention to the inherent danger of such movements and to the urgent need for precautions against the spread of animal disease.

While the department has the responsibility for safeguarding our livestock and poultry by the prevention and control of domestic diseases as well as those of foreign origin it is not the intention to interfere unduly with the normal movement of specimens to diagnostic laboratories and research institutions or the transfer of organisms and viruses indigenous to this country between research workers and laboratories. For example permits are not required for the interstate transportation of members of the *Listerella* group *Salmonella* group *Clostridium* group Newcastle disease virus of mild virulence the viruses of such diseases as fowl pox hog cholera and swine influenza or of canine distemper. Permits are required in advance for the interstate movement of such agents as Venezuelan equine encephalomyelitis virus highly virulent strains of Newcastle disease virus *Plasmodium burgheri* the viruses of blue tongue or scrapie vesicular disease viruses and any other foreign animal disease viruses organisms or vectors that may have gained entrance into the United States accidentally or otherwise. Permits are required for the importation of any organisms or vectors that may transmit diseases to our animal and poultry population.

It is evident that absolute protection against the introduction and dissemination of foreign diseases is virtually impossible however every reasonable precaution should be taken toward this end. In order to protect the nation's livestock industry the Secretary of Agriculture is empowered to promulgate rules and regulations which may be modified in accordance with changing conditions and requirements.²⁵

References

- 1 MEYER K F 1953 Animal disease and human welfare *Advances in Veterinary Science* 1 1-40 Academic Press New York N Y
- 2 WAGENFUR K 1957 Zoonosen anthroozoonosen zooanthroponosen Berlin u Munch tierarztl Wochschr 70 12-13
- 3 MIDDLETON W S 1956 Zoonosis—a serious medical problem *Wisconsin Med J* 55 1189-1196
- 4 SHOPE R E 1955 Epizootiology of virus diseases *Advances in Veterinary Science* 2 32-33 Academic Press New York N Y
- 5 FELDMAN W H & H MOSES 1941 Human tuberculosis in a bovine case report of a spontaneous infection in an adult bovine *Am Rev Tuberc* 43 418-424
- 6 NASSAL J 1956 Humane Tuberkuloseinfektion beim Rind mit Ausscheidung der Erreger durch das unveränderte Futter *Rindertuberkulose* 5 189-198
- 7 HARING C M & J TRAUM 1913-1917 *Repts Calif Agr Expt Station*
- 8 COBBETT L 1917 The danger of immunizing milch cows with living tubercle bacilli *In The Causes of Tuberculosis* 674-679 Cambridge Univ Press Cambridge England
- 9 BRANDLY C A 1952 Newcastle disease *In Diseases of Poultry* H F Biester and L H Schwarte Fds 531-568 Iowa State Coll Press Ames Iowa
- 10 BLOOD B D 1950 Epidemiology of Newcastle disease *Bull Pan Am Sanit Bur* 29 78-49
- 11 SABIN A B 1941a The filtrable micro organisms of the pleuropneumonia group *Bacteriol Revs* 5 1
- 12 SABIN A B 1941b The filtrable micro organisms of the pleuropneumonia group (appendix on classification and nomenclature) *Bacteriol Revs* 5 331
- 13 EDWARD D G 1954 The pleuropneumonia group of organisms a review together with some new observations *J Gen Microbiol* 10 27-64
- 14 CORDY D R H E ADLER & J BERG 1957 The pathogenicity for swine of pleuropneumonia like organisms from goats *Cornell Vet In press*
- 15 CORDY D R & H E ADLER 1957 The pathogenesis of the encephalitis in turkeys produced by a neurotropic pleuropneumonia like organism *Avian Dis* 1 235-245
- 16 ADLER H E & R YAMAMOTO 1957 Pathogenic and nonpathogenic pleuropneumonia like organisms in infectious sinusitis of turkeys *Am J Vet Research* 18 655-656
- 17 MOULTON J E & H E ADLER 1957 Pathogenesis of arthritis in chicken embryos caused by a pleuropneumonia like organism *Am J Vet Research* 18 731-734
- 18 ADLER H E & R YAMAMOTO & J BERG 1957 Strain differences of pleuropneumonia like organisms of avian origin *Avian Dis* 1 19-76
- 19 MERCHANT I A & R A PACKER 1956 *Veterinary Bacteriology and Virology* 5th ed Iowa State Coll Press Ames Iowa
- 20 STUMOLLER P I C KRANEVELD & A VAN DER SCHAAF 1957 Melioidosis (pseudomalleus) in sheep goats and pigs on Aruba (Netherlands Antilles) *J Am Vet Med Assoc* 120 415-417
- 21 DYER R E 1955 Q fever *Diseases of Animals Transmitted from Animals to Man* 597-607 T G Hull Ed 4th ed Thomas Springfield Ill
- 22 MEYER K F C M HARING & B HOWITT 1930 The etiology of epizootic encephalomyelitis of horses in the San Joaquin Valley *Science* 74 227-228
- 23 BECK C E & I W G WYCKOFF 1938 Venezuelan equine encephalomyelitis *Science* 88 530
- 24 REPORT OF COMMITTEE ON EXOTIC DISEASES 1954 U S Livestock San As oc Trenton N J
- 25 SMITHBURN K C A J HADDOW & W H R LUMSDEN 1949 Rift Valley fever transmission of virus by mosquitoes *Brit J Exptl Pathol* 30 35-47
- 26 SABIN A B & K W BLUMBERG 1947 Human infection with Rift Valley fever virus and immunity twelve years after single attack *Proc Soc Exptl Biol Med* 64 385-389
- 27 WEISS K E D A HAIG & R A ALEXANDER 1956 Wesselsbron virus—a virus not previously described associated with abortion in domestic animals *Onderstepoort J Vet Research* 27 183-195
- 28 RIVERS T M 1952 *Viral and Rickettsial Infections of Man* 2nd ed Lippincott Philadelphia Pa
- 29 BRENTS E G C NEWBAUM & F W HIRST 1949 Another case of louping ill in man Isolation of the virus *Lancet* 256 689-691
- 30 LAWSON J H W G MANDERSON & E W HIRST 1949 Louping ill meningoencephalitis a further case and a serological survey *Lancet* 257 696-699

- 31 TRAUM J 1952 Foot and Mouth Disease A Textbook of Medicine 49-51 Cecil and Loeb Eds 8th ed Saunders Philadelphia Pa
- 32 VETTERLEIN W 1954 Das klinische Bild der Maul und Klauenseuche beim Menschen aufgestellt aus den bisher experimentell gesicherten Erkrankungen Arch Exptl Vet Med 8 540-564
- 33 REPORT OF THE DEPARTMENTAL COMMITTEE ON FOOT AND MOUTH DISEASE 1952-1954 1956 Ministry of Agriculture and Fisheries H M Stationery Off London England
- 34 RATNER S I C P KOROLEV G V GUBIN & R P KOMOLOVA 1956 A case of lingering foot and mouth disease in man (trans from Russian) Klin Med 34 10-77
- 35 HORSTMAN D M 1952 Experiments with Teschen disease (virus encephalomyelitis of swine) J Immunol 69 319-394
- 36 HAGAN W A 1947 The relation of diseases in the lower animals to human welfare Ann N Y Acad Sci 48(6) 353-359
- 37 NORDQUIST A V & C H PALS 1956 Economic losses from animal diseases and parasites Yearbook Agr U S Dept Agr 11 14
- 38 U S DEPT AGR Mimeo CIRC SFR 1957 ZV Oct 8 (revised May 12 1953)
- 39 GOOPING C L 1956 Quarantining Yearbook Agr U S Dept Agr 62 70

References

- 1 MEYER K F 1953 Animal disease and human welfare *Advances in Veterinary Science* 1 1-40 Academic Press New York N Y
- 2 WAGENER K 1957 Zoonosen anthroozoonosen zooanthropozosen Berlin u Munch tierarztl Wochschr 70 12-13
- 3 MIDDLETON W S 1956 Zoonosis—a serious medical problem *Wisconsin Med J* 55 1189-1196
- 4 SHOPE R E 1955 Epizootiology of virus diseases *Advances in Veterinary Science* 2 32-33 Academic Press New York N Y
- 5 FELDMAN W H & H MOSES 1941 Human tuberculosis in a bovine case report of a spontaneous infection in an adult bovine *Am Rev Tuberc* 43 418-424
- 6 NASSAL J 1956 Humane Tuberkuloseinfektion beim Rind mit Ausscheidung der Erreger durch das unveränderte Euter Rindertuberkulose 5 189-198
- 7 HARING C M & J TRAUM 1913-1917 Repts Calif Agr Expt Station
- 8 COBBETT L 1917 The danger of immunizing milch cows with living tubercle bacilli *In The Causes of Tuberculosis* 674-679 Cambridge Univ Press Cambridge England
- 9 BRANDLY C A 1952 Newcastle disease *In Diseases of Poultry* H F Biester and L H Schwarte Eds 531-568 Iowa State Coll Press Ames Iowa
- 10 BLOOD B D 1950 Epidemiology of Newcastle disease *Bull Pan Am Sanit Bur* 29 28-49
- 11 SABIN A B 1941a The filtrable micro organisms of the pleuropneumonia group *Bacteriol Revs* 5 1
- 12 SABIN A B 1941b The filtrable micro organisms of the pleuropneumonia group (appendix on classification and nomenclature) *Bacteriol Revs* 5 331
- 13 EDWARD D G 1954 The pleuropneumonia group of organisms a review together with some new observations *J Gen Microbiol* 10 27-64
- 14 CORDY D R H E ADLER & J BERG 1957 The pathogenicity for swine of pleuropneumonia like organisms from goats *Cornell Vet In press*
- 15 CORDY D R & H E ADLER 1957 The pathogenesis of the encephalitis in turkeys produced by a neurotropic pleuropneumonia like organism *Avian Dis* 1 235-245
- 16 ADLER H E & R YAMAMOTO 1957 Pathogenic and nonpathogenic pleuropneumonia like organisms in infectious sinusitis of turkeys *Am J Vet Research* 18 655-656
- 17 MOULTON J E & H I ADLER 1957 Pathogenesis of arthritis in chicken embryos caused by a pleuropneumonia like organism *Am J Vet Research* 18 731-734
- 18 ADLER H E R YAMAMOTO & J BERG 1957 Strain differences of pleuropneumonia like organisms of avian origin *Avian Dis* 1 19-26
- 19 MERCHANT I A & R A PACKER 1956 *Veterinary Bacteriology and Virology* 5th ed Iowa State Coll Press Ames Iowa
- 20 SUTMOLLER P F C KRAVEVELD & A VAN DER SCHAAF 1957 Melioidosis (pseudomalleus) in sheep goats and pigs on Aruba (Netherlands Antilles) *J Am Vet Med Assoc* 130 415-417
- 21 DYER R F 1955 Q fever *Diseases of Animal Transmitted from Animals to Man* 597-602 T G Hull Ed 4th ed Thomas Springfield Ill
- 22 MEYER K F C M HARING & B HOWITT 1930 The etiology of epizootic encephalomyelitis of horses in the San Joaquin Valley *Science* 74 221-228
- 23 BECK C F & R W G WICKOFF 1938 Venezuelan equine encephalomyelitis *Science* 88 530
- 24 REPORT OF COMMITTEE ON EXOTIC DISEASES 1954 U S Livestock San Assoc Trenton N J
- 25 SMITHBURN K C A J HADDOW & W H R LUMSDEN 1949 Rift Valley fever transmission of virus by mosquitoes *Brit J Exptl Pathol* 30 35-47
- 26 SABIN A B & R W BLUMBERG 1947 Human infection with Rift Valley fever virus and immunity twelve years after single attack *Proc Soc Exptl Biol Med* 64 385-389
- 27 WEISS K F D A HAIG & R A ALEXANDER 1956 Wesselsbron virus—a virus not previously described associated with abortion in domestic animals *Onderstepoort J Vet Research* 27 183-193
- 28 RIVERS T M 1952 *Viral and Rickettsial Infections of Man* 2nd ed Lippincott Philadelphia Pa
- 29 BREWIS F G C NEWBALER & F W HURST 1949 Another case of louping ill in man Isolation of the virus *Lancet* 256 689-691
- 30 LAWSON J H W G MANDERSON & F W HURST 1949 Louping ill meningoencephalitis a further case and a serological survey *Lancet* 257 696-699

tions caused by *Listeria* in ruminants, swine and men are meningoencephalitis, septicemia and abortion.

Since Nyfeldt isolated *Listeria* from humans with a disease indistinguishable from infectious mononucleosis,⁴ numerous other cases have been reported.⁴⁵⁻⁴⁹ *Listeria* infections usually do not give a positive Paul Bunnell reaction and produce a relatively high infant mortality, while infectious mononucleosis is usually benign and the affected individual recovers without specific treatment.

Conjunctivitis with or without mononucleosis is another manifestation of *Listeria* infection. This has occurred as a result of laboratory infection⁵ and a granulomatous type has been reported in a group of poultry processors.⁶ The ability of *Listeria* to produce conjunctivitis in rabbits and guinea pigs makes it a useful diagnostic tool.⁹

Septicemia is probably the most commonly occurring form of listeriosis in man.⁴¹ The resulting involvement of the lymph glands and the reticuloendothelial system stimulates the production of monocytes, thus causing confusion in differentiating the disease from infectious mononucleosis. Monocytosis is not always present and in these cases a fever, leukocytosis and swollen liver and spleen are predominant symptoms.

A disease of the newborn described as a granulomatous septicemia has been described by Potel.⁶⁻⁴² It is highly fatal to the infant but curiously enough it does not seriously affect the mother. Often the mothers deny any symptoms compatible with listeriosis. Thorough questioning frequently establishes some influenzalike symptoms a few days before occurrence of the abortion or premature birth. Slight chills, light lumbar pains and a slight elevation of temperature usually are all that can be ascribed to the infection in these mothers. If the infants survive the first septicemic stage they may develop meningitis.⁴⁴ While the majority of *Listeria* meningitis cases occur in the young, it is by no means confined to children.⁴⁵ Kaplan has listed numerous cases of meningitis due to *Listeria* and many are in the adult age group.⁴⁶

Therapeutic agents have been evaluated in both animals and man.^{1, 28, 47-49} and the following assumptions can be made from the results obtained and reported in recent literature. The antibiotics of the tetracycline group are the drugs of choice for the treatment of listeriosis, but penicillin and the sulfonamides in combination proved to be almost as effective. An excellent summary of human *Listeria* cases treated with various therapeutic agents has been published by Seeliger.⁴⁵ The *in vitro* sensitivity test gives some indication of the agent of choice. Certainly chemotherapy and antibiotic therapy have increased the odds in favor of recovery from malignant listeriosis when such treatment is begun early and with sufficient amounts to bring blood levels up to an effective listeristatic or listeriocidal concentration. Untreated cases of meningitis, polyserositis and overt septicemic disease usually are fatal.

Vaccination of animals against listeriosis has been attempted.⁵⁰⁻⁵¹ The results of these trials have not met with any great degree of success. The economic value of sheep and cattle populations together with public health considerations indicate strongly that these investigations should be continued.

LISTERIOSIS A POTENTIAL PUBLIC HEALTH PROBLEM

By John F. Winn, William B. Cherry and Elizabeth O. King

Laboratory Branch Communicable Disease Center Public Health Service Atlanta Ga

Listeriosis is a little known infrequently recognized but widespread bacterial disease of man and other warm blooded creatures. The causative agent, *Listeria monocytogenes* has been recovered from man other mammals and birds in the following countries Australia¹ Belgium² Brazil³ Canada⁴ Ceylon⁵ Denmark⁶ Germany⁷ Uruguay⁸ Austria⁹ Argentina¹⁰ England¹¹ Finland¹² France¹³ The Netherlands¹⁴ India¹⁵ Israel¹⁶ Japan¹⁷ New Zealand¹⁸ Norway¹⁹ Poland²⁰ Sweden²¹ Switzerland²² the Union of Soviet Socialist Republics²³⁻²⁵ South Africa²⁶ Czechoslovakia²⁷ Turkey²⁸ and the United States²⁹. This indicates that, geographically, this is certainly a ubiquitous organism. Many species of animals have served as host for the bacterium. *Listeria monocytogenes* has been isolated most often from sheep¹⁻³ cattle⁷⁻⁹ and man^{20,29} but occasionally from foxes³⁰ chinchillas⁴⁰ lemmings⁴¹ rabbits¹¹ swine¹¹ horses¹⁵ canaries⁴ hares⁴ chickens⁴³⁻⁴⁴ goats²⁹ ducks⁴⁵ geese⁵ eagles⁴⁶ turkeys⁴⁷ heath cocks⁴⁸ dogs⁴⁹⁻⁵⁰ deer⁵¹ cats⁵¹ field mice⁵ ferrets⁵² house mice⁵³ voles¹⁶ gerbils³ skunks⁴⁴ and raccoons⁵⁵. Laboratory animals such as the white mouse⁵ and guinea pig³ also have been shown to be susceptible to *Listeria*.

Murray Webb and Swann isolated the organism in 1924 from laboratory rabbits with a fatal septicemic disease. Because of the monocytosis produced in this species these investigators tentatively called the organism *Bacterium monocytogenes*¹¹. Three years later Pirie found a similar disease in South African rodents that was caused by the same type of organism which he called *Listerella hepatolytica*³. English investigators agreed to call the organism *Listerella monocytogenes* but because this name is a homonym of a mycetozoon Pirie suggested *Listeria monocytogenes* which is now the accepted name⁵⁷.

Morphologically the organism is a Gram positive nonacid fast nonspore forming rod. There is a tendency for the organisms to arrange themselves into V's or into a palisade formation when the smear is made from solid media. The bacteria stain evenly and are 0.5 to 0.6 by 1.0 to 2.0 μ in size. Short but sometimes long threadlike chains may be observed in the rough phase of growth. Capsules are not demonstrable. Motility can be demonstrated in young (6-hour) glucose broth cultures but is more pronounced at 25°C after 24 hours incubation. The peritrichously flagellated organisms have a characteristic tumbling motion.

An interesting and important cultural characteristic of *Listeria* is its frequent failure to grow when cultured from fresh tissue. This unusual behavior may account to some extent for failures to diagnose listeriosis in human disease. It has been found that refrigeration of the original necropsy specimens and periodic subculturing will increase the number of isolations obtained⁴⁶.

Listeriosis in sheep and cattle usually is recognized readily and the etiological agent is frequently recovered without difficulty. The main clinical manifesta-

spread might be venereal in nature cannot be overlooked. *Listeria* has been recovered from the urethra of some young adult males²⁴ and we must not forget that the meconium of infected prematurely born infants is swarming with *Listeria* and that isolations have been made from the placenta, the cervix and the vagina of human and animal females^{25, 26, 27, 28, 29}.

Two recent reports of human infections following direct contact with animals moribund with listeriosis strongly suggest that this mode of transmission is possible^{30, 31}. It may be, however, that the source of infection for the individuals concerned is as common to both the livestock and the animal husbandry man.

Nonpasteurized milk may contain *Listeria*³². Seeliger points out that in East Germany, where milk was still being rationed in 1955, unpasteurized milk was frequently illegally purchased directly from farmers. Human cases of listeriosis are frequent in this area, as are animal infections. The sale of unpasteurized milk on the black market may be a source of infection for humans³³.

Another interesting observation was a number of cases among shoe repair men³⁴. These individuals might have become infected from the dirt and manure brought in on shoes to be repaired. One can speculate that in this instance the disease was contracted via the airborne route by means of the dust created during the repair process. It is entirely possible that unclean hands could have contaminated food eaten by the individuals, or that tacks customarily carried in the mouths of repairmen might be a means of spreading contamination.

There are four serotypes of *Listeria*, and among these are several biotypes^{35, 36}. All serotypes and biotypes can and do attack man, poultry and mammals. Serotype 4 has been divided into 4a and 4b³⁷. The most prevalent serotypes are 1 and 4b³⁸. There does not appear to be any association of serotypes either to the species of animal or to the clinical type of disease. The primary value of serotyping and biotyping is for epidemiological purpose. For example, serologic surveys show more rural children with agglutinating antibodies than are present in urban children³⁹.

Agglutinating antibodies against *Listeria* differ in amount. For example, Lang demonstrated that about 60 per cent of several hundred children 3 to 10 years of age had antibodies against somatic or flagellar antigens with titers ranging from 1:40 to 1:160. Only one third of older children had such titers⁴⁰. The specificity of agglutinating titers is open to question, since some sera spontaneously agglutinate antigen and *Listeria* shares antigens with other bacteria such as the staphylococci and *Streptococcus fecalis*⁴¹. This being the case, agglutinating titers must be interpreted with caution and preferably compared with another type of serologic test using a different antibody system such as the complement fixation test.

The identification of any organism found in the spinal fluid or blood of patients is desirable; likewise, cultures from necropsy specimens should be identified. *Listeria* cultures are sometimes discarded because the bacteriologist, being unfamiliar with the organism, occasionally considers it to be a diphtheroid.

Whether or not an infectious disease constitutes a public health problem depends upon its appearance within a population the severity of the disease, and the nature of its epidemiology. Therefore by definition listeriosis constitutes such a problem because we do not know its frequency of occurrence nor do we understand the epidemiology of the disease. We do know that it occurs frequently in livestock that it is often fatal to both animals and man and that its role in human infections requires clarification.

The geographical distribution and host range of the disease have been mentioned. Other factors such as season sex and age distribution persistence and serotypes are of interest in solving the problem.

There is a definite seasonal distribution in animal infections^{26 7 9 77}. The greatest number of cases occurs during the winter and spring months. It would appear that this is not true of human infections. *Listeria* cultures referred to the Communicable Disease Center laboratory for identification or verification were slightly more numerous during the cool months. A breakdown by months gives the following figures: January 8 February 2 March 6 April 4 May 4 June 3 July 4 August 4 September 6 October 10 November 7 and December 7. Case reports published^{30 33 35 37 65 66 77} that give the dates of admission to the hospital show the following distribution: January 0 February 3 March 0 April 0 May 1 June 0 July 0 August 3 September 2 November 1 and December 1.

It is possible that neither the sex distribution nor the age distribution of the disease in animals and man is significantly different. It must be emphasized however that in human infection mothers frequently have no definite symptoms pointing to listeriosis and consequently may be overlooked as a source of infection for the infant. It may be that sex does not influence the course of the disease but that age is a factor in determining whether the course of the disease is fatal that is the younger the animal or human the less favorable the prognosis. The sex distribution of listeriosis in livestock is influenced by husbandry practices. For example the number of male animals employed usually is 1 bull for 25 to 50 cows and 1 ram to 25 to 35 ewes. Therefore the small number of cases in noncastrated male animals is easily explained.

Direct contact seems to play a part in the spread of infection among animals. Those animals that shared a common feed rack but that were separated by a fence from infected animals did not become infected while those in the same pen did become diseased. Attempts to recover the organism from straw litter drinking water and feed were unsuccessful. However it was found that *Listeria* could survive for long periods of time under barn conditions. The shortest period of survival was six weeks on straw and wood shavings and the longest time was twenty six weeks on rabbit food pellets⁷⁷. Other studies have shown that it is possible to infect animals through seeding their drinking water with cultures of *Listeria*⁷⁷.

Human infections while usually sporadic have occurred as epidemics. The exact mechanism of spread other than by direct contact was not apparent. *Listeria monocytogenes* was recovered from the blood and also from the throat washings of 8 of 26 individuals with overt illness⁷³. The possibility that human

spread might be venereal in nature cannot be overlooked. *Listeria* has been recovered from the urethra of some young adult males⁷⁴ and we must not forget that the meconium of infected prematurely born infants is swarming with *Listeria* and that isolations have been made from the placenta, the cervix and the vagina of human and animal females.⁷⁵ " " " "

Two recent reports of human infections following direct contact with animal moribund with listeriosis strongly suggest that this mode of transmission is possible.⁷⁶⁻⁷ It may be however that the source of infection for the individuals concerned was common to both the livestock and the animal husbandry man.

Nonpasteurized milk may contain *Listeria*.¹⁹ Seeliger points out that in East Germany where milk was still being rationed in 1955 unpasteurized milk was frequently illegally purchased directly from farmers. Human cases of listeriosis are frequent in this area as are animal infections. The sale of unpasteurized milk on the black market may be a source of infection for humans.⁴

Another interesting observation was a number of cases among shoe repairmen.⁴⁵ These individuals might have become infected from the dirt and manure brought in on shoes to be repaired. One can speculate that in this instance the disease was contracted via the airborne route by means of the dust created during the repair process. It is entirely possible that unclean hands could have contaminated food eaten by the individual or that tacks customarily carried in the mouths of repairmen might be a means of spreading contamination.

There are four serotypes of *Listeria* and among these are several biotypes.⁷⁸⁻⁷⁹ All serotypes and biotypes can and do attack man, poultry and mammals. Serotype 4 has been divided into 4a and 4b.⁴⁵ The most prevalent serotypes are 1 and 4b.⁴⁵ There does not appear to be any association of serotypes either to the species of animal or to the clinical type of disease. The primary value of serotyping and biotyping is for epidemiological purposes. For example serologic surveys show more rural children with agglutinating antibodies than are present in urban children.⁸¹

Agglutinating antibodies against *Listeria* differ in amount. For example Lang demonstrated that about 65 per cent of several hundred children 3 to 10 years of age had antibodies against somatic or flagellar antigens with titers ranging from 1:40 to 1:160. Only one third of older children had such titers.⁸¹ The specificity of agglutinating titers is open to question since some sera spontaneously agglutinate antigen and *Listeria* shares antigens with other bacteria such as the staphylococci and *Streptococcus fecalis*.⁸ This being the case agglutinating titers must be interpreted with caution and preferably compared with another type of serologic test using a different antibody system such as the complement fixation test.

The identification of any organism found in the spinal fluid or blood of patients is desirable likewise cultures from necropsy specimens should be identified. *Listeria* cultures are sometimes discarded because the bacteriologist being unfamiliar with the organism occasionally considers it to be a diphtheroid.

Whether or not an infectious disease constitutes a public health problem depends upon its appearance within a population the severity of the disease and the nature of its epidemiology. Therefore by definition listeriosis constitutes such a problem because we do not know its frequency of occurrence nor do we understand the epidemiology of the disease. We do know that it occurs frequently in livestock that it is often fatal to both animals and man and that its role in human infections requires clarification.

The geographical distribution and host range of the disease have been mentioned. Other factors such as season, sex and age distribution, persistence and serotypes are of interest in solving the problem.

There is a definite seasonal distribution in animal infections.^{6, 7, 23, 24} The greatest number of cases occurs during the winter and spring months. It would appear that this is not true of human infections. *Listeria* cultures referred to the Communicable Disease Center laboratory for identification or verification were slightly more numerous during the cool months. A breakdown by months gives the following figures: January 8, February 2, March 6, April 4, May 4, June 3, July 4, August 4, September 6, October 10, November 7 and December 7. Case reports published^{20, 22, 25, 27, 45, 46, 7} that give the dates of admission to the hospital show the following distribution: January 0, February 3, March 0, April 0, May 1, June 0, July, 0, August, 3, September 2, November 1 and December 1.

It is possible that neither the sex distribution nor the age distribution of the disease in animals and man is significantly different. It must be emphasized however that in human infection mothers frequently have no definite symptoms pointing to listeriosis and consequently may be overlooked as a source of infection for the infant. It may be that sex does not influence the course of the disease but that age is a factor in determining whether the course of the disease is fatal; that is, the younger the animal or human, the less favorable the prognosis. The sex distribution of listeriosis in livestock is influenced by husbandry practices. For example, the number of male animals employed usually is 1 bull for 25 to 50 cows and 1 ram to 25 to 35 ewes. Therefore the small number of cases in noncastrated male animals is easily explained.

Direct contact seems to play a part in the spread of infection among animals. Those animals that shared a common feed rack but that were separated by a fence from infected animals did not become infected, while those in the same pen did become diseased. Attempts to recover the organism from straw, litter, drinking water and feed were unsuccessful. However, it was found that *Listeria* could survive for long periods of time under barn conditions. The shortest period of survival was six weeks on straw and wood shavings and the longest time was twenty-six weeks on rabbit food pellets.²⁷ Other studies have shown that it is possible to infect animals through seedling, their drinking water with cultures of *Listeria*.⁷

Human infections, while usually sporadic, have occurred as epidemics. The exact mechanism of spread other than by direct contact was not apparent. *Listeria monocytogenes* was recovered from the blood and also from the throat washings of 8 of 26 individuals with overt illness.²³ The possibility that human

- 8 PORZECALSKI B & C DE BAIGORRIA 1938 A proposito de su constatacion en el Uruguay de una meningocefalitis humana Arch soc Biol Montevideo 9 99
- 9 ANTON W 1934 Kritisch experimenteller Beitrag zur Biologie seiner Beziehung zur infektiösen Mononukleose des Menschen Zentr Bakteriell Parasitenk Abt I 131 89-103
- 10 SAVINO F 1940 Observacion de listeriosis humana y accion terapeutica del yage nan Semana medica Buenos Aires 1 336-340
- 11 MURRAY F D C R A WFB & M B R SWANN 1976 A disease of rabbits characterized by a large mononuclear leukocytosis caused by a hitherto undescribed bacillus *Bacterium monocyti* J Pathol Bacteriol 29 40-439
- 12 KOINE P A RAITIO & U YAROKAARA 1953 Listeria infection in the guinea pig caused by feeding with Aureomycin Nature 172 61-68
- 13 FORCET P C TRICHE A STROUB & R LARRY 1941 Bull Acad Vet 14 195 (Cited by Seeliger⁴⁴)
- 14 DEBLIECK L & J JANSEN 1943 Listeriose bei Tieren J Mikrobiol Serol 9 93-103
- 15 VISWANATHAN C K & V AYYAR 1950 Indian Vet J 26 395 (Cited by Seeliger⁴⁴)
- 16 LEVI M L 1948 *Listeria monocytogenes* in voles Vet J 104 310-317
- 17 TAJIMA M Y FUJIMOTO & I HIGURO 1951 J Vet Med Japan 56 15 (Cited by Seeliger⁴⁴)
- 18 GRINF O 1943 *Listerella monocytogenes* som arsak til septicopyemi hos fall Norsk Vet Tidsskr 55 9-104
- 19 WRAMBY B O 1944 Om *Listerella monocytogenes* bakterier og om forkomst av *Listerella* infektion hos dyr Skand Vet Tidsskr 34 277-29
- 20 SCHLEGEL & FEY (Quoted by Seeliger⁴⁴)
- 21 SLABOSPILSKY T J & SYNGEN 1939 Zhur Mikrobiol Epidemiol Immun Biol 6 123 (Cited by Seeliger⁴⁴)
- 22 GUDKOWA, E. I & I P SACKAROFF 1946 Zhur Eksp Biol i Med 22 54 (Cited by Seeliger⁴⁴)
- 23 IIRIE J H H 1977 A new disease of wild rodents Publs S African Inst Med Research 3 163-186
- 24 PATOCKA F J BENDA & J STARKY 1953 Listerose infekce novorzenku Ceskoslov hyg epidemiol mikrobiol imunol 2 375-340
- 25 OZCEBE I (Quoted by Seeliger⁴⁴)
- 26 BUSTER H F & L H SCHWARTZ 1939 Studies on *Listeria* infection in sheep J Infectious Diseases 64 135-144
- 27 GRAY M I H J SLAFSETH & F THORP 1951 A four year study of Listeriosis in Mich J Am Vet Med Assoc 118 247-252
- 28 MURRAY F D C R 1955 A characterization of listeriosis in man and other animals Can Med Assoc 72 99-103
- 29 GRAHAM R N D LEVINE & C C MORRILL 1943 Listeriosis in domestic animals Bull 499 Agr Fpt Stat Univ Ill Urbana Ill
- 30 LINE F G & W B CHERRY 1952 Meningitis due to *Listeria monocytogenes* J Am Med Assoc 148 366-369
- 31 LANG K 1955 *Listeria* infektion als mögliche Ursache früh erworbener Cerebralschaden Z Kinderheilk 76 378-339
- 32 JAEGER R F & D M MYERS 1954 *Listeria monocytogenes* a study of two strains isolated from human listeriosis Can J Microbiol 1 12-21
- 33 REED R W W F GAVIN, CROSLEY & P DOBSON 1955 Listeriosis in Man Can Med Assoc J 73 400-407
- 34 JOHNSON W H S A MORTON M H WONG & T E ROY 1955 Septicemia of the newborn due to *Listeria monocytogenes* Can Med Assoc J 73 407-405
- 35 LINE F G & F G APPLETON 1952 *Listeria meningitis* in a premature infant J Pediatr 41 97-99
- 36 WILLIAMS R & J E HORMBURG 1954 *Listeria monocytogenes* in human infections Am J Clin Pathol 24 962-964
- 37 SCHULZE M L G H WARLE & J B WHITE 1953 Meningitis due to *Listeria monocytogenes* a case of disseminated lupus erythematosus Am J Clin Pathol 23 1078-1030
- 38 FINEGOLD S M J G BRADLEY M K CAMPBELL & A J GREENBERG 1954 *Listeria monocytogenes* meningitis A M A Arch Internal Med 93 515-527
- 39 PEFFRAAMP C W A N & J JANSEN 1947 Listeriose bij de kip Tijdschr Diergeneesk 72 389-390
- 40 SHOLKOP W T 1950 *Listeria monocytogenes* isolated from chinchillas J Am Vet Med Assoc 116 447-449

contaminant. Sometimes especially in smears of centrifuged spinal fluid *Listeria* will decolorize or stain unevenly and this may lead to confusion with *Hemophilus influenzae*.

Listeriosis is not a reportable disease and most cases are made known through scientific publications but three human cases were nevertheless reported through the weekly *Morbidity and Mortality Report* of the National Office of Vital Statistics Washington D C during 1956. Eighty three cultures isolated mostly from the spinal fluid or blood of humans have been received by the Communicable Disease Center laboratory for identification since 1950 in 1950 5 1951 4 1952 3 1953 4 1954 6 1955 17, 1956 27 and (through July) 1957 17 cultures. By states Louisiana has sent in 24 Georgia 6 Illinois 5 North Carolina 5 Maryland 4 Florida 4 New York 3 Missouri 3 Pennsylvania 3 Washington D C 3 West Virginia 2, Michigan 2 Texas 2 Virginia 2 Tennessee 2 and Alabama Oklahoma Washington New Mexico Rhode Island California Indiana and Ohio 1 each.

The numbers of animal cases reported to official agencies and compiled by the Communicable Disease Center are as follows 1954 837 1955 668 1956 601 and (January through June) 1957 411. Some of these represent individual herds rather than isolated cases so the number of animals affected exceeds these figures. The totals by month verify the fact that this is a winter and spring disease in livestock. Seventy two and two tenths (72.2) per cent of 2106 reported cases occurred during winter and spring. These reports represent a fraction of the actual occurrence of the disease in livestock and humans for only 10 of the 48 states have sent in information. This is a reflection on the interest in and the facilities for both diagnosis and reporting.

We may conclude therefore that listeriosis is a public health problem in the United States and elsewhere. It is necessary that the physician and medical bacteriologist be constantly aware that *Listeria* is widespread that the premature infant and the aborted fetus may be due to a congenitally acquired agent that does not always manifest itself in an overt illness of the mother that all ages and both sexes are susceptible that animal contact is not necessarily a factor in the epidemiology of listeriosis that the older the individual the greater is the chance of survival from this particular disease except in the ages 60 years or more and that the tetracyclines are efficacious in the treatment if started early.

References

- 1 GILL D A 1937 Ovine bacterial encephalitis and the bacterial genus *Listerella*. Australian Vet J 13 46
- 2 GLEEDEN L M G & A DEVOS 1954 Listeriosis in calves. Vlaamse diergeneesk Tijdschr 23 57
- 3 OLAFSON I 1940 *Listerella* encephalitis of sheep, cattle and goats. Cornell Vet 30 590-593
- 4 BIRLAND C H 1950 A report on the isolation of *Listeria* organisms from a canary and a chicken in the province of Alberta. Can J Comp Med Vet Sci 14 319-324
- 5 BANDARANAYAKE A 1953 Ceylon Vet J 40 (Cited by Seliger¹⁴)
- 6 NYFFELDT A 1929 Étude sur la mononucléose infectieuse. Compt rend soc sci 101 590-597
- 7 FALKSKE C 1941 *Listerella* Infektion bei Hühnern in Deutschland. Berlin u Münch tierärztl Wochschr 441-445

- 8 PORZECANSKI B & C DE BAYGORRIU 1938 A proposito de su constatacion en el Uruguay de una meningitis febril humana Arch soc Biol Montevideo 9 93
- 9 ANTON W 1934 Kritisch experimenteller Beitrag zur Biologie seiner Beziehung zur infektiösen Mononukleose des Menschen Zentr Bakteriell Parasitenk Abt I 131 89-103
- 10 SAVINO F 1940 Observacion de listeriosis humana y accion terapeutica del la e nan Sema a med Buenos Aires 1 336-340
- 11 MURRAY F D G R A WEBB & M B R SWANN 1936 A disease of rats characterized by a large mononuclear leukocytosis caused by a hitherto undescribed bacillus *Bacterium monocytogenes* J Pathol Bacteriol 29 40-439
- 12 ROINE P A RAITIO & L VARKKJAARA 1933 Listeria infection in the guinea pig caused by feeding with Aureomycin Nature 172 67-68
- 13 FORGEOT J C TRUCHE A STROUB & R LARRY 1941 Bull Acad Vet 14 195 (Cited by Seeliger⁴⁰)
- 14 DEBLICKA I & J JANEN 1943 Listeriose bei Tieren J Mikrobiol Serol 9 93-103
- 15 VISWANATHAN G R & V AYYAR 1950 Indian Vet J 26 395 (Cited by Seeliger⁴⁰)
- 16 LEVY M L 1948 *Listeria monocytogenes* in voles Vet J 104 310-312
- 17 TAJIMA M A FUJIMOTO & I HIGUCHI 1951 J Vet Med Japan 66 15 (Cited by Seeliger⁴⁰)
- 18 CRINE O 1943 *Listerella monocytogenes* som årsak til septikemi hos fall Norsk Vet Tidskr 55 9-104
- 19 WRAMBY B O 1944 Om *Listerella monocytogenes* bakterier og om forekomst av listerella infektion hos dyr Skand Vet Tidskr 34 2 1-29
- 20 SCHLEGEL & FEY (Quoted by Seeliger⁴⁰)
- 21 SLABOSPILSKY T P & SVINCEN 1939 Zhur Mikrobiol Epitemi i Immunol 6 123 (Cited by Seeliger⁴⁰)
- 22 GEDKOWA F I & P P SACKAROFF 1946 Zhur Ekptl Biol i Med 22 54 (Cited by Seeliger⁴⁰)
- 23 PIRIE J H H 192 A new disease of wild rodents Publ S Afric Inst Med Research 3 163-186
- 24 PATOCKA F J BENDA & J STARKY 1933 Listerose infekce novorozenku Českoslov hyg etlemlol mikrol Immunol 2 325-340
- 25 OZCEBE I (Quoted by Seeliger⁴⁰)
- 26 BIESTER H F & L H SCHWARTZ 1939 Studies on *Listeria* infection in sheep J Infectious Diseases 64 135-144
- 27 GRAY M L H J SLAFETH & F THORP 1951 A four year study of Listeriosis in Michigan J Am Vet Med Assoc 118 247-252
- 28 MURRAY F D G 1935 A characterization of listeriosis in man and other animals Can Med Assoc 72 99-103
- 29 GRAHAM R N D LEVINE & C C MORRILL 1943 Listeriosis in domestic animals Bull 49 Agr Expt Stat Univ Ill Urbana Ill
- 30 LINE F C & W B CHERRY 1952 Meningitis due to *L. tera monocytogenes* J Am Med Assoc 148 366-369
- 31 LANG K 1935 Listeria Infektion als mögliche Ursache früh erworbener Cerebralschaden Z Kinderheilk. 76 378-339
- 32 JAEGER R F & D M MYERS 1954 *Listeria monocytogenes* a study of two strains isolated from human listeriosis Can J Microbiol 1 12-21
- 33 REED R W W F GAVIN J CROSLY & I DOBSON 1955 Listeriosis in Man Can Med Assoc J 73 400-407
- 34 JOHN ON W H S A MORTON M H WONG & T F ROY 1955 Septicemia of the newborn due to *Listeria monocytogenes* Can Med Assoc J 73 407-409
- 35 LINE F G & F G APPLETON 1952 *Listeria meningitis* in a premature infant J Pediat 41 97-99
- 36 WILLIAMS R & J E HORNBERG 1954 *Listeria monocytogenes* in human infections Am J Clin Pathol 24 967-964
- 37 SCHULZE M L C H WAXLE & J B WHITE 1953 Meningitis due to *Listeria monocytogenes*: a case of disseminated lupus erythematosis Am J Clin Pathol 23 1078-1030
- 38 FINEGOLD S M J G BRADLEY M K CAMPBELL & A J GREENBERG 1954 *Listeria monocytogenes* meningitis A M A Arch Int Med 93 515-57
- 39 PEPEKAMP C W A N & J JANSEN 1947 Listeriosis bij de kip Tijdschr Dierg neek 72 389-340
- 40 SHOLKOP W T 1950 *Listeria monocytogenes* isolated from chinchillas J Am Vet Med Assoc 116 447-449

contaminant. Sometimes especially in smears of centrifuged spinal fluid *Listeria* will decolorize or stain unevenly, and this may lead to confusion with *Hemophilus influenzae*.

Listeriosis is not a reportable disease and most cases are made known through scientific publications but three human cases were nevertheless reported through the weekly *Morbidity and Mortality Report* of the National Office of Vital Statistics Washington D C during 1956. Eighty three cultures isolated mostly from the spinal fluid or blood of humans have been received by the Communicable Disease Center laboratory for identification since 1950. In 1950 5 1951 4 1952 3 1953 4 1954 6 1955 17 1956 21 and (through July) 1957 17 cultures. By states, Louisiana has sent in 24 Georgia 6 Illinois 5 North Carolina 5 Maryland 4 Florida 4 New York 3 Missouri 3 Pennsylvania 3 Washington D C 3 West Virginia 2 Michigan 2 Texas 2 Virginia 2 Tennessee 2 and Alabama Oklahoma Washington New Mexico Rhode Island California Indiana and Ohio 1 each.

The numbers of animal cases reported to official agencies and compiled by the Communicable Disease Center are as follows 1954 837 1955 668 1956 601 and (January through June) 1957 411. Some of these represent individual herds rather than isolated cases so the number of animals affected exceeds these figures. The totals by month verify the fact that this is a winter and spring disease in livestock. Seventy two and two tenths (72.2) per cent of 2106 reported cases occurred during winter and spring. These reports represent a fraction of the actual occurrence of the disease in livestock and humans for only 10 of the 48 states have sent in information. This is a reflection on the interest in and the facilities for both diagnosis and reporting.

We may conclude therefore that listeriosis is a public health problem in the United States and elsewhere. It is necessary that the physician and medical bacteriologist be constantly aware that *Listeria* is widespread that the premature infant and the aborted fetus may be due to a congenitally acquired agent that does not always manifest itself in an overt illness of the mother that all ages and both sexes are susceptible that animal contact is not necessarily a factor in the epidemiology of listeriosis that the older the individual the greater is the chance of survival from this particular disease except in the ages 60 years or more and that the tetracyclines are efficacious in the treatment if started early.

References

1. GILL D A 1937 Ovine bacterial encephalitis and the bacterial genus *Listerella*. *Australian Vet J* 13: 46.
2. GUERDEN L M G & A DEVOS 1954 Listeriosis in calves. *Vlaam diergeneesk Tijdschr* 23: 57.
3. OLAFSON P 1940 *Listerella* encephalitis of sheep cattle and goats. *Cornell Vet* 30: 590-597.
4. BIGLAND C H 1950 A report on the isolation of *Listeria* organisms from a canary and a chicken in the Province of Alberta. *Can J Comp Med Vet Sci* 14: 319-324.
5. BANDARANAYAKE A 1953 *Ceylon Vet J* 40 (Cited by Seeliger).
6. NAFFLDT A 1929 Etologie de la mononucleose infectieuse. *Compt rend soc bio* 101: 550-597.
7. PALLASKE G 1941 *Listerella* Infektion bei Huhnern in Deutschland. *Berlin u Munch tierarztl Wochschr* 441-445.

- 76 GRAY M I S CLINTAMANI & F THORP 1956 Abortion and prior postnatal death of young due to *Listeria monocytogenes*. III. Studies in ruminants. Am J Vet Research 17 510-520
- 77 ØDEGAARD B R CRELLAND & S D HENRIKSEN 1957 A case of *Listeria* infection in man transmitted from sheep. Acta Med Scand 162 231-235
- 78 PATERSON J S 1940 The antigenic structure of the genus *Listerella*. J Pathol Bacteriol 51 427-436
- 79 SEFLIGER H 1955 Serologische Kreuzreaktionen zwischen *Listeria monocytogenes* und Enterokokken. Z Hyg Infektionskrankh 141 15-24
- 80 LINZENMEIER C & H SEELIGER 1954 Die *in vitro* Empfindlichkeit von *Listeria monocytogenes* gegen Sulfonamide und Antibiotica. Zentr Bakteriell Parasitenk Mitt 1 160 543-549
- 81 OEBOLD J W & M T SAWYER 1955 Listeriosis—factors in immunity and pathogenesis. Proc Book Am Vet Med Assoc 189-195

- 41 BARRALES D 1953 Listeriosis in lemmings Can J Public Health 44 180-184
- 42 VALLÉE A 1952 Ann inst Pasteur 83 832 (Cited by Seeliger³)
- 43 HURT A H & D LEVINE & R GRAHAM 1941 Isolation of *Listeria* from the chicken Am J Vet Research 2 279-280
- 44 SEGRÉ D C OLSON C I MARSH & I C BLORE 1956 Attempts to influence listeriosis in chickens using antimetabolites Am J Vet Research 17 299-307
- 45 SEELIGER H 1955 Listeriose Barth Leipzig Germany
- 46 SCHULZE W 1950 Listeriose beim Blauadler Monatsh Veterinarmed 5 200-201
- 47 LATERSON J S 1931 *Listerella* infection in fowls Preliminary note on its occurrence in East Anglia Vet Record 49 1533-1534
- 48 ILLERGEEN K 1947 Svensk Vet Tidskr 56 (Cited by Seeliger⁴)
- 49 COX R F 1945 Listerellosis in dogs Auburn Vet 98-99
- 50 CHAPMAN M P 1947 Listerellosis in a dog a field case North Am Veterinarian 28 532-538
- 51 REH T 1955 (Quoted by Seeliger⁴)
- 52 MUSTEJEV N S & O S IMELJANOWA 1954 Zhur Mikrobiol Epidemiol Immuno biol 36 (Cited by Seeliger⁵)
- 53 MORRIS J & M C NORMAN 1950 The isolation of *Listeria monocytogenes* from ferrets J Bacteriol 59 313-314
- 54 OSBOLD J W G SCHULTZ & E W JAMESON 1957 An epizootiological study of Listeriosis J Am Vet Med Assoc 130 471-475
- 55 GIFFORD A & I JUCHERR 1947 Listeriosis in Connecticut with particular reference to a case in a wild raccoon Cornell Vet 37 39-48
- 56 GRAY M L H J STAFSETH F THORP L B SKOLL & W F RILEY 1948 A new technique for isolating *Listerellae* from the bovine brain J Bacteriol 56 471-476
- 57 IRIE J H 1940 *Listeria* change of name for a genus of bacteria Nature 146 264
- 58 KAPLAN M M 1945 Listeriosis New Engl J Med 232 755-759
- 59 GIRARD K F & E G D MURRAY 1951 *Listeria monocytogenes* as the cause of disease in man and animals and its relation to infectious mononucleosis from an etiological and immunological aspect Am J Med Sci 221 343-357
- 60 FELSENFELD O 1951 Diseases of poultry transmissible to man Iowa State Coll Vet 13 89-92
- 61 BLUM C G 1936 Clinical and pathological features of infection caused by a new pathogen of genus *Listeria* Am J Pathol 12 341-348
- 62 PÖTEL J 1952 Weitere Untersuchungen über den Erreger der Granulomatosis infantiseptica Zentr Bakteriell Parasitenk Abt I 169 86-87
- 63 PÖTEL J 1953 Ätiologie der Granulomatosis infantiseptica Wiss Z Martin Luther Univ 3 341-364
- 64 PÖTEL J 1955 6 Jahre Listeriose Forschung und Diagnose mit besonderer Berücksichtigung der Neugeborenen Listeriose Arch Hyg u Bakteriol 139 245-264
- 65 WINKLER C H M F JAMES & R P CARTER 1952 *Listeria meningitis* a case report Southern Med J 45 1181-1183
- 66 HOEPRICH I O & H M CHERNOFF 1955 Subacute bacterial endocarditis due to *Listeria monocytogenes* Am J Med 19 488-494
- 67 ZING A G C DEMELLO & R I BURHART 1951 Listeriosis field and laboratory studies and aureomycin activity Am J Vet Research 12 194-197
- 68 NORMAN M C D LONGFELLOW & J K LEVIN 1951 Effects of Aureomycin and chloramphenicol on two recently isolated strains of *Listeria monocytogenes* Proc Soc Exptl Biol Med 76 435
- 69 PORTER J R & W H HALF 1939 Effect of Sulfanilamide and sulfapyridine on experimental infections with *Listerella* and *Erysipel thrix* in mice Proc Soc Exptl Biol Med 42 47-50
- 70 OLSON C R H COOK & V BAUDONA 1951 An attempt to immunize sheep during an outbreak of listeriosis Am J Vet Research 12 306-312
- 71 JENSEN K & D R MACKEN 1947 Listerellosis in sheep and cattle J Am Vet Med Assoc 114 420-424
- 72 MORBIDITY AND MORTALITY WEEKLY REPORT 1950 1957 Natl Off Vital Stat 6 13 36 39 6 21
- 73 URBACH H & G SCHABINSKI 1955 Zur Listeriose des Menschen Z Hyg Infektionskrankh 141 239-248
- 74 WENCKEBACK C K 1953 Züchtung von *Listeria monocytogenes* aus der Harnröhre des Mannes Proc 6th Intern Congr Microbiol 2 406-407
- 75 BINDER M A C DIFHL J WEISS & H RAY 1953 *Listeria meningitis* Ann Internal Med 38 1315-1319

- 6 CRAY M L S CLINTAMANI & F THORP 1956 Abortion and prior postnatal death of young due to *Listeria monocytogenes* III Studies in ruminants Am J Vet Research 17 510-520
- 77 ØDEGAARD B R CRILLAND & S D HENRIKSEN 1952 A case of *Listeria* infection in man transmitted from sheep Acta Med Scand 142 231-235
- 8 PATERSON J S 1940 The antigenic structure of the genus *Listerella* J Pathol Bacteriol 61 421-436
- 79 SEFLIGER H 1955 Serologische Kreuzreaktionen zwischen *Listeria monocytogenes* und Enterokokken Z Hyg Infektionskrankh 141 15-24
- 80 LINZENMEIER C & H SEELIGER 1954 Die in vitro Empfindlichkeit von *Listeria monocytogenes* gegen Sulfonamide und Antibiotica Zentr Bakteriell Parasitenk Abt I 160 543-559
- 81 OSBOLD J W & M T SAWYER 1955 Listeriosis—factors in immunity and pathogenesis Proc Book Am Vet Med Assoc 169-195

THE PUBLIC HEALTH IMPORTANCE OF ANIMAL TUBERCULOSIS

By R. J. Anderson

Communicable Disease Center Public Health Service Atlanta Ga

Although tuberculosis is common to a wide range of animals mammals birds and even fish and reptiles only cattle are important hosts where human health is at risk. Tuberculosis in swine is a problem to the farmer to be sure for his hogs may be condemned at slaughter as unfit for human consumption. Pet animals sometimes have tuberculosis but the likelihood of their transmitting it to human beings is rather remote. Since cow's milk is a staple of the diet in the United States however the bovine type of tuberculosis which humans can contract by ingesting milk from tuberculous cows has long been a concern of public health in this country.

The control of tuberculosis in cattle has had a remarkable history in the United States. Controversy surrounded the advocacy of tuberculin testing and slaughter of reactors forty years ago. There was both skepticism about the communicability of bovine tuberculosis to animals or to man and a wide belief that the destruction of tuberculin reacting cattle would be ruinous to the agricultural economy. In spite of these obstacles the proponents of tuberculin testing and slaughter of reactors were able to establish a national campaign in 1917 supported by a congressional appropriation. The campaign rapidly grew to full force and in the ensuing forty years hundreds of millions of cattle have been tested and the reactors destroyed.

The success of this work in reducing the percentage of tuberculin reactors among cattle tested is a matter of record. From a high of almost 5 per cent in 1918 reacting cattle reached a low in 1952 of 0.11 per cent. Slightly higher rates in recent years (0.15 per cent in 1956) are attributed to more aggressive attempts to find sources of tuberculosis by tracing reactors to their herds of origin and testing those herds. Even 15 reactors per 1000 cattle tested constitutes a very low rate compared with those found in many other areas of the world. In some countries the control of bovine tuberculosis is only beginning.

As the practice of pasteurizing milk has become more widespread it has given added protection to the population against exposure to bovine tuberculosis. Today about 95 per cent of the milk consumed by the urban population of the United States is pasteurized. Nine states and 64 cities require pasteurization of the milk that is sold in their jurisdiction.

The effect of these programs in lowering the prevalence of bovine tuberculosis in humans is difficult to measure. Statements that the bovine type of tuberculosis now plays an insignificant role in this country are common in current medical literature but no data are available to substantiate them because presence of the causative strain of bacillus is not required in tuberculosis reporting in the United States. In the 1920s and 1930s a number of authors studied the relative frequency of bovine infection and reported widely ranging percentages in the various groups of patients. In discussing B. Mollers summary of the literature up to 1927 from which he found that 11.1 per cent of

2277 cases studied by various authors were due to bovine infection. Rich¹ points out that such a composite must be considered in light of the fact that the incidence of bovine infection varies widely from area to area that it has been greatly reduced in some places since 1927 and that studies usually have been made where there was some reason to believe that the incidence of bovine infection was high a set of conditions that would tend to inflate the figures on bovine infection. Since the United States has always had a lower rate of tuberculosis in cattle than most of the rest of the world application of Mollers 11.1 per cent or of Price's 10 per cent (based on a 1939 study of 18 000 cases reported in the literature from all over the world) would be inappropriate here.

Since bovine infection affects children more often than adults and usually lodges in parts of the body other than the lungs deaths from extrapulmonary tuberculosis among children are often used as a rough index of the incidence of bovine infection. Price² points out that the reduction in mortality from extrapulmonary forms of tuberculosis parallels the eradication of tuberculosis in cattle. Such a parallel however is less convincing when one considers that the reduction in mortality from extrapulmonary forms has also paralleled the reduction in mortality from pulmonary tuberculosis. In the absence of exact data the question of whether we have less tuberculosis of the spine bones and joints and less scrofula in the United States because of the control of tuberculosis in cattle or because of factors that have brought about lowered pulmonary tuberculosis death rates remains a matter of speculation. Better social and economic conditions and control measures such as more aggressive case finding the isolation of patients in sanatoria and better medical care all would certainly be expected to bring about a decline in the incidence of the nonpulmonary as well as the pulmonary forms of the disease.

I do not wish to give the impression that I am minimizing the importance of the excellent work that has been done in the United States to eradicate tuberculosis in cattle. I do believe however that we are now at a point where assessment of its public health importance is in order. Consideration of the situation raises these challenging questions for both medical and veterinary epidemiologists. To what extent are bovine type tubercle bacilli infecting humans and causing disease in them? What is the source and the mechanism of transmission? Is the program of eradication of tuberculosis in cattle to be justified in terms of its effect on human tuberculosis or is its importance related chiefly to the agricultural economy? Indeed is a program that is sufficient for the agricultural economy adequate for the control of human tuberculosis? What indices are valid for decisions about the relaxation cessation or intensification of present programs?

When a disease is fairly widespread effective control programs can be based upon empirical judgments of where disease exists and how much of it there is. With success in lowering prevalence comes the necessity for focusing efforts directly on the remaining problem in order to use control facilities economically.

Our first question is perhaps the most important. To what extent are bovine type tubercle bacilli infecting humans and causing disease in them? The culturing and strain typing of bacilli are now routine practice in many

THE PUBLIC HEALTH IMPORTANCE OF ANIMAL TUBERCULOSIS

By K. J. Anderson

Communicable Disease Center Public Health Service Atlanta Ga

Although tuberculosis is common to a wide range of animals mammals birds and even fish and reptiles only cattle are important hosts where human health is at risk. Tuberculosis in swine is a problem to the farmer to be sure for his hogs may be condemned at slaughter as unfit for human consumption. Pet animals sometimes have tuberculosis but the likelihood of their transmitting it to human beings is rather remote. Since cow's milk is a staple of the diet in the United States however the bovine type of tuberculosis which humans can contract by ingesting milk from tuberculous cows, has long been a concern of public health in this country.

The control of tuberculosis in cattle has had a remarkable history in the United States. Controversy surrounded the advocacy of tuberculin testing and slaughter of reactors forty years ago. There was both skepticism about the communicability of bovine tuberculosis to animals or to man and a wide belief that the destruction of tuberculin reacting cattle would be ruinous to the agricultural economy. In spite of these obstacles the proponents of tuberculin testing and slaughter of reactors were able to establish a national campaign in 1917 supported by a congressional appropriation. The campaign rapidly grew to full force and in the ensuing forty years hundreds of millions of cattle have been tested and the reactors destroyed.

The success of this work in reducing the percentage of tuberculin reactors among cattle tested is a matter of record. From a high of almost 5 per cent in 1918, reacting cattle reached a low in 1952 of 0.11 per cent. Slightly higher rates in recent years (0.15 per cent in 1956) are attributed to more aggressive attempts to find sources of tuberculosis by tracing reactors to their herds of origin and testing those herds. Even 1.5 reactors per 1000 cattle tested constitutes a very low rate compared with those found in many other areas of the world. In some countries the control of bovine tuberculosis is only beginning.

As the practice of pasteurizing milk has become more widespread it has given added protection to the population against exposure to bovine tuberculosis. Today about 95 per cent of the milk consumed by the urban population of the United States is pasteurized. Nine states and 64 cities require pasteurization of the milk that is sold in their jurisdiction.

The effect of these programs in lowering the prevalence of bovine tuberculosis in humans is difficult to measure. Statements that the bovine type of tuberculosis now plays an insignificant role in this country are common in current medical literature but no data are available to substantiate them because presence of the causative strain of bacillus is not required in tuberculosis reporting in the United States. In the 1920s and 1930s a number of authors studied the relative frequency of bovine infection and reported widely ranging percentages in the various groups of patients. In discussing B. Mollers summary of the literature up to 1921 from which he found that 11.1 per cent of

not announce itself in sudden dramatic onset but often develops slowly and can go unrecognized for long periods. Its public health importance must be evaluated by some means more definite than opinion. The relatively much greater prevalence of tuberculosis of human origin overshadows and obscures the absolute indices of the extent of bovine tuberculosis infection. This denies to us the use of the mass of data on tuberculosis in humans that is readily available in assessing the present-day importance and nature of bovine type infection. We are therefore forced into specific inquiry if such is our objective.

In the history of any disease-control program particularly those directed against widespread infections the later phases during which eradication is almost in sight often present unusual difficulties. In the beginning when a method of control is first available enthusiasm is high and this enthusiasm will continue as long as the method produces demonstrable results. As the problem diminishes control measures may either be neglected or be carried on routinely without critical examination of their effectiveness. Too often re-evaluation and remeasurement of the problem and of the methods in use against it remain long overdue. I suggest that the time is ripe for such a re-evaluation of the public health importance of animal tuberculosis.

References

1. RICH, A. R. 1951. The Pathogenesis of Tuberculosis. 56-59. Thomas, Springfield, Ill.
2. PRICE, R. M. 1939. The bovine tubercle bacillus in human tuberculosis. *Am J Med Sci* 197: 413.

tuberculosis hospitals and clinics, especially in connection with studies of the atypical acid fast bacteria (the so called yellow bacilli and Battey type organisms) that are being recovered in many cases diagnosed as tuberculosis. However, since both tuberculosis hospitals and public health clinics are concerned chiefly with pulmonary disease even when routine typing is done it probably can give little information about the extent of bovine infection. Canvassing of general hospitals, children's hospitals, and physicians such as orthopedists or dermatologists who may encounter tuberculosis of the bones or skin would be a necessary additional productive line of inquiry. In every case where bovine bacilli are isolated, thorough investigation of the source of infection might uncover otherwise unsuspected remaining reservoirs of the disease or throw light on the mechanism of transmission and on the significance of cattle infection for humans.

From a practical standpoint, this type of investigation would be most appropriate in communities where tuberculin testing of cattle is sporadic. Objective review of the figures on cattle testing by states suggests that there is considerable variation in coverage both geographically and in point of time. Furthermore, in many rural areas where dairy farming is not the major emphasis of agriculture, families customarily keep one or several cows to supply their own needs. How often are these family cows tested? Is their milk customarily pasteurized before the family drinks it? It seems to me that the possibility that bovine tuberculosis is still harbored in such areas would bear investigation not only through examination of hospital records and inquiry about physicians' experience but also by use of the tuberculin test on the children in such areas. By and large, human tuberculin sensitivity has been found to be low in the rural districts of the United States, as we should expect from the fact that fewer cases of disease are found there. However, some children in almost every rural area do react to tuberculin, and it would seem both practical and profitable to investigate carefully their opportunities for exposure to bovine infection.

In any discussion of bovine tuberculosis the suggestion always arises that this form of the disease occurs more often than we think, and that it goes unrecognized. The occasional instance in which bovine tuberculous infection of the gastrointestinal or genitourinary organs is diagnosed raises in the epidemiologist's mind the question: are other cases occurring that are never diagnosed accurately? Are we so unsuspicious of the existence of bovine infection that we fail to notice it? Does it produce subclinical illness that rarely receives medical attention? This too is a matter not for speculation or for argument but for investigation. The tuberculin test and history of possible exposure to bovine tuberculosis could perhaps provide leads in the differential diagnosis of obscure abdominal complaints.

The public health importance of animal tuberculosis it seems to me cannot be measured with the information we now have. We think that bovine tuberculosis in humans is rare. The cases that have come to the attention of the Communicable Disease Center since 1950, cases confirmed by laboratory study, can be counted on the fingers of one hand. However, this communicable disease which exists in animal hosts closely associated with man does

not announce itself in sudden dramatic onset but often develops slowly and can go unrecognized for long periods. Its public health importance must be evaluated by some means more definite than opinion. The relatively much greater prevalence of tuberculosis of human origin overshadows and obscures the absolute indices of the extent of bovine tuberculosis infection. This denies to us the use of the mass of data on tuberculosis in humans that is readily available in assessing the present day importance and nature of bovine type infection. We are therefore forced into specific inquiry if such is our objective.

In the history of any disease-control program particularly those directed against widespread infections the later phases during which eradication is almost in sight often present unusual difficulties. In the beginning when a method of control is first available enthusiasm is high and this enthusiasm will continue as long as the method produces demonstrable results. As the problem diminishes control measures may either be neglected or be carried on routinely without critical examination of their effectiveness. Too often re evaluation and remeasurement of the problem and of the methods in use against it remain long overdue. I suggest that the time is ripe for such a re evaluation of the public health importance of animal tuberculosis.

References

- 1 RICH A R 1951 The Pathogenesis of Tuberculosis 56-59 Thomas Springfield Ill
- 2 PRICE R M 1939 The bovine tubercle bacillus in human tuberculosis Am J Med Sci 197 413

tuberculosis hospitals and clinics, especially in connection with studies of the atypical acid fast bacteria (the so-called yellow bacilli and Battey type organisms) that are being recovered in many cases diagnosed as tuberculosis. However, since both tuberculosis hospitals and public health clinics are concerned chiefly with pulmonary disease even when routine typing is done it probably can give little information about the extent of bovine infection. Canvassing of general hospitals, children's hospitals, and physicians such as orthopedists or dermatologists who may encounter tuberculosis of the bones or skin would be a necessary additional productive line of inquiry. In every case where bovine bacilli are isolated, thorough investigation of the source of infection might uncover otherwise unsuspected remaining reservoirs of the disease or throw light on the mechanism of transmission and on the significance of cattle infection for humans.

From a practical standpoint, this type of investigation would be most appropriate in communities where tuberculin testing of cattle is sporadic. Objective review of the figures on cattle testing by states suggests that there is considerable variation in coverage, both geographically and in point of time. Furthermore, in many rural areas where dairy farming is not the major emphasis of agriculture, families customarily keep one or several cows to supply their own needs. How often are these family cows tested? Is their milk customarily pasteurized before the family drinks it? It seems to me that the possibility that bovine tuberculosis is still harbored in such areas would bear investigation, not only through examination of hospital records and inquiry about physicians' experience, but also by use of the tuberculin test on the children in such areas. By and large, human tuberculin sensitivity has been found to be low in the rural districts of the United States, as we should expect from the fact that fewer cases of disease are found there. However, some children in almost every rural area do react to tuberculin, and it would seem both practical and profitable to investigate carefully their opportunities for exposure to bovine infection.

In any discussion of bovine tuberculosis, the suggestion always arises that this form of the disease occurs more often than we think, and that it goes unrecognized. The occasional instance in which bovine tuberculous infection of the gastrointestinal or genitourinary organs is diagnosed raises in the epidemiologist's mind the question: are other cases occurring that are never diagnosed accurately? Are we so unsuspicious of the existence of bovine infection that we fail to notice it? Does it produce subclinical illness that rarely receives medical attention? This, too, is a matter not for speculation or for argument, but for investigation. The tuberculin test and history of possible exposure to bovine tuberculosis could perhaps provide leads in the differential diagnosis of obscure abdominal complaints.

The public health importance of animal tuberculosis, it seems to me, cannot be measured with the information we now have. We think that bovine tuberculosis in humans is rare. The cases that have come to the attention of the Communicable Disease Center since 1950, cases confirmed by laboratory study, can be counted on the fingers of one hand. However, this communicable disease, which exists in animal hosts closely associated with man, does

morbidity*. In the rural population of the United States it is estimated that approximately 10 to 80 per cent of fungus infections involving the exposed parts of the body are caused by zoophilic dermatophytes with animals serving as vectors of these diseases. In the urban population it is estimated that 10 to 30 per cent of such infections are produced by zoophilic dermatophytes with animals playing the role of vectors of infection.

It is evident that animal ringworm presents definite public health implications. Recognizing the importance of this problem and the many voids in our over all knowledge of this entity the Mycology Laboratory, Communicable Disease Center, Chamblee, Ga. in cooperation with the Veterinary Public Health Section of the Center initiated a long range animal ringworm study in 1953. An integral part of this study is a survey¹⁰ to determine among other things the fungi involved and the prevalence of these infections in our lower animal population. This survey has been conducted with the cooperation of veterinarians from all parts of the United States who submit clinical materials from animals with skin lesions for mycologic study. In addition a companion survey was carried out on wild animals trapped in southwest Georgia by personnel of the Newton Field Station of the Communicable Disease Center. It is noteworthy that the wild animals did not show any manifest cutaneous abnormalities at the time the specimens were collected.

The purpose of this report is to present and to discuss our findings in these studies for the 18 month period from January 1956 to June 1957.

MATERIALS AND METHODS

The clinical materials received for study consisted of hairs and/or skin scrapings. Upon arrival at the Mycology Laboratory each specimen was examined under a Wood's light for evidence of fluorescence. This procedure was followed by a direct microscopic examination to detect the presence of dermatophytes. The microscopy was carried out in 10 per cent potassium hydroxide mounts. Each specimen was then cultured on a special selective isolation medium¹¹ containing the antibiotics cycloheximide*, penicillin and streptomycin and incubated at room temperature (25°C). For the isolation of *Trichophyton verrucosum* the medium was enriched with thiamine and cultures held at 31°C¹. The cycloheximide was omitted from the medium for the isolation of *Trichosporon cutaneum* since that organism was found to be inhibited by this antibiotic. Each specimen was inoculated into a minimum of three tubes of media and all cultures were held for one month before reporting any as negative.

RESULTS

Domestic Animal Ringworm Study

As noted on TABLE 1 during the period covered by this report specimens from 2361 domestic animals with skin lesions were examined. These materials were submitted by veterinarians from 35 states in all sections of the country.

Actidione manufactured by The Upjohn Co. Kalamazoo Mich. Trade names are used for identification purposes only. Mention of them does not constitute an endorsement by the Public Health Service.

RECENT DEVELOPMENTS IN ANIMAL RINGWORM AND THEIR PUBLIC HEALTH IMPLICATIONS

By William Kaplan Lucille K. Georg and Libero Ajello
Communicable Disease Center Public Health Service Atlanta Ga

INTRODUCTION

The ability to break down keratin is an important characteristic of the dermatophytes. *In vivo* these fungi are parasites of the keratinized tissues of the body such as the epidermis the hair, and the nails and they produce alterations of the structures invaded. These alterations together with attendant immunological reactions are clinically classified as ringworm.

Both man and the lower animals are susceptible to infection by ringworm fungi. However the individual dermatophyte species show varying degrees of host specificity. Some are parasites of man and these rarely if ever infect lower animals, others are primarily animal pathogens that are found to produce human disease on occasion. Thus on the basis of host preference or host adaptation, the ringworm fungi have been divided into two groups. In one category have been placed the human pathogens or anthropophilic organisms and in the other the animal pathogens or zoophilic dermatophytes. While it is true that some of the human and animal pathogens have been recovered from the contaminated environment of infected hosts^{1, 2} such apparent saprophytism appears to be secondary to their parasitism of man or animals. Until recently nearly all of the known dermatophytes have been placed in one or the other of these two groups.

Recent findings however point to the necessity of establishing still another category of dermatophytes one that would draw its members from among the keratinophilic soil inhabiting fungi that have the ability under certain conditions to invade the keratinized tissues of human and animal hosts and there to produce disease. Members of this group are essentially saprophytes and probably serve along with other keratinophilic organisms in the microbiological breakdown in soil of keratin into simple elements. To date only two species *Microsporum gypsum*^{3, 4} and *Keratinomyces ajelloi*^{5, 6} have been demonstrated to belong in this third category. Thus from an epidemiological standpoint it is logical at this time to consider the existence of these three groups of ringworm fungi the anthropophilic or human type organisms the zoophilic or animal type organisms and the geophilic or soil organisms.

Members of the three groups of dermatophytes are involved in human disease and each type presents distinct public health problems. As might be anticipated the vast majority of ringworm infections in man are caused by anthropophilic organisms. Such conditions as athlete's foot and ringworm of the groin and nails are common conditions affecting large segments of our population. They are caused by anthropophilic dermatophytes and animals have not been shown to play a role in the epidemiology of these diseases. However if one excludes these common infections and considers only ringworm of the exposed parts of the body such as the scalp face neck and extremities then the zoophilic organisms appear as important contributors to human

morbidity.⁹ In the rural population of the United States it is estimated that approximately 70 to 80 per cent of fungus infections involving the exposed parts of the body are caused by zoophilic dermatophytes with animals serving as vectors of these diseases. In the urban population it is estimated that 10 to 30 per cent of such infections are produced by zoophilic dermatophytes with animals playing the role of vectors of infection.

It is evident that animal ringworm presents definite public health implications. Recognizing the importance of this problem and the many voids in our over-all knowledge of this entity, the Mycology Laboratory, Communicable Disease Center, Chamblee, Ga., in cooperation with the Veterinary Public Health Section of the Center, initiated a long range animal ringworm study in 1953. An integral part of this study is a survey¹⁰ to determine among other things the fungi involved and the prevalence of these infections in our lower animal population. This survey has been conducted with the cooperation of veterinarians from all parts of the United States who submit clinical materials from animals with skin lesions for mycologic study. In addition a companion survey was carried out on wild animals trapped in southwest Georgia by personnel of the Newton Field Station of the Communicable Disease Center. It is noteworthy that the wild animals did not show any manifest cutaneous abnormalities at the time the specimens were collected.

The purpose of this report is to present and to discuss our findings in these studies for the 18 month period from January, 1956 to June, 1957.

MATERIALS AND METHODS

The clinical materials received for study consisted of hairs and/or skin scrapings. Upon arrival at the Mycology Laboratory each specimen was examined under a Wood's light for evidence of fluorescence. This procedure was followed by a direct microscopic examination to detect the presence of dermatophytes. The microscopy was carried out in 10 per cent potassium hydroxide mounts. Each specimen was then cultured on a special selective isolation medium¹¹ containing the antibiotics cycloheximide,* penicillin and streptomycin and incubated at room temperature (25° C). For the isolation of *Trichophyton verrucosum* the medium was enriched with thiamine and cultures held at 37° C.¹² The cycloheximide was omitted from the medium for the isolation of *Trichosporon cutaneum* since that organism was found to be inhibited by this antibiotic. Each specimen was inoculated into a minimum of three tubes of media and all cultures were held for one month before reporting any as negative.

RESULTS

Domestic Animal Ringworm Study

As noted on TABLE 1 during the period covered by this report specimens from 2361 domestic animals with skin lesions were examined. These materials were submitted by veterinarians from 35 states in all sections of the country.

* *Acto-d-o* manufactured by The Upjohn Co., Kalamazoo, Mich. Trade names are used for identification purpose only. Mention of them does not constitute an endorsement by the Public Health Service.

TABLE 1
DOMESTIC ANIMAL RINGWORM SURVEY JANUARY 1936 TO JUNE 1937
DERMATOPHYTES ISOLATED BY HOST SPECIES

Host animal species	Number specimens examined	Organisms isolated										Total positive	Percentage
		<i>M. canis</i>	<i>M. gypseum</i>	<i>M. audouinii</i>	<i>M. distortum</i>	<i>T. tonsurans</i>	<i>T. equinum</i>	<i>T. verrucosum</i>	<i>T. rubrum</i>	<i>Keratinomyces</i>	<i>Trichosporon</i>		
Dog	1659	252	68	1	1	41			1	3		368	22.1
Cat	524	193	4			3						200	38.1
Cattle	58							29				29	50.0
Horse	56	2	12				4					18	32.1
Monkey	21	10			4						1	15	71.4
Chinchilla	12	1										3	25.0
Guinea pig	9					6						6	66.0
Rabbit	8											0	0.0
Donkey	2						1	1				2	100.0
Sheep	5											0	0.0
Malabar squirrel	1									1		1	100.0
Chicken	1											0	0.0
Parakeet	4											0	0.0
Chimpanzee	1											0	0.0
Total	2301	358	84	1	5	52	5	30	1	1 + 3*	1	642	27.2

* *A. ajelloi* strains not shown to be pathogenic

† *T. cutaneum*

Pathogenic fungi were recovered from animals residing in 29 different states. Of the total number of specimens studied 642 (27.2 per cent) were found to be positive by culture.

As shown on TABLE 1, 9 dermatophyte species and a *Trichosporon* were isolated and identified from the clinical materials submitted. These were *Microsporum canis*, *M. gypseum*, *M. audouinii*, *M. distortum*, *Trichophyton mentagrophytes*, *T. equinum*, *T. verrucosum*, *T. rubrum*, *Keratinomyces ajelloi*, and *Trichosporon cutaneum*. As might be expected, the majority of these dermatophytes are specialized zoophilic parasites of lower animals. However, 2 of the species, *M. gypseum* and *A. ajelloi*, are soil organisms and 2 others, *M. audouinii* and *T. rubrum*, are essentially human pathogens. As a background to the presentation of our findings relative to the individual host animal species, it would be well to discuss briefly each of the organisms encountered.

Microsporum canis (*M. felinum*, *M. equinum*, *M. lanosum*). This fungus has been shown by a number of investigators both in the United States and elsewhere to be the common cause of ringworm in cats and dogs. Our findings further confirm this fact and, in addition, reveal that *M. canis* is also an important cause of ringworm in monkeys. Although *M. canis* is primarily a zoophilic dermatophyte, human infections by this fungus are not uncommon. The organism is readily transmitted to man by animals, particularly by cats and dogs, and is responsible for the majority of the animal type ringworm infections in the urban human population of the United States¹². In rural areas this agent presents less of a public health problem. Person-to-person

transmission of this fungus can occur however the organism appears to lose its infectivity after a few such human passages. Thus large scale epidemics in man are not caused by *M canis*. Rather isolated cases or group infections form the pattern observed the total number of human cases being a reflection of the morbidity in animals. The Wood's light is a useful tool in the detection of this infection in both man and animals. However this diagnostic aid does have its limitations as evidenced by our findings that only 30 per cent of the *M canis* infected hairs fluoresced.

Microsporum gypsum (*M fulvum* Ichoriou *gypsum*) *M gypsum* has been shown to be a common soil fungus. Ajello⁴ reported the recovery of this agent from 31 (31.9 per cent) of 116 soil samples collected from various sites in Tennessee and Georgia. Subsequent analyses of foreign and domestic soils by several workers⁵⁻⁸⁻¹⁴ have resulted in equally frequent isolations. Although *M gypsum* is infectious for man it does not appear to be an important human pathogen. In several surveys¹⁵⁻¹⁷ of human ringworm this fungus has been found to be responsible for only 0.7 (2.1 per cent) of the cases studied. Until recently animal infections have been considered to be equally uncommon. A review of the world literature by Ajello⁴ in 1953 revealed that up to that time only 61 authenticated lower animal infections by this organism had been recorded. However the results of our studies would indicate that *M gypsum* is a not uncommon cause of ringworm in animals. As noted on TABLE I *M gypsum* was isolated from 84 animals with clinical ringworm during the period covered by this report. This fungus is often referred to as a zoophilic dermatophyte in the belief that animals serve as reservoirs of infection for man. It is possible that occasional animal to-human transmission does occur. However in those instances where the infection has been diagnosed concurrently in man and his animal contacts the possibility that soil served as a common source of infection for the hosts involved was not considered or investigated. The wide spread distribution of this fungus in soil together with its demonstrated ability to exist in nature as a free living agent¹⁸ would indicate that *M gypsum* is essentially a saprophyte that can under certain conditions assume a parasitic existence. Consequently this mold would fall into the category of geophilic dermatophytes and should not be regarded as an animal type fungus. Although some workers have reported fluorescence of *M gypsum* infected human hairs under the Wood's light we have never observed this phenomenon in our animal cases. Thus the Wood's lamp does not appear to be an appropriate tool for the detection of this infection in animals.

Microsporum audouinii This fungus is considered to be a human pathogen with animals playing no role in the epidemiology of the infection. *M audouinii* is the agent most commonly implicated in epidemic ringworm of the scalp in children in the United States. In recent years outbreaks involving thousands of school children have been reported from all sections of the country. Although this agent is essentially a human pathogen there are recorded in the literature¹⁹⁻²³ five authenticated infections in lower animals. In three of these cases dogs in one a monkey and in one a guinea pig were involved. These few cases of course do not indicate that animals serve as unsuspected reservoirs of this infection yet they are significant to the extent that lower animals

have been proved to be susceptible to this agent. It is possible although perhaps unlikely that future studies may change our concept of the part played by animals in the natural history of this agent. As noted on TABLE 1 we recovered *M. audouinii* from a dog with clinical ringworm. The animal involved was a 10-week old boxer that had been acquired by a family that had two children under dermatologic care for tinea capitis. Infected hairs collected from this dog were observed to fluoresce under the Wood's light. Subsequent to the period covered by this report we again isolated *M. audouinii* from a dog with clinical ringworm.⁴ This animal was a 6 month old terrier that had been recently purchased by a family that included a young boy under treatment for ringworm of the scalp. In this case *M. audouinii* was also recovered from the human. It would appear that in these two episodes man transmitted the infection to the lower animals.

Microsporum distortum. This new species of *Microsporum* was described in 1954 by Di Menna and Marples⁵ as a cause of human disease in New Zealand. The epidemiological features of the outbreaks in that country suggested to these workers that animals were the source of the human infections. However these investigators did not recover this fungus from lower animals. As shown on TABLE 1 we isolated *M. distortum* in four instances from monkeys and once from a dog that had been in contact with one of the infected simians.²⁸ All five animals had clinical ringworm. These recoveries demonstrated that *M. distortum* does indeed infect lower animals in nature and that it is present in the New World. The monkeys involved had been recently imported from Central America. While conclusive proof was not obtained to show that *M. distortum* had been brought into the United States by the monkeys the evidence is sufficiently suggestive in nature to warrant further investigations into this possible avenue of importation. The public health implications of this infection are evidenced by the fact that six humans that had been in contact with three of the simians were reported to have developed ringworm. The demonstrated ability of this agent to infect a variety of host species points to the possibility of its establishment in the lower animal population of this country. Hairs invaded by *M. distortum* fluoresce under the Wood's lamp.

Trichophyton mentagrophytes var. *granular* (*T. gypsum*, *T. asteroides*, *T. granulosum*). This variety must be differentiated from the downy type of *T. mentagrophytes* that is the human pathogen commonly found to cause athlete's foot.²⁷ The granular variety is a zoophilic dermatophyte that is transmitted by animals to man. Human infections with this agent are encountered principally in rural areas. In one study,⁹ *T. mentagrophytes* var. *granular* was shown to be responsible for approximately one half of the cases of suppurative ringworm in humans in rural areas of Michigan. This fungus is pathogenic for a diversity of animal species both domestic and wild including rodents and larger mammals.⁹

Trichophyton equinum. As the species name would imply this fungus is primarily a pathogen of horses. Epizootics of equine ringworm caused by this fungus have been described from the United States² and other parts of the world.²⁹⁻³² In some of the outbreaks infections have been reported among human contacts. However such cases have been few in number a fact that

indicates minimal pathogenicity for man. In this country *T. equinum* has not been generally recognized as a distinct species. In the manual of Conant *et al*³³ it is listed as a synonym of *T. mentagrophytes*. However nutritional and morphologic studies³⁴ recently carried out in our laboratory have led us to recognize that *T. equinum* is a valid species and is not a variant of *T. mentagrophytes*.

Trichophyton terreus (*T. sativum*, *T. album*, *T. discoides*, *T. ochraceum*) This dermatophyte is the principal cause of cattle ringworm in the United States and other parts of the world.^{35, 36} Occasional infections occur in other animal species.⁹ Man is susceptible to infection by this fungus and *T. terreus* appears to be an important cause of suppurative ringworm in humans residing in rural areas. In the study carried out in Michigan³⁵ approximately one half of the cases of suppurative ringworm in rural residents were shown to be caused by this fungus.

Trichophyton rubrum (*T. purpureum*) This dermatophyte is an important cause of ringworm of the feet and nails of man in the United States and in other parts of the world. In recent years there have been indications that human infections with this agent have been increasing.³⁷ The reasons for this trend are not too well understood and have been the subject of considerable speculation. There is no evidence, however, that animals are involved in the epidemiology of this disease. Although *T. rubrum* is generally regarded as an obligate anthropophilic dermatophyte, natural infections in animals can occur. Chakraborty, Ghosh, and Blank³⁸ reported the first recovery of this agent from lower animals. As noted on TABLE 1, during the period covered by this report we isolated *T. rubrum* from a dog with clinical ringworm. The animal involved was a one year old boxer the owner of which had a mild case of tinea pedis of several years duration. The owner had the habit of rubbing the dog with his bare feet. *T. rubrum* was recovered from the owner's feet. Thus it would appear that in this case man transmitted the infection to a lower animal.³⁹

Keratinomyces ajelloi This fungus was first described in 1952 by Van Breuseghem,⁶ who recovered the organism from Belgian soil. Subsequent studies^{4, 6} carried out in the United States and elsewhere have shown this agent to be a common soil inhabiting fungus. Although man and lower animals are constantly exposed to this agent in nature, there had been no reports of infection with this species until 1956. During that year we recovered *K. ajelloi* from a case of ringworm in a Malabar squirrel.⁴⁰ In this instance we were able to demonstrate tissue invasion and were successful in reproducing the disease in guinea pigs with the strain isolated. As noted on TABLE 1, this fungus was also recovered from three dogs with skin lesions. However tissue invasion was not demonstrated, which would indicate that in these instances the organism most probably was present on the skin as a contaminant. To date there have been no reports of human infections with this interesting fungus. Although cases of ringworm due to *K. ajelloi* are apparently very rare, it is possible that future studies may show this fungus to be an occasional cause of disease.

Trichosporon cutaneum Members of this genus are not dermatophytes. They are the cause of a fungus disease of the hair known as white piedra. In

fectured hairs are characterized by the formation of nodules along their shafts. Most of the cases in man have been reported from South America, Asia, and Europe.³³ The paucity of reported human cases from the United States⁴⁴ would indicate that white piedra is rare there. Piedra has been reported in lower animals. This condition has been observed in a horse⁴⁵ and in clinical materials from chimpanzees in Germany.⁴¹ The chimpanzee cases, however, appear to be black piedra, a disease caused by an ascomycete *Piedraia hortai*. The organism responsible for the equine case is not known. As noted in TABLE 1 we recovered a *Trichosporon* from a monkey⁴⁴ with clinical white piedra. This is the first such authenticated infection to be reported from a simian. The animal involved had recently been imported from Latin America. This interesting finding suggests the possibility that piedra may not be uncommon in the simians of Latin America.

It is evident from the above comments that the dermatophytes encountered in our survey show varying degrees of host specificity. Some are pathogenic for a diversity of animals; others parasitize only a few species. It is interesting to note that, with the exception of *A. ajelloi*, all fungi to date have been shown capable of infecting man. With these points in mind let us examine some of our findings with respect to the individual host animal species.

Infections in dogs. As noted on TABLE 1, specimens from 1659 dogs were received for laboratory study. Of this number 368 (22.1 per cent) were found to be positive by culture. In order of frequency, the three principal dermatophytes encountered were *M. canis* in 252 cases (69 per cent of the infections), *M. gypsum* in 68 cases (18 per cent of the infections), and *T. mentagrophytes* in 41 cases (11 per cent of the infections). The dermatophytes *M. audouinii*, *M. distortum*, and *T. rubrum* each were recovered once, and the soil organism *A. ajelloi* was recovered three times.

There was no apparent difference in the clinical picture produced by the various dermatophytes encountered. Approximately 80 per cent of the infections occurred in dogs less than two years of age. The most common lesions observed by the attending veterinarians in this series were circular, scaly areas of alopecia, with or without crust formation, located on the head and extremities. Kerion development was reported in some cases, and in others the clinical picture simulated eczema.

Of the 368 dogs shown to have ringworm, there was a history of skin lesions among members of 36 owner families. In 30 of these episodes *M. canis* was recovered from the dogs involved. The dermatophytes *M. audouinii*, *M. distortum*, *M. gypsum*, and *T. rubrum* were each isolated in one instance from the other dogs. *T. mentagrophytes* was recovered in two cases from the pets involved. Unfortunately, in only five episodes did the attending physicians culture the human cases or submit clinical materials to a public health laboratory for mycologic study. Consequently, evidence that the humans were suffering from the same infection as their pets must be considered presumptive.

Infections in cats. Specimens from 524 cats were submitted for laboratory study (TABLE 1). Of this number 200 (38.1 per cent) were found to be positive by culture. The principal dermatophyte encountered in this series was *M. canis*. The fungus was isolated in 193 cases (96.5 per cent of the cases). *M.*

gypseum and *T. mentagrophytes* were recovered in 4 cases and 3 cases respectively.⁴⁵⁻⁴⁸ Approximately 65 per cent of the infections occurred in cats less than one year of age, approximately 50 per cent of the infections occurred in felines less than two years of age. The lesions observed by the attending veterinarians varied from scattered loss of hair to diffuse scaling and alopecia. The areas of the body involved were principally the head and the extremities.

The public health importance of feline ringworm was emphasized by the fact that among the owners of the 200 cats found to be infected there was a history of skin lesions among members of 63 of the families. In all such cases *M. canis* was recovered from the felines involved. As far as could be determined in 12 instances in which lesions were reported among human contacts the cats involved had been recently purchased from breeding establishments; in 10 instances the animals involved were strays that had been recently brought into the home; in 2 episodes the animals had been recently obtained from friends; and in 14 cases the felines had been permitted to run out of doors at will. In the remaining cases it was not possible to obtain additional information. It should be pointed out that in only 3 of these episodes were the humans cultured by the attending physicians and *M. canis* was recovered by the dermatologists from these patients. Therefore evidence that in the remaining instances the humans were infected by the same agent as the cats must be considered presumptive.

Infections in cattle. Clinical materials for laboratory study were received from 58 head of cattle including dairy and beef breeds (TABLE 1). Of this number 29 (50.0 per cent) were found to be positive on culture. The only dermatophyte encountered was *T. verrucosum*. Although ringworm in cattle is most commonly observed during the fall and winter months when cattle are kept in barns, three outbreaks were reported in the summer months. The lesions most commonly observed were raised, crusted areas, circular in outline, located principally on the head, neck, and tail regions.

In 3 outbreaks of cattle ringworm lesions were reported among members of the owner families. Unfortunately the attending physicians did not culture the human cases. On the basis of clinical appearance, however, the lesions were diagnosed as animal type ringworm. In one such episode a farmer, his wife, and their 4 children, ranging in age from 6 months to 10 years, developed skin lesions. It is of interest to note that the 6 month old infant, who had never been in contact with the animals, was the first to become infected, and that the father, who had the greatest contact with the infected stock, was the last to develop lesions. This episode again demonstrates the possibility of indirect spread of this infection.⁴⁹

Infections in horses. During the year 1956 clinical materials from 56 horses were examined (TABLE 1). Of this number 18 (32.1 per cent) were found to be positive on culture. Three different dermatophyte species were isolated. They were *M. canis*, *M. gypseum*, and *T. equinum*.

The first organism, *M. canis*, is an important cause of equine ringworm in some parts of the world.⁴⁵ In this country, however, little information is available concerning the prevalence of this infection in horses. A review of the literature shows only one previously reported case of *M. canis* ringworm in

a horse in the United States⁴⁹ Our two isolations of this dermatophyte represent the second and third times that *M. canis* has been demonstrated to cause equine ringworm in the United States

The second organism, *M. gypsum*, does not appear to be a common cause of equine ringworm A review of the world literature by Ajello⁴ in 1953 disclosed that up to that time only 50 such infections had been recorded All of the cases had been described from Europe, with the exception of 2 from Madagascar Prior to 1956 this disease had not been reported in horses in this country Since that time we have recovered *M. gypsum* from 12 horses with clinical ringworm Eight of the animals were involved in an epizootic of ringworm⁵⁰ and 4 appeared as isolated cases Thus *M. gypsum* does not appear to be an uncommon equine pathogen in this country

The third fungus *T. equinum* was recovered from horses in four instances

In our series the clinical picture produced by the three dermatophytes was basically the same The appearance of the lesions varied with the stage of the disease Early lesions appeared as raised crusted areas that tended to be circular in outline As the disease progressed the crusts fell off leaving scaly areas of alopecia Where widespread loss or breakage of hair had occurred the affected areas presented a moth eaten appearance

Although there was close contact between the infected horses and humans over a long period of time there was no evidence of transmission of disease from equines to man

Infections in monkeys Specimens from 21 monkeys were received for study (TABLE 1) All the simians were of species indigenous to Central and South America Ten of the animals were found to be infected with *M. canis* four with *M. distortum* and one with *Trichosporon cutaneum*

The lesions most commonly seen were scaliness of the skin with varying degrees of alopecia In two cases a more severe tissue reaction was observed as evidenced by the development of kerions The parts of the body most commonly involved were the extremities and tail

Five of the simians shown to be infected with *M. canis* and three with *M. distortum* were reported to have transmitted the disease to human contacts However the attending physicians did not culture the human cases to determine the etiological agent The diagnoses of ringworm were made on the basis of clinical appearance More intensive studies are indicated to determine the full public health importance of ringworm in monkeys

Infections in chinchillas Specimens from twelve chinchillas were received for laboratory study (TABLE 1) The dermatophytes *M. canis* and *T. mentagrophytes* were isolated in one and two cases respectively

Blank⁴¹ presented evidence that would indicate that *T. mentagrophytes* is a cause of fur slipping a skin disorder of economic importance to chinchilla breeders In one case where we recovered *T. mentagrophytes* the animal involved was a member of a group of three hundred chinchillas the majority of which were reported to have skin lesions The infected animals had scaly patches of alopecia located most commonly on the nose behind the ears and around the base of the tail In this outbreak the owner and her son were reported to have developed skin lesions as a result of handling the animals

Infections in guinea pigs Nine guinea pigs maintained at the Communicable Disease Center were cultured and *T. mentagrophytes* was isolated in six instances (TABLE 1). This agent had previously been incriminated in an epizootic of ringworm involving several thousand guinea pigs at the Communicable Disease Center.⁶² In our few cases the animals showed scaly erythematous patches of alopecia on the tip of the nose around the eyes and at the base of the tail. None of the individuals handling these animals developed skin lesions.

Infections in other animal hosts *Keratinomyces apellii* was isolated from a Malabar squirrel. Clinical material was received from 2 donkeys (TABLE 1). One of the animals was found to be infected with *T. equinum* and the other with *T. terrucosum*. In addition specimens from 8 rabbits, 3 sheep, 1 chicken, 4 parakeets and 1 chimpanzee were cultured. In none of these cases were ringworm fungi recovered.

Wild Animal Ringworm Study

During the period covered by this report hair specimens from 2350 wild animals were cultured (TABLE 2). Twenty-one different animal species including rodents and larger mammals are included in this series. Of the total number of specimens examined *M. gypsum* was recovered in 3 instances, once

TABLE 2
WILD ANIMAL RINGWORM SURVEY, JANUARY 1956 TO JUNE 1957
FUNGI ISOLATED BY HOST SPECIES

Host animal sp.	No. specimens examined	Fungi isolated		
		<i>M. gypsum</i>	<i>T. mentagrophytes</i>	<i>M. dermatophytes</i>
Opossum (<i>Didelphis marsupialis</i>)	319		6	11
Raccoon (<i>Procyon lotor</i>)	352			1
Skunk (<i>Mephitis mephitis</i>)	246			2
Gray fox (<i>Urocyon cinereoargenteus</i>)	87			
Cottontail rabbit (<i>Sylvilagus floridanus</i>)	205			1
Cotton rat (<i>Sigmodon hispidus</i>)	336		1	57
House mouse (<i>Mus musculus</i>)	91			1
Gray squirrel (<i>Sciurus carolinensis</i>)	3			
Red fox (<i>Vulpes fulva</i>)	18			
Cottontail mouse (<i>Peromyscus polionotus</i>)	18			
Florida squirrel (<i>Sciurus niger</i>)	51			
Marsh rabbit (<i>Sylvilagus palustris</i>)	4			
Domestic cat (feral) (<i>Felis domesticus</i>)	26			
Beaver (<i>Lynx rufus</i>)	61			
Old field mouse (<i>Peromyscus polionotus</i>)	61	2	1	
Harvest mouse (<i>Reithrodontomys sp.</i>)	4			
Roof rat (<i>Rattus attus</i>)	76			
Golden mouse (<i>Peromyscus nuttalli</i>)	1			
Wood rat (<i>Neotoma floridana</i>)	1			
Otter (<i>Lutra canadensis</i>)	1			
Norway rat (<i>Rattus norvegicus</i>)	129	1	1	1
Total	2350	3	9	74

A new species of *Mus sporum*. Pathogenicity never demonstrated.

from a Norway rat (*Rattus norvegicus*), and twice from old field mice (*Peromyscus polionotus*). The dermatophyte *T. mentagrophytes* was isolated from 6 opossums (*Didelphis marsupialis*), 1 cotton rat (*Sigmodon hispidus*), 1 old field mouse (*P. polionotus*), and 1 Norway rat (*R. norvegicus*). In addition as shown on TABLE 2 we obtained 74 isolates of a fungus that at this time is simply referred to as the 'red *Microsporum*'. This organism was recovered from 11 opossums (*D. marsupialis*), 1 raccoon (*Procyon lotor*), 2 skunks (*Mephitis mephitis*), 1 cottontail rabbit (*Sylvilagus floridanus*), 57 cotton rats (*S. hispidus*), 1 house mouse (*Mus musculus*) and 1 Norway rat (*R. norvegicus*). The red *Microsporum* is considered to be a new dermatophyte species²⁴ and a manuscript describing this fungus is now being prepared for publication. A similar organism has been recovered from soil in Idaho and Washington by Cooke²⁴ and in Michigan by Aiello.²⁵ It should be pointed out again that none of the animal hosts from which this organism was recovered showed any skin lesions. In addition the Wood's light and direct microscopic examination of the clinical materials failed to reveal any evidence of infection. Furthermore all experimental attempts to demonstrate pathogenicity of the red *Microsporum* for man and animals have been unsuccessful. Thus the potential public health implications of this new fungus species remain to be determined.

It is of interest to note that the pathogen *Trychophyton mentagrophytes* was recovered from wild rodents and opossums. In a previous report²⁶ similar findings have been recorded. The fact that feral animals, particularly rodents of the type commonly found on farm premises, have been repeatedly demonstrated to be carriers of this dermatophyte may have important epidemiological implications. Large numbers of rodents are commonly found in barns, feed storage bins, and other sites on farms, and rodent carriers of *T. mentagrophytes* may serve as indirect sources of human infections by contaminating the environment with elements of this fungus.

DISCUSSION AND CONCLUSIONS

It is obvious that the term 'ringworm' denotes a clinical entity rather than a specific infection. A number of distinct fungus species can cause dermatophytoses in animals as well as man. Each organism presents its own epidemiological and epizootiological implications. Thus measures designed to control outbreaks of ringworm in human and animal populations must be based upon knowledge of the organisms involved and their ecology. Therefore reliable mycologic support would be imperative for any well founded control program.

Our studies have shown that ringworm in animals is a common dermatological condition with definite public health implications. Nevertheless this entity has been largely ignored by public health workers in the United States to date. To a great degree this situation reflects the neglect of the whole field of medical mycology in the past. In recent years, however, there has been a marked increase of interest in the fungus diseases of man and animals, and it is encouraging to note the many advances made in our over-all knowledge of these infections. It is to be expected that as our understanding of animal ringworm infections accumulates and their public health implications are fully

appreciated control programs will be developed in the United States. Such programs already have been initiated in Great Britain.²⁷

SUMMARY

Based on recent findings a modified classification of the dermatophytes from an epidemiological standpoint is presented and described. This classification includes three groups: the anthropophilic or human type dermatophytes, the zoophilic or animal type, and the geophilic or soil inhabiting fungi. Members of the third group are free living keratinophilic fungi, commonly found in soil which have the ability, under certain conditions, to parasitize human and lower animal hosts and produce disease. The public health importance of these three groups of organisms is reviewed.

The results of the animal ringworm survey conducted during the period of January 1956 to June 1957 are presented and discussed. Clinical materials from 2361 domestic animals with skin lesions were examined. Of these animals 642 (27.2 per cent) were found to be positive by culture. Nine dermatophyte species and a *Trichosporon* caused these infections. They were *Microsporum canis*, *M. gypsum*, *M. audouinii*, *M. distortum*, *Trichophyton mentagrophytes*, *T. equinum*, *T. verrucosum*, *T. rubrum*, *Keratinomyces ajelloi* and *Trichosporon cutaneum*.

Clinical materials from 2350 wild animals trapped in southwestern Georgia also were cultured. These animals did not show any apparent cutaneous abnormalities. The dermatophyte *M. gypsum* was recovered in 3 cases, *T. mentagrophytes* in 9 instances, and a new species of *Microsporum* in 74 other cases.

References

1. MCINDIE I & P. WEBB. 1937. Ringworm fungus growing as a saprophyte under natural conditions. *Arch. Dermatol. and Syphilol.* 36: 98-990.
2. AJELLO L. & M. E. GETZ. 1954. Recovery of dermatophytes from shoes and shower stalls. *J. Invest. Dermatol.* 22: 17-21.
3. WALKER J. 1955. Possible infection of man by indirect transmission of *Trichophyton discoid*. *Brit. Med. J.* 1955(II): 1430-1433.
4. AJELLO L. 1953. The dermatophyte *Microsporum gypsum* as a saprophyte and parasite. *J. Invest. Dermatol.* 21: 157-171.
5. AJELLO L. 1956. Soil as a natural reservoir for human pathogenic fungi. *Science* 123: 876-879.
6. VANBREUSEGHEM R. 1952. Intérêt théorique et pratique d'un nouveau dermatophyte isolé du sol *Keratinomyces ajelloi* gen. nov. sp. nov. *Bull. acad. roy. med. Belg.* 38: 1068-1077.
7. DANIELS G. 1954. Isolation of *Keratinomyces ajelloi* from soils in Great Britain. *Nature* 174: 224-226.
8. FREY D. M. & E. B. DURYE. 1956. The isolation of keratinophilic fungi including *Microsporum gypsum* from Australian soil. *Australian J. Exptl. Biol. Med. Sci.* 34: 199-204.
9. GEORG L. K. 1954. The diagnosis of ringworm in animals. *Vet. Med.* 49: 157-166.
10. MORGES R. W. & L. K. GEORG. 1955. Animal ringworm study. *Vet. Med.* 50: 293-297.
11. GEORG L. K., L. AJELLO & C. PAPAGEORGE. 1954. Use of cycloheximide in the selective isolation of fungi pathogenic to man. *J. Lab. Clin. Med.* 44: 422-428.
12. GEORG L. K. 1950. The relation of nutrition to the growth and morphology of *Trichophyton fasciforme*. *Mycologia* 42: 683-697.
13. GEORG L. K. 1956. The role of animals as vectors of human fungal diseases. *Trans. N. Y. Acad. Sci. Ser. II* 18(7): 639-647.
14. AJELLO L. 1954. Occurrence of *Penicillium asperulatum* and other human pathogenic molds in Panamanian soil. *Am. J. Trop. Med. Hyg.* 3: 897-904.

- 15 LEWIS G M & M E HOPPER 1948 An Introduction to Medical Mycology Year book Chicago Ill
- 16 SWARTZ J H E M ROCKWOOD & E A GLICKLICH 1949 A survey of tinea capitis including favus Arch Dermatol and Syphilol 60 486-494
- 17 BENEDEK T & I M FELSHER 1944 Epidemiology of tinea capitis. I A study of tinea capitis in a dispensary Arch Dermatol and Syphilol 49 120-123
- 18 GORDON M A L AJELLO & I K GEORG 1952 *Microsporium gypseum* and *Histo plasma capsulatum* pores in soil and water Science 116 208
- 19 SABOURAUD R 1908 The dermatomycoses common to man and animals Brit Med J 2 1089-1094
- 20 MURRELL T W 1951 *Microsporium audouinii* isolated from a dog Arch Dermatol and Syphilol 63 638
- 21 SCULLY J P & A M KLIGMAN 1951 Coincident infection of a human and an anthropoid with *Microsporium audouinii* Arch Dermatol and Syphilol 64 495-498
- 22 KAPLAN W & I K GEORG 1957 Isolation of *Microsporium audouinii* from a dog J Invest Dermatol 28 313-315
- 23 VOGEL R A & A M TIMPE 1957 Spontaneous *Microsporium audouinii* infection in a guinea pig J Invest Dermatol 28 311-312
- 24 MYCOLOGY UNIT COMMUNICABLE DISEASE CENTER Chamblce Ga Unpublished data
- 25 DI MENNA M & M J MARPLES 1954 *Microsporium distortum* sp nov from New Zealand Trans Brit Mycol Soc 37 372-374
- 26 KAPLAN W I K GEORG S L HENDRICKS & P A LEEFFER 1957 Isolation of *Microsporium distortum* from animals in the United States J Invest Dermatol 28 449-453
- 27 GEORG L K 1954 The relationship between the downy and granular forms of *Trichophyton mentagrophytes* J Invest Dermatol 23 123-141
- 28 GEORG L K E A HAND & R A MENGES 1956 Observations on rural and urban ringworm J Invest Dermatol 27 335-353
- 29 BATTE E G & W S MILLER 1953 Ringworm in horses and its control J Am Vet Med Assoc 123 111-114
- 30 BROQU ROUSSEAU D A URBAIN & J BAROTTE 1927 Étude des teignes du cheval et l'immunité dans les teignes expérimentales Ann inst Pasteur 41 513-531
- 31 NEFFS & GILLMAN 1931 Contribution à l'étude de la teigne Ann med vét 76 193-209
- 32 MACLANNON J E 1936 Epizootica de tina en caballos y contaminación de hombre Arch uruguay med cir y especial Montevideo 8 498-502
- 33 COVANT N F D T SMITH R D BAKER J L CALLAWAY & D S MARTIN 1954 Manual of Clinical Mycology 2nd ed Saunders Philadelphia Pa
- 34 GEORG L K W KAPLAN & L B CAMP 1951 *Trichophyton equinum*—A re evaluation of its taxonomic status J Invest Dermatol 29 27-37
- 35 BLANK F 1953 Ringworm of cattle due to *Trichophyton discoides* and its transmission to man Can J Comp Med Vet Sci 17 277-281
- 36 WINSWORTH C C & P K C ALSTWICK 1955 A survey of animal mycoses in Britain general aspects Vet Record 67 88-97
- 37 MASKIN D L C L TASCHDJIAN & A C FRANK 1957 The etiology of dermatophytosis Shift from *Trichophyton mentagrophytes* to *Trichophyton rubrum* 1935-1954 A M A Arch Dermatol 76 66-69
- 38 CHAKRABORTY A N S GHOSH & F BLANK 1954 Isolation of *Trichophyton rubrum* (Castellani) Sabouraud 1911 from animals Can J Comp Med Vet Sci 18 436-438
- 39 KAILAN W I K GEORG & R H GLIMP *Trichophyton rubrum* infection in a dog In press
- 40 GEORG L K W KAILAN I AJELLO & W M WILLIAMSON The parasitic nature of the soil fungus *Acremonium ajelloi* In press
- 41 DALY J F 1957 Piedra in Vermont A M A Arch Dermatol 75 584
- 42 MIGLENS M P 1952 Fiebra blanca europea Primer caso descrito en España Actas Dermosifiliograficas Santiago de Compostela 6 1-9
- 43 LOCHTE T 1937 Über das Vorkommen der Fiebra beim Schimpansen und über die Beziehungen der tierischen Fiebra zur Menschlichen Arch Dermatol u Syphilis 175 107-113
- 44 MYCOLOGY UNIT COMMUNICABLE DISEASE CENTER Chamblce Ga Unpublished data
- 45 KAPLAN W L K GEORG & C I BROMLEY 1957 Ringworm in cats caused by *Microsporium gypseum* Vet Med 52 347-349

- 46 GEORG I K C S I OBERTS I W MENGES & W KAPLAN 1957 *Trichophyton mentagrophytes* infections in dogs and cats J Am Vet Med Assoc 130 427-432
- 47 ROOK A J & W FRANK BELL 1954 Cattle ringworm Brit Med J 1954(ii) 1198-1200
- 48 GEORG I K W KAPLAN & L B CAMP 1957 Equine ringworm with special reference to *Trichophyton equinum* Am J Vet Res 18 793-810
- 49 HOERLEIN A B 1945 Studies on animal dermatomycoses I Clinical studies Cornell Vet 35 287-298
- 50 KAPLAN W J L HOPPING & I K GEORG 1957 An outbreak of ringworm in horses caused by the dermatophyte *Microsporum gypsum* J Am Vet Med Assoc 131 329-332
- 51 BLANK F J I BYRNE P J G PILMER & R J AVERY 1953 Isolation of *Trichophyton granulosum* Sabouraud 1907 from chinchillas showing fur slipping Can J Comp Med Vet Sci 17 396-402
- 52 MENGES R W & I K GEORG 1956 An epizootic of ringworm among guinea pigs caused by *Trichophyton mentagrophytes* J Am Vet Med Assoc 128 395-398
- 53 AJELLO L In preparation
- 54 COOK W B 1952 Western fungi II 3 Species from eastern Washington and adjacent Idaho Mycologia 44 245-61
- 55 AJELLO L Unpublished data
- 56 SMITH W W R W MENGES & I K GEORG 1957 Ecology of ringworm fungi on commensal rats from rural premises in southwestern Georgia Am J Trop Med Hyg 6 81-85
- 57 LA TOUCHE C J 1955 The importance of the animal reservoir of infection in the epidemiology of animal type ringworm in man Vet Record 67 666-669

- 15 LEWIS G M & M E HOPPER 1948 An Introduction to Medical Mycology Year book Chicago Ill
- 16 SWARTZ J H L M ROCKWOOD & E A GLICKLICH 1949 A survey of tinea capitis including favus Arch Dermatol and Syphilol 60 486-494
- 17 BENEDEK, T & I M IELSHER 1944 Epidemiology of tinea capitis. 1 A study of tinea capitis in a dispensary Arch Dermatol and Syphilol 49 120-123
- 18 GORDON M A I AJELLO & I K GEORG 1952 *Microsporium gypsum* and *Microplasma capsulatum* spores in soil and water Science 116 208
- 19 SABOURAUD R 1908 The dermatomycoses common to man and animals Brit Med J 2 1089-1094
- 20 MURRELL T W 1951 *Microsporium audouinii* isolated from a dog Arch Dermatol and Syphilol 63 638
- 21 SCULLY J I & A M KLIKMAN 1951 Coincident infection of a human and an anthropoid with *Microsporium audouinii* Arch Dermatol and Syphilol 64 495-498
- 22 KAPLAN W & L K GEORG 1957 Isolation of *Microsporium audouinii* from a dog J Invest Dermatol 28 313-315
- 23 VOGEL R A & A M TIMPE 1957 Spontaneous *Microsporium audouinii* infection in a guinea pig J Invest Dermatol 28 311-312
- 24 MYCOLOGY UNIT COMMUNICABLE DISEASE CENTER Chamblee Ga Unpublished data
- 25 DI MENNA M & M J MARILES 1954 *Microsporium distortum* sp nov from New Zealand Trans Brit Mycol Soc 37 372-374
- 26 KAPLAN W L K GEORG S L HENDRICKS & I A LEEIER 1957 Isolation of *Microsporium distortum* from animals in the United States J Invest Dermatol 28 449-453
- 27 GEORG I K 1954 The relationship between the downy and granular forms of *Trichophyton mentagrophytes* J Invest Dermatol 23 123-141
- 28 GEORG I K E A HAND & R A MENGES 1956 Observations on rural and urban ringworm J Invest Dermatol 27 335-353
- 29 BATTE E G & W S MILLER 1953 Ringworm in horses and its control J Am Vet Med Assoc 123 111-114
- 30 BROCC ROUSSEAU D A URBAIN & J BAROTTE 1927 Étude des teignes du cheval et l'immunité dans les teignes expérimentales Ann inst Pasteur 41 513-551
- 31 NEEDS & GILLAIN 1931 Contribution à l'étude de la teigne Ann med vét 76 193-209
- 32 MACANNOV J E 1936 Epizootica de tina en caballos y contaminacion de hombre Arch uruguay med cir y especial Montevideo 8 498-502
- 33 CONANT N I D T SMITH J D BAKER J L CALLAWAY & D S MARTIN 1954 Manual of Clinical Mycology 2nd ed Saunders Philadelphia Pa
- 34 GEORG I K W KAPLAN & L B CAMP 1957 *Trichophyton equinum*—A re evaluation of its taxonomic status J Invest Dermatol 29 27-37
- 35 BLANK F 1953 Ringworm of cattle due to *Trichophyton discoides* and its transmission to man Can J Comp Med Vet Sci 17 277-281
- 36 AINSWORTH G C & P K C ALSTWICK 1955 A survey of animal mycoses in Britain general aspects Vet Record 67 88-97
- 37 MASKIN D L C I TASCHDJIAN & A C FRANK 1957 The etiology of dermatophytosis Shift from *Trichophyton mentagrophytes* to *Trichophyton rubrum* 1935 1954 A.M.A Arch Dermatol 75 66-69
- 38 CHAARABORTY A N S GHOSH & F BLANK 1954 Isolation of *Trichophyton rubrum* (Castellani) Sabouraud 1911 from animals Can J Comp Med Vet Sci 18 436-438
- 39 KAPLAN W L K GEORG & R H GUMP *Trichophyton rubrum* infection in a dog In press
- 40 GEORG I K W KAPLAN L AJELLO & W M WILLIAMSON The parasitic nature of the soil fungus *Aeratinomyces ajelloi* In press
- 41 DALY J F 1957 Fiedra in Vermont A.M.A Arch Dermatol 75 584
- 42 MIGLENS M P 1952 Fiedra blanca europea Primer caso descrito en España Actas Dermosifilograficas Santiago de Compo tela 6 1-9
- 43 LOCHTE T 1937 Über das Vorkommen der Fiedra beim Schimpan en und über die Beziehungen der tierischen Fiedra zur Menschlichen Arch Dermatol u Syphilis 176 10, 113
- 44 MYCOLOGY UNIT COMMUNICABLE DISEASE CENTER Chamblee Ca Unpublished data
- 45 KAPLAN W L K GEORG & C L BROMLEY 1957 Ringworm in cats caused by *Microsporium gypsum* Vet Med 52 347-349

Etiology

In contrast to the modern trend in virology in which numerous agents are found in nature without diseases to call their own cat scratch fever is somewhat old fashioned. In this case we have a definite disease but no established cause for it although by processes of elimination a virus of some type is the most likely suspect.

Mollaret *et al*²² reported the isolation of a virus as the cause of the disease but confirmation of this work has not been possible to date. Numerous attempts to isolate such an agent in tissue culture by animal injection and by chick embryo investigation have failed with great regularity.

The consensus has been that the agent is probably a member of the psittacosis-lymphogranuloma group. Three factors have lent seeming credibility to this hypothesis. First the disease is characterized by subacute granulomatous lesions of the lymph nodes; second the sera of patients who have recovered from the disease frequently show the presence of complement fixing (CF) antibodies for lymphogranuloma antigen; and third one of the most common viruses present in the domestic feline is the agent of feline pneumonitis, a member of the psittacosis-lymphogranuloma group. Admittedly these criteria form an extremely shaky base for the theoretical selection of an etiological factor but the nature of the agent has been assumed by the majority of investigators.

Recently however Armstrong *et al*²³ have suggested that there is no reason to assume that the causal agent of cat scratch fever is a member of the psittacosis-lymphogranuloma group of viruses. They report that the reaction to lymphogranuloma CF antigen is not a significant factor in diagnosis. Many cases have no rise in titer and some individuals in the older age groups may have a titer with no history of clinical disease. This contention is also emphasized by the work of others^{24, 25}

The presence of inclusion or elementary bodies has been discussed in detail by Mollaret *et al*²². The finding of basophilic intracytoplasmic bodies was considered by these workers as evidence of specific cellular reaction to the agent of benign lymphoreticulosis or direct virus growth. Intracellular bodies were found in both the peripheral primary lesion and in affected lymph nodes. Basophilic inclusion bodies are commonly found in the macrophages of the lesion²⁶ but are also present in the same pathological tissues of other diseases including tularemia, tuberculosis and bacterial adenitis. The absence of inclusion bodies has been noticed in cases of cat scratch fever by Kalter, Prier and Prior²⁷ and it is now generally accepted that there are no specific intracellular bodies associated with the lesions.

The possibility that cat scratch fever may be a manifestation of some other specific disease has been considered but all evidence to date leads to the conclusion that it is a complete and separate pathological entity. Beaumont and Gill¹ have suggested that the disease was the same as rat bite fever but the organisms associated with the latter have not been isolated from active lesions.

CAT SCRATCH FEVER

By J. E. Prier

Merck Sharp & Dohme West Point 1

The close relationship existing between the human and the domestic animals and the interdependence of these species are matters of record in the sociological and scientific literature that forms the history of civilization. At one time such association was close physically as well as economically. Man shared his living quarters, his water and in some instances even his food with the species of agricultural importance, and this cooperative existence frequently resulted in equal sharing of some diseases.

Only a few remnants of this type of existence remain today. It is reported at least through that medium of modern electronics that has become an integral part of existence in the United States that the cowboy still lives with his horse, the shepherd with his flock, and Tarzan with the beasts of the jungle. For the most part, however, modern husbandry dictates that the livestock upon which our way of life depends be maintained separate and apart from the human domicile. Most of the diseases that were common to the species have been brought under control. Progress has been made in these areas.

However, in the relationships of man to animals other than agricultural species, the situation is somewhat different. The home, which was well on its way to becoming a human sanctuary, has been invaded by a curious array of nonhuman species. Also, the communities of the world are overrun with the domestic feline, a species of dubious value but definite detriment.

The most common disease transmitted to man by the domestic cat, aside from insomnia, is a condition that has been given such names as nonbacterial regional lymphadenitis, benign inoculation lymphoreticulosis, and cat bite fever, and L. Foshay of the University of Cincinnati, Cincinnati, Ohio, is credited with the name, cat scratch fever. As a recognized clinical entity, this disease is new, the majority of cases having been reported within the past six years. The number of diagnosed cases is increasing, although there is no direct evidence that the disease actually occurs more frequently now than in previous years. Some cases have been tentatively diagnosed in retrospect one as early as 1925.

It is generally conceded that the disease was first recognized in the United States, although this was unpublished by Foshay in 1932. In 1935 the disease was described in Great Britain by Beaumont and Gill¹ and by Brown.² Following the descriptions by Debre *et al* in 1950³ and by Mollaret *et al* in France, and Creer and Keefer (1951)¹² in the United States, numerous case histories and research reports appeared in the literature of the United States and other countries.¹⁴⁻¹⁹ There is little doubt that the disease is one of clinical and public health importance and that probably the present incidence will persist or increase until more information regarding etiology and prophylaxis becomes available.

Etiology

In contrast to the modern trend in virology in which numerous agents are found in nature without diseases to call their own cat scratch fever is some what old fashioned. In this case we have a definite disease but no established cause for it although by processes of elimination a virus of some type is the most likely suspect.

Mollaret *et al*²² reported the isolation of a virus as the cause of the disease but confirmation of this work has not been possible to date. Numerous attempts to isolate such an agent in tissue culture by animal injection and by chick-embryo investigation have failed with great regularity.

The consensus has been that the agent is probably a member of the psittacosis-lymphogranuloma group. Three factors have lent seeming credibility to this hypothesis. First the disease is characterized by subacute granulomatous lesions of the lymph nodes; second the sera of patients who have recovered from the disease frequently show the presence of complement fixing (CF) antibodies for lymphogranuloma antigen; and third one of the most common viruses present in the domestic feline is the agent of feline pneumonitis, a member of the psittacosis-lymphogranuloma group. Admittedly these criteria form an extremely shaky base for the theoretical selection of an etiological factor but the nature of the agent has been assumed by the majority of investigators.

Recently however Armstrong *et al*²³ have suggested that there is no reason to assume that the causal agent of cat scratch fever is a member of the psittacosis-lymphogranuloma group of viruses. They report that the reaction to lymphogranuloma CF antigen is not a significant factor in diagnosis. Many cases have no rise in titer and some individuals in the older age groups may have a titer with no history of clinical disease. This contention is also emphasized by the work of others^{24, 25}.

The presence of inclusion or elementary bodies has been discussed in detail by Mollaret *et al*²⁴. The finding of basophilic intracytoplasmic bodies was considered by these workers as evidence of specific cellular reaction to the agent of benign lymphoreticulosis or direct virus growth. Intracellular bodies were found in both the peripheral primary lesion and in affected lymph nodes. Basophilic inclusion bodies are commonly found in the macrophages of the lesion²⁵ but are also present in the same pathological tissues of other diseases including tularemia, tuberculosis and bacterial adenitis. The absence of inclusion bodies has been noticed in cases of cat scratch fever by Kalter, Prier and Prior²³ and it is now generally accepted that there are no specific intracellular bodies associated with the lesions.

The possibility that cat scratch fever may be a manifestation of some other specific disease has been considered but all evidence to date leads to the conclusion that it is a complete and separate pathological entity. Beaumont and Gill⁹ have suggested that the disease was the same as rat bite fever but the organisms associated with the latter have not been isolated from active lesions.

Source of Infection

The most frequent predisposing factor to the clinical case of cat scratch fever is a skin injury produced by the claw of a cat. The age of the offending animal is not important. Cat bites and, occasionally, dog bites also have been incriminated. At least one case has been attributed to contact with urine. Other sources of infection have been a porcupine quill, wooden splinters, thorns, insect bites, and abrasions.^{23, 24} It is not known, of course, whether such vectors were contaminated with the infecting agent by cats, either directly or indirectly.

It is reasonable to assume that cats may carry a transmissible agent in their upper respiratory system or oral cavity. The feline habit of frequently licking the paws results in almost constant contamination of the claws with saliva. It remains to be demonstrated, however, that clinically normal cats are carriers of a viral agent. With the present knowledge of orphan viruses in many species, it becomes plausible to assume that such a possibility exists.

Cats that have been incriminated definitely in the transmission of cat scratch fever have been examined in detail. It is usual that the animals involved are normal, even to detailed clinical and laboratory examination, but such normal cats are capable of subsequent transmission of the disease; even multiple family outbreaks have been noted.^{25, 27} Thompson and Miller²⁸ in discussing a case of cat scratch fever encephalitis, noted that an involved cat died several days after onset of the primary lesion. There were insufficient facts available to assume that this death was related to the capacity of the animal to transmit cat scratch fever.

Detailed post mortem examinations of cats involved in the production of cat scratch fever cases have been done by the author. In no case has there been identified a lesion or disease syndrome that could be connected with the animal's ability to transmit disease. In at least one case it was demonstrated that the animal was capable of inducing cat scratch fever at the time of post mortem inspection. This cat, received from a patient who had recently developed the disease, accidentally scratched an individual at the time of examination. As a result, a mild but typical case of cat scratch fever developed, including initial lesion, lymphadenopathy, and subsequent development of a positive reaction to a skin test antigen.

In general, therefore, the sources of infection shed no light on the problem of etiology. In fact, there is no evidence indicating that the cat acts other than merely as a common mechanical carrier of the disease agent.

Transmission to Experimental Animals

Information to date tends to indicate that man is peculiar in his susceptibility to benign lymphoreticulosis. Mollaret²⁹ described the transmission of the disease to monkeys. He injected a suspension of lymph node obtained from a human case and produced both a local lesion and regional lymphadenopathy. Unfortunately, however, sufficient confirmation of this work has never been presented.

Peterman³⁰ in discussing a probable case that occurred in 1925, noted that materials from the lesions of the patient were injected into guinea pigs. The

injected animals developed lung lesions that reportedly bore a histological similarity to those of the naturally infected human

For the most part the use of laboratory animals in the study of cat scratch fever has been entirely unsatisfactory. Mice guinea pigs rats rabbits chicks suckling animals and hamsters have all been used in attempts to elicit a regular response to injection of supposedly infective materials. Dogs and cats have also been used without success. Studies employing schemes of blind passage in guinea pigs were futile.

It is perhaps of particular interest that cats of any age are totally refractory to the injection of purulent material obtained from the lesions of acute cases. Such animals do not develop and do not naturally carry antibodies against the lymphogranuloma group of agents. Also they do not react to the skin test antigen which is a result of diagnostic significance in the human patient.

The Clinical Disease

The primary lesion. The majority of cases develop a lesion at the site of inoculation. Although the time of appearance is variable in the extreme it usually appears 3 to 14 days after the injury. One case¹⁰ had an apparent incubation period of 62 days. It is a relatively benign lesion being red slightly raised and papular in appearance. It may proceed to a minor ulceration with a small necrotic core. Small multiple papular eruptions have been observed in some cases and infrequently a transient localized measles like eruption occurs.

Systemic symptoms. The majority of patients exhibit a febrile period some temperatures exceeding 104° F. The course of the fever is quite variable. With some an undulating curve is noted others may have a low fever of short duration (1 or 2 days) still others may have a relatively high fever that persists for 4 days or longer. About one third of the patients are afebrile.

Other general symptoms include anorexia malaise weakness chills and aches. Many cases are hospitalized although this acute phase subsides within a few days in most instances.

Lymphadenopathy. Aside from the primary skin lesion the most common change associated with the disease is a regional lymphadenopathy. Varying degrees of swelling are noted and nodes of nearly any area may be involved. Most typically the immediate draining lymph gland is affected first and subsequently the secondary nodes become involved. Thus in a scratch or bite about the hand swelling of the epitrochlear node is seen after several days and then the axillary nodes are involved. Submandibular suprascapular preauricular and cervical nodes are enlarged following injuries about the face head and neck. Such nodes may also be secondarily involved following inoculations of the upper limbs. Hilar and mesenteric nodes have been involved in a few cases but the pathogenesis of such conditions is not clear.

The affected nodes are swollen and produce varying degrees of pain and tenderness. As the disease progresses the enlargement may cease and the node becomes indurated. In other instances the nodes enlarge suppurate and frequently require surgical drainage. The pus derived from such lesions is thick gray and bacteriologically sterile. Nodes recede in size in 2 to 6

weeks, though some may require a longer period to return completely to normal. Surgical removal of the affected node does not seem indicated commonly.

Involvement of the nervous system By far the most serious sequela of cat scratch fever is involvement of the central nervous system (CNS). Such complications were first brought to light by Stevens²⁴. In describing the disease in a 13 year old boy, he recorded convulsions, stupor, coma, and other evidence of CNS disturbance. In the same year Debre²⁵ reported similar symptoms in a case in France. These cases were actually diagnosed in retrospect. Thompson and Miller²³ recognized the first case during the acute stages of the disease. The central nervous symptoms did not become apparent until one month after the initial injury. Sufficient cases of this type have not been recognized to ascertain without doubt that CNS involvement is an important and not altogether rare result of infection with the agent of benign lymphoreticulosis.

Other sequelae In the examination of patients presumably ill of cat scratch fever, other involvements have been observed. In some cases, however, it is not quite clear that the symptoms and lesions are referable to the primary disease. Rather, they may be coincidental disease syndromes.

Belber, Davis and Epstein¹ reported the association of thrombocytopenic purpura with a typical regional lymphadenopathy type of cat scratch disease. Five weeks after the initial injury, and following a course of regional lymphadenopathy, generalized petechiae in the skin, hematuria, bleeding gums, and blood in the stools characterized a well defined disease process. The condition was sufficiently severe to require splenectomy. After recovery, a cat scratch skin test was done and supposedly due to an allergic response, a brief episode of purpura occurred.

At least two cases of bone involvement have been reported due to, or at least associated with, cat scratch fever.²⁶ One patient developed a granulomatous osteomyelitis on a digit following a severe cat bite. The second individual developed a lesion on the right ileum, a presumptive diagnosis of eosinophilic granuloma, which healed spontaneously in nine months. The latter case had a typical syndrome of lymphadenopathy preceding development of the bone lesion.

Lesions of the eye have been reported, though these are not of frequent occurrence. Daniels and MacMurray²⁷ have described a conjunctivitis and preauricular adenitis due, possibly, to entrance of the infection via the nasopharynx. Parinaud's oculoglandular syndrome has been discussed by Casady and Culbertson^{15, 28} and others^{22, 29} as a probable result of direct conjunctival infection. This syndrome, nonspecific in etiology, may also result from infection with the tubercle bacillus, yeast *Pasteurella tularensis*, viruses, and *Leptothrix*. Pronounced enlargement of the regional lymph nodes is associated with a severe acute conjunctivitis.

Pathology

The specific pathology is associated either with the primary lesion or the involved lymph node. Grossly, the primary lesion is a papular eruption

microscopically it is a superficial circumscribed ulceration with an underlying acute inflammatory process²²⁴

Lymph nodes appear either as small indurated specimens or enlarged firm masses encapsulated and with or without evidence of suppuration. On section white necrotic nodules are apparent beneath the capsule bearing a distinct resemblance to tubercles. Cut surfaces usually have a pearl gray to tan background.

The histopathological appearance of lymph nodes has been classified into three groups by Kalter, Prier, and Prior²²⁵ (1) acute caseous (2) acute necrotizing and (3) epithelioid cell. In all types a capsular thickening with marked chronic inflammatory cell reaction is noted. Large numbers of eosinophils frequently are present among the plasma cells and lymphocytes in the capsular region and giant cells are sometimes present in various parts of the lesions.

The acute caseous lesion is essentially a soft tuberculoid granuloma. There is a large central amorphous caseous mass around which is found a peripheral zone of epithelioid cells. Foreign body type giant cells also may be present in this area. Lymphocytes are likewise present and many of the epithelioid cells show degenerative changes. These lesions are not readily distinguishable from reactions induced by the tubercle bacillus.

Acute necrotizing lesions are characterized by a central core of polymorphonuclear leukocytes. There is no well-defined layer of epithelioid cells at the periphery and no definite demarcation from surrounding tissue. A marked tendency for fusion of adjacent lesions is observed.

Epithelioid cell lesions are essentially collections of epithelioid cell granuloma. No evidence of caseation or cellular degeneration is present. The cells are large with vesicular nuclei, acidophilic nucleoli and a relatively large portion of cytoplasm. Marked germinal center hyperplasia is also observed in this type of involvement.

Diagnosis

Tentative diagnosis is possible by consideration of the history of a cat bite or scratch followed by the development of the typical picture of lymphadenopathy and gradual spontaneous regression of symptoms and lesions. More difficulty is encountered of course when atypical syndromes occur.

I do not suggest however that full attention need not be given to complete examination for diseases that may produce similar clinical pictures such as tularemia, brucellosis, lymphogranuloma venereum, tuberculosis and neoplastic disease of the lymphatic system. Experience indicates that overenthusiasm in the diagnosis of cat scratch fever may be as misleading as lack of consideration of the malady. Consequently an adequate laboratory diagnostic procedure may include in addition to complete hematological studies, serological examination for *Brucella*, tularemia, infectious mononucleosis and lymphogranuloma and skin tests for histoplasmosis, blastomycosis and tuberculosis.

Supporting evidence for diagnosis can be obtained by histological examination of lymph nodes if surgical intervention is indicated.

If positive serologic examination for antibodies to lygranum antigen may support the diagnosis (TABLE 1). It must be emphasized however that

weeks though some may require a longer period to return completely to normal. Surgical removal of the affected node does not seem indicated commonly.

Involvement of the nervous system By far the most serious sequela of cat scratch fever is involvement of the central nervous system (CNS). Such complications were first brought to light by Stevens³⁴. In describing the disease in a 13 year old boy he recorded convulsions, stupor, coma, and other evidence of CNS disturbance. In the same year, Debre³⁵ reported similar symptoms in a case in France. These cases were actually diagnosed in retrospect. Thompson and Miller³⁶ recognized the first case during the acute stages of the disease. The central nervous symptoms did not become apparent until one month after the initial injury. Sufficient cases of this type have not been recognized to ascertain without doubt that CNS involvement is an important and not altogether rare result of infection with the agent of benign lympho-reticulosis.

Other sequelae In the examination of patients presumably ill of cat scratch fever other involvements have been observed. In some cases however it is not quite clear that the symptoms and lesions are referable to the primary disease. Rather, they may be coincidental disease syndromes.

Belber, Davis and Epstein⁴ reported the association of thrombocytopenic purpura with a typical regional lymphadenopathy type of cat scratch disease. Five weeks after the initial injury and following a course of regional lymphadenopathy generalized petechiae in the skin, hematuria, bleeding gums and blood in the stools characterized a well defined disease process. The condition was sufficiently severe to require splenectomy. After recovery a cat scratch skin test was done and supposedly due to an allergic response a brief episode of purpura occurred.

At least two cases of bone involvement have been reported due to or at least associated with cat scratch fever⁴³. One patient developed a granulomatous osteomyelitis on a digit following a severe cat bite. The second individual developed a lesion on the right ileum a presumptive diagnosis of eosinophilic granuloma which healed spontaneously in nine months. The latter case had a typical syndrome of lymphadenopathy preceding development of the bone lesion.

Lesions of the eye have been reported though these are not of frequent occurrence. Daniels and MacMurray⁴⁴ have described a conjunctivitis and preauricular adenitis due possibly to entrance of the infection via the nasopharynx. Parinaud's oculoglandular syndrome has been discussed by Cassidy and Culbertson⁴⁵ and others^{3, 46} as a probable result of direct conjunctival infection. This syndrome nonspecific in etiology may also result from infection with the tubercle bacillus, yeast *Pasteurella tularensis*, viruses and *Leptothrix*. Pronounced enlargement of the regional lymph nodes is associated with a severe acute conjunctivitis.

Pathology

The specific pathology is associated either with the primary lesion or the involved lymph node. Grossly the primary lesion is a papular eruption

microscopically it is a superficial circumscribed ulceration with an underlying acute inflammatory process²⁵

Lymph nodes appear either as small indurated specimens or enlarged firm masses encapsulated and with or without evidence of suppuration. On section white necrotic nodules are apparent beneath the capsule bearing a distinct resemblance to tubercles. Cut surfaces usually have a pearl gray to tan background.

The histopathological appearance of lymph nodes has been classified into three groups by Kalter, Prier, and Prior²⁶ (1) acute caseous (2) acute necrotizing and (3) epithelioid cell. In all types a capsular thickening with marked chronic inflammatory cell reaction is noted. Large numbers of eosinophils frequently are present among the plasma cells and lymphocytes in the capsular region and giant cells are sometimes present in various parts of the lesions.

The acute caseous lesion is essentially a soft tuberculoid granuloma. There is a large central amorphous caseous mass around which is found a peripheral zone of epithelioid cells. Foreign body type giant cells also may be present in this area. Lymphocytes are likewise present and many of the epithelioid cells show degenerative changes. These lesions are not readily distinguishable from reactions induced by the tubercle bacillus.

Acute necrotizing lesions are characterized by a central core of polymorphonuclear leukocytes. There is no well-defined layer of epithelioid cells at the periphery and no definite demarcation from surrounding tissue. A marked tendency for fusion of adjacent lesions is observed.

Epithelioid cell lesions are essentially collections of epithelioid cell granuloma. No evidence of caseation or cellular degeneration is present. The cells are large with vesicular nuclei, acidophilic nucleoli and a relatively large portion of cytoplasm. Marked germinal center hyperplasia is also observed in this type of involvement.

Diagnosis

Tentative diagnosis is possible by consideration of the history of a cat bite or scratch followed by the development of the typical picture of lymphadenopathy and gradual spontaneous regression of symptoms and lesions. More difficulty is encountered of course when atypical syndromes occur.

I do not suggest however that full attention need not be given to complete examination for diseases that may produce similar clinical pictures such as tularemia, brucellosis, lymphogranuloma venereum, tuberculosis and neoplastic disease of the lymphatic system. Experience indicates that overenthusiasm in the diagnosis of cat scratch fever may be as misleading as lack of consideration of the malady. Consequently an adequate laboratory diagnostic procedure may include in addition to complete hematological studies, serological examination for *Brucella*, tularemia, infectious mononucleosis and lymphogranuloma and skin tests for histoplasmosis, blastomycosis and tuberculosis.

Supporting evidence for diagnosis can be obtained by histological examination of lymph nodes if surgical intervention is indicated.

If positive serologic examination for antibodies to lymph node antigen may support the diagnosis (TABLE 1). It must be emphasized however that

weeks though some may require a longer period to return completely to normal. Surgical removal of the affected node does not seem indicated commonly.

Involvement of the nervous system By far the most serious sequela of cat scratch fever is involvement of the central nervous system (CNS). Such complications were first brought to light by Stevens²⁴¹. In describing the disease in a 13 year old boy he recorded convulsions, stupor, coma, and other evidence of CNS disturbance. In the same year, Debre²⁴² reported similar symptoms in a case in France. These cases were actually diagnosed in retrospect. Thompson and Miller²⁴³ recognized the first case during the acute stages of the disease. The central nervous symptoms did not become apparent until one month after the initial injury. Sufficient cases of this type have not been recognized to ascertain without doubt that CNS involvement is an important and not altogether rare result of infection with the agent of benign lymphoreticulosis.

Other sequelae In the examination of patients presumably ill of cat scratch fever other involvements have been observed. In some cases however it is not quite clear that the symptoms and lesions are referable to the primary disease. Rather they may be coincidental disease syndromes.

Belber, Davis, and Epstein⁴² reported the association of thrombocytopenic purpura with a typical regional lymphadenopathy type of cat scratch disease. Five weeks after the initial injury, and following a course of regional lymphadenopathy, generalized petechiae in the skin, hematuria, bleeding gums, and blood in the stools characterized a well defined disease process. The condition was sufficiently severe to require splenectomy. After recovery, a cat scratch skin test was done and supposedly due to an allergic response, a brief episode of purpura occurred.

At least two cases of bone involvement have been reported due to, or at least associated with, cat scratch fever⁴³. One patient developed a granulomatous osteomyelitis on a digit following a severe cat bite. The second individual developed a lesion on the right ileum, a presumptive diagnosis of eosinophilic granuloma, which healed spontaneously in nine months. The latter case had a typical syndrome of lymphadenopathy preceding development of the bone lesion.

Lesions of the eye have been reported though these are not of frequent occurrence. Daniels and MacMurray⁴⁴ have described a conjunctivitis and preauricular adenitis due possibly to entrance of the infection via the nasopharynx. Parinaud's oculoglandular syndrome has been discussed by Casady and Culbertson^{45, 246} and others^{3, 49} as a probable result of direct conjunctival infection. This syndrome, nonspecific in etiology, may also result from infection with the tubercle bacillus, yeast *Pasteurella tularensis*, viruses, and *Leptothrix*. Pronounced enlargement of the regional lymph nodes is associated with a severe acute conjunctivitis.

Pathology

The specific pathology is associated either with the primary lesion or the involved lymph node. Grossly the primary lesion is a papular eruption

microscopically it is a superficial circumscribed ulceration with an underlying acute inflammatory process²³⁵

Lymph nodes appear either as small indurated specimens or enlarged firm masses encapsulated, and with or without evidence of suppuration. On section white necrotic nodules are apparent beneath the capsule bearing a distinct resemblance to tubercles. Cut surfaces usually have a pearl gray to tan background.

The histopathological appearance of lymph nodes has been classified into three groups by Kalter Prier and Prior²³¹ (1) acute caseous (2) acute necrotizing and (3) epithelioid cell. In all types a capsular thickening with marked chronic inflammatory cell reaction is noted. Large numbers of eosinophils frequently are present among the plasma cells and lymphocytes in the capsular region and giant cells are sometimes present in various parts of the lesions.

The acute caseous lesion is essentially a soft tuberculoid granuloma. There is a large central amorphous caseous mass around which is found a peripheral zone of epithelioid cells. Foreign body type giant cells also may be present in this area. Lymphocytes are likewise present and many of the epithelioid cells show degenerative changes. These lesions are not readily distinguishable from reactions induced by the tubercle bacillus.

Acute necrotizing lesions are characterized by a central core of polymorphonuclear leukocytes. There is no well defined layer of epithelioid cells at the periphery and no definite demarcation from surrounding tissue. A marked tendency for fusion of adjacent lesions is observed.

Epithelioid cell lesions are essentially collections of epithelioid cell granuloma. No evidence of caseation or cellular degeneration is present. The cells are large with vesicular nuclei, acidophilic nucleoli and a relatively large portion of cytoplasm. Marked germinal center hyperplasia is also observed in this type of involvement.

Diagnosis

Tentative diagnosis is possible by consideration of the history of a cat bite or scratch followed by the development of the typical picture of lymphadenopathy and gradual spontaneous regression of symptoms and lesions. More difficulty is encountered of course when atypical syndromes occur.

I do not suggest however that full attention need not be given to complete examination for diseases that may produce similar clinical pictures such as tularemia, brucellosis, lymphogranuloma venereum, tuberculosis and neoplastic disease of the lymphatic system. Experience indicates that overenthusiasm in the diagnosis of cat scratch fever may be as misleading as lack of consideration of the malady. Consequently an adequate laboratory diagnostic procedure may include in addition to complete hematological studies, serological examination for *Brucella*, tularemia, infectious mononucleosis and lymphogranuloma and skin tests for histoplasmosis, blastomycosis and tuberculosis.

Supporting evidence for diagnosis can be obtained by histological examination of lymph nodes if surgical intervention is indicated.

If positive serologic examination for antibodies to lymph node antigen may support the diagnosis (TABLE 1). It must be emphasized however that

weeks though some may require a longer period to return completely to normal. Surgical removal of the affected node does not seem indicated commonly.

Involvement of the nervous system By far the most serious sequela of cat scratch fever is involvement of the central nervous system (CNS). Such complications were first brought to light by Stevens.⁴¹ In describing the disease in a 13 year old boy he recorded convulsions, stupor, coma, and other evidence of CNS disturbance. In the same year Debre³⁶ reported similar symptoms in a case in France. These cases were actually diagnosed in retrospect. Thompson and Miller³⁸ recognized the first case during the acute stages of the disease. The central nervous symptoms did not become apparent until one month after the initial injury. Sufficient cases of this type have not been recognized to ascertain without doubt that CNS involvement is an important and not altogether rare result of infection with the agent of benign lymphoreticulosis.

Other sequelae In the examination of patients presumably ill of cat scratch fever other involvements have been observed. In some cases however it is not quite clear that the symptoms and lesions are referable to the primary disease. Rather they may be coincidental disease syndromes.

Belber, Davis, and Epstein⁴² reported the association of thrombocytopenic purpura with a typical regional lymphadenopathy type of cat scratch disease. Five weeks after the initial injury, and following a course of regional lymphadenopathy, generalized petechiae in the skin, hematuria, bleeding gums, and blood in the stools characterized a well defined disease process. The condition was sufficiently severe to require splenectomy. After recovery, a cat scratch skin test was done and, supposedly due to an allergic response, a brief episode of purpura occurred.

At least two cases of bone involvement have been reported due to, or at least associated with, cat scratch fever.⁴³ One patient developed a granulomatous osteomyelitis on a digit following a severe cat bite. The second individual developed a lesion on the right ileum, a presumptive diagnosis of eosinophilic granuloma, which healed spontaneously in nine months. The latter case had a typical syndrome of lymphadenopathy preceding development of the bone lesion.

Lesions of the eye have been reported though these are not of frequent occurrence. Daniels and MacMurray⁴⁴ have described a conjunctivitis and preauricular adenitis due possibly to entrance of the infection via the nasopharynx. Parinaud's oculoglandular syndrome has been discussed by Cassidy and Culbertson^{24, 45} and others^{3, 46} as a probable result of direct conjunctival infection. This syndrome, nonspecific in etiology, may also result from infection with the tubercle bacillus, yeast *Pasteurella tularensis*, viruses, and *Leptothrix*. Pronounced enlargement of the regional lymph nodes is associated with a severe acute conjunctivitis.

Pathology

The specific pathology is associated either with the primary lesion or the involved lymph node. Grossly the primary lesion is a papular eruption

TABLE 2
SKIN REACTIONS TO CAT SCRATCH ANTIGEN IN A POPULATION

Total tested	Skin test reaction	No.	Percentage	Scratched by cat		Injection with cat		Known test with cat		Allergy	
				No.	Percentage	No.	Percentage	No.	Percentage	No.	Percentage
Patients 250	Positive	210	84.0	135	65.0	10	33.0	5	2.0	210	100
	Negative†	25	10.0	5	20.0	15	60.0	5	2.0	25	100
	Doubtful	15	6.0	15	100.0	—	—	—	—	15	100
Controls 94	Positive	3	3.2	2	66.7	1	33.3	—	—	—	0
	Negative	83	88.3	56	67.5	7	8.5	20	24.0	—	0
	Doubtful	8	8.5	6	75.0	1	12.5	1	12.5	—	0

Reproduced by permission from *Annals of Internal Medicine* ¹⁸

† Clinical picture suggested cat scratch fever. Several were patients with abscesses or other pathological conditions (proved later)

value has been demonstrated. Brand and Linkel¹⁷ report some benefit to systemic symptoms from the use of chloramphenicol, chlortetracycline and oxytetracycline, but such compounds do not affect the course of the lymphadenitis. Others^{18, 19} report negative results with antibiotics.

If it is true, as some have suggested²⁰ that the cat is the only vector of the disease, the control procedure becomes clear. On the other hand, if cats are merely mechanical vectors, the solution of the problem of prevention is the discovery of the real source of the causal agent.

And so it becomes clear that the playful kitten and the gentle companionable house cat have taken on a new role, namely, that of purveyors of an insidious and sometimes dangerous disease process. Such news must come as a shock to many who look to the domestic feline as a symbol of homely comfort synonymous with the warmth and cheer of the hearthstone. But who would not be disturbed when the last retreat in a troubled world is invaded by an unseen enemy and, as if by a terrible twist of fate, the entrance is facilitated by the second best friend of man?

References

1. BEALMONT G. F. & A. M. GILL. 1935. A case of cat scratch fever. *Brit. Med. J.* 1935 (i) 382.
2. BROWN R. J. 1935. A case of septicaemia resulting from a cat bite. *Can. Med. Assoc. J.* 32: 182-183.
3. DERRÉ R. M., LAMY M. L., JAMMET L., COSTIL & P. MOZZICONACCI. 1950. La maladie des griffes du chat. *Sémin. hóp.* 96: 1895.
4. DERRÉ R. 1950. La maladie des griffes du chat (diseases from cat scratches). *Marseille m. d.* 67: 375-378.
5. DERRÉ R. M., LAMY M. J., JAMMET L., COSTIL & P. MOZZICONACCI. 1950. La maladie des griffes du chat. *Bull. mém. soc. méd. hóp. Paris* 66: 6-9.
6. MOLLARET P. J., REILLY R., BASTIN & P. TOLMANER. 1950. Sur une adénopathie régionale subaiguë et spontanément curable avec intradermoréaction et lésions ganglionnaires particulières. *Bull. mém. soc. m. d. hóp. Paris* 66: 424.
7. MOLLARET P. J., REILLY R., BASTIN & P. TOLMANER. 1951. L'lymphaticulose bénigne d'oculation et de monstration de l'inoculabilité au singe. *Bull. mém. soc. méd. hóp. Paris* 67: 687.

TABLE 1*
RESULTS OF COMPLEMENT FIXATION TESTS WITH LAGRANUM CF ANTIGEN SERA FROM PATIENTS AND CATS

No sera (patients)	No negative	Complement fixing titers (positives)					
		1:5	1:10	1:20	1:40	1:80	1:160
Single 20	10	4	1	1	?	1	1
Paired 2†	0	1 A x	—	—	—	—	—
		B x	—	—	—	—	—
		2 A x	x	x	—	—	—
		B x	x	x	x	x	—
Cats 4	4	—	—	—	—	—	—

Reproduced by permission from *Annals of Internal Medicine*²³

† The two paired sera were obtained approximately two weeks apart. A = acute phase; B = convalescent phase. Both patients were ill several weeks prior to the obtaining of the acute specimen.

such results are not specific, and are negative in over 50 per cent of the cases. A positive CF test has less significance in mature and elderly individuals since members of these age groups normally carry such CF antibodies.

It was demonstrated by Foshay that a specific skin test antigen could be prepared from the exudate of the affected lymph node. Material is treated in a manner similar to that used in the preparation of the Frei antigen for lymphogranuloma diagnosis. Pus is collected in a sterile manner and diluted 1:1 with sterile saline or buffer. The material is heated to 60° C for 2 hours. It is then stored for 24 hours in the refrigerator and reheated to 60° C for an additional hour. Bacterial sterility tests are run and with the advent of newer techniques tests in tissue culture for the presence of cytopathogenic viral agents.

One tenth ml of the antigen is injected intradermally in the forearm and the results of the tests are read 48 hours later though some reactions are clear at 24 hours. An area of erythema with or without induration about the site of injection constitutes a positive reaction however areas of only 1 to 3 mm in diameter are considered doubtful.

The efficacy of skin testing in the diagnosis of cat scratch fever has been considered by many workers (TABLE 2). The consensus is that it is of a high order of specificity although some persons react without prior history of disease such false positives are few.

It is known that ability to develop a positive reaction to the antigen may persist for many years. One case is reported in which the reaction was positive 28 years after clinical disease. Subsequent secondary subclinical infection of course may be a factor in the persistence of the reactive state.

Prevention and Therapy

No adequate measure for prevention by use of biological products is available. Many therapeutic agents have been used but it is doubtful that real

TABLE 2
SKIN REACTIONS TO CAT SCRATCH ANTIGEN IN A POPULATION

Total tested	Skin test reaction	N	Percent	Scratched by cat		I contacted with cats		I know contacts with cats		All together	
				N	Percent	N	Percent	N	Percent	N	Percent
Patients 250	Positive	210	84.0	135	63.0	0	33.0	5	2.0	210	100
	Negative	25	10.0	5	20.0	15	60.0	5	2.0	25	100
	Doubtful	15	6.0	15	100.0	—	—	—	—	15	100
Controls 94	Positive	3	3.2	2	66.7	1	33.3	—	—	—	0
	Negative	83	88.3	56	67.5	7	8.5	20	24.0	—	0
	Doubtful	8	8.5	6	75.0	1	12.5	1	12.5	—	0

Reproduced by permission from *Annals of Internal Medicine*²⁸

† Clinical picture suggested cat scratch fever. Several were patients with abscesses or other pathological conditions (proved later)

value has been demonstrated. Brand and Finkel²⁹ report some benefit to systemic symptoms from the use of chloramphenicol, chlortetracycline and oxytetracycline, but such compounds do not affect the course of the lymphadenitis. Others²⁸⁻³⁰ report negative results with antibiotics.

If it is true, as some have suggested²⁸ that the cat is the only vector of the disease, the control procedure becomes clear. On the other hand, if cats are merely mechanical vectors, the solution of the problem of prevention is the discovery of the real source of the causal agent.

And so it becomes clear that the playful kitten and the gentle companionable house cat have taken on a new role, namely, that of purveyors of an insidious and sometimes dangerous disease process. Such news must come as a shock to many who look to the domestic feline as a symbol of homely comfort, synonymous with the warmth and cheer of the hearthstone. But who would not be disturbed when the last retreat in a troubled world is invaded by an unseen enemy and, as if by a terrible twist of fate, the entrance is facilitated by the second best friend of man?

References

1. BEALMONT G. E. & A. M. GILL. 1935. A case of cat bite fever. *Brit. Med. J.* 1935 (ii) 587.
2. BROWN R. J. 1935. A case of septicæmia resulting from a cat bite. *Can. Med. Assoc. J.* 32: 182-183.
3. DEBRÉ R., M. LAMY, M. L. JAMMET, L. COSTIL & P. MOZZICONACCI. 1950. La maladie des griffes du chat. *Semaine hôp.* 26: 1895.
4. DEBRÉ R. 1950. La maladie des griffes du chat (disease of man cat scratches). *Mar-seille m. d.* 87: 315-378.
5. DEBRÉ R., M. LAMY, M. L. JAMMET, L. COSTIL & P. MOZZICONACCI. 1950. La maladie des griffes du chat. *Bull. m. m. soc. méd. hôp. Paris* 66: 16-9.
6. MOLLARET P., J. REILLY, R. BASTIN & P. TOURNIER. 1950. Sur une adénopathie ganglionnaire papuleuse et spontanément curable avec intradermo-inoculation et lésions locales. *Bull. mém. soc. m. d. h. p. Paris* 66: 424.
7. MOLLARET P., J. REILLY, R. BASTIN & P. TOURNIER. 1951. La lymphatite localisée bénigne d'inoculation et démonstration de l'inoculabilité au singe. *Bull. mém. soc. m. d. hôp. Paris* 67: 687.

TABLE 1
RESULTS OF COMPLEMENT FIXATION TESTS WITH LAGRANUM CF ANTIGEN SERA FROM PATIENTS AND CATS

No sera (patients)	No. negative	Complement fixation titer (positive)					
		1:5	1:10	1:20	1:40	1:80	1:160
Single 20	10	4	1	1	2	1	1
Paired 2†	0	1 A	—	—	—	—	—
		B	—	—	—	—	—
		2 A	+	+	—	—	—
		B	+	+	+	+	—
Cats 4	4	—	—	—	—	—	—

Reproduced by permission from *Annals of Internal Medicine*²³

† The two paired sera were obtained approximately two weeks apart. A = acute phase, B = convalescent phase. Both patients were ill several weeks prior to the obtaining of the acute specimen.

such results are not specific and are negative in over 50 per cent of the cases. A positive CF test has less significance in mature and elderly individuals, since members of these age groups normally carry such CF antibodies.

It was demonstrated by Foshay that a specific skin test antigen could be prepared from the exudate of the affected lymph node. Material is treated in a manner similar to that used in the preparation of the Frei antigen for lymphogranuloma diagnosis. Pus is collected in a sterile manner and diluted 1:1 with sterile saline or buffer. The material is heated to 60° C for 2 hours. It is then stored for 24 hours in the refrigerator and reheated to 60° C for an additional hour. Bacterial sterility tests are run and, with the advent of newer techniques, tests in tissue culture for the presence of cytopathogenic viral agents.

One tenth ml of the antigen is injected intradermally in the forearm and the results of the tests are read 48 hours later, though some reactions are clear at 24 hours. An area of erythema, with or without induration, about the site of injection constitutes a positive reaction; however, areas of only 1 to 3 mm in diameter are considered doubtful.

The efficacy of skin testing in the diagnosis of cat scratch fever has been considered by many workers (TABLE 2). The consensus is that it is of a high order of specificity, although some persons react without prior history of disease; such false positives are few.

It is known that ability to develop a positive reaction to the antigen may persist for many years. One case is reported in which the reaction was positive 28 years after clinical disease. Subsequent secondary subclinical infection, of course, may be a factor in the persistence of the reactive state.

Prevention and Therapy

No adequate measure for prevention by use of biological products is available. Many therapeutic agents have been used, but it is doubtful that real

TABLE 2^a
SKIN REACTIONS TO CAT SCRATCH ANTIGEN IN A POPULATION

Total no. tested	Skin test result	No.	Per cent	Scratched by cat		Total contacts with cats		No. known contact with cat		Adenopathy	
				No.	Per cent	No.	Per cent	No.	Per cent	No.	Per cent
Patients 250	Positive	210	84.0	135	63.0	10	33.0	5	2.0	210	100
	Negative	25	10.0	5	20.0	15	60.0	5	2.0	25	100
	Doubtful	15	6.0	15	100.0	—	—	—	—	15	100
Controls 93	Positive	3	3.2	2	66.7	1	33.3	—	—	—	0
	Negative	83	88.3	36	67.5	7	8.5	20	24.0	—	0
	Doubtful	8	8.5	6	5.0	1	12.5	1	12.5	—	0

Reproduced by permission from *Annals of Internal Medicine*²²

^a Clinical picture suggested cat scratch fever. Several were patients with abscesses or other pathological conditions (proved later).

value has been demonstrated. Brand and Finkel⁴ report some benefit to systemic symptoms from the use of chloramphenicol, chlortetracycline and oxytetracycline, but such compounds do not affect the course of the lymphadenitis. Other^{24, 25} report negative results with antibiotics.

If it is true, as some have suggested²⁷ that the cat is the only vector of the disease, the control procedure becomes clear. On the other hand, if cats are merely mechanical vectors, the solution of the problem of prevention is the discovery of the real source of the causal agent.

And so it becomes clear that the playful kitten and the gentle companionable house cat have taken on a new role, namely, that of purveyors of an insidious and sometimes dangerous disease process. Such news must come as a shock to many who look to the domestic feline as a symbol of homely comfort synonymous with the warmth and cheer of the hearthstone. But who would not be disturbed when the fast retreat in a troubled world is invaded by an unseen enemy and as if by a terrible twist of fate, the entrance is facilitated by the second best friend of man.

References

1. BRACOMO T. G. F. & A. M. GILL. 1935. A case of cat bite fever. *Brit. Med. J.* 1935 (i) 532.
2. BROWN R. J. 1935. A case of septicaemia resulting from a cat bite. *Can. Med. Assoc. J.* 32: 187-193.
3. DEBRÉ K., M. LAMY, I. L. JAMMET, L. COSTIL & P. MOZZICONACCI. 1950. La maladie des griffes du chat. *Semaine méd.* p. 26 (1950).
4. DEBRÉ K. 1950. La maladie des griffes du chat (ch. 2. Les formes from cat scratches). *Mar. soc. mé.* 87: 375-378.
5. DEBRÉ K., I. LAMY, M. L. JAMMET, L. COSTIL & P. MOZZICONACCI. 1950. La maladie des griffes du chat. *Bull. mem. soc. méd. hôp. Paris* 66: 6-9.
6. MOLLAPET P., J. REILLY, R. BASTIN & P. TOLRNER. 1950. Sur une adénopathie régionale aiguë et spontanément curable avec intradermo-réaction et lésions ganglionnaires particulières. *Bull. mem. soc. mé.* hôp. Paris 66: 424.
7. MOLLAPET P., J. REILLY, R. BASTIN & P. TOLRNER. 1951. Lymphaticulose chronique d'inoculation et démonstration de l'inoculabilité au singe. *Bull. mem. soc. mé.* hôp. Paris 67: 687.

- 8 MOLLARET P J REILLY R BASTIN & P TOURNIER 1951 II Lymphoreticulose benigne d'inoculation et caractérisation du virus par sérologie et colorations Bull m m soc méd hôp Paris 67 693
- 9 MOLLARET P J REILLY R BASTIN & P TOURNIER 1950 Réaction de fixation du complément dans la lymphoreticulose benigne d'inoculation Compt rend soc biol 144 1493
- 10 MOLLARET P J REILLY R BASTIN & P TOURNIER 1950 Recherches virologiques dans la lymphoreticulose benigne d'inoculation Compt rend soc biol 144 1495
- 11 MOLLARET P J REILLY R BASTIN & P TOURNIER 1950 Une maladie ganglionnaire nouvelle Adénopathie régionale subaiguë spontanément curable avec intra-dermoréaction et lésions histologiques particulières (A new lymphatic disease Regional subacute adenopathy spontaneously curable with intradermal reaction and unusual histologic lesions) Presse med 58 287-283
- 12 MOLLARET P J REILLY R BASTIN & P TOURNIER 1950 Documentation nouvelle sur l'adénopathie régionale subaiguë et spontanément curable décrite en 1950 la lymphoreticulose benigne d'inoculation (New report on subacute regional adenopathy spontaneously curable described in 1950 benign inoculation lymphoreticulosis) Presse med 58 1353-1355
- 13 GREER W E R & C S KEEFER 1951 Cat scratch fever disease entity New Engl J Med 244 545-548
- 14 ANVINO R 1951 Benign lymphoreticulosis due to inoculation (cat scratch disease) ca e Pediatrics 40 591-597
- 15 BENNETT I L JR & J T MELTON 1951 Cat scratch disease J Med Assoc Georgia 40 466-468
- 16 BENZECRY E 1951 La linforeticulosis benigna de inoculación una nueva enfermedad descubierta recientemente de Francia (Benign lymphoreticulosis due to inoculation a new disease discovered recently in France) Vida nueva 68 140
- 17 BLATTNER R J 1951 Cat scratch fever J Pediat 39 123-124
- 18 BOLGERT M M LE SOURD & M SOULÉ 1950 So called cat scratch disease Bull soc franç dermatol syphilig 57 511-513
- 19 BONNIN H F G MORETTI & P PENE 1951 Sur un cas de lymphoreticulose benigne d'inoculation (maladie par griffes de chat (Case of benign inoculation lymphoreticulosis cat scratch disease) J med Bordeaux et Sud Ouest 128 857
- 20 CACHIN M & J L BEAUMONT 1951 Benign inoculation lymphoreticulosis (cat scratch) with ocular port of entry Paris méd 41 294 295
- 21 CALZETTI G 1951 Una nuova malattia l'adenopatia regionale subacuta o malattia da unghie di gatto o linforeticulosis benigna da inoculazione (New disease regional subacute adenopathy or cat scratch disease benign lymphoreticulosis due to inoculation) Clin Pel hat 33 515-518
- 22 CAT SCRATCH FEVER 1951 Lancet 1951 (i) 958
- 23 CLEMENT R A COMBES HAMELLE & B MEYER 1951 Forme oculaire de la lymphoreticulose benigne dite des griffes de chat (Ocular form of benign lymphoreticulosis or so-called cat scratch disease) Bull m m soc méd hôp Paris 67 1108
- 24 DANIELS W B & F G MACMURRAY 1951 Cat scratch disease non bacterial regional lymphadenitis A M A Arch Internal Med 88 735-736
- 25 DANIELS W B & F C MACMURRAY 1951 Cat scratch disease non bacterial regional lymphadenitis Trans Assoc Am Physicians 64 137-146
- 26 DUERRAT B 1951 Pseudovenereal form of cat scratch disease Bull m m soc méd hôp Paris 67 848 850
- 27 GARNIER G 1950 Subacute regional adenopathy due to cat scratch case Bull soc franç dermatol syphilig 57 395-396
- 28 CSFEL O K FORSTER & I KLAUS 1951 Virus disease caused by cat scratch (lymphadenitis benign lymphoreticulosis) Schweiz med Wochschr 81 699-704
- 29 KAPLAN M P CRUMBACH P BOUVATTIER & M MONTOUCHET 1950 Subacute regional adenopathy spontaneously curable following scratches cat scratch disease Bull m m soc méd hôp Paris 66 852 860
- 30 LANGE H L 1951 Cat scratch fever a report of three cases J Pediat 39 431-434
- 31 LEBOURG M M BURFAU & J COURTIER 1951 Adénopathies cervico-faciales subaiguës spontanément curables Lymphoreticulose d'inoculation (Subacute spontaneously curable cervicofacial adenopathy Inoculation lymphoreticulosis) Rev stomat 52 801-806
- 32 MADURO R P MOLLARET & L G CHEVANCE 1951 Une forme nouvelle de lymphoreticulose benigne d'inoculation la forme d'entrée et pharyngée (New form of benign inoculation lymphoreticulosis port of entry at pharynx) Ann oto laryng 1 68 510-514

- 33 MASHOUEAL D (LEMPERIFRE) & C LEBOW 1950 Subacute polyal nonathy possibly due to scratching by cat case Bull m m soc m d h p Paris 66 410-412
- 34 MERALEY, F I & K WATTEBLFD 1950 Benign inoculation lymphoreticulosis due to cat bite Bull soc franc dermatol syphil - 67 558-559
- 35 MEULEMAN H K 1951 Ein neues stomatologisches Krankheitsbild die Virus Kraz lymphoeritis (A new stomatologic syndrome cat scratch disease) Schweiz Monatsschr Zahnheilk 61 896-901
- 36 ORLANDI F 1950 La malattia delle unghie di gatto (Cat scratch disease) Lat tante 21 610-672
- 37 PIERRE O D 1951 Una nuova enfermedad linforeticulosis benigna de inoculación (A new disease benign inoculation lymphoreticulosis) Irensa pediat 2 139 140
- 38 LACES F, A MAC F & J CAPTEVILLE 1951 Cat scratch disease (benign inoculation lymphoreticulosis) coexistent with primary tuberculous infection Bull mem soc méd h p Paris 67 213-216
- 39 RICHARD J 1951 Disease due to cat scratches curable subacute regional a leucopathy Acta Paediat Belg 5 40-57
- 40 SCARZELLA M & A FORTINA 1951 Su di un caso di linforeticulosis benigna da inoculazione (Case of cat scratch disease) Minerva med 42 904-907
- 41 SICCHER F J J WELT & J SPBAOV 1951 Benign lymphoreticulosis of inoculation due to cat scratches or thorn pricks etiology study of 4 cases Bull m m soc m d h p Paris 67 511-576
- 42 STEHELEN B 1951 Cat scratch disease 3 cases observed in Mediterranean region Bull mem soc méd h p Paris 67 810-834
- 43 TRASSAC F J & J MAROT 1951 So-called cat scratch disease benign lymphoreticulosis due to inoculation J m d Bordeaux et Sud Ouest 128 540-546
- 44 LESTER C & C HEDINGER 1951 Über die maulhe des griffes de chat (On cat scratch disease) Schweiz med Wochschr 81 221 223
- 45 WEILL J & P LÉVY 1950 Cat scratch disease Bull mém soc méd h p Paris 66 967-969
- 46 ABEL F P MICHON P DORNIER & G LASCOMBES 1953 Lymphoreticulose bénigne d'inoculation sousmaxillaire bilatérale chez un adolescent atteint d'ictère hémolytique familial (Bilateral submaxillary benign inoculation lymphoreticulosis in an adolescent with familial hemolytic jaundice) Rev méd Nancy 78 818 931
- 47 ANDRE P & B BREVES 1952 Adénopathies subaiguës par griffure de chat pasturelleuse (Subacute lymph node disease caused by cat scratches pasturellous) Bull mem soc m d h p Paris 68 157 160
- 48 ANDREU LERRA J 1953 La fiebre de la garra del gato (Cat scratch fever) Medicina 30 457-461
- 49 ARAGAO P M DE & E DAVIDOVICH 1953 Doença de Foshay Mollaret (Foshay Mollaret disease) Hospital Rio de Janeiro 43 751-763
- 50 BEDONČEK B 1952 Benign virus lymphoeritis (Cat scratch disease) Časopis lékařů český 91 550-563
- 51 DE LAVERGNE A C THOMAS & B ALGAN 1951 La forme oculo-ganglionnaire de la lymphoreticulose bénigne d'inoculation Bull mem soc méd h p Paris 67 985
- 52 DROLET F L G FAIVRE & B CRALC 1950 Premiers cas lorrains de la maladie des griffures de chat Rev m d Nancy 76 487
- 53 WEGMANN T C LESTER & C HEDINGER 1951 Die sogenannte Katzenkrankheit eine benignen Virus lymphadenitis Schweiz m d Wochschr 81 853
- 54 PARAF J & E PERÉZ 1951 Sur un nouveau cas de maladie des griffes du chat et de l'importance du diagnostic dans certaines circonstances Arch fran pediat 8 31
- 55 LEMAIRE A & J DES AX 1950 Sur une épidémie familiale d'adénite subaiguë bénigne de nature inconnue Bull mém soc méd h p Paris 66 375
- 56 PRIER J F & S S KALTER 1954 Cat scratch fever—a public health problem Vet Med 49 337
- 57 CAT SCRATCH A D ILLARFSLIA 1950 Lancet 1950/11 817
- 58 INÉLIN F & R MARTIN DU PAN 1951 Nos observations of diseases due to cat scratch Paris 67 40 74
- 59 LESME I & P MARCHAND 1950 Sur un nouveau type de conjonctivite infectieuse expérimentalement transmise par un chat J m d Bordeaux et Sud Ouest 127 127 131
- 60 BENIGN INOCULATION LYMPHORETICULOSIS A NEW DISEASE ENTITY 1952 What's New Abbott Labs 166 8-14
- 61 PERARDIVELLI W 1953 Lympho-réticulose bénigne d'inoculation ou maladie des griffes de chat (Benign lymphoreticulosis by inoculation or cat scratch disease) Presse méd 61 1104 1105

- 62 BERARDINELLI W 1953 Fluronicose ou doenca da arranhadura de gato (Ailuronicosis or cat scratch disease) *Revista clin cient São Paulo* 22 120-128
- 63 BERCHER J 1952 Les formes stomatologiques de la lympho-réticulose bénigne d'inoculation maladie des griffes du chat ou maladie de Debré et Mollaret (The oral forms of benign inoculation lymphoreticulosis or cat scratch disease or Debré Mollaret disease) *Rev odontol stomatol* 8 20-22
- 64 BETKE K 1952 Zur Klinik und Cytodagnostik chronischer Lymphknotenerkrankungen die Viruskratzympadenitis benigne Inokulationslymphoretikulose (Symptoms and cytological diagnosis of chronic diseases of the lymph node viral scratch lymphadenitis benign inoculation lymphoreticulosis) *Klin Wochschr* 30 583-588
- 65 BETTLEY F R & F A FAIRBURN 1953 Cat scratch disease the intradermal test in controls *Lancet* 1953(i) 520-521
- 66 BIJLMER J & D GRANAAT 1952 Benign lymphoreticulosis (Cat scratch disease case) *Ned Tijdschr Geneesk* 96 1158-1159
- 67 BORIOLI F & F BADARACCO 1952 A propos d'un cas de maladie dite des griffes de chat (On a case of so called cat scratch disease) *Traxis* 41 263-264
- 68 BREIJER H B & R GISPEN 1953 Lymphoreticulosis benigna als kattenkrabziekte (Benign lymphoreticulosis or cat scratch disease) *Acta leiden* 23 52-58
- 69 BREIJER H B & R GISPEN 1953 Lymphoreticulosis benigna al kattenkrabziekte (Benign lymphoreticulosis as cat scratch disease) *Ned Tijdschr Geneesk* 97 1849-1853
- 70 CAT SCRATCH FEVER 1952 *Lancet* 1952(i) 302
- 71 CAT SCRATCH DISEASE 1952 *J Am Med Assoc* 148 746-747
- 72 CAT SCRATCH DISEASE 1953 *Med World* 78 280-281
- 73 CHAIA J & N TALEB 1953 A propos du premier cas libanais de lymphoréticulose bénigne d'inoculation maladie des griffes du chat (First case of cat scratch disease in Lebanon) *Sang* 24 635-637
- 74 CHARBONNEAU H 1952 Quatre cas de maladie des griffes du chat à Montréal lymphoréticulose aigue bénigne d'inoculation (Four cases of the cat scratch disease in Montreal acute benign inoculation lymphoreticulosis) *Union méd Canada* 81 123-128
- 75 CHIEMIN J & J P IMBERT 1952 A propos d'une adénopathie cervicale à évolution subaigue lymphoréticulose bénigne d'inoculation (Subacute evolution of a disease of a lymph node of the neck benign inoculation lymphoreticulosis) *Rev odontol stomatol* 8 23-25
- 76 CHIARI G 1952 Su di un caso di linforeticulosis infettiva benigna malattia da unghie di gatto (A case of benign infectious lymphoreticulosis cat scratch disease) *Lattante* 23 159-176
- 77 CITTERIO L 1952 Linforeticulosis benigna per morso di cane (Benign lymphoreticulosis due to dog bite) *Arch ital chir* 76 331-346
- 78 CUTLER T D 1952 Benign lymphoreticulosis of inoculation (cat scratch fever) *Calif Med* 76 404-406
- 79 DAESCHNER C W G W SALMON & I M HEYS 1953 Cat scratch fever *J Pediatr* 43 371-384
- 80 DE JONGH J 1953 Benign virus lymphadenitis (cat scratch disease) case *Ned Tijd chr Geneesk* 97 1910-1911
- 81 DENECHAL D 1952 Un nouvel exemple de maladie de griffes de chat excision du foyer et du ganglion infectés guérison spectaculaire (A new example of cat scratch disease excision of the focus and infected lymph node spectacular recovery) *Bull mun soc méd hôp Paris* 68 34-349
- 82 DEPAILLAT A & A CONDAT 1952 A propos d'un cas de thrombophlébite cérébrale au cours d'une lymphoréticulose bénigne d'inoculation (A case of cerebral thrombophlebitis in the course of benign inoculation lymphoreticulosis) *Arch franç pediat* 9 744-745
- 83 DREOSTI A F & J F MURRAY 1953 Cat scratch disease non bacterial regional lymphadenitis or benign lymphoreticulosis of inoculation *S African Med J* 27 81-83
- 84 EUSTEFIN I 1952 Cat scratch fever *A M A Arch Dermatol Syphilol* 66 210-213
- 85 FERGUSON G 1952 Acute non specific mesenteric lymphadenitis possible mechanism of pain illustrated by two cases *A M A Arch Surg* 65 906-911

- 86 FITZPATRICK T J & L W WOODRUFF 1952 Cat scratch disease. *Ill Med J* 102 384-395
- 87 FLOROS A 1953 Die benigne durch Inokulation übertragbare Lymphoretikulose Morbus Petzetakis oder Katzenkratzkrankheit (Benign lymphoreticulosis contagious by inoculation—Morbus Petzetakis or cat scratch disease) *Wien klin Wochschr* 64 963-964
- 89 FOLBERTH S 1952 Die Katzenkratzkrankheit eine neue Virus lymphadenitis (Cat scratch disease a new viral lymphadenitis?) *Deut med Wochschr* 77 943-945
- 90 GENNES I DE & H BRICAIRE 1953 Sur quatre observations d'infections traitées par l'érythromycine (Erythromycin treatment of four infections) *Bull mcm soc med hôp Paris* 69 377-384
- 91 GIFFORD H M BROCKBANK & J MONTMAN 1952 Cat scratch fever. *Stanford Med Bull* 10 157-161
- 97 CROPP C 1952 Su un caso di linforeticulosis benigna da inoculazione malattia la grassia di gatto cat fever (Benign inoculated lymphoreticulosis cat scratch disease cat scratch fever case) *Ciorn mal infettive e parasit* 4 308-310
- 93 CROSSIORD A A WIMBEN & M SELIGMAN 1952 Infectious myelitis aiguë et lymphoreticulose bénigne d'inoculation maladie des griffures du chat (Subacute encephalomyelitis and benign inoculation lymphoreticulosis cat scratch disease) *Bull mcm soc med hôp Paris* 68 671-674
- 94 HAHN A 1952 Benigna virusna limfadenitis (Benign virus lymphadenitis) *Med glasnik (Beogr)* 6 170-172
- 95 HEDINGER C 1952 Die histologischen Veränderungen bei der sog Katzenkratzkrankheit einer benignen Virus lymphadenitis (Histologic changes in so-called cat scratch disease Benign virus lymphadenitis) *Arch pathol Anat u Physiol Verhows* 322 159-174
- 97 HEDINGER C C LESTER & T WEGMANN 1952 Der kutane Primäraffekt der sogenannten Katzenkratzkrankheit einer benignen Virus lymphadenitis (Cutaneous primary affection of the so-called cat scratch disease a benign virus lymphadenitis) *Dermatologica* 104 101-10
- 98 HEGG L 1952 Recueil de faits lymphoreticulose l'origine d'inoculation maladie des griffures du chat (Benign inoculation lymphoreticulosis cat scratch disease) *Dermatologica* 106 198-200
- 99 HENRY M 1952 Leptothricosis conjunctivae (Parinaud's conjunctivitis) report of 13 cases relationship to cat scratch disease. *Trans Pacific Coast Oto-Ophthalm Soc* 33 173-196
- 100 HERBELL W F A BALOWS & B F ROACH 1953 Cat scratch fever report of a case. *J Kentucky State Med Assoc* 53 486-488
- 101 JAEGER H & J DELACRYAZ 1953 Lymphoreticulose et cgu ou maladie des griffures de chat a caractère d'entéropénie (Benign lymphoreticulosis or cat scratch disease with primary penile lesion) *De matologica* 107 260-261
- 102 JAMBOUR J & E EMURA 1953 Benign inoculated lymphoreticulosis (cat scratch disease) report of two cases with positive skin tests. *Am J Arch Dermatol Syphil* 67 437-442
- 103 KALKOFF K W 1952 Zur Ätiologie der Lymphadenitis n follicularis benigna cutis on Hautmanifestationen der chronischen Lymphadenose (Differentiation of lymphadenitis n follicularis benigna cutis from skin manifestation of chronic lymphadenosis) *Dermatol Wochschr* 126 1146-1156
- 104 KAMPMEIER R H 1952 Cat scratch disease. *J Tenn State Med Assoc* 45 494-495
- 105 KASER D F 1953 Katzenkrankheit (cat scratch fever) maladie des griffures de chat) *Ned Tijdsch Geneesk* 97 1958-1961
- 106 KEYSER J I 1953 Een geval van kattekratz ziekte (A case of cat scratch disease) *Monatssch k n iergene sk* 21 341-345
- 107 KRAEPI D 1953 Suppurative lymphadenitis concomitant with infectious mononucleosis. *N Y State J Med* 52 2043
- 108 LAHAM F 1953 Cas cliniques de lymphoreticulose régionale et d'une maladie par griffures de chat (A case of benign regional lymphoreticulosis in Lebanon—cat scratch disease) *Soc med Moyen Orient* 10 391-397
- 109 LANGE H I 1953 Cat scratch fever. *Am J Nursing* 53 832-833
- 110 LATALLA J 1952 Choroba Defektu Molaret chorobu kociego pazusa (Cat scratch disease) *Peliat polska* 27 1109-1112

- 62 BERARDINELLI W 1953 Eluronicose ou doenca da arranhadura de gato (Ailuronicosis or cat scratch disease) *Resenha clin cient Sao Paulo* 22 120-128
- 63 BERCHER, J 1952 Les formes stomatologiques de la lympho-réticulose bénigne d'inoculation maladie des griffes du chat ou maladie de Debré et Mollaret (The oral forms of benign inoculation lymphoreticulosis or cat scratch disease or Debré Mollaret disease) *Rev odontol stomatol* 8 20-22
- 64 BETAE K 1952 Zur Klinik und Cytodagnostik chronischer Lymphknotenerkrankungen die Viruskratzympfadentitis benigne Inokulationslymphoretikulose (Symptoms and cytological diagnosis of chronic diseases of the lymph node viral scratch lymphadenitis benign inoculation lymphoreticulosis) *Klin Wochschr* 30 583-588
- 65 BETTLEY I R & I A FAIRBURN 1953 Cat scratch disease the intradermal test in controls *Lancet* 1953(i) 520-521
- 66 BIJLMER J & D GRANAAT 1952 Benign lymphoreticulosis (Cat scratch disease case) *Ned Tijdschr Geneesk* 96 1158-1159
- 67 BORIOLI F & F BADARACCO 1952 A propos d'un cas de maladie dite des griffes de chat (On a case of so called cat scratch disease) *Praxis* 41 263-264
- 68 BREIJER H B & R GISPEN 1953 Lymphoreticulosis benigna als kattenkrabziekte (Benign lymphoreticulosis or cat scratch disease) *Acta leiden* 23 52-58
- 69 BREIJER H B & R GISPEN 1953 Lymphoreticulosis benigna als kattenkrabziekte (Benign lymphoreticulosis as cat scratch disease) *Ned Tijdschr Geneesk* 97 1849-1853
- 70 CAT SCRATCH FEVER 1952 *Lancet* 1952(i) 302
- 71 CAT SCRATCH DISEASE 1952 *J Am Med Assoc* 148 146-147
- 72 CAT SCRATCH DISEASE 1953 *Med World* 78 280-281
- 73 CHAIA J & N TALEB 1953 A propos du premier cas libanais de lymphoréticulose bénigne d'inoculation maladie des griffes du chat (First case of cat scratch disease in Lebanon) *Sang* 24 635-637
- 74 CHARBONNEAU H 1952 Quatre cas de maladie des griffes du chat à Montréal lymphoreticulose aigue benigne d'inoculation (Four cases of the cat scratch disease in Montreal a cute benign inoculation lymphoreticulosis) *Union méd Canada* 81 123-128
- 75 CHFMIV J & J P IMBERT 1952 A propos d'une adenopathie cervicale à evolution subaigue lymphoreticulose bénigne d'inoculation (Subacute evolution of a disease of a lymph node of the neck benign inoculation lymphoreticulosis) *Rev odontol stomatol* 8 23-25
- 76 CHIARI G 1952 Su di un caso di linforeticulosis infettiva benigna malattia da unghie di gatto (A case of benign infectious lymphoreticulosis cat scratch disease) *Lattante* 23 159-176
- 77 CITTERIO L 1952 Linforeticulosis benigna per morso di cane (Benign lymphoreticulosis due to dog bite) *Arch ital chir* 76 331-346
- 78 CUTTLE T D 1952 Benign lymphoreticulosis of inoculation (cat scratch fever) *Calif Med* 76 404-406
- 79 DAESCHNER C W G W SALMON & F M HEYS 1953 Cat scratch fever *J Pediat* 43 371-384
- 80 DE JONGH J 1953 Benign virus lymphadenitis (cat scratch disease) case *Ned Tijdschr Geneesk* 97 1910-1911
- 81 DENECHAU D 1952 Un nouvel exemple de maladie de griffes de chat excision du foyer et du ganglion infectés guérison spectaculaire (A new example of cat scratch diseases excision of the focus and infected lymph node spectacular recovery) *Bull mem soc med hop Paris* 68 347-349
- 82 DEPAILLAT A & A CONDAT 1952 A propos d'un cas de thrombophlébite cérébrale au cours d'une lymphoréticulose bénigne d'inoculation (A case of cerebral thrombophlebitis in the course of benign inoculation lymphoreticulosis) *Arch franç pédiat* 9 744-745
- 83 DREOSTI A E & J F MURRAY 1953 Cat scratch disease non bacterial regional lymphadenitis or benign lympho reticulosis of inoculation *S African Med J* 27 81-83
- 84 EPSTEIN E 1952 Cat scratch fever *A M A Arch Dermatol Syphilol* 66 240-243
- 85 FERGUSON G 1952 Acute nonspecific mesenteric lymphadenitis possible mechanism of pain illustrated by two cases *A M A Arch Surg* 65 906-911

- 86 FITZPATRICK T J & L W WOODRUFF 1953 Cat scratch disease. Ill Med J 102 394-395
- 87 FLORES A 1953 Die benigne durch Inokulation übertragbare Lymphoretikulose Morbus Petzetakis oder Katzenkratzkrankheit (Benign lymphoreticulosis contagiosa by inoculation—Morbus Petzetakis or cat scratch disease) Wien klin Wochschr 64 963-964
- 88 FORBETH S 1952 Die Katzenkratzkrankheit eine neue Virus lymphadenitis? (Cat scratch disease a new viral lymphadenitis) Deut med Wochschr 77 943-945
- 89 GENNES L DE & H BRICURE 1953 Sur quatre observations d'infections traitées par l'érythromycine (Erythromycin treatment of four infections) Bull. m m soc méd hôp Paris 69 377-381
- 90 GIFFORD H M BROCKBANK & J MONZMAN 1953 Cat scratch fever Stanford Med Bull 10 157-161
- 91 GROPPI C 1953 Su un caso di linforeticulosi benigna da inoculazione malattia da griffi di gatto cat fever (Benign inoculation lymphoreticulosis cat scratch disease) Giorn mal infettive e parasit 4 308-310
- 92 CROSSKOPF A A WIMPHIE & M SELLMANN 1953 Étiologie étiologique et lymphoreticulose bénigne d'inoculation maladie des griffures du chat (Subacute encephalomyelitis and benign inoculation lymphoreticulosis cat scratch disease) Bull. m m soc méd hôp Paris 68 611-624
- 93 HAY A 1953 Benign virus lymphadenitis (Benign virus lymphadenitis) Med gla u (Beogr) 16 10-172
- 94 HEDINGER C 1952 Die histologischen Veränderungen bei der sog Katzenkratzkrankheit einer benignen Virus lymphadenitis (Histologic changes in so-called cat scratch disease benign virus lymphadenitis) Arch pathol Anat u Physiol Virchow s 322 152-154
- 95 HEDINGER C C LESTER & T WEGMANN 1952 Der kutane Primäraffekt der sogenannten Katzenkratzkrankheit einer benignen Virus lymphadenitis (Cutaneous primary affection of the so-called cat scratch disease a benign virus lymphadenitis) Dermatologica 104 101-107
- 96 HELG L 1952 Recueil de faits lymphoreticulose bénigne d'inoculation maladie des griffures du chat (Benign inoculation lymphoreticulosis cat scratch disease) Dermatologica 105 198-200
- 97 HENRY M 1952 Leptothrix conjunctivae (Parinaud's conjunctivitis) report of 13 cases relationship to cat scratch disease Trans Pacific Coast Oto-Ophthalm Soc 33 173-196
- 98 HERRELL W E A BALOWS & B F ROACH 1953 Cat scratch fever report of a case J Kentucky State Med Assoc 53 486-488
- 99 JAEGER H & J DELACRETAZ 1953 Lymphoreticulose benigne ou maladie des griffures du chat avec porte d'entrée au pénis (Benign lymphoreticulosis or cat scratch disease with primary penile lesion) Dermatologica 107 260-261
- 100 JAMES J & E EMURA 1953 Benign inoculation lymphoreticulosis (cat scratch disease) report of two cases with positive skin tests A M A Arch Dermatol Syphilol 67 439-447
- 101 KALLOFF K W 1953 Zur Abgrenzung der Lymphadenitis non follicularis benigna cutis on Hautmanifestationen der chronischen Lymphadenitis (Differentiation of lymphadenitis non follicularis der gna cutis from skin manifestation of chronic lymphadenitis) Dermatol Wochschr 126 1146-1150
- 102 KAMMEIER R H 1953 Cat scratch disease J Tenn State Med Assoc 45 441-49
- 103 KEISER D P 1953 Fatt krabziekte (cat scratch fever maladie des griffures de chat) Ned Tijdschr Geneesk 97 1958-1961
- 104 KLEYER J J 1953 Een geval van kattenkrabziekte (A case of cat scratch disease) Maatsch k vergezond 21 341-345
- 105 KRAPIN D 1952 Suppurative lymphadenitis concomitant with infectious mononucleosis N Y State J Med 52 2043
- 106 LAHAM T 1953 Cas litana s de lymphoreticulose regionale benigne maladie par griffures de chat (A case of benign regional lymphoreticulosis in Lebanon—cat scratch disease) Rev med Moson Orient 10 391-397
- 107 LANGE H L 1953 Cat-scratch fever Am J Nurs 53 837-833
- 108 LATALLA Z 1953 Choroba Dobre i Miarec choroba kociego pazura (Cat scratch disease) Pe lat pol a 27 1109-1112

- 111 PEDIATRIA 1952 Linforeticolosi benigna da inoculazione malattia da graffi di gatto (Benign inoculation lymphoreticulosis cat scratch disease) 60 385-389
- 112 IOVISATO I 1953 Linforeticolosi benigna da inoculazione (Inoculation benign lymphoreticulosis) Kass giuliana med 9 49-81
- 113 MACMURRAY I C M C SANFORD & T WINSHIP 1952 Cat scratch disease simulating sarcoma of the neck Am J Surg 84 483-485
- 114 MARCHINI I 1952 Prime osservazioni su di un caso di cosiddetta linforeticolosi benigna da inoculazione o malattia da graffi di gatto (A case of so called benign inoculation lymphoreticulosis or cat scratch disease) Minerva dermatol 27 13-17
- 115 MARCZYNSKA ROBOWSKA M 1952 Przypadek choroby kożnego pazurau dziewczynki 12 letniej (Case of cat scratch disease in a 12 year old girl) I ediat pol ka 27 859-860
- 116 MASSHOFF W 1953 Eine neuartige Form der mesenterialen Lymphadenitis (A recently discovered form of mesenteric lymphadenitis) Deut med Wochschr 78 532-535
- 117 MOLLARET I 1952 Eimen neue lymphatische Erkrankung und ein neues Virus die gutartige Impflymphoreticulosis (A new lymphatic disease and a new virus benign inoculation lymphoreticulosis) Wien klin Wochschr 64 497-500
- 118 MOLLARET P 1952 La lymphoreticulose benigne d inoculation (Benign inoculation lymphoreticulosis) Med hyg, Genève 10 307-308
- 119 MOLLARET I 1952 La lymphoreticulose benigne d inoculation (The benign inoculation lymphoreticulosis) Helv Med Acta 19 316-326 (Abstr in Biol Abstr 27 10243 1953)
- 120 MONCRIEFF A 1952 Cat scratch fever Proc Roy Soc Med 45 456
- 121 MORDANT H 1952 Un cas de lymphoreticulose benigne d inoculation ou maladie par griffure de chat (A case of cat scratch disease) Arch belges dermatol syphi 18 240-243
- 122 NINNI M 1952 La linforeticolosi benigna da inoculazione malattia dei graffi del chat (Cat scratch disease) Progr in Med Naples 8 685 688
- 123 OFFRET G 1952 Benign inoculation lymphoreticulosis so-called cat scratch disease (A propos d un cas lymphoreticulose benigne d inoculation dite maladie des griffes du chat) Bull soc ophtalmol France 537-541
- 124 PAVLANIS V 1952 Sur le virus de la lymphoreticulose benigne d inoculation (The virus of benign inoculation lymphoreticulosis) Union med 81 129-131
- 125 PAYAN H R MURATORE & H GASTALT 1952 A propos du diagnostic histologique de la lymphoreticulose benigne d inoculation maladie des griffes du chat (Histological diagnosis of benign inoculation lymphoreticulosis cat scratch disease) Toulouse med 63 925-927
- 126 PELOUX Y 1952 Un cas de lymphoreticulose benigne d inoculation au Tchad (A case of benign inoculation lymphoreticulosis in Tchad) Méd trop 12 336-338
- 127 PEPLE W I JR & F W GILL 1952 Mesenteric adenitis complicated by multiple segmental intussusception of the small intestine Virginia Med Monthly 79 673 675
- 128 QUINN F L 1953 Cat scratch disease J Michigan State Med Soc 62 71-72
- 129 RAMIOLI H 1952 Lymphoreticulose benigne d inoculation à localisation palpébrale sans adénopathie (Benign lymphoreticulosis from palpebral localization and without adenopathy) Rev med Liège 7 789-800
- 130 REGINSTER A 1953 Un nouveau cas de maladie des griffes du chat (A recent case of cat scratch disease) Rev med Liège 8 205 207
- 131 REGINSTER A & L F DUMONT 1952 Deux cas de lymphoreticulose benigne d inoculation (Two cases of benign lymphoreticulosis from inoculation) Rev med Liège 7 786-787
- 132 ROCET J & R FAU (BEALDOIN) 1952 Encéphalite clinique confirmée par l'électroencéphalographie contemporaine d'une lymphoreticulose benigne (Clinical encephalitis confirmed by electroencephalography appearing during benign lymphoreticulosis) Bull mem soc med hôp Paris 68 710-712
- 133 ROSENBLUM H 1953 Cat scratch disease cat scratch fever benign lymphoreticulosis of inoculation Am Practitioner and Dig Treatment 4 431-437
- 134 ROUX J A 1952 Un cas d'érythème noueux dans la maladie des griffes du chat (Erythema nodosum during cat scratch disease case) J dermatol 41 238-239
- 135 SCHUERMANN H & H REICH 1952 Katzenkratzerkrankheit (Maladie des griffes de chat) in Deutschland (Cat scratch disease in Germany) Hautarzt 3 225-227
- 136 SCHUERMANN H & H REICH 1952 Katzenkratzerkrankheit (Maladie des griffes de chat) in Deutschland (Cat scratch disease in Germany) Klin Wochschr 30 366-367

- 137 SCHWARTZ J R GEIGER & M MARGROFF 1953 Maladie des griffes du chat (Cat scratch disease) *Strasbourg med* 4 390-402
- 138 SIGLIER I 1952 Une traversée mouvementée *Presse m* 1 60 179
- 139 SIGLIER F J J WELTI & P LOMBROSO 1952 Pseudo venereal forms of cat scratch disease (Nouvelles observations de formes pseudovenériennes de la maladie des griffes du chat) *Bull mém soc méd hôp Paris* 68 216-219
- 140 SOARES A D & T GOLVEIA 1952 Linforeticulose benigna de inoculacao (Benign inoculation lymphoreticulosis) *Ann inst med trop* 9 145-159
- 141 SOMMER C 1953 Über die Katzenkratzkrankheit (On cat scratch disease) *Zentr Chir* 78 890-897
- 142 STARCICH R 1953 Le forme atipiche della cosiddetta malattia da graffi di gatto (Atypical forms of so called cat scratch disease) *Giorn clin med* 34 1499-1507
- 143 STIGTER W 1953 Forms of scabies in dogs or cats transmissible to humans *Ned Tijdschr Geneesk* 97 1958 1961
- 144 THEISSING G 1953 Beitrag zur Frage der benignen Impflymphoretikulose (Problem of the benign vaccination lymphoreticulosis) *Arch Ohren Nasen u Kehlkopf heilk ver Z Hals Nasen u Ohrenheilk* 163 499-505
- 145 THIERS H T 1952 Erythème noueux au cours de l'évolution d'une lymphoréticulose bénigne d'inoculation (Erythema nodosum during the evolution of benign inoculation lymphoreticulosis) *Icon méd* 187 10-11
- 146 THOMAS C J CORDIER & B ALGAN 1952 La lymphoréticulose bénigne oculoganglionnaire (Benign oculoglandular lymphoreticulosis) *Ophthalmologica* 123 129-144
- 147 TONGE J I J A INGLIS & E H DERRICK 1953 Regional non bacterial suppurative lymphadenitis and its relationship to cat scratch disease *Med J Australia* 40 81-85
- 148 TURIAT J & Y JEANJEAN 1953 Le syndrome pneumo-ganglionnaire de la pneumonie atypique de la lymphoreticulose bénigne d'inoculation maladie des griffes du chat (The pneumoganglionic syndrome of atypical pneumonia due to benign inoculation lymphoreticulosis cat scratch disease) *Bull mém soc méd hôp Paris* 69 914-970
- 149 TURIAT J & Y JEANJEAN 1954 La pneumonie atypique de la lymphoreticulose bénigne d'inoculation maladie des griffes du chat (Atypical pneumonia due to cat scratch disease) *J française de méd et chir thorac* 8 141-147
- 150 TZANCK A G R MELLI & R TEMIME 1957 Un cas de maladie de la griffe du chat (Case of cat scratch disease) *Bull soc franç dermatol syphilis* 59 216-217
- 151 USTERI C T WEGMANN & C HEDINGER 1952 Über atypische Formen der sog Katzenkratzkrankheit einer benignen Virus-Lymphadenitis (Atypical forms of the so-called cat scratch disease a benign virus lymphadenitis) *Schweiz med Wochschr* 82 1287-1290
- 152 VEELAN A W VAN & P D STIBBE 1953 Conjunctivitis van Parinaud en kattekrabziekte (Parinaud's conjunctivitis and cat scratch disease) *Ned Tijdschr Geneesk* 97 1203-1205
- 153 VIVELL O 1952 Die Viruskratzlymphadenitis Gutartige Impflymphoretikulosis (Viral scratch lymphadenitis benign inoculation lymphoreticulosis) *Deut med Wochschr* 77 945-946
- 154 WATERS W J S S KALTER & J T PRIOR 1952 Cat scratch syndrome *Pediatrics* 10 311-318
- 155 WATRIN J J BELREY R LEDUC & A MFRAND 1952 Lymphoreticulose bénigne d'inoculation forme génitale (Benign lymphoreticulosis of inoculation genital form) *Bull soc franç dermatol syphilis* 59 399-400
- 156 WATRIN J J BEUREY R LEDUC & A MERAND 1957 Lymphoréticulose bénigne d'inoculation forme génitale (Benign inoculation lymphoreticulosis genital form) *Bull soc franç dermatol syphilis* 59 431-432
- 157 WEILL J & F BLAS 1951 Erythème noueux et griffure du chat (Erythema nodosum and cat scratch disease) *Bull m m soc m d hôp Paris* 67 1141-1144
- 158 WIJERS P J 1953 Een geval van kattkrabziekte (Case of cat scratch disease) *Ned Tijdschr Geneesk* 97 2085-2086
- 159 WINSHIP T 1953 Cat scratch disease in children a report of five cases *Clin Proc Child Hosp* 9 244-247
- 160 BERARDINELLI W 1954 Galeomucose de enca de Delré Mollaret Reilly Foshay (Cat scratch disease) *Gaz med port* 7 20-22
- 161 BONNET I 1956 Conjunctivite folliculaire géante unilatérale avec adénopathie prétragénne et cervicale conjonctivite de Parinaud inoculée par le chat (Unilateral giant follicular conjunctivitis with adenopathy pretragenic and cervical conjunctivitis of Parinaud inoculated by the cat)

- 111 PEDIATRIA 1952 Linforeticulosis benigna da inoculazione malattia da graffi di gatto (Benign inoculation lymphoreticulosis cat scratch disease) 60 395-389
- 112 LOVISATO L 1953 Linforeticulosis benigna da inoculazione (Inoculation benign lymphoreticulosis) I ass giuliana med 9 19-81
- 113 MACMURRAY I G M C SAINFORD & T WINSHIP 1952 Cat scratch disease simulating sarcoma of the neck Am J Surg 84 483-485
- 114 MARCHINI E 1952 Prime osservazioni su di un caso di cosiddetta linforeticulosis benigna da inoculazione o malattia da graffi di gatto (A case of so-called benign inoculation lymphoreticulosis or cat scratch disease) Minerva dermatol 27 13-17
- 115 MARCZYNSKA ROBOWSKA M 1952 Przypadek choroby kociego pazurazu dziewczynki 12 letniej (Case of cat scratch disease in a 12 year old girl) Pediatr polska 27 859-860
- 116 MASSHOFF W 1953 Eine neuartige Form der mesenterialen Lymphadenitis (A recently discovered form of mesenteric lymphadenitis) Deut med Wochschr 78 532-535
- 117 MOLLARET I 1952 Einen neue lymphatische Erkrankung und ein neues Virus die gutartige Impflymphoreticulosis (A new lymphatic disease and a new virus benign inoculation lymphoreticulosis) Wien klin Wochschr 64 497-500
- 118 MOLLARET P 1952 La lymphoreticulose benigne d inoculation (Benign inoculation lymphoreticulosis) Méd hyg Genève 10 307-308
- 119 MOLLARET I 1952 La lymphoreticulose benigne d inoculation (The benign inoculation lymphoreticulosis) Helv Med Acta 19 316-326 (Abstr in Biol Abstr 27 10243 1953)
- 120 MONCRIEFF A 1952 Cat scratch fever Proc Roy Soc Med 45 456
- 121 MORDANT H 1952 Un cas de lymphoréticulose benigne d inoculation ou maladie par griffure de chat (A case of cat scratch disease) Arch belges dermatol syphig 8 240-243
- 122 NINNI M 1957 La linforeticulosis benigna da inoculazione malattia dei griffe del chat (Cat scratch disease) Progr in Med Naples 8 685-688
- 123 OFFRET G 1952 Benign inoculation lymphoreticulosis so-called cat scratch disease (A propos d un cas lymphoreticulose benigne d inoculation dite maladie des griffes du chat) Bull soc ophtalmol France 537-541
- 124 PAVLANIS V 1952 Sur le virus de la lymphoreticulose benigne d inoculation (The virus of benign inoculation lymphoreticulosis) Union m d 81 129-131
- 125 PAYAN H R MURATORE & H GASTALT 1957 A propos du diagnostic histologique de la lymphoreticulose benigne d inoculation maladie des griffes du chat (Histological diagnosis of benign inoculation lymphoreticulosis cat scratch disease) Toulouse med 53 975-977
- 126 PELOUX Y 1952 Un cas de lymphoreticulose benigne d inoculation au Tchad (A case of benign inoculation lymphoreticulosis in Tchad) M d trop 12 336-338
- 127 PEPLE W L JR & F W GILL 1952 Mesenteric adenitis complicated by multiple segmental intussusception of the small intestine Virginia Med Monthly 79 673-675
- 128 QUINN E L 1953 Cat scratch disease J Michigan State Med Soc 52 71-72
- 129 RAMIOLL H 1952 Lymphoreticulose benigne d inoculation a localisation palpébrale sans adenopathie (Benign lymphoreticulosis is from palpebral localization and without adenopathy) Rev med Liege 7 789-800
- 130 REGINSTER A 1953 Un nouveau cas de maladie des griffes du chat (A recent case of cat scratch disease) Rev m d Liège 8 205-207
- 131 REGINSTER A & L E DUMONT 1952 Deux cas de lymphoreticulose benigne d inoculation (Two cases of benign lymphoreticulosis from inoculation) Rev med Liege 7 766-787
- 132 ROGET J & R FAU (BEALDOIN) 1952 Encephalite clinique confirmée par l'électroencephalographie contemporaine d'une lymphoréticulose benigne (Clinical encephalitis confirmed by electroencephalography appearing during benign lymphoreticulosis) Bull mem soc med h p Paris 68 710-712
- 133 ROSENBLUM H 1953 Cat scratch disease cat scratch fever benign lymphoreticulosis of inoculation Am Practitioner and Dig Treatment 4 431-437
- 134 ROUX J A 1952 Un cas d'érythème noueux dans la maladie des griffes du chat (Erythema nodosum during cat scratch disease case) Pediatre 41 238-239
- 135 SCHUERMAN H & H REICH 1952 Katzenkratzerkrankheit (Maladie des griffes de chat) in Deutschland (Cat scratch disease in Germany) Hautarzt 3 775-727
- 136 SCHUERMAN H & H REICH 1952 Katzenkratzerkrankheit (Maladie des griffes de chat) in Deutschland (Cat scratch disease in Germany) Klin Wochschr 30 366-367

- 137 SCHWARTZ J & GEIGER & M MARGROFF 1953 Maladie des griffes du chat (Cat scratch disease) *Strasbourg med* 4 399-402
- 138 SICLIER F 1952 Une traversée mouvementée *Presse m* 1 60 179
- 139 SIGLER J J J WELT & P LOMBRIO 1952 Pseudo venereal forms of cat scratch disease (Nouvelles observations de formes pseudovenériennes de la maladie les griffes du chat) *Bull mem soc m'd hôp Paris* 68 216-219
- 140 SOARES A D & E COLAIEA 1952 Inforeticulose benigna de inoculacao (Benign inoculation lymphoreticulosis) *Ann inst mei trop* 9 145-159
- 141 SOMMER C 1953 Über die Katzenkratzkrankheit (On cat scratch disease) *Zentr Chir* 78 890-891
- 142 STARCHI R 1953 Le forme atipiche della cosiddetta malattia da graffi di gatto (Atypical forms of so-called cat scratch disease) *Giorn clin mei* 34 1499-1507
- 143 STIGTER W 1953 Forms of scabies in dogs or cats transmissible to humans *Ned Tijdschr Geneesk* 97 1958 1961
- 144 THIELING G 1953 Beitrag zur Frage der benignen Impflymphoretikulose (Problem of the benign vaccinal lymphoreticulosis) *Arch Ohren Nasen u Kehlkopf heilk ver Z Hals Nasen u Ohrenheilk* 163 499-505
- 145 THIERS H T 1952 Erythème noueux au cours de l'évolution d'une lymphoréticulose bénigne d'inoculation (Erythema nodosum during the evolution of benign inoculation lymphoreticulosis) *Lyon mei* 187 10-11
- 146 THOMAS C J CORDIER & B ALGAN 1952 La lymphoréticulose bénigne oculoganglionnaire (Benign oculoglandular lymphoreticulosis) *Ophthalmologica* 123 179-184
- 147 TONGE J I J A INGLIS & F H DERRICK 1953 Regional non bacterial suppurative lymphadenitis and its relationship to cat scratch disease *Med J Australia* 40 81 85
- 148 TURIAU J & Y JEANJEAN 1953 Le syndrome pneumo-ganglionnaire de la pneumonie atypique de la lymphoréticulose bénigne d'inoculation maladie des griffes du chat (The pneumoganglionic syndrome of atypical pneumonia due to benign inoculation lymphoreticulosis cat scratch disease) *Bull mem soc m'd hôp Paris* 69 914-920
- 149 TURIAU J & Y JEANJEAN 1954 La pneumonie atypique de la lymphoréticulose bénigne d'inoculation maladie des griffes du chat (Atypical pneumonia due to cat scratch disease) *J franç mei et chir tho ac* 8 141 147
- 150 TZANCA A G R MELLI & R TEMME 1952 Un cas de maladie de la griffe du chat (Case of cat scratch disease) *Bull soc franç dermatol syphilig* 59 216-217
- 151 USTERI C T WEGMANN & C HEDINGER 1952 Über atypische Formen der sog Katzenkratzkrankheit einer benignen Virus-Lymphadenitis (Atypical forms of the so-called cat scratch disease a benign virus lymphadenitis) *Schweiz med Wochschr* 82 1281 1290
- 152 VEELEN A W VAN & P D STIBBE 1953 Conjunctivitis van Parinaud en kattekrabziekte (Parinaud's conjunctivitis and cat scratch disease) *Ned Tijdschr Geneesk* 97 1203-1205
- 153 VINELL O 1952 De Virus-Kratzlymphadenitis Cutartige Impflymphoretikulosis (Viral scratch lymphadenitis benign inoculation lymphoreticulosis) *Deut med Wochschr* 77 945-946
- 154 WATERS W J S S KALTER & J T IRIOR 1952 Cat scratch syndrome *Pediatrics* 10 311 318
- 155 WATRIN J J BELREY R LEDUC & A MERAND 1952 Lymphoreticulose bénigne d'inoculation forme génitale (Benign lymphoreticulosis of inoculation genital form) *Bull soc franç dermatol syphilig* 59 399-400
- 156 WATRIN J J BEUREY R LEDUC & A MERAND 1952 Lymphoréticulose bénigne d'inoculation forme génitale (Benign inoculation lymphoreticulosis genital form) *Bull soc franç dermatol syphilig* 59 431-432
- 157 WEILL J & F BLAS 1951 Erythème noueux et griffure du chat (Erythema nodosum and cat scratch disease) *Bull m'm soc m'd hôp Paris* 67 1141 1144
- 158 WIJERS J J 1953 Een geval van kattekrabziekte (Case of cat scratch disease) *Ned Tijdschr Geneesk* 97 2085 2086
- 159 WINSHIP T 1953 Cat scratch disease in children a report of five cases *Clin Proc Child Hosp* 9 244 247
- 160 BERARDINELLI W 1954 Galeoncosi della de Del ré Molla et Reilly Foshay (Cat scratch disease) *Gaz med port* 7 20-29
- 161 BONNET I 1956 Cojctivite folliculaire géante unilatérale avec adénopathie prétragienne et cervicale conjonctivite de Parinaud inoculation par le chat (Uni

- 111 PEDIATRIA 1952 Linforeticulosis benigna da inoculazione malattia de graffi di gatto (Benign inoculation lymphoreticulosis cat scratch disease) 60 345-389
- 112 LOVISATO L 1953 Linforeticulosis benigna de inoculazione (Inoculation benign lymphoreticulosis) Rass giuliana med 9 79-81
- 113 MACMURRAY F G M C SANFORD & T WINSHIP 1952 Cat scratch disease simulating sarcoma of the neck. Am J Surg 84 483-485
- 114 MARCHINI E 1957 Prime osservazioni su di un caso di cosiddetta linforeticulosis benigna da inoculazione o malattia da graffi di gatto (A case of so called benign inoculation lymphoreticulosis or cat scratch disease) Minerva dermatol 27 13-17
- 115 MARCZYNSKA ROBOWSKA M 1952 Przypadek choroby kociego pazurau dziewczynki 12 letniej (Case of cat scratch disease in a 12 year old girl) Pediat polska 27 859-860
- 116 MASSHOFF W 1953 Eine neuartige Form der mesenterialen Lymphadenitis (A recently discovered form of mesenteric lymphadenitis) Deut med Wochschr 78 532-533
- 117 MOLLARET P 1952 Einen neue lymphatische Erkrankung und ein neues Virus die gutartige Impflymphoreticulosis (A new lymphatic disease and a new virus benign inoculation lymphoreticulosis) Wien klin Wochschr 64 497-500
- 118 MOLLARET P 1952 La lymphoreticulose benigne d inoculation (Benign inoculation lymphoreticulosis) Med hyg Genève 10 307-308
- 119 MOLLARET P 1952 La lymphoreticulose benigne d inoculation (The benign inoculation lymphoreticulosis) Helv Med Acta 19 316-326 (Abstr in Biol Abstr 27 10243 1953)
- 120 MONCRIEFF A 1952 Cat scratch fever Proc Roy Soc Med 45 456
- 121 MORDANT H 1952 Un cas de lymphoreticulose benigne d inoculation ou maladie par griffure de chat (A case of cat scratch disease) Arch belges dermatol vph 118 240-243
- 122 NINNI M 1952 La linforeticulosis benigna da inoculazione maladie des griffes du chat (Cat scratch disease) Progr in Med Naples 8 685-688
- 123 OFFRET G 1952 Benign inoculation lymphoreticulosis so-called cat scratch disease (A propos d'un cas lymphoreticulose benigne d inoculation dite maladie des griffes du chat) Bull soc ophtalmol France 537-541
- 124 PAVILANIS V 1957 Sur le virus de la lymphoreticulose benigne d inoculation (The virus of benign inoculation lymphoreticulosis) Union med 81 129-131
- 125 PAYAN H R MURATORE & H GASTAUT 1952 A propos du diagnostic histologique de la lymphoreticulose benigne d inoculation maladie des griffes du chat (Histological diagnosis of benign inoculation lymphoreticulosis cat scratch disease) Toulouse med 53 925-927
- 126 PELOU Y 1952 Un cas de lymphoreticulose benigne d inoculation au Tchad (A case of benign inoculation lymphoreticulosis in Tchad) Med trop 12 336-338
- 127 PEPLE W L JR & F W GILL 1952 Mesenteric adenitis complicated by multiple segmental intussusception of the small intestine Virginia Med Monthly 79 673-675
- 128 QUINN E L 1953 Cat scratch disease J Michigan State Med Soc 52 71-72
- 129 RAMIOUL H 1952 Lymphoreticulose benigne d inoculation a localisation palpebrale sans adenopathie (Benign lymphoreticulosis from palpebral localization and without adenopathy) Rev med liege 7 789-800
- 130 REGINSTER A 1953 Un nouveau cas de maladie des griffes du chat (A recent case of cat scratch disease) Rev med Liege 8 205-207
- 131 REGINSTER A & L E DUMONT 1957 Deux cas de lymphoreticulose benigne d inoculation (Two cases of benign lymphoreticulosis from inoculation) Rev med Liege 7 786-787
- 132 ROGET J & R FAU (BEAUDOIN) 1957 Encephalite clinique confirmee par l'electroencephalographie contemporaine d'une lymphoreticulose benigne (Clinical encephalitis confirmed by electroencephalography appearing during benign lymphoreticulosis) Bull mem soc med hyp Paris 68 710-712
- 133 ROSENBLUM H 1953 Cat scratch disease cat scratch fever benign lymphoreticulosis of inoculation Am Practitioner and Dig Treatment 4 431-437
- 134 ROUX J A 1952 Un cas d'erytheme noueux dans la maladie des griffes du chat (Erythema nodosum during cat scratch disease case) Pediatrie 41 238-239
- 135 SCHUERMANN H & H REICH 1957 Katzenkratzkrankheit (Maladie des griffes de chat) in Deutschland (Cat scratch disease in Germany) Hautarzt 3 225-227
- 136 SCHUERMANN H & H REICH 1952 Katzenkratzkrankheit (Maladie des griffes de chat) in Deutschland (Cat scratch disease in Germany) Klin Wochschr 30 366-367

- 137 SCHWARTZ J R GEFER & M MARCROFF 1953 Maladie des griffes du chat (Cat scratch disease) *Strasbourg méd* 4 399-407
- 138 SIGUIER F 1952 Une traversée mouvementée *Presse méd* 60 179
- 139 SIGUIER F J J WELT & P LOMBROSO 1952 I seu lo-veneraal forms of cat scratch disease (Nouvelles observations de formes pseudovenériennes de la maladie des griffes du chat) *Bull mem soc méd hôp Paris* 58 216-219
- 140 SOARES A D & T COLVEIA 1952 Inforeticulose benigna de inoculacao (Benign inoculation lymphoreticulosis) *An in t med trop* 9 145-159
- 141 SOMMER G 1953 Über die Katz kratzkrankheit (On cat scratch disease) *Zentr Chir* 78 840-897
- 142 STARCHI R 1953 Le forme atipiche della cosiddetta malattia da graffi di gatto (Atypical forms of so-called cat scratch disease) *Giorn clin med* 34 1499-1507
- 143 STEIGER W 1953 Forms of scabies in dogs or cats transmissible to humans *Ned Tijdschr Geneesk* 97 1958 1961
- 144 THIFFESSING C 1953 Beitrag zur Frage der benignen lymphoretikulose (Problem of the benign vaccinator lymphoreticulosis) *Arch Ohren Nasen u Kehlkopf heilk* 7 Hals Nasen u Ohrenheilk 163 499-505
- 145 THIERS H T 1952 Erythème noueux au cours de l'évolution d'une lymphoreticulose bénigne d'inoculation (Erythema nodosum during the evolution of benign inoculation lymphoreticulosis) *Lyon méd* 187 10-11
- 146 THOMAS C J CORDIER & B ALGAN 1952 La lymphoreticulose bénigne oculoganglionnaire (Benign oculoglandular lymphoreticulosis) *Ophthalmologica* 123 129-144
- 147 TONGE J I J A INGLIS & E H DERRICK 1953 Regional non-lactating suppurative lymphadenitis and its relationship to cat scratch disease *Med J Australia* 40 81-85
- 148 TURIAP J & Y JEANJEAN 1953 Le syndrome pneumo-ganglionnaire de la pneumonie atypique de la lymphoreticulose bénigne d'inoculation maladie des griffes du chat (The pneumoganglionic syndrome of atypical pneumonia due to benign inoculation lymphoreticulosis cat scratch disease) *Bull mem soc méd hôp Paris* 69 914-970
- 149 TURIAP J & Y JEANJEAN 1954 La pneumonie atypique de la lymphoreticulose bénigne d'inoculation maladie des griffes du chat (Atypical pneumonia due to cat scratch disease) *J franç méd et chir thorac* 8 141-147
- 150 TZANCK A G R MEIKI & R TEMIME 1952 Un cas de maladie de la griffe du chat (Case of cat scratch disease) *Bull soc franç dermatol syphilis* 59 216-217
- 151 USTERI C T WEFMANN & C HEDINGER 1952 Über atypische Formen der sog Katzenkratzenkrankheit einer benignen Virus Lymphadenitis (Atypical forms of the so-called cat scratch disease a benign virus lymphadenitis) *Schweiz med Wochschr* 82 1287-1290
- 152 VEELEN A W VAN & I D STIBBE 1953 Conjunctivitis van Parinaud en kattekrabziekte (Parinaud's conjunctivitis and cat scratch disease) *Ned Tijdschr Geneesk* 97 1203-1205
- 153 VIVELL O 1952 Die Virus Kratzlymphadenitis Gutartige Impflymphoretikulosis (Viral scratch lymphadenitis benign inoculation lymphoreticulosis) *Deut med Wochschr* 77 945-946
- 154 WATERS W J S S KALTER & J T PRIOR 1952 Cat scratch syndrome *I ed atnes* 10 311-318
- 155 WATRIN J J BELREY R LEDUC & A MFRAND 1957 Lymphoreticulose bénigne d'inoculation forme génitale (Benign lymphoreticulosis of inoculation genital form) *Bull soc franç dermatol syphilis* 59 399-400
- 156 WATRIN J J BELREY R LEDUC & A MERAND 1952 Lymphoreticulose bénigne d'inoculation forme génitale (Benign inoculation lymphoreticulosis genital form) *Bull soc franç dermatol syphilis* 59 431-432
- 157 WEILL J & F BLAS 1951 Erythème noueux et griffure du chat (Erythema nodosum and cat scratch disease) *Bull mem soc méd hôp Paris* 67 1141-1144
- 158 WIJERS P J 1953 Een geval van kattekrabziekte (Case of cat scratch disease) *Ned Tijdschr Geneesk* 97 2085-2086
- 159 WINSHIP T 1953 Cat scratch disease in children a report of five cases *Clin Proc Child Hosp* 9 244-247
- 160 BERARDINELLI W 1954 Clinica del cat scratch (Cat scratch disease) *Gaz med port* 7 20-29
- 161 BONNET I 1956 Clinique folliculaire générale unilatérale avec adenopathie prétragienne et cervicale conjonctive de Parinaud inoculation par le chat (Unilateral follicular general adenopathy pretragial and cervical conjunctive of Parinaud inoculation by the cat) (Un

- 111 PEDIATRIA 1952 Linforeticolosi benigna da inoculazione malattia da graffi di gatto (Benign inoculation lymphoreticulosis cat scratch disease) 60 385-389
- 112 IOVISATO I 1953 Linforeticolosi benigna da inoculazione (Inoculation benign lymphoreticulosis) Rass giuliana med 9 49-81
- 113 MACMURRAY T G M C SANFORD & T WINSHIP 1952 Cat scratch disease simulating sarcoma of the neck Am J Surg 84 483-485
- 114 MARCHINI I 1952 Prime osservazioni su di un caso di cosiddetta linforeticolosi benigna da inoculazione o malattia da graffi di gatto (A case of so-called benign inoculation lymphoreticulosis or cat scratch disease) Minerva dermatol 27 13-17
- 115 MARCZYŃSKA ROBOWSKA M 1952 Przypadek choroby kociego pazurau dziewczynki 12 letnicy (Case of cat scratch disease in a 12 year old girl) Ieiat polska 27 859-860
- 116 MASSHOFF W 1953 Eine neuartige Form der mesenterialen Lymphadenitis (A recently discovered form of mesenteric lymphadenitis) Deut med Wochschr 78 532-535
- 117 MOLLARET P 1952 Eimen neue lymphatische Erkrankung und ein neues Virus die gutartige Impflymphoreticulosis (A new lymphatic disease and a new virus benign inoculation lymphoreticulosis) Wien klin Wochschr 64 497-500
- 118 MOLLARET I 1952 La lymphoreticulose benigne d inoculation (Benign inoculation lymphoreticulosis) Med hyg Geneve 10 307-308
- 119 MOLLARET P 1952 La lymphoreticulose benigne d inoculation (The benign inoculation lymphoreticulosis) Helv Med Acta 19 316-326 (Abstr in Biol Abstr 27 10243 1953)
- 120 MONCRIEFF A 1952 Cat scratch fever Proc Roy Soc Med 45 456
- 121 MORDANT H 1952 Un cas de lymphoreticulose benigne d inoculation ou maladie par griffure de chat (A case of cat scratch disease) Arch belges dermatol syph 8 240-243
- 122 NINNI M 1952 La linforeticolosi benigna da inoculazione malattia dei griffi del chat (Cat scratch disease) Progr in Med Naples 8 685 688
- 123 OFFRET G 1952 Benign inoculation lymphoreticulosis so-called cat scratch disease (A propos d un cas lymphoreticulose benigne d inoculation dite maladie des griffes du chat) Bull soc ophtalmol France 537-541
- 124 PAVILANIS V 1952 Sur le virus de la lymphoreticulose benigne d inoculation (The virus of benign inoculation lymphoreticulosis) Union med 81 129-131
- 125 PAYAN H R MURATORE & H GASTAUT 1952 A propos du diagnostic histologique de la lymphoreticulose benigne d inoculation maladie des griffes du chat (Histological diagnosis of benign inoculation lymphoreticulosis cat scratch disease) Toulouse med 53 975-977
- 126 PELOUX Y 1952 Un cas de lymphoreticulose benigne d inoculation au Tchad (A case of benign inoculation lymphoreticulosis in Tchad) Méd trop 12 336-338
- 127 PEILE W L JR & F W GILL 1952 Mesenteric adenitis complicated by multiple segmental intussusception of the small intestine Virginia Med Monthly 79 673-675
- 128 QUINN E L 1953 Cat scratch disease J Michigan State Med Soc 52 71-72
- 129 RAMOLL H 1952 Lymphoreticulose benigne d inoculation à localisation palpébrale sans adénopathie (Benign lymphoreticulosis from palpebral localization and without adenopathy) Rev med Liege 7 789-800
- 130 PEGINSTER A 1953 Un nouveau cas de maladie des griffes du chat (A recent case of cat scratch disease) Rev méd Liege 8 205-207
- 131 PEGINSTER A & L E DUMONT 1952 Deux cas de lymphoreticulose benigne d inoculation (Two cases of benign lymphoreticulosis from inoculation) Rev m d Liege 7 786-787
- 132 ROGET J & R FAU (BEAUDOIN) 1952 Encéphalite clinique confirmée par l'électroencéphalographie contemporaine d'une lymphoreticulose benigne (Clinical encephalitis confirmed by electroencephalography appearing during benign lymphoreticulosis) Bull mém soc med hop Paris 68 710-712
- 133 ROSENBLUM H 1953 Cat scratch disease cat scratch fever benign lymphoreticulosis of inoculation Am Practitioner and Dig Treatment 4 431-437
- 134 ROUX J A 1952 Un cas d'erythème noueux dans la maladie des griffes du chat (Erythema nodosum during cat scratch disease case) Pédiatrie 41 238 239
- 135 SCHLIERMANN H & H REICH 1952 Katzenkratzkrankheit (Maladie des griffes de chat) in Deutschland (Cat scratch disease in Germany) Hautarzt 3 225-227
- 136 SCHLIERMANN H & H REICH 1952 Katzenkratzkrankheit (Maladie des griffes de chat) in Deutschland (Cat scratch disease in Germany) Klin Wochschr 30 366-367

- 13 SCHWARTZ J & GEIGER & M MARGROFF 1953 Maladie des griffes du chat (Cat scratch disease) Strassbourg med 4 399-407
- 138 SIGUIER F 1952 Une traversée mouvementée Ires-c-m 1 60 1-9
- 139 SIGUIER F J J WELTI & P LOMBROSO 1957 I seculo-venereali forms of cat scratch disease (Nouvelles observations de formes pseudovenériennes de la maladie des griffes du chat) Bull m m soc m d h p Paris 68 216-219
- 140 SOARES A D & F COUVEIA 1952 Linforeticulose benigna de inoculacao (Benign inoculation lymphoreticulosis) Ann inst med trop 9 145-155
- 141 SOMMER C 1953 Über die Katzenkratzkrankheit (On cat scratch disease) Zentr Chir 78 890-897
- 142 STARCICH R 1953 Le forme atipiche della cosiddetta malattia da griffa di gatto (Atypical forms of so-called cat scratch disease) Giorn clin med 34 1499-1507
- 143 STIGTER W 1953 Forms of scabies in dogs or cats transmissible to humans Ned Tijdschr Geneesk 97 1958 1961
- 144 THEISSING C 1953 Beitrag zur Frage der benignen lymphoreticulose (Problem of the benign vaccinal lymphoreticulosis) Arch Ohren Nasen u Kehlkopf heilk ver Z Hals Nasen u Ohrenheilk 163 49 505
- 145 THIERS H T 1952 Erythème nouveau au cours de l'évolution d'une lymphoreticulose benigne d'inoculation (Erythema nodosum during the evolution of benign inoculation lymphoreticulosis) Lyon m d 187 10-11
- 146 THOMAS C J CORDIER & B ALGAN 1952 La lymphoreticulose benigne oculoganglionnaire (Benign oculoglandular lymphoreticulosis) Ophthalmolog ca 123 139-144
- 147 TONGE J I J A INGLIS & F H DERRICK 1953 Regional non bacterial suppurative lymphadenitis and its relationship to cat scratch disease Med J Australia 40 81 85
- 148 TURIAT J & A JEANJEAN 1953 Le syndrome pneumoganglionnaire de la pneumonie atypique de la lymphoreticulose benigne d'inoculation maladie des griffes du chat (The pneumoganglionic syndrome of atypical pneumonia due to benign inoculation lymphoreticulosis cat scratch disease) Bull m m soc m d h p Paris 69 914-920
- 149 TURIAT J & A JEANJEAN 1954 La pneumonie atypique de la lymphoreticulose benigne d'inoculation maladie des griffes du chat (Atypical pneumonia due to cat scratch disease) J franç m l et chir tho ac 8 141 147
- 150 TZANCK A G R MELI & R TEMME 1952 Un cas de maladie de la griffe du chat (Case of cat scratch disease) Bull soc franç dermatol syphil 69 216-217
- 151 USTERI C T WEGMANN & C HEDINGER 1952 Über atypische Formen der sog Katzenkratzkrankheit einer benignen Virus Lymphadenitis (Atypical forms of the so-called cat scratch disease a benign virus lymphadenitis) Schweiz med Wochschr 82 1287-1290
- 152 VEELAN A W VAN & P D STIBBE 1953 Conjunctivitis an Parinaud en kattekrabzkte (Parinaud's conjunctivitis and cat scratch disease) Ned Tijdschr Geneesk 97 1203-1205
- 153 VIVELL O 1952 Die Virus-Kratzlymphadenitis Gutartige Impflymphoretikulosis (Viral scratch lymphadenitis benign inoculation lymphoreticulosis) Deut med Wochschr 77 945-946
- 154 WATERS W J S S KALTER & J T PRIOR 1952 Cat scratch syndrome Pediatrics 10 311 318
- 155 WATRIN J J BELREY R LEDUC & A MERAND 1952 Lymphoreticulose benigne d'inoculation forme genitale (Benign lymphoreticulosis of inoculation genital form) Bull soc franç dermatol syphil 69 399-400
- 156 WATRIN J J BELREY R LEDUC & A MERAND 1952 Lymphoreticulose benigne d'inoculation forme genitale (Benign inoculation lymphoreticulosis genital form) Bull soc franç dermatol syphil 69 431-432
- 157 WEILL J & F BLAS 1951 Erythème nouveau et griffure du chat (Erythema nodosum and cat scratch disease) Bull m m soc med hôp Paris 67 1141-1144
- 158 WIJERS P J 1953 Een geval van kattekratz ziekte (Case of cat scratch disease) Ned Tijdschr Geneesk 97 2085-2086
- 159 WINSTE T 1953 Cat scratch disease in children a report of five cases Clin Proc Child Hosp 9 244-247
- 160 BERARDINELLI W 1954 Caloicose do ca de D. Ir. M. Haret Reilly Foshay (Cat scratch disease) Gaz med port 7 20-29
- 161 BONNEY I 1956 Conjunctivite folliculaire gauche unilatérale avec adénopathie prétragienne et cervicale conjonctivite de Parinaud inoculation par le chat (Uni-

- lateral giant follicular conjunctivitis with pretragal and cervical adenopathy (Parinaud's conjunctivitis inoculation by the cat) Bull soc ophtalmol France 2 325-326
- 162 CASABAN D 1956 Keratoderme localisée des doigts consécutive à une atteinte de lymphoréticulose bénigne d'inoculation (Keratoderma localized in the fingers subsequent to an attack of benign inoculation lymphoreticulosis) Bull soc franç dermatol syphilig 1 115-116
- 163 CAT SCRATCH FEVER 1956 Med J Australia 1 802
- 164 CAT SCRATCH DISEASE 1956 Diagnostic procedures for virus and rickettsial diseases 470-471 Am Public Health Assoc New York N Y
- 165 CHAIA J & N TALEB 1952 Lymphoréticulose bénigne d'inoculation ou maladie des griffes du chat (Benign inoculation lymphoreticulosis or cat scratch disease) J méd libanais 5 268-277 (Biol Abstr 28 11122 1954)
- 166 CONNOLLY V P 1955 Cat scratch disease Can Service Med J 11 10-21
- 167 CREYX I I J DAVID CHASSE & R CEINDRE 1955 Reticulose bénigne d'inoculation par piqûres d'arêtes de morue à début cutané du type érythème polymorphe (Benign inoculation reticulosis caused by codfish bone scratches with polymorphous erythema) Bull soc franç dermatol syphilig 4 433-434
- 168 CREYX L L J DAVID CHASSE A SERRES & R CEINDRE 1955 Lymphoréticulose bénigne d'inoculation par piqûre d'arête de morue maladie professionnelle (Benign inoculation lymphoreticulosis from codfish bone puncture occupational disease) J méd Bordeaux et Sud Ouest 132 420-421
- 169 DEBRÉ R 1954 Die Katzenkratzkrankheit (Cat scratch disease) Med Klin Berlin 49 945-950
- 170 DEBRÉ R 1954 La maladie des griffes de chat (Cat scratch disease) Biliotheca Paediat Suppl Ann Paediat No 58 465-475
- 171 DEBRÉ R & J JOB 1954 Cat scratch fever Acta paediat 43 1-86 (Abstr in AM J Disca Children 89 107 1955)
- 172 DE SAN MARTIN A M & R J CALIGARI 1956 La enfermedad por rascuno de gato (maladie de la griffe du chat or cat scratch fever) Dia med 28 69-71
- 173 DEVITO J J 1954 Cat scratch disease J Florida Med Assoc 40 638-640
- 174 FAIVRE G P LAMY G J AUBER & A LARCAN 1956 Lymphoréticulose bénigne d'inoculation avec méningisme splénomégalie et évolution récurrente (Benign inoculation lymphoreticulosis with meningism splenomegaly and recurrence) Rev méd Nancy 81 11-15
- 175 FOX R A 1952 So-called cat scratch fever AM J Arch Pathol 54 75-83
- 176 FRANK D E & H I HARDER 1954 Cat scratch disease in the Los Angeles area Calif Med 80 48-49
- 177 FRICK P G 1954 Cat scratch disease associated with encephalitis and herpes zoster Minn Med 37 815-817
- 178 GIFFORD H 1955 Skin test reactions to cat scratch disease among veterinarians AM J Arch Internal Med 95 828-833
- 179 GIFFORD H 1955 Cat scratch disease Pediat Clin N Am 33-40
- 180 GOODMAN D H 1955 Cat scratch disease Ariz Med 12 197-194
- 181 GROFF S 1954 Tod an Katzenkratzkrankheit Felineose (Fatal scratch disease felineosis) Monatsschr Kinderheilk 102 232-237
- 182 GUBNITSKY A & J M WAGHELSTEIN 1955 Cat scratch disease manifested by generalized lymphadenopathy Case report Maryland Med J 4 407-408
- 183 GUTTMAN P H 1955 Pathology of cat scratch disease Calif Med 82 25-31
- 184 HOFSTETTER J & J P GIRARD 1956 Deux cas atypiques de lymphoréticulose bénigne d'inoculation (LBI) chez des conjoints (Two atypical cases of benign inoculation lymphoreticulosis in husband and wife) Praxis 45 265-267
- 185 HORING F O & T ZWISLER 1954 Über eine kleine Epidemie von Katzenkratzkrankheit (Minor epidemic of cat scratch disease—lymphadenitis caused by virus) Verhandl deut Ges inn Med 60 624-630
- 186 HRAZDRA C L 1955 Nemoc kociciko skralnuti s encefalitickow komplikaci (Cat scratch disease with encephalic complication) Vnitri Lkafst 1 81-86
- 187 HUNNICUTT T N JR & C F EVANS JR 1954 Cat scratch disease—a regional lymphadenitis of non bacterial etiology Virginia Med Monthly 81 538-540
- 188 IRELAND P E & T J MOLONY 1956 Unusual virus diseases in otolaryngology Ann Otol Rhinol & Laryngol 65 442-443
- 189 KAMPMEIER R H & O A COUCH JR 1954 Cat scratch disease J Tenn Med Assoc 47 246-83
- 190 KEIL C 1956 Maladie des griffes du chat (Cat scratch disease) Rev méd Liège 11 3-4

- 191 KINLAW W B 1954 Cat scratch disease: report of a case representing hepatosplenomegaly Conn Med J 18 960-962
- 192 IOVENITAL K 1954 Cat scratch disease N Y State J Med 54 2090-2094
- 193 LISAUGHIT J N 1954 Benign inoculation lymphoreticulosis cat scratch disease J Oklahoma Med Assoc 47 131-132
- 194 MARETSKAIA M I 1955 Infektsionnoe vyzvanoe tsarajm i ili ukusom koshki (Infectious disease caused by cat bite or scratch) Sovet M d 19 45-48
- 195 MARQUÉZY R A & C BACH 1955 Erythème noueux et lymphoreticulose bénigne deux observations (Erythema nodosum and benign lymphoreticulosis two observations) Bull mem soc med hop Paris 71 647-644
- 196 MARSHALL C F 1956 Cat scratch fever Can Med Assoc J 76 724-732
- 197 MCGOVERN J J L J KUNZ & F M BLODGETT 1955 Nonarterial regional lymphadenitis (cat scratch fever) New Engl J Med 252 166-177
- 198 MEYER A & P FRÉOUR 1954 Allonopathies manifestes transitoires coexistent avec une réaction positive à l'antigène de la lymphoreticulose (Transitory manifestations of lymphopathies associated with positive antigen reactions of cat scratch disease) J franç m l et chir thorac 8 163-168
- 199 MEYER A J C HESSE & J C TABARY 1955 Erythème noueux et lymphoreticulose bénigne d'inoculation avec allonopathie cervicale (Erythema nodosum and benign inoculation lymphoreticulosis with cervical adenopathy) Bull m m soc med hôp Paris 71 577-581
- 200 MISIL F 1955 Onemocnění Kočičího Skrabnutí (Cat scratch disease) Rozhledy chir 34 439-446
- 201 NETTER R Y CHANALET & J BERNOT 1956 Premier cas de lymphoreticulose bénigne d'inoculation au Vietnam (First case benign inoculation lymphoreticulosis in Viet Nam) Bull soc pathol exotique 49 23-25
- 202 ORBETON F A 1954 Benign inoculation lymphoreticulosis Cat scratch disease J Maine Med Assoc 45 223-225
- 203 O REAR H B 1954 Cat scratch disease J Med Assoc Georgia 43 869-870
- 204 PARNAS J 1954 Choroba kocia (Cat scratch disease) Wiadomości Lekarskie 7 57-60
- 205 PATZEROWA T 1954 Przypadek choroby Debré Mollaret u 10 letniej dziewczynki w przebiegu przewlekłego zapalenia wyrostka robaczkowego (Case of Debré Mollaret disease in a 10-year old girl during the course of chronic appendicitis) Pediatria polska 29 290-291
- 206 PELNER L 1955 Cat scratch disease or benign lymphoreticulosis Am Practitioner and Dig Treatment 6 1164-1166
- 207 PENDLETON T L 1954 Cat scratch disease on the San Francisco peninsula Calif Med 80 46-48
- 208 PHILLIPS W R 1954 Cat scratch syndrome N Y State J Med 54 169-171
- 209 PHILPOTT O S A R WOODBURN & J A PHILPOTT JR 1955 Cat scratch fever Rocky Mt Med J 52 789-790
- 210 POLAYES S H 1954 Cat scratch fever Brooklyn Hosp J 12 50-51
- 211 RATKOWSKI B 1954 Cat scratch disease Harefuah 47 31-37
- 212 REILLY J 1955 La lymphoreticulose bénigne d'inoculation dite maladie des griffes du chat (Benign inoculation lymphoreticulosis or so-called cat scratch disease) Bruxelles med 35 213-223
- 213 RISTIC L 1953 Takozvana bolest usled ogrebatina macke (The so-called cat scratch disease) Spski arh cel lek 81 400-403 (Abstr in Biol Abstr 28 11497 1954)
- 214 ROSEMAN R B H BRODY & J A STERLING 1954 Cat scratch disease case report J Allert Einstein Med Center 2 67-65
- 215 RUDMAN I & J R CUMMINGS 1955 Cat scratch disease Ill Med J 108 120
- 216 SCHMID K O 1956 Ein Beitrag zur sogenannten Katzenkratzenkrankheit (So-called cat scratch disease) Wien klin Wochschr 68 568-573
- 217 SEBASTIANI F 1956 A proposito di un caso clinico di linforeticulosi benigna da inoculazione nota di aggiornamento (Clinical case of benign lymphoreticulosis caused by inoculation an up-to-date note) Gio ital dermatol e sifilol 92 173-180
- 218 SEMELAIGNE P J HUREZ E BRAULT & M BATTISTELLI 1956 Maladie des griffes du chat pandémie familiale coïncidant avec des épizooties à virus (Cat scratch disease family epidemic coincidence with viral epizootic diseases) Arch franç pédiat 13 67-69
- 219 SLATER S 1954 Cat scratch disease N Y State J Med 54 2486-2487

- 220 STALLINGS J H JR 1955 Cat scratch disease report of two cases *Chn Trac Child Hosp* 11 222-224
- 221 STERLING J A R B ROSEMAN & H BRODY 1954 Axillary adenopathy due to cat scratch *Am Surgeon* 20 1012-1014
- 222 TAUBENHAUS L J 1954 Cat scratch disease clinical case report *N Carolina Med J* 15 136
- 223 VALLEDOR T I E EXPÓSITO L BARBELL A CUERRA CHABAU & A GARCÍA PALACIOS 1955 Enfermedad por arañazos de gato una epidemia familiar en tres hermanos primeras observaciones reportadas en Cuba (Cat scratch disease familial epidemic in three siblings first cases reported in Cuba) *Rev cubana pediat* 27 125-142
- 224 WARWICK W J A PAGE & R A GOOD 1956 Passive transfer with circulating leukocytes of delayed hypersensitivity to cat scratch antigen *Proc Soc Exptl Biol Med* 93 253-255
- 225 WATSON J R H 1956 Cat scratch disease in Western Australia A report of 3 cases *Med J Australia* 2 20-21
- 226 WEEKLY CLINICOPATHOLOGICAL EXERCISES CASE 40052 1954 *New Engl J Med* 250 217-220
- 227 WEINSTEIN I & R H MEAD 1955 III The neurological manifestations of cat scratch disease *Am J Med Sci* 229 500-505
- 228 WILLCOX R R 1955 Skin tests for psittacosis and cat scratch fever in non gonococcal urethritis *Acta Dermato-Venerol* 35 174-19
- 229 MOLLARET P J REILLY R BASTIN & P TOURNIER 1951 La découverte du virus de la lymphoréticulose bénigne d'inoculation II Inoculation expérimentale au singe et colorations *Presse med* 59 681-682
- 230 ARMSTRONG C W B DANIELS F G MACMURRAY & H C TURNER 1956 Complement fixation in cat scratch disease employing Lygranum CF as antigen *J Am Med Assoc* 161 149-150
- 231 CAMPBELL F S K & C E W WHEATON 1952 Cat scratch fever *Lancet* 1952(i) 975
- 232 DANIELS W B & F G MACMURRAY 1952 Cat scratch disease non bacterial regional lymphadenitis a report of 60 cases *Ann Internal Med* 37 697-713
- 233 KALTER S S J E PRIER & J T PRIOR 1955 Recent studies on the diagnosis of cat scratch fever *Ann Internal Med* 42 562-573
- 234 MOLLARET P J REILLY R BASTIN & P TOURNIER 1951 La découverte du virus de la lymphoréticulose bénigne d'inoculation I Caractérisation sérologique et un munologique *Presse m d* 59 681-682
- 235 WINSHIP T 1953 Pathologic changes in so-called cat scratch fever review of findings in lymph node of 29 patients and cutaneous lesions of 2 patients *Am J Clin Pathol* 23 1012-1018
- 236 DEBRÉ R L VAN BOGAERT S THIEFFRY & M ARTHUS 1952 Accidents nerveux de la maladie des griffes du chat (Nervous complications in cat scratch disease) *Bull acad natl med* 136 454-459
- 237 MOLLARET P J REILLY R BASTIN & P TOURNIER 1951 La lymphoréticulose bénigne d'inoculation (Benign inoculation lymphoreticulosis) *Acta Med Ital Mal Infett* 6 173-182
- 238 THOMSON T E JR & K F MILLER 1953 Cat scratch encephalitis *Ann Internal Med* 39 146-151
- 239 PETERMAN M G 1954 Cat scratch disease report of a case in 1925 *J Pediat* 44 563-564
- 240 CAMPBELL W N & T G ANDERSON 1953 Cat scratch disease *Penn Med J* 56 188-190
- 241 STEVENS H 1952 Cat scratch fever encephalitis *AMA J Diseases Children* 84 218-222
- 242 BELBER J P A E DAVIS & E H EPSTEIN 1954 Thrombocytopenic purpura associated with cat scratch disease response of cat scratch disease to steroid hormones *AMA Arch Internal Med* 94 371-375
- 243 ADAMS W C & S M HINDMAN 1954 Cat scratch disease associated with an osteolytic lesion *J Pediat* 44 665-669
- 244 DANIELS W B & F G MACMURRAY 1954 Cat scratch disease report of one hundred sixty cases *J Am Med Assoc* 164 1241-1251
- 245 CASSADY J V & C S CULBERTSON 1953 Cat scratch disease and Parinaud's oculoglandular syndrome *AMA Arch Ophthalmol* 50 68-74

- 246 CAT SCRATCH DISEASE AND FARINATO'S OCULOCILLIARY SYNDROME 1953 Editorial J Am Med Assoc 162 1717
- 247 BRAND T A & K C FINKEL 1956 Cat scratch fever Brit Med J 1956(ii) 88-91
- 248 CARAI D F 1952 Cat scratch fever a case in England Lancet 1952(ii) 646
- 249 BLA A H & C RAKE 1955 Viral and Rickettsial Diseases of the Skin Eye and Mucous Membranes of Man 208 Little Brown & Co Boston Mass.
- 250 DANIELS W B & F G MacMURRAY 1953 Cat scratch disease a report of 169 cases Trans Am Climat Clin Assoc 65 99 115

- 220 STALLINGS J H JR 1955 Cat scratch disease report of two cases Clin Proc Child Hosp 11 222-224
- 221 STERLING J A P B ROSEMAN & H BRODY 1954 Axillary adenopathy due to cat scratch Am Surgeon 20 1012-1014
- 222 TAUBENHAUS I J 1954 Cat scratch disease clinical case report N Carolina Med J 15 136
- 223 VALLEDOR T I F EXPOSITO L BARBELLA A GUERRA CHABAU & A GARCIA PALACIOS 1955 Enfermedad por arañazos de gato una epidemia familiar en tres hermanos primeras observaciones reportadas en Cuba (Cat scratch disease famihial epidemic in three siblings first cases reported in Cuba) Rev cubana pediat 27 125-142
- 224 WARWICK W J A PAGE & R A GOOD 1956 Passive transfer with circulating leukocytes of delayed hypersensitivity to cat scratch antigen Proc Soc Exptl Biol Med 93 253-255
- 225 WATSON J R H 1956 Cat scratch disease in Western Australia A report of 3 cases Med J Australia 2 20-21
- 226 WEEKLY CLINICOPATHOLOGICAL EXERCISES CASE 40052 1954 New Engl J Med 250 217-220
- 227 WEINSTEIN L & R H MEADE 1955 III The neurological manifestations of cat scratch disease Am J Med Sci 229 500-505
- 228 WILLCOX P R 1955 Skin tests for psittacosis and cat scratch fever in non gonococcal urethritis Acta Dermato Venereol 35 174-179
- 229 MOLLARET P J REILLY R BASTIN & P TOURNIER 1951 La decouverte du virus de la lymphoréticulose bénigne d'inoculation II Inoculation expérimentale au singe et colorations Presse med 59 681-682
- 230 ARMSTRONG C W B DANIELS F G MACMURRAY & H C TURNER 1956 Complement fixation in cat scratch disease employing Lygranum CF as antigen J Am Med Assoc 161 149-150
- 231 CAMPBELL F S K & C E W WHEATON 1952 Cat scratch fever Lancet 1952(i) 975
- 232 DANIELS W B & F G MACMURRAY 1952 Cat scratch disease non bacterial regional lymphadenitis a report of 60 cases Ann Internal Med 37 697-713
- 233 KALTER S S J E PRIER & J T PRIOR 1955 Recent studies on the diagnosis of cat scratch fever Ann Internal Med 42 562-573
- 234 MOLLARET P J REILLY R BASTIN & P TOURNIER 1951 La decouverte du virus de la lymphoréticulose benigne d'inoculation I Caractérisation sérologique et immunologique Presse méd 59 681-682
- 235 WINSHIP T 1953 Pathologic changes in so-called cat scratch fever review of findings in lymph node of 29 patients and cutaneous lesions of 2 patients Am J Clin Pathol 23 1012-1018
- 236 DEBRÉ R L VAN BOGAERT S THIEFFRY & M ARTHUS 1952 Accidents nerveux de la maladie des griffes du chat (Nervous complications in cat scratch disease) Bull acad natl med 136 454-459
- 237 MOLLARET P J REILLY R BASTIN & P TOURNIER 1951 La linforeticulose benigna d'inoculazione (Benign inoculation lymphoreticulosis) Acta Med Ital Mal Infett 6 173-182
- 238 THOMPSON T F JR & K F MILLER 1953 Cat scratch encephalitis Ann Internal Med 39 146-151
- 239 IETERMAN M G 1954 Cat scratch disease report of a case in 1975 J Pediat 44 563-564
- 240 CAMPBELL W N & T G ANDERSON 1953 Cat scratch disease Penn Med J 56 188-190
- 241 STEVENS H 1952 Cat scratch fever encephalitis AMA J Diseases Children 84 218-222
- 242 BELBER J P A E DAVIS & F H EPSTEIN 1954 Thrombocytopenic purpura associated with cat scratch disease response of cat scratch disease to steroid hormones AMA Arch Internal Med 94 321-375
- 243 ADAMS W C & S M HINDMAN 1954 Cat scratch disease associated with an osteolytic lesion J Pediat 44 665-669
- 244 DANIELS W B & F G MACMURRAY 1954 Cat scratch disease report of one hundred sixty cases J Am Med Assoc 164 1247-1251
- 245 CASSADY J A & C S CULBERTSON 1953 Cat scratch disease and Parinaud's oculoglandular syndrome AMA Arch Ophthalmol 50 68-74




FIGURE 1 Animal plague in the United States from 1903 through 1940

Numerous species of wild rodents have been implicated in the epizootiology of plague ever since the first suggestive evidence of an epizootic in ground squirrels *Citellus beecheyi* was found in Contra Costa County Calif from 1903 to 1905 (Meyer 1942b). When plague was conclusively demonstrated in ground squirrels by McCoy (1908) and by Wherry (1908) the attention of investigators was focused upon the larger colonial rodents. Thus Eskey and Haas (1940) in their classic study of plague in the western part of the country indicated that the primary reservoirs of plague were ground squirrels *Citellus* spp, wood rats *Neotoma* spp, and prairie dogs *Cynomys* spp. Occasional findings of plague in native mice such as *Peromyscus* spp mentioned by these and other authors were considered by them to be unimportant accidental infections acquired by contact with ground squirrels or other important reservoirs.

Colonial rodents best served the purpose of the early plague surveys which were conducted primarily to demonstrate the geographic extent of the disease. The adoption in 1936 of routine inoculation of pooled fleas in addition to tissue inoculations by Eskey and Haas (1940) increased the efficiency of surveys in delineating the rodent and flea reservoirs of plague. Consequently tissue and flea pools from ground squirrels and prairie dogs constitute the major source of plague positive records thus far accumulated on wild rodent plague in this country.

In more recent years there has been a growing conviction among investigators of plague that rodents other than the ground squirrel and the large conspicuous

NEW KNOWLEDGE ON THE ECOLOGY OF SYLVATIC PLAGUE

By Leo Kartman, Frank M. Prince, Stuart F. Quan, and
Harold E. Stark

*San Francisco Field Station, San Francisco, Calif., Communicable Disease Center,
U. S. Public Health Service*

There is every reason to believe that an epidemic of plague is a self-regulating phenomenon which is capable of description provided enough of its ecology is known. Indeed, it would remove a great deal of the fearsome unknown if one could forecast the peregrinations of plague which in man are merely accidental sequels to a broad rodent parasitism in his environment. —Karl F. Meyer (1942a)

INTRODUCTION

The history of plague is an eloquent example of changing concepts of investigation in which numerous hypotheses have been advanced, their limitations described, their positive values extracted, and their remaining skeletons buried in favor of new ideas. Known in classical antiquity, in the Middle Ages, and in the pandemics of the latter Nineteenth and early Twentieth centuries, plague has been studied primarily as an epidemic disease causing widespread mortality in human populations. Until comparatively recently, plague was thought to be fundamentally a disease of domestic rats in which rat fleas were the major vectors. By 1928 it was clear that sylvatic or wild rodent plague represented an independent epizootiologic entity, and since that time it has been recognized that sylvatic plague may transfer to domestic rodents under certain conditions (Pollitzer 1954).

Detailed historical and ecologic views and reviews of plague in the western United States have been written by Meyer (1942a and 1942b), while the works of Pollitzer (1954) and Macchiavello (1954) document the vast world literature. The world distribution of sylvatic plague has been summarized by Garnham (1949). Sylvatic plague is now known to be established in wild rodents and their fleas in the western region of the United States and in western Canada. Recently, the presence of wild rodent plague was shown in Mexico with the isolation of the organism from prairie dogs in the northern state of Coahuila (Varela and Vasquez 1954). FIGURE 1 shows the distribution of plague in animals as a result of extensive surveys, especially from 1936 to 1950, conducted by the United States Public Health Service and by certain states. The pre-dominance of wild rodent over rat-borne plague is shown in FIGURE 2. Wild rodent plague foci have been found in 131 counties of 15 western states in surveys from the Pacific Coast to the 100th meridian, an area comprising about 40 per cent of the continental United States. The evidence shows that the disease in wild rodents forms a vast enzootic reservoir as a potential threat to humans. Pollitzer (1954) lists 91 plague cases and 57 deaths in 7 western states during the period 1908 to 1951 as having been contracted from wild rodent sources. In June 1956 a fatal human case occurred in Ventura County, Calif.; this was apparently contracted from ground squirrel fleas. FIGURE 3 shows the distribution of human plague cases resulting from contact with wild rodents, with the exception of the case shown in Michigan, which is a laboratory infection.




FIGURE 3 United States counties in which human plague has occurred

mice *Microtus* spp and *Dipodomys* spp are resistant to experimental plague infection

It is important to note that some of these small rodents are generally more closely associated with domestic rats *Rattus* spp in the immediate environs of human habitations than are the ground squirrels and other colonial species. The potential danger is obvious and in the words of Meyer (1933) "The increase in the population of California has given rise to two situations that must be taken into account. The building of suburbs ultimately makes a habitat unsuitable for maintaining appreciable wild rodent populations but there is at least an initial period of joint tenancy by people and wild and commensal rodents—a condition theoretically ideal for the propagation of plague. The wilderness recreational areas of California have been crowded with more people every year. This too has increased the opportunity for close contact between man and reservoir animals and ectoparasites."

It is known that native field mice and domestic rats coexist and live in newly suburbanized areas around refuse disposal sites on ranches in recreational areas and even in city vacant lots. If plague infective fleas from field mice transfer to rats causing plague then infective rat fleas are introduced into the human environment. The dangers inherent in the close association of wild rodents and domestic rats were recognized over 40 years ago by McCoy (1910). Eskey (1938) found the wild rodent flea *Xenopsylla conformis* on Norway rats in San Francisco. From an analysis of data based on 5783 fleas




FIGURE 2 Rat-borne and sylvatic plague in the United States from 1900 through 1949

species may be more usual reservoirs of *Pasteurella pestis*. The colonial rodents are like man perhaps only occasionally included in plague outbreaks initiated by epizootics among native field mice (Meyer 1955). In California since 1908 investigations have shown that plague in wild rodents was enzootic in three-fourths of the state and by 1952 the following 10 genera of native wild rodents and 2 genera of lagomorphs and/or their fleas were implicated with plague: *Citellus*, *Neotoma*, *Peromyscus*, *Futamias*, *Tamiasciurus*, *Glaucomys*, *Marmota*, *Thomomys*, *Microtus*, *Reithrodontomys*, *Lepus*, and *Sylvilagus* (Jones and Street 1956). A perusal of the California records shows that plague-positive fleas have been taken from the small native field mice on numerous occasions. For instance during 1943 in Monterey County meadow mice (*Microtus californicus*), white-footed mice (*Peromyscus* spp.) and their fleas were discovered to be infected with plague in an area where ground squirrels were thought to have been virtually eradicated by intensive control operations (Jones and Street 1956).

Epizootiologic studies in various parts of the world have indicated that the mammalian plague reservoirs of primary importance are not necessarily the highly susceptible conspicuous species. Some native mice that are relatively plague-resistant may constitute the more usual reservoirs for the persistence of plague as has been found in Africa (Heisch *et al.* 1953 and Davis 1953). Preliminary findings in California (Quan and Kartman 1956) and in New Mexico (Holdenried and Quan 1956) suggest that certain strains of native field




FIGURE 3 United States counties in which human plague has occurred

mice *Microtus* spp and *Dipodomys* spp are resistant to experimental plague infection

It is important to note that some of these small rodents are generally more closely associated with domestic rats *Rattus* spp in the immediate environs of human habitations than are the ground squirrels and other colonial species. The potential danger is obvious and in the words of Meyer (1955) "The increase in the population of California has given rise to two situations that must be taken into account. The building of suburbs ultimately makes a habitat unsuitable for maintaining appreciable wild rodent populations but there is at least an initial period of joint tenancy by people and wild and commensal rodents—a condition theoretically ideal for the propagation of plague. The wilderness recreational areas of California have been crowded with more people every year. This too has increased the opportunity for close contact between man and reservoir animals and ectoparasites."

It is known that native field mice and domestic rats coexist and live in newly suburbanized areas around refuse disposal sites on ranches in recreational areas and even in city vacant lots. If plague infective fleas from field mice transfer to rats causing plague then infective rat fleas are introduced into the human environment. The dangers inherent in the close association of wild rodents and domestic rats were recognized over 40 years ago by McCoy (1910). Eskey (1938) found the wild rodent flea *Xaloraeus telchium* on Norway rats in San Francisco. From an analysis of data based on 5785 fleas




FIGURE 2 Rat borne and sylvatic plague in the United States from 1900 through 1949

species may be more usual reservoirs of *Pasteurella pestis*. The colonial rodents are like man perhaps only occasionally included in plague outbreaks initiated by epizootics among native field mice (Meyer 1955). In California since 1908 investigations have shown that plague in wild rodents was enzootic in three fourths of the state and by 1952 the following 10 genera of native wild rodents and 2 genera of lagomorphs and/or their fleas were implicated with plague: *Citellus*, *Neotoma*, *Peromyscus*, *Futamias*, *Tamiasciurus*, *Glaucomys*, *Marmota*, *Thomomys*, *Microtus*, *Reithrodontomys*, *Lepus* and *Sylvilagus* (Jones and Street 1956). A perusal of the California records shows that plague positive fleas have been taken from the small native field mice on numerous occasions. For instance during 1943 in Monterey County meadow mice (*Microtus californicus*), white footed mice (*Peromyscus* spp.) and their fleas were discovered to be infected with plague in an area where ground squirrels were thought to have been virtually eradicated by intensive control operations (Jones and Street 1956).

Epidemiologic studies in various parts of the world have indicated that the mammalian plague reservoirs of primary importance are not necessarily the highly susceptible conspicuous species. Some native mice that are relatively plague resistant may constitute the more usual reservoirs for the persistence of plague as has been found in Africa (Heisch *et al.* 1953 and Davis 1953). Preliminary findings in California (Quan and Kartman 1956) and in New Mexico (Holdenried and Quan 1956) suggest that certain strains of native field

suggest what the present position is in regard to various ecologic problems and to control. Each item will be discussed under appropriate headings.

The history of sanitary legislation has been a reflection of the evolution of theories and knowledge of the epidemiology and control of the relevant communicable diseases.—H. S. Gear and L. Deutschman (1956)

PLAGUE IN THE UNITED STATES AND THE INTERNATIONAL SANITARY REGULATIONS

The fascinating history of quarantine practices has been told recently by Hirst (1953) and briefly summarized by Gear and Deutschman (1956). Plague is one of the so-called quarantinable diseases against which legislation has been primarily directed from the time of the early sanitary conventions to the present International Sanitary Regulations (World Health Organization 1951). In modern practice quarantine regulations on plague deal with rodent control on ships and in ports, the isolation of human contacts or actual plague cases, and disinsectionization of ships and aircraft. Article 51 of the International Sanitary Regulations specifically states: "Each state shall employ all means in its power to diminish the danger from the spread of plague by rodents and their ectoparasites. Its health administration shall keep itself constantly informed by systematic collection and regular examination of rodents and their ectoparasites of the conditions of any local area, especially any port or airport infected or suspected of being infected by rodent plague."

The theoretical basis for these rules is rapidly becoming obsolete. The extent and severity of plague epidemics has significantly declined throughout the world, and it should not be long before most ports will feel secure as the result of the reduction of rat populations and because of adequate medical facilities. Nevertheless, there is a growing cognizance that even when the danger from rats has been lowered, the danger of carryover from wild rodent sources is still a fact to be considered seriously. Thus the World Health Organization Expert Committee on Plague (1953) stated: "Taking into consideration that the relationship between wild rodent and domestic rodent plague has not been fully elucidated, it was further recommended that countries in which plague is still endemic should undertake systematic studies on *ecologic* lines to establish the comparative importance of wild and domestic rodent species in the maintenance of the plague reservoir, and that particular attention should be given to a comprehensive investigation of the vector ability and survival of plague infected fleas under different conditions of temperature and humidity, both in the laboratory and in the field."

In the above quotation we have italicized the word *ecologic* since this is the area of investigation least understood in modern concepts of the plague. As will be seen in other sections of this paper, plague ecology is now the chief concern of the San Francisco Field Station, even though routine surveillance of formerly plague infected port cities is still maintained. It has been suggested (Gear and Deutschman 1956) that plague in rural rodents need not be given very much consideration in international traffic. However, it appears obvious that as long as wild rodent plague represents a possible source of the infection to urban areas, it must be considered in the philosophy of quarantine. We

from 4188 domestic rats caught in 13 western states Prince (1943) showed that these rats harbored 15 different species of wild rodent fleas Meyer and Holdenried (1949) established the existence of plague in rats *Rattus r. rattus* and *R. norvegicus* and in ground squirrels *Citellus beecheyi* on a ranch in Ventura County Calif Rat fleas were not found on the squirrels but 25 per cent of the fleas taken from the rats were squirrel fleas *Diplospyllus montanus* and *Hoplospyllus anomalus* These authors concluded that plague was transferred to the rats by these wild rodent ectoparasites

Although this introduction has merely presented a few of the more important facts concerning plague in the western region of the United States it is sufficient to indicate that studies of plague in this country have only scratched the surface of the formidable epizootiological complex represented by the phrase 'wild rodent plague' There is a definite need for intensive ecologic studies to elucidate the host ectoparasite disease relationships in plague foci especially those found near human populations The hazard of plague to the public health at present cannot be accurately evaluated therefore and only ill defined recommendations can be made for safeguarding humans against the enzootic plague reservoir It is generally agreed that while it is impracticable to attempt the control of wild rodent plague on a wide scale natural plague foci should not be permitted to exist where they threaten human populations The potentially dangerous nature of natural focal localizations or 'nidi' of diseases transmissible to man have been described in detail by Pavlovsky (1955) It is now somewhat vaguely suggested that plague free zones should be maintained by the elimination of foci around populated areas or places temporarily inhabited by man This concept is confronted at the outset by the paucity of techniques for determining rapidly the presence or absence of plague in a given area Furthermore the currently meager knowledge of the ecology of plague precludes a sound judgment of such basic considerations as the weakest link in the infection chain the seasons of prevalence and quiescence practical protective zones and the types of habitats reservoirs and wild domestic rodent contacts that may harbor plague locally and present a threat to humans

Pioneering work on plague ecology in California has been done by a group of scientists formerly at the University of California Hooper Foundation Berkeley Calif (1939 to 1946), under the direction of K F Meyer These studies did much to show the direction in which modern concepts of plague ecology must travel More recently the Communicable Disease Center (CDC) of the Public Health Service established a field station at Santa Fe N M (Holdenried 1951) where studies of plague ecology in small native rodents were carried out during a three year period During the last six years the San Francisco Field Station of the CDC has initiated field and laboratory studies on the ecology and control of plague in the San Francisco Bay area (Kartman Miles and Prince 1958 Miles Kinney and Stark 1957) and on the control of enzootic plague in Hawaii (Kartman and Loneragan 1955a and 1955b) It would be beyond the scope of the present paper to discuss sylvatic plague on a world wide basis We shall therefore attempt to give an account of our own investigations in the western United States and to

families in the United States. All these rodents are diurnal and fairly large and are therefore more frequently observed by man. *Cynomys* noted for its colonial habits is most frequently involved in the spectacular die offs in which whole prairie dog towns covering hundreds of acres have been virtually extirpated. Among other rodents cricetine and murine species have yielded plague positive fleas and tissues on a number of occasions but no spectacular die offs have been observed as in the case of prairie dogs. Most of these animals are nocturnal and all are small and inconspicuous thus permitting far less chance for gross observation. Eskey and Haas (1940) reported plague in epizootic form among *Neotoma* the pack rats. Field workers noticed many deserted nests of this animal. When opened these nests revealed dead animals. Although plague was not isolated from their tissues the organism was found in fleas removed from the nests.

Recently new views have developed as to which rodents constitute the basic plague reservoir. Multimammate mice in South Africa are considered almost entirely responsible for transfers of wild rodent plague to *Rattus* (Davis 1953). Heisch *et al* (1953) in East Africa and Baltazard *et al* (1952) in Persia consider resistant rodents to be basic plague reservoirs. Holdenrud and Quan (1956) found that various rodents in New Mexico were resistant to experimental infections of plague. Data presented below show a correlation of plague resistance and the actual occurrence of plague in some small rodent populations in the San Francisco Bay area of California.

One of the first objectives in ecologic studies conducted in San Francisco was to determine areas of association between domestic rats and wild rodents and to detect exchange of fleas between these wild rodents and domestic rats. Hooper (1944) recorded 43 species and subspecies of native rodents from 9 ecologic associations in the San Francisco Bay area. In our studies 12 native rodents were found in association with 3 species of domestic rodents in 4 ecologic associations as follows: (1) *Salicornia* marsh association, (2) bunch grass meadow association, (3) sage coyote brush association, and (4) riparian association. Of the 12 rodents at least 4 were found to commingle with rats in significant numbers. These were *Merotus californicus* (2 subspecies), *Reithrodontomys megalotis* (1 subspecies), *Ieromyscus maniculatus* (1 subspecies) and *Citellus beecheyi* (2 subspecies). All ecologic associations encountered were variously modified by suburbanization, industrialization or farming. The trap sites may be characterized as follows: dumps, hog farms, dairies, parks, highway developments, factories, feed mills, and food packing plants. All of these were located close to the bay or in mountains yet unclaimed by suburban developments. Five mammal groups all belonging to orders other than Rodentia were encountered. Only one group *Sorex* (3 species) was taken in significant numbers. All of the individuals in the remaining 4 groups constituted about 1 per cent of the total capture. TABLE 1 presents a list of the most important mammals trapped in descending order of frequency of capture. The data presented are from a study by Miles *et al* (1957) and from subsequent and more detailed studies made in some of the areas where plague was found. The 7 mammals listed constituted virtually 100 per cent of the total captures. The remaining 10 mammals not shown were captured

have isolated plague infected rat fleas from the port of Tacoma Wash on at least two occasions in recent years. Surveys in that port have shown that microtine rodents live in close contact with domestic rats in vacant lots and other areas of the city. Thus there exists the strong possibility until proved otherwise that rats had acquired plague from wild rodent sources. Actual transfer of plague from native field mice to rats has been found in our studies in the San Francisco area (Kartman *et al* 1958). The fact that transfer of plague actually does take place under modern conditions of sanitation emphasizes the thin tissue of protection now afforded humans living in certain areas. It would probably take only a slight breakdown in present conditions to allow an increase in plague to spread to domestic rats. Under disaster or emergency conditions such a spread could be explosive. With present rapid means of travel an infected person could move thousands of miles away from the sylvatic plague source before the onset of clinical symptoms.

We can conclude only that the international danger of the spread of plague from wild to domestic reservoirs is at present not subject to clear definition. More intensive ecologic studies are needed for an exact delimitation of wild rodent plague and for an intelligent assessment of its epidemic potential. It would seem that until such knowledge is available the international sanitary code must give proper recognition to sylvatic plague as an international menace and not treat it merely as a disagreeable local phenomenon.

The ecology of wild rodent plague is most complex to unravel it is like following the different voices in a Bach fugue except that in plague the basic design is unknown.—P. C. C. Garnham (1949)

SOME ECOLOGIC FACTORS IN SYLVATIC PLAGUE

Rodent Species Involved

It is recognized (Pollitzer 1954; Macchiavello 1954) that both mammals and fleas may serve either as reservoirs or vectors of plague. The important factor in the occurrence and transmission of plague is not simply whether or not the mammal or flea is a reservoir or a vector but the degree to which it serves in either capacity and what species is involved. Macchiavello (1954) lists 344 mammal and 2 bird species that have been reported from various parts of the world as being experimentally infected as naturally infected or from which plague infected fleas were removed. The great majority of these mammals are rodents of which 67 species occur in the Nearctic region. Tables presented by Pollitzer (1954) and by Macchiavello (1954) include certain rodents that were easily infected experimentally with plague and thus presumed to be of importance. However as will be pointed out later some rodents that have been proved to be ecologically implicated with plague are quite resistant when tested in the laboratory. These rodents may actually be of greater importance to the perpetuation of the disease though they do not exhibit spectacular die off rates.

Rodents of the family Scuridae are of great importance in plague. They have been involved in contacts with humans under sylvatic conditions and have transmitted plague directly to man more often than have other rodent

families in the United States. All these rodents are diurnal and fairly large and are therefore more frequently observed by man. *Cynomys* noted for its colonial habits is most frequently involved in the spectacular die-offs in which whole prairie dog towns covering hundreds of acres have been virtually extirpated. Among other rodents cricetine and microtine species have yielded plague positive fleas and tissues on a number of occasions but no spectacular die-offs have been observed as in the case of prairie dogs. Most of these animals are nocturnal and all are small and inconspicuous thus permitting far less chance for gross observation. Eskey and Haas (1940) reported plague in epizootic form among *Neotoma* the pack rats. Field workers noticed many deserted nests of this animal. When opened these nests revealed dead animals. Although plague was not isolated from their tissues the organism was found in fleas removed from the nests.

Recently new views have developed as to which rodents constitute the basic plague reservoir. Multimammate mice in South Africa are considered almost entirely responsible for transfers of wild rodent plague to *Rattus* (Davis 1953). Hensch *et al* (1953) in East Africa and Baltazard *et al* (1952) in Persia consider resistant rodents to be basic plague reservoirs. Holdenried and Quan (1956) found that various rodents in New Mexico were resistant to experimental infections of plague. Data presented below show a correlation of plague resistance and the actual occurrence of plague in some small rodent populations in the San Francisco Bay area of California.

One of the first objectives in ecologic studies conducted in San Francisco was to determine areas of association between domestic rats and wild rodents and to detect exchange of fleas between these wild rodents and domestic rats. Hooper (1944) recorded 43 species and subspecies of native rodents from 9 ecologic associations in the San Francisco Bay area. In our studies 12 native rodents were found in association with 3 species of domestic rodents in 4 ecologic associations as follows: (1) *Salicornia* marsh association, (2) bunch grass meadow association, (3) sage-coyote brush association, and (4) riparian association. Of the 12 rodents at least 4 were found to commingle with rats in significant numbers. These were *Microtus californicus* (2 subspecies), *Reithrodontomys megalotis* (1 subspecies), *Peromyscus maniculatus* (1 subspecies) and *Citellus beecheyi* (2 subspecies). All ecologic associations encountered were variously modified by suburbanization, industrialization or farming. The trap sites may be characterized as follows: dumps, hog farms, dairies, parks, highway developments, factories, feed mills, and food packing plants. All of these were located close to the bay or in mountains yet unclaimed by suburban developments. Five mammal groups all belonging to orders other than Rodentia were encountered. Only one group, *Sorex* (3 species) was taken in significant numbers. All of the individuals in the remaining 4 groups constituted about 1 per cent of the total capture. TABLE 1 presents a list of the most important mammals trapped in descending order of frequency of capture. The data presented are from a study by Miles *et al* (1951) and from subsequent and more detailed studies made in some of the areas where plague was found. The 7 mammals listed constituted virtually 100 per cent of the total captures. The remaining 10 mammals not shown were captured

TABLE 1

WILD MAMMALS TRAPPED FROM THE SAN FRANCISCO BAY REGION OF CALIFORNIA IN 13 AREAS WHERE DOMESTIC RODENTS WERE PRESENT

Species collected	Number trapped	Percent of total	Fleas (all species)	Approximate difference between (ft)
<i>Microtus californicus</i>	2442	60.0	5.3	<100
<i>Peromyscus maniculatus</i>	528	12.0	2.7	>100
<i>Reithrodontomys megalotis</i>	454	11.0	0.3	>100
<i>Rattus norvegicus</i>	273	7.0	2.3	>500
<i>Mus musculus</i>	237	6.0	0.2	>100
<i>Sorex</i> sp.	110	3.0	0.4	not known
<i>Citellus beecheyi</i>	23	0.5	38.2	>500

* A total of 37 individuals of 10 additional mammal species were trapped or about 0.9 per cent of all animals taken

incidentally. Either they were not in close association with feral rats or they were present in very limited numbers.

It will be noted that the ground squirrel *Citellus beecheyi* has the greatest number of fleas per host. However, squirrel fleas are usually highly host specific, thus tending to minimize the importance of commingling with domestic rodents. *Microtus* accounted for the greatest number of fleas taken from all areas. *Peromyscus* and *Rattus* each having lesser concentrations of fleas. *Sorex*, *Reithrodontomys*, and *Mus* had very low concentrations of fleas.

The first conclusive evidence of plague transfer from wild rodents to rats was reported by Meyer and Holdenried (1949) in an ecologic study of a plague epizootic in a feral colony of rats closely associated with *C. beecheyi*. The authors indicated that squirrel fleas were involved in the transfer. The work by Miles *et al.* (1957) suggested that in suburban areas there may be frequent flea interchange between smaller rodent species and rats. *Microtus californicus* appeared to be of greatest importance as regards flea exchange in the areas studied. In areas where the concentrations of both rats and small rodents were fairly equal, the index of fleas on alternative hosts was considerably higher than in areas where one or the other host predominated.

Blair (1940) has shown that the home range of small rodents such as *Microtus* probably is quite restricted, most individuals spending an entire lifetime in an area smaller than one acre. Grids of live traps were used in our more detailed studies of certain areas. The interval that seemed to have the best results was 30 feet. Our findings indicated that the ranges of the small rodents were quite limited. The range of *Peromyscus* extended relatively farther than that of other native rodents. Hayne (1950) discussed the relative merits of grid trapping, and the subject will not be pursued further.

In the San Francisco Bay area, *Microtus* was definitely confined to meadow-bunch grass association and rapidly gave way to other species as the ecologic association changed. In one area, rats and mice (*Mus*) were located only in *Salicornia* association bordering the bay, while *Microtus* and *Reithrodontomys* were in an adjoining grassy area. The line of demarcation was both definite and constant. In most other areas where domestic rats were encountered

Microtus and other wild rodents were repeatedly observed to coexist in the same niche however the rats dominated their particular home territory

Rats usually build burrows underground and within trash piles if these are available and they usually persist in a given area until environmental pressures force them to move. Kartman *et al* (1958) noted for a brief period the presence of rats along a small stream a short distance from a hog farm. Following a poisoning campaign at the environs of the hog farm by the owners and by county health authorities the rats disappeared from the stream location perhaps returning to the hog farm where the pressure from the previously larger population was eliminated. *Microtus* usually were observed to build their nests either above ground in a protected place or about 12 inches beneath the ground somewhere in their intricate tunnel systems. It may be that *Microtus* like other rodents builds two types of nests a resting nest built by all individuals and a breeding nest built only by females for rearing young. It appears therefore that the location of *Microtus* nests is a product of the type of area. In loose sandy soils the nests were invariably located underground in heavy claylike or soggy soils they were built within dense clumps of vegetation (usually in *Juncus* sp.) or at the base of shrubs (*Baccharis*). *Peromyscus* and *Reithrodontomys* nests were difficult to find they seemed to be located mainly above ground in shrubs. The predominant rodent of the sage coyote brush association was *Peromyscus*. The heaviest concentration of fleas occurred on *Microtus* this may be due primarily to the nest building habits of this rodent.

The unexplained complexity of sylvatic plague ecology is exemplified in the following account of recent observations in one of our study areas near San Francisco. During November 1956 it was evident that a spectacular disappearance of *Microtus* was taking place this apparently had its beginning in mid October. Six rodents were taken from the dwindling population of the area and autopsied to determine the possible presence of disease. Nothing was detected in the blood in histological sections and animal inoculations of brain liver spleen heart or lungs of the rodents or from flea inoculations of laboratory mice. The *Microtus* population remained low until the beginning of April when trapping was extended to include all areas plague positive in previous years in northern San Mateo County (south of San Francisco). The first trappings produced plague positive findings simultaneously in 5 areas and in 2 more weeks a total of 7 positive areas had been located. After the first plague finding new trapping grids were established in a section of the area that looked most promising for a study of the physical extent of wild rodent plague. The predominant types of flora and fauna in this area were grasses and few forbs the California vole *Microtus californicus* and the fleas *Xenopsylla conformis*, *Hystrihopysylla insulata* and *Athyphloceras multidentatus*. *M. californicus* constituted more than 90 per cent of the rodent hosts.

From April 16 through July 3 the capture of *Microtus* per trap day rose from 6 to more than 60. Plague was detected each week during that period by inoculations of flea pools into laboratory mice by plate culture of individual fleas and by cultures from occasional *Microtus* found dead in the field. No generally adverse effect on the *Microtus* population was noted. *Microtus*

nests were removed at intervals and some *Microtus* were taken for tissue and serologic tests. Fleas from 38 nests were examined for plague and 5 contained *P. pestis*. Eight *Microtus* were found with plague positive tissues. The study showed that plague was not limited to a single ecologic niche, but was found in various types of biotic communities in which *Microtus* and its fleas were present.

Three significant findings were made during this study.

(1) Plague occurred after the decimation of a 'stable' *Microtus* population. No plague had been detected from October 1955 to April 1957 at which time it became evident in a *Microtus* population that had dropped below the level of associated rodent species that are usually considerably less numerous than the predominant vole population.

(2) The *Microtus* population increased rapidly at a time when plague was being found in about 1 per cent of the fleas taken from the animals and from their nests and in the tissues of some of the rodents.

(3) In view of the fact that neither plague nor any other disease agent was detected in the *Microtus* during the October-April period, there is at present no explanation for the extreme reduction of the population at a time when *Microtus* populations in other areas appeared stable.

Other reports of reductions in vole populations have been attributed to extreme competition for food. However in the present case there was an ample supply of food and nesting material at all times. It is notable that plague was found in all 7 areas within a span of 3 weeks without any obvious warning of its presence. Thus it became apparent that plague appeared at a level readily demonstrable in the laboratory but without creating a noticeable effect on the *Microtus* or other small wild rodents over a wide area. As noted above, plague appeared after a decline and during a rise in the *Microtus* population. Heisch (1956) notes in his studies of sylvatic plague in Africa that outbreaks usually coincided with an increase in the wild rodents. However unlike the *Microtus* in our studies the African rodents were reported to decline suddenly at times due to plague. On the basis of present knowledge it would be difficult to speculate on this mechanism of epizootic plague and impossible to explain it. We feel that the highly resistant *Microtus* population and its fleas constitutes the *modus vivendi* of wild rodent plague in this area. However the mode of operation of the complex is still obscure.

The ecologic information gathered thus far points to the importance of the occurrence of small rodents as related to the perpetuation of plague in an area. Furthermore a given area is able to support greater numbers of small individuals than of large ones. The smaller native rodents may thus be a basic link in the plague chain serving as reservoirs of the organism during periods of apparent quiescence and supplying the source of infected fleas that may transfer the disease to domestic rats and then to man.

Rodent Susceptibility

The rodent hosts of *Pasteurella pestis* not only vary in epizootiological importance but also show marked differences in ecology. Some of the factors underlying these variations are evident and known, one is the experimental

susceptibility to infection. Correlation of human infection with sylvatic plague in Transbaikalia and Mongolia was well recognized in 1850 according to references cited by Pollitzer (1934). At that time the susceptible Siberian marmot was implicated in plague epidemics. A hypothesis pointing to wild rodents as the reservoir of sylvatic plague was proposed in 1910 by McCoy (quoted by Pollitzer 1934 p. 263). After finding insusceptible ground squirrels in plague foci in California he concluded that this partially resistant race of rodents will if not vigorously attacked perpetuate the disease for many years. Later from an extensive series of investigations on California ground squirrels Meyer (1942a) explained that at the apparent termination of the epizootics both susceptible and resistant rats or squirrels may survive and interbreed furnishing sufficient hosts to maintain the infection in a smoldering non readily recognizable state. Recently Baltazard and his associates (1952) were satisfied that in Iranian Kurdistan the *Meriones* subspecies which were somewhat resistant to infection with *Y. pestis* were an important reservoir of the disease. Some investigators (Meyer 1942b, Tinker and Kalabukhov 1934) found that young squirrels were highly susceptible to laboratory plague infections.

That rodent species somewhat resistant to infection are important reservoirs of sylvatic plague has been confirmed in our laboratory and we believe that the highly prevalent and resistant meadow vole *Microtus californicus* is a significant factor in the sylvatic plague reservoir in the San Francisco Bay area. However it is not yet known what this resistance is or how it fits into the ecologic puzzle. Also it is not known what part is played by other rodents some of which are highly susceptible but may have an increased resistance during estivation or hibernation. Some species have been found infected with *Y. pestis* and thus may serve as temporary reservoirs. The relative susceptibility to the plague bacillus has been determined for only a few of these rodent species.

Recent studies were directed toward elucidating the susceptibility to *Y. pestis* infection of all the species of rodents within relatively small areas. In Santa Fe N. M. (Holdenried and Quan 1956) about 400 individuals of 21 wild rodent species were compared with a highly susceptible strain of white laboratory mice by intracutaneous inoculation of 0.05 ml. of aqueous 1 per cent peptone dilutions of a virulent strain (Alexander) isolated earlier from a human case in New Mexico (Link 1950b). In the San Francisco Bay area (Quan unpublished data) about 1100 animals of 5 wild rodent and 2 feral domestic species were identically compared but in San Francisco guinea pigs and white laboratory rats (Sprague Dawley strain) were used as controls in addition to white mice. The intracutaneous route of inoculation was employed in these tests because the investigators felt that this route best simulated the bite of the flea, a natural method of plague infection.

Although the rodent species tested in New Mexico and in California were different in the main one fact was outstanding in the results both areas were inhabited by rodent species varying from highly susceptible to completely resistant individuals (TABLE 2). Holdenried and Quan (1956) have divided these into 4 groups as follows:

TABLE 2
APPROXIMATE PLAGUE LD₅₀S OF SOME OF THE SPECIES OF RODENTS TESTED
IN NEW MEXICO AND CALIFORNIA

Rodent species	Santa Fe N M		San Francisco Calif	
	Not tested	Approx LD	No tested	Approx LD
<i>Citellus variegatus</i>	9	>10 ⁴	0	—
<i>Dipodomys ordii</i>	22	R	0	—
<i>D. spectabilis</i>	13	R	0	—
<i>Microtus californicus</i>	0	—	461	>10 ⁴ <10 ⁵
<i>M. longicaudus</i>	15	>10 ⁴	0	—
<i>Neotoma albigula</i>	32	<10	0	—
<i>Onychomys leucogaster</i>	10	>10 ⁴	0	—
<i>Peromyscus boylii</i>	11	<10 ⁴	0	—
<i>P. leucopus</i>	49	<10	0	—
<i>P. maniculatus</i>	53	<10 ²	273	>10 ⁴
<i>P. truei</i>	74	<10	2	>10 ⁴
<i>Reithrodontomys megalotis</i>	8	<10	148	<10
<i>Thomomys talpoides</i>	9	>10 ⁴	0	—

* R = refractory to intracutaneous inoculations of about 10⁴ *Pasteurella pestis*

Highly susceptible Approximately the same susceptibility as white laboratory mice having an LD₅₀ of less than 10 virulent plague bacilli by intracutaneous inoculation for example *Reithrodontomys megalotis* in both New Mexico and California

Susceptible to slightly resistant The majority died from 1 to 1000 mouse LD₅₀s (mouse ID₅₀ about 10 organisms) for example *Peromyscus maniculatus rufinus*, *P. boylii rouleyi* of New Mexico and *Rattus norvegicus* of California

Moderately resistant to resistant Some deaths with doses greater than 1000 mouse LD₅₀s for example *Microtus longicaudus mordax* and *Onychomys leucogaster pallescens* of New Mexico and *Microtus californicus* and *Peromyscus maniculatus* of California

Highly resistant to refractory No deaths with doses of about 1 million virulent *Pasteurella pestis* for example *Dipodomys ordii medius* in New Mexico. The susceptibility of a rodent species could be estimated and classified in one of the above groups by intracutaneous inoculation of as few as 2 or 3 animals each with undiluted 10⁻³ and 10⁻⁶ dilutions of a 24 hour second transfer brain heart infusion broth culture of a virulent *P. pestis* strain grown at 28° C. Such a culture usually has 100 million to 300 million organisms per milliliter.

An observation of importance made in Santa Fe and in San Francisco was the lack of noticeable die-off of rodents during plague epizootics. When *P. pestis* was found in New Mexico (Holdenried and Morlan 1935) no decrease in rodents per trap per night was detectable. In the San Francisco Bay area epizootic (Kartman *et al.* 1938) *Microtus* were primarily involved but here again there was no decrease in the *Microtus* count.

The finding of more resistant ground squirrels in California plague foci than in plague free areas (Meyer 1942b) is well known. In an attempt to check this variance at Santa Fe four species of rodents (*Neotoma albigula*, *Peromyscus truei*, *P. leucopus* and *P. maniculatus*) from areas with and without plague

TABLE 3

SUSCEPTIBILITY OF *MICROTUS CALIFORNICUS* FROM VARIOUS LOCATIONS NEAR
SAN FRANCISCO CALIF. TO INTRACUTANEOUS INOCULATION OF ABOUT ONE
MILLION PLAGUE ORGANISM *PASTEURRELLA PESTIS*

Host locality	Plague found	Number tested			Results		
		Males	Females	Total	Males	Females	Total
San Francisco game refuge	+	12	18	30	12.5	12.8	12.6
Central Hog Farm	+	40	40	80	12.5	10.3	11.2
Colma Creek	?	19	31	50	15.8	9.7	12.0
Assoc. Hog Farm	?	11	19	30	21.2	21.1	23.4
South San Francisco	—	49	54	103	6.5	64.8	66.0

were compared. No difference was observed in the degree of susceptibility of the comparative groups of these rodent species. It is yet impossible to explain precisely these contradictory results. One reason may be the difference in species employed. Another possible explanation is that plague free or plague positive areas in Santa Fe could not be delineated exactly. The importance of geographic separation is indicated (TABLE 3) by the finding that in a well isolated area in south San Francisco where *Pasteurella pestis* has not been found *Microtus* were susceptible to plague infection (Quan and Kartman 1956). It should also be pointed out that in the California plague foci studied by Meyer (1942b) and by the present authors *P. pestis* has been found many times and probably has a continuous existence while the status of plague in northern New Mexico has not been determined.

In spite of reports of varying degrees of infection in *Meriones* (Iobanov and Federov 1938) and in *Suslik* (Gaiski 1926) during different times of the year Davis (1948) stated that rodent epizootics whether in wild or domestic species do not show any marked seasonal incidence. Our data (TABLE 4) substantiate this observation since no significant seasonal differences were noted in tests with three species of wild rodents at different times of the year. On the other hand studies at the Hooper Foundation (Meyer unpublished data) have shown that guinea pigs were more resistant to experimental plague from October to March than at other times. Evidence of hibernation as a factor in the prolongation or modification of *P. pestis* infections in rodents also needs further elucidation (see Pollitzer 1954 for review). The general importance and specific effects of season on wild rodent plague is but poorly understood at present.

The factors of age and sex of host also need further study. Meyer (1942b) reported that young ground squirrels were most susceptible, adult males less so, and adult females least so. Pollitzer (1954) felt that the higher susceptibility in young wild rodents was more apparent than real since it was possible that the higher resistance of adults was due to hibernation. Also in contrast to the above Tinker and Kalabukhov (1934) found adult males more resistant than adult females. Our own results (TABLE 5) with *Microtus* do not show any marked differences either in the age (based on body length) or in the sex of the

TABLE 2
APPROXIMATE PLAGUE LD₅₀S OF SOME OF THE SPECIES OF RODENTS TESTED
IN NEW MEXICO AND CALIFORNIA

R d t s p c	Santa Fe N M		S F Calif	
	N t t e l	Appro LD	N t t d	App ox LD ₅₀
<i>Citellus variegatus</i>	9	>10 ⁴	0	—
<i>Dipodomys ordii</i>	22	R*	0	—
<i>D. spectabilis</i>	13	R	0	—
<i>Microtus californicus</i>	0	—	461	>10 ⁴ <10 ⁵
<i>M. longicaudus</i>	15	>10 ⁴	0	—
<i>Neotoma albigula</i>	32	<10	0	—
<i>Onychomys leucogaster</i>	10	>10 ⁴	0	—
<i>Peromyscus boylii</i>	11	<10 ⁴	0	—
<i>P. leucopus</i>	49	<10	0	—
<i>P. maniculatus</i>	53	<10	273	>10 ⁴
<i>P. truei</i>	74	<10	2	>10 ⁴
<i>Reithrodontomys megalotis</i>	8	<10	148	<10
<i>Thomomys talpoides</i>	9	>10 ⁴	0	—

* R = refractory to intracutaneous inoculations of about 10⁴ *Pasteurella pestis*

Highly susceptible Approximately the same susceptibility as white laboratory mice having an LD₅₀ of less than 10 virulent plague bacilli by intracutaneous inoculation for example *Reithrodontomys megalotis* in both New Mexico and California

Susceptible to slightly resistant The majority died from 1 to 1000 mouse LD₅₀s (mouse LD₅₀, about 10 organisms) for example *Peromyscus maniculatus rufinus* *P. boylii rouleyi* of New Mexico and *Rattus norvegicus* of California

Moderately resistant to resistant Some deaths with doses greater than 1000 mouse LD₅₀s for example *Microtus longicaudus mordax* and *Onychomys leucogaster pallescens* of New Mexico and *Microtus californicus* and *Peromyscus maniculatus* of California

Highly resistant to refractory No deaths with doses of about 1 million virulent *Pasteurella pestis* for example *Dipodomys ordii medius* in New Mexico. The susceptibility of a rodent species could be estimated and classified in one of the above groups by intracutaneous inoculation of as few as 2 or 3 animals each with undiluted 10⁻³ and 10⁻⁶ dilutions of a 24 hour second transfer brain heart infusion broth culture of a virulent *P. pestis* strain grown at 28° C. Such a culture usually has 100 million to 300 million organisms per milliliter.

An observation of importance made in Santa Fe and in San Francisco was the lack of noticeable die off of rodents during plague epizootics. When *P. pestis* was found in New Mexico (Holdenried and Morlan 1955) no decrease in rodents per trap per night was detectable. In the San Francisco Bay area epizootic (Kartman *et al.* 1958) *Microtus* were primarily involved but here again there was no decrease in the *Microtus* count.

The finding of more resistant ground squirrels in California plague foci than in plague free areas (Meyer 1942b) is well known. In an attempt to check this variance at Santa Fe four species of rodents (*Neotoma albigula* *Peromyscus truei* *P. leucopus* and *P. maniculatus*) from areas with and without plague

are presented by Iollitzer (1954) and by Machiavello (1954) who list locality hosts and epidemiologic findings. There are certain important ecologic relationships that affect a flea's importance in connection with plague and these relationships are observable mainly in the field. Some of these are (1) the natural plague infection rate of different flea species (2) flea host relationships (3) flea nest relationships (4) abundance of different species seasonally on hosts and in nests (5) geographic location of fleas and finally (6) the natural systematic relationships of flea complexes or individual species to plague vector capacity.

On a world wide basis the following are some of the fleas thought to be important in plague (Costa Lima and Hathaway 1946; Jellison and Cood 1942; Jellison *et al* 1953; Machiavello 1954; Pollitzer 1954): *Dinopsyllus ellobius*, *Neopsylla setosa*, *Lepopsylla segnis*, *Diamanus montanus*, *Xosopsyllus fasciatus*, *Orchopeas sexdentatus*, *Iolygenis greyii*, *Ctenocephalides felis*, *Echidnophaga gallinacea* and in the genera *Thrassis* (particularly *acamantis* and *bacchi*), *Oropsylla* (particularly *idahoensis* and *silantieri*), *Opisocrostitis*, *Malaraeus*, *Monopsyllus* (particularly *anisus* and *cagneri*) and *Xenopsylla* (particularly *cheopis astia* and *brasiliensis*). Recently Blanc (1956) has restated the case for *Pulex irritans* as an important vector in certain regions. This list contains but a fraction of the number reported in laboratory and field investigations concerned with plague. It would probably be cut in half and then again enlarged by the addition of many different fleas if the actual ecologic importance of each flea to plague could be considered and compared. Kartman (1954) has proposed a guiding model in this direction by suggesting a method that combines field and laboratory findings to provide a single comparative rating. Although detailed data are required for such information they are justified for those flea species implicated in wild rodent plague ecology.

During recent work in the ecologically and geographically limited area around San Francisco certain species were studied quite exhaustively in relation to plague. Some of the findings are presented in detail by Miles *et al* (1951) dealing with a preliminary study of 13 areas in the vicinity of San Francisco Bay and by Stark *et al* (1958) reporting an intensive ecologic study of one area where plague was found. TABLE 6 (compiled from data of these studies) shows flea relationships with the hosts from which they were taken. It should be noted that these data present figures on the percentage of occurrence of a given flea species (Miles *et al* 1957). This was defined as the index of a given flea on a particular host as the percentage of all flea host indices of that flea. The resultant numerical values compensate for the disproportionate numbers of hosts obtained thereby modifying the significance of the usual index (mean number of fleas per host) and in addition may be somewhat diagnostic of the host preference of the flea. A simple listing of host flea records (as for instance *Microtus californicus* with ten-odd species of fleas) only serves to clutter current literature without providing significant information. It is felt that the number of species of fleas on a given host is directly proportionate to the frequency of capture of the host involved and this must weight the literature considerably. Of concern to plague ecology studies are the fleas that appear in large numbers. Vertical and horizontal

TABLE 4
MORTALITY FROM PLAGUE OF 3 SPECIES OF RODENTS TESTED AT DIFFERENT MONTHS OF
THE YEAR IN SAN FRANCISCO CALIF

Month y of test	Number dying/ number tested and death rate among indicated species							
	Mortality				Percentage		Risk of mortality	
	moderately		susceptible		moderately		susceptible	
February 1955	8/44	0.18	—	—	2/17	0.12	3/4	0.75
January 1957	2/31	0.06	6/7	0.86	5/25	0.20	8/11	0.73
March 1957	11/43	0.25	—	—	9/49	0.18	3/9	0.33
April 1955	5/24	0.21	—	—	1/9	0.11	7/10	0.70
May 19 4 1955	5/19	0.26	17/28	0.61	1/34	0.03	20/36	0.56
June 1955 1956	14/42	0.33	30/39	0.72	—	—	8/13	0.67
July 1956	3/19	0.16	4/4	1.00	1/10	0.10	4/12	0.33
August 1955	2/8	0.25	10/21	0.48	7/32	0.22	3/7	0.43
October 1954	5/42	0.12	—	—	4/46	0.09	22/25	0.88
November 1954	2/17	0.12	—	—	2/9	0.22	2/8	0.25
December 1956	23/68	0.34	4/5	0.80	2/42	0.05	6/13	0.46
Total	80/357	0.22	71/104	0.68	34/273	0.12	86/148	0.56

All rodents received 0.05 ml intracutaneously containing varying numbers of *Pasteurella pestis*

TABLE 5
SUSCEPTIBILITY TO EXPERIMENTAL PLAGUE OF *MICROTUS CALIFORNICUS* ACCORDING TO
SEX AND AGE

Sex	Body length (mm) from top of ear to tip of tail														
	105 to 130			131 to 150			151 to 170			171 to 190			All length		
	N	dying	Percent dying	N	dying	Percent dying	N	dying	Percent dying	N	dying	Percent dying	N	dying	Percent dying
Male	13	4	12.2	68	20	12.1	90	20	10.6	27	10	34.4	198	54	12.2
Female	20	5	15.3	98	19	11.5	99	31	16.4	2	1	3.4	219	56	13.4
Totals	33	9	27.2	166	39	23.4	189	51	26.9	29	11	37.9	417	110	26.3

* As determined by body length

rodent. However, some of the groups tested had too few individuals to furnish conclusive results.

Flea Species Involved

a priori, every flea is a potential vector, whether a particular species of flea is actually a significant vector is another matter (Jordan 1948). Tables of flea species implicated with plague in nature and in the laboratory

are presented by Iollitzer (1954) and by Machiavello (1954) who list locality hosts and epidemiologic findings. There are certain important ecologic relationships that affect a flea's importance in connection with plague and these relationships are observable mainly in the field. Some of these are (1) the natural plague infection rate of different flea species (2) flea host relationships (3) flea nest relationships (4) abundance of different species seasonally on hosts and in nests (5) geographic location of fleas and finally (6) the natural systematic relationships of flea complexes or individual species to plague vector capacity.

On a world wide basis the following are some of the fleas thought to be important in plague (Costa Lima and Hathaway 1946; Jellison and Cood 1942; Jellison *et al* 1953; Macchiavello 1954; Pollitzer 1954): *Dinopsyllus ellobius*, *Xenopsylla setosa*, *Leptopsylla segnis*, *Diamanus montanus*, *Xenopsyllus fasciatus*, *Orchopeas sergentatus*, *Iolygenis gwyni*, *Ctenocephalides felis*, *Echidnophaga gallinacea* and in the genera *Thraassis* (particularly *acamantis* and *bacchi*), *Oropsylla* (particularly *idahoensis* and *silantietti*), *Opisocrostitis*, *Malaraeus*, *Monopsyllus* (particularly *anisus* and *agneri*) and *Xenopsylla* (particularly *cheopis asiatica* and *brasiliensis*). Recently Blanc (1956) has restated the case for *Pulex irritans* as an important vector in certain regions. This list contains but a fraction of the number reported in laboratory and field investigations concerned with plague. It would probably be cut in half and then again enlarged by the addition of many different fleas if the actual ecologic importance of each flea to plague could be considered and compared. Kartman (1951) has proposed a guiding model in this direction by suggesting a method that combines field and laboratory findings to provide a single comparative rating. Although detailed data are required for such information they are justified for those flea species implicated in wild rodent plague ecology.

During recent work in the ecologically and geographically limited area around San Francisco certain species were studied quite exhaustively in relation to plague. Some of the findings are presented in detail by Miles *et al* (1957) dealing with a preliminary study of 13 areas in the vicinity of San Francisco Bay and by Stark *et al* (1958) reporting an intensive ecologic study of one area where plague was found. TABLE 6 (compiled from data of these studies) shows flea relationships with the hosts from which they were taken. It should be noted that these data present figures on the percentage of occurrence of a given flea species (Miles *et al* 1951). This was defined as the index of a given flea on a particular host as the percentage of all flea host indices of that flea. The resultant numerical values compensate for the disproportionate numbers of hosts obtained thereby modifying the significance of the usual index (mean number of fleas per host) and in addition may be somewhat diagnostic of the host preference of the flea. A simple listing of host flea records (as for instance *Microtus californicus* with ten odd species of fleas) only serves to clutter current literature without providing significant information. It is felt that the number of species of fleas on a given host is directly proportionate to the frequency of capture of the host involved and this must weight the literature considerably. Of concern to plague-ecology studies are the fleas that appear in large numbers. Vertical and horizontal

TABLE 6

HOST RELATIONSHIPS OF FLEAS COLLECTED FROM MAMMALS IN 13 AREAS IN THE SAN FRANCISCO BAY AREA

Host class	S p		Citellus b h y		Reithrodontomys m g f c s		Peromyscus m l a t s		Microtus f v c s		Reithrodontomys m g f c s		Microtus m c f	
Number examined	110		23		454		53		244		273		23	
Flea species	F l i n d e x		P e r c e n t o c c u r r e n c e		F l i n d e x		P e r c e n t o c c u r r e n c e		F l i n d e x		P e r c e n t o c c u r r e n c e		F l i n d e x	
	F l i n d e x	P e r c e n t o c c u r r e n c e	F l i n d e x	P e r c e n t o c c u r r e n c e	F l i n d e x	P e r c e n t o c c u r r e n c e	F l i n d e x	P e r c e n t o c c u r r e n c e	F l i n d e x	P e r c e n t o c c u r r e n c e	F l i n d e x	P e r c e n t o c c u r r e n c e	F l i n d e x	P e r c e n t o c c u r r e n c e
<i>Hystrichopsylla linsdalei</i>					0 02	3 0 08 14	0 43 74	0 05 9 0						
<i>Athyloceras m multidentatus</i>					0 08	14 0 09 16	0 27 47	0 14 24 0						
<i>Corrodopsylla c obliuata</i>	0 14 91													
<i>Catallagia wymani</i>	0 10 8	0 04	3 0 07	6 0 38 31	0 48 38	0 16 13 0 0 01	1 0							
<i>Leptopsylla segnis</i>						0 01 3 0 01	3 0 25 76	0 06 18 0						
<i>Diamanus montanus</i>		32 2 100												
<i>Xenopsyllus fasciatus</i>	0 01 1	0 09	6 0 03	2 0 07 5 0 07	5 1 10 76	0 07 5 0								
<i>Opis dasys k nesiotus</i>	0 04 2		0 05	3 1 60 92	0 03 2	0 3								
<i>Monopsyllus x phidius</i>				0 08 64	0 01 8 0 03 24	0 4 0								
<i>Malaraeus telchitum</i>	0 16 3	0 4	1 0 08	1 0 57 11 4 0	4 0 48 9 0 0 07	1 0								
<i>Hoplopsyllus anomalus</i>		5 8 100												
<i>Xenopsylla cheopis</i>							3 0 16 94 0	3 0						

Value less than 0.005 never more than 1 flea involved

comparisons of the percentage of occurrence (TABLE 6) provide an indication of host specificity of the flea (or the degree of acceptability of the host) and numbers of flea species with high occurrences. The percentage of occurrence varied slightly when data from different areas were compared. There was however a definite limit to the range of the percentage of occurrence for example that of *Opisodasys nesiotus* on *Peromyscus* was always observed to be 90+.

Specific host flea relationships in the San Francisco Bay region may now be discussed. Fleas such as *Clenocephalides felis felis* and *Dactylopsylla ignota franciscana* were taken in low numbers but since their normal hosts (that is the domestic cat and the pocket gopher) were not considered for computations presented here confusing values would be obtained on hosts that are presented. For this reason they are not included in TABLE 6.

Hystrichopsylla linsdalei. This flea was found in great concentration on *Microtus* in all areas studied and from field and laboratory findings it was considered to be the most important flea in wild rodent epizootics in the San Francisco Bay area. TABLE 7 shows it to have the highest percentage of plague positives in the field. In the laboratory this flea was found to have an experimental vector index of 0.02 as compared with 0.06 for *Xenopsylla cheopis* (Kartman and Prince 1956). Stark et al. (1958) present a vector potential computed from the vector index field infection index and

TABLE 7

NATURAL INFECTION RATES OF VARIOUS FLEA SPECIES DURING A PLAGUE
EPIZOOTIC AMONG FIELD MICE MARCH 23 TO APRIL 30 1954

Chest week N	Infestation of field mice													
	<i>Malpue</i>		<i>Hoplit</i>		<i>Allyphl</i>		<i>C. llyt</i>		<i>Op d y</i>		Oth		All pec	
	N	Per t pos	N	P t pos	N	P t pos	N	P t p	N	P t pos	N	P t pos	N	P t pos
1	316	10.1	42	23.8	26	0	26	3.8	12	25.0	3	0	475	10.8
2	421	5.7	61	9.8	30	3.3	45	0	1	0	0	—	55	5.6
3	257	4.2	41	14.6	21	0	25	0	9	0	0	—	353	4.8
4	190	4.4	28	7.1	17	5.8	30	0	1	0	0	—	256	4.3
5	132	0	28	10.7	16	0	19	0	10	0	1	0	206	1.5
6	251	0	47	2.1	36	2.7	53	0	13	0	4	0	404	0.5
Totals	1557	4.8	247	11.3	146	2.0	198	0.5	46	6.5	8	0	2207	5.0

Mainly Nosopsyllus fasciatus

field prevalence index which was greater than that of any of the other fleas encountered. This flea appeared less likely to transfer to other hosts such as rats than did some of the other flea species.

Allyphloceras m. multidentatus This species was fairly abundant and showed a field infection index below that of *Hystrichopsylla* and *Malariaeus* (TABLE 7). However these data were based on a six week period. Studies carried on for a whole year with additional plague findings gave *m. multidentatus* a higher rate than *Malariaeus*.

Catallagia wymani In one study the number of individuals of this flea exceeded that of *H. linsdalei*. However this flea was restricted to a more limited type of environment than other species and was entirely absent in some areas where plague appeared. The percentage of occurrence for *C. wymani* indicated that this flea had the widest range of acceptable hosts; therefore it may be of considerable importance in transfer between rodents. The field infection rate was exceedingly low and laboratory findings (Eskey and Haas 1940) indicated that it was probably not a vector of importance.

Diamanus montanus This flea has been found to be an efficient vector of plague (Eskey and Haas 1940; Wheeler and Douglas 1941; Burroughs 1947; Kartman and Prince 1956). Its high index on *Citellus* not only in the San Francisco Bay area but in 11 western states indicates that this flea is of importance in plague epizootics among the ground squirrels. However its importance may be restricted because of its high degree of host specificity. Even in areas where opportunity for transfer was afforded (for example to *Rattus* which is an acceptable host to many flea species) the percentage of occurrence and high index (TABLE 6) indicated a rather unusual host preference. Nevertheless the work of Meyer and Holdenried (1949) shows that this species on occasion may transfer plague to domestic rats.

Nosopsyllus fasciatus The northern rat flea is a fairly efficient experimental

vector of plague (Eskey and Haas 1940 Burroughs 1947), being approximately equal in this respect to *H. linsdalei*. In the areas studied in the vicinity of San Francisco its numbers were low, and it was not found to be infected in the plague areas where it occurred. However, because of its wide range of acceptable hosts as indicated by its percentage of occurrence it may be potentially of importance in any part of the San Francisco Bay area, where a build up of the rat population would increase the incidence of this flea. In a recent survey of the city of San Francisco a general index of 1.5 *X. fasciatus* per rat was evident. However, the purpose of this survey was to avoid a general index and to seek out rather the index in relation to specific factors such as season, habitat of the rat and section of the city. Surprisingly high indices were found in some sections of the city which indicated that favorable conditions still persist in some areas for the propagation of these fleas and their rat hosts despite recent socioeconomic improvements.

Opisodasys keeni nestolus. This flea was of some importance in the plague picture since it is a known experimental vector and because it had a high field infection rate (TABLE 7). However Stark *et al* (1958) reported a rather low vector potential due to its scarcity in the areas studied. Like *D. montanus* it appeared restricted by a strong host specificity. It may be of importance to plague among small rodents whenever its principal host *Peromyscus* sp. is present in fairly large numbers.

Monopsyllus agneri ophidius. The very low index of this flea in the San Francisco Bay area was rather surprising since it is a major flea of several rodents over much of the United States. This finding adds one more factor to the uniqueness of this section of Pacific Coast flora and fauna.

Malariaeus telchinum. This species was the most abundant flea in all areas studied. Its principal host *Microtus* occurred in rather limited ecologic environments and the flea appeared to be restricted by the host's distribution. In other parts of the United States *Peromyscus* is the principal host. Populations of *Microtus* are always quite dense and therefore are conducive to a high flea population. The range of acceptable hosts and the degree of host specificity of *M. telchinum* were similar to those of *Histricopsylla linsdalei*. Although laboratory findings indicated that these fleas are poor individual vectors at best our field studies revealed a fairly high infection rate. This fact taken in conjunction with evidence of mass transmission in the laboratory (Burroughs 1944 1947) suggests that this species may be an important secondary vector in the field.

Hoplopsyllus anomalus. This flea was found to be naturally infected with plague in New Mexico (Holdenried and Morlan 1955) and is able to transmit plague (Eskey and Haas 1940) but is probably a poor vector (Wheeler and Douglas 1941). Like *Diamanus montanus* it was restricted to its host because of a high degree of specificity.

The remaining fleas *Corrodopsylla curiata obtusata* and *Xenopsylla cheopis* were taken in very low numbers on their preferred hosts *Sorex* and *Rattus* respectively. Whether or not these two species of fleas are of importance to sylvatic plague cannot be ascertained from present studies. The occurrence of these two species in sylvan areas may be of significance. *X. cheopis* oc.

curted in large numbers on urban rats in San Francisco being a very close second to *Xenopsyllus fasciatus*. A recent survey of San Francisco rat fleas substantiated previous findings that *X. cheopis* occurred more frequently under strictly urban conditions while the reverse was true for *X. fasciatus*.

In general there seemed to be very little correlation between the seasonal incidence of fleas on their hosts and the occurrence of plague. *H. linsdalei* had a uniform incidence on its preferred host *Microtus californicus* throughout the year. The incidence of *Malariaeus telchinum* followed fairly closely that of its host *Microtus* as indicated by trapping results. Increases in flea indices followed shortly behind the increase of host prevalence. The peak of *M. telchinum* occurred during and following the warmer and drier season of the bay region. In contrast *Catallagia wymani* had its greatest incidence during winter and spring and its lowest following the drier season. There were insufficient data to consider seasonal incidence for other fleas.

The numbers of fleas on the host tell very little about what occurs in the host's nest where the adult fleas seem to spend a good deal of their time and on which the immature stages are completely dependent. Bacot and Martin (1914) when writing up their remarkable discovery of blocking in fleas theorized on the effects of adverse climatic conditions on blocked fleas. Other workers ascribed the trend of plague seasons to the influence of climatic conditions on fleas (Petrie 1939). However studies of these effects on fleas and plague have not progressed very far. Hurst (1926) and Grubbs (1924) have postulated that even though an infection is present there is little danger of epidemic spread as long as the *Xenopsylla cheopis* index remains below 1. Stewart and Evans (1941) considered that the value of flea surveys in the accepted sense of collecting fleas from the rodents themselves might be enhanced by obtaining the composition of the flea population at the openings of wild rodent burrows. Davis (1939) and Macchiavello (1948) have recommended that the total or absolute numbers of fleas should be determined for surveys by the examination of burrows and nests. Consequently some workers have advocated the use of a formula in which the absolute flea index is obtained by dividing the total number of rats into the total number of fleas found on rodents and in their nests. It must be emphasized here that ecologic studies of plague in the field go beyond the survey method. Other types of data are needed especially in relation to the life of the flea vector in the host's nest. For instance we have found that the optimum quantity of moisture in nesting material was between 50 and 100 per cent as indicated by numbers of fleas per nest. In nature nests most frequently showed the lowest moisture range (10 to 30 per cent) and least frequently the highest range (70 to 90 per cent). *Catallagia wymani* flourished under relatively high moisture conditions while *Malariaeus telchinum* prevailed in drier nests. This type of micro-environmental investigation may reveal important factors in seasonal plague and the vector potentials of fleas and lead to the development of better methods for flea and rodent control and the laboratory culture of wild rodent fleas.

Finally let us consider briefly the importance of systematics. Some of the material presented in this paper illustrates the need for exact taxonomic identification of fleas as has been repeatedly demonstrated in the past. The

vector of plague (Eskey and Haas 1940 Burroughs 1947) being approximately equal in this respect to *H. linsdalei*. In the areas studied in the vicinity of San Francisco its numbers were low, and it was not found to be infected in the plague areas where it occurred. However, because of its wide range of acceptable hosts as indicated by its percentage of occurrence it may be potentially of importance in any part of the San Francisco Bay area where a build up of the rat population would increase the incidence of this flea. In a recent survey of the city of San Francisco a general index of 15 *X. fasciatus* per rat was evident. However the purpose of this survey was to avoid a general index and to seek out rather the index in relation to specific factors such as season habitat of the rat and section of the city. Surprisingly high indices were found in some sections of the city which indicated that favorable conditions still persist in some areas for the propagation of these fleas and their rat hosts despite recent socioeconomic improvements.

Opisodasys keeni nestotus. This flea was of some importance in the plague picture since it is a known experimental vector and because it had a high field infection rate (TABLE 7). However Stark *et al* (1958) reported a rather low vector potential due to its scarcity in the areas studied. Like *D. montanus* it appeared restricted by a strong host specificity. It may be of importance to plague among small rodents whenever its principal host *Peromyscus* sp. is present in fairly large numbers.

Monopsyllus uagneri ophidius. The very low index of this flea in the San Francisco Bay area was rather surprising since it is a major flea of several rodents over much of the United States. This finding adds one more factor to the uniqueness of this section of Pacific Coast flora and fauna.

Malaraeus telchinum. This species was the most abundant flea in all areas studied. Its principal host *Microtus* occurred in rather limited ecologic environments and the flea appeared to be restricted by the host's distribution. In other parts of the United States *Peromyscus* is the principal host. Populations of *Microtus* are always quite dense and therefore are conducive to a high flea population. The range of acceptable hosts and the degree of host specificity of *M. telchinum* were similar to those of *Histricopsylla linsdalei*. Although laboratory findings indicated that these fleas are poor individual vectors at best our field studies revealed a fairly high infection rate. This fact taken in conjunction with evidence of mass transmission in the laboratory (Burroughs 1944 1947) suggests that this species may be an important secondary vector in the field.

Hoplopsyllus anomalus. This flea was found to be naturally infected with plague in New Mexico (Holdenried and Morlan 1955) and is able to transmit plague (Eskey and Haas 1940) but is probably a poor vector (Wheeler and Douglas 1941). Like *Diamanus montanus* it was restricted to its host because of a high degree of specificity.

The remaining fleas *Corrodopsylla curvata obtusata* and *Xenopsylla cheopis* were taken in very low numbers on their preferred hosts *Sorex* and *Rattus* respectively. Whether or not these two species of fleas are of importance to sylvatic plague cannot be ascertained from present studies. The occurrence of these two species in sylvan areas may be of significance. *X. cheopis* oc




FIGURE 4 The cumulative blocking rate of fleas *Xenopsylla vexabilis hawaiiensis* and *X. cheopis* which were infected with *Pasteurella pestis* at various days after the infectious blood meal

More recently Kartman Prince and Quan (1956) and Kartman and Prince (1956) have utilized an additional component the blocking survival potential to derive a vector index. It is based on a ratio of the mean day of death of fleas after blocking over the mean day of blocking after the infectious meal. This potential was suggested by the fact that the speed of blocking and the survival of the flea after blocking are critical factors affecting biological transmission of plague (FIGURES 4 and 5). The shorter the incubation period and the longer the survival after blocking the greater the blocking survival potential. The vector index is the product of the vector efficiency and the blocking survival potential (or the product of the three potentials of Wheeler and Douglas and the blocking survival potential). As an example TABLE 8 presents data from experimental tests with the native Hawaiian flea compared




FIGURE 5 The longevity of fleas after blocking with *Pasteurella pestis*

Plague Commission in India (1904 to 1907) found that rat fleas were involved in transmission of plague, but was unable to determine that two prevalent but similar species (*Xenopsylla cheopis* and *X. astia*) were present in the endemic plague area. Fluctuations in the occurrence of human plague were not explained until these fleas had been identified and their nature determined. Of great significance was the fact that these fleas differed greatly in their ability to transmit and in their seasonal occurrence. Today confusion still exists as to the identity of *Thrassitis howelli* and *T. stanfordi* and as to which species was actually involved in plague transmission experiments (reported in Eskey and Haas 1940 Macchiavello 1954 and Pollitzer 1954). In our field studies a flea of the genus *Hystriechopsylla* was encountered that had a higher natural infection rate and was a more efficient transmitter of plague than any other wild rodent flea. Exact taxonomic evaluation of this species showed that it had previously entered the literature under a misnomer (Burroughs 1944 Snodgrass 1946 Hubbard 1947). The flea was in fact a new species and observations were made on it for more than four years before its specific determination as *Hystriechopsylla linsdalei* was made in 1957 (Holland 1957).

Vector Efficiency*

During the course of investigations of plague by many workers the concept of vector capacity of separate flea species became of paramount concern. When the importance of strict taxonomic classification of the flea species had been realized the role of these species as vectors became a critical problem in studies of plague epidemiology. The classic work of the Plague Research Commission in India (1906 to 1915) constitutes the basic knowledge of the transmission of plague by the rat flea. The work of Bacot and Martin (1914) gave a conclusive demonstration of a mechanism of the infectious process in the flea and a mode of plague transmission by this insect.

Wheeler and Douglas (1941-1945) attempted to assess the importance of several factors that when combined would give a more exact picture of the vector efficiency of individual fleas of a single species. They set up a mathematical model in which the factors could be combined in a product symbolic of plague vector capacity. These authors postulated three potentials for deriving vector efficiency: (1) the infection potential is the proportion of fleas becoming infected after feeding on an animal with plague septicemia; (2) the vector potential is the proportion of infected fleas that become infective or blocked (Kartman *et al.* 1956) and (3) the transmission potential is the mean number of transmissions accomplished by the infective fleas each of which is fed daily upon a laboratory mouse. The vector efficiency is the product of these three potentials and represents the mean number of plague transmissions per flea. It should be noted in the words of Wheeler and Douglas (1945) that the infection and vector potentials actually do not enter into the determination of vector efficiency; however they are included to emphasize the fallacy of estimating the efficiency of a vector solely on the basis of the number that became infected or infective.




FIGURE 4 The cumulative blocking rate of fleas *Venopsylla vexabilis hawaiiensis* and *X. cheppis* which were infected with *Pasteurella pestis* at various days after the infectious blood meal

More recently Kartman, Prince, and Quan (1956) and Kartman and Prince (1956) have utilized an additional component—the blocking-survival potential—to derive a vector index. It is based on a ratio of the mean day of death of fleas after blocking over the mean day of blocking after the infectious meal. This potential was suggested by the fact that the speed of blocking and the survival of the flea after blocking are critical factors affecting biological transmission of plague (FIGURES 4 and 5). The shorter the incubation period and the longer the survival after blocking, the greater the blocking-survival potential. The vector index is the product of the vector efficiency and the blocking-survival potential (or the product of the three potentials of Wheeler and Douglas and the blocking-survival potential). As an example, TABLE 8 presents data from experimental tests with the native Hawaiian flea compared




FIGURE 5 The longevity of fleas after blocking with *Pasteurella pestis*

TABLE 8

SUMMARY DATA FOR THE DETERMINATION OF PLAGUE VECTOR EFFICIENCY INDICES

Flea species	Y. <i>psyll</i> <i>g</i> <i>h</i> <i>u</i> <i>e</i> <i>s</i>			Y. <i>he</i> <i>p</i> <i>s</i>		
	M	I	M + I	M	I	M + I
Sex						
Number used	29	31	60	21	9	30
Number infected	23	28	51	21	8	29
Number blocked	21	23	44	16	1	23
Number of transmissions	9	11	20	12	9	21
Blocking survival ratio†	3 5/14 5	3 3/22 4	3 4/18 9	3 9/20 3	3 0/30 9	3 7/23 5
Infection potential	0 79	0 90	0 85	1 00	0 88	0 96
Blocking potential	0 91	0 82	0 86	0 76	0 87	0 79
Transmission potential	0 47	0 47	0 45	0 75	1 28	0 91
Vector efficiency‡	0 30	0 35	0 33	0 51	0 99	0 69
Blocking survival potential	0 24	0 14	0 17	0 18	0 09	0 15
Vector index	0 07	0 05	0 05	0 11	0 09	0 10

From Kartman *et al* 1956

† Mean day of death after blocking/mean day of blocking after infectious meal

‡ Vector efficiencies calculated according to the method of Wheeler and Douglas 1945

with the classic vector *Xenopsylla cheopis* (Kartman *et al* 1956). The vector indices suggest the differences in plague vector capacity of the two flea species under experimental conditions. When atmospheric conditions plague strains and the fleas used are carefully controlled, replications of this type of laboratory test will result in fairly consistent data.

The central problem lies in the circumstance that the vector efficiency and the vector index as now used represent only a limited number of the factors actually involved in a total concept of vector capacity. The actual need is for an index that will include pertinent data from field studies. Few investigators would cavil at the suggestion that knowledge of the total ecology of a flea species would aid in evaluating its role as a vector of plague. That is an ideal toward which workers with many arthropod borne diseases aim. Nevertheless, knowledge of detailed ecologic processes accumulates but slowly and the worker must deal with the level of known facts at any particular moment.

Certain types of ecologic data are now fairly easy to obtain and they may be included in the evaluation of plague vector capacity. The prevalence of various flea species or the flea index based on numbers of fleas per rodent is almost a routine observation in field studies. Considerably less attention has been devoted to the flea index in rodent nests. Nevertheless the most realistic flea index is one that combines data from both the rodent and its nest. Furthermore no attempt has been made in plague ecology studies to obtain flea infection rates based upon bacteriological tests of individual fleas taken from rodents and their nests. In recent studies we have paid particular attention to these factors and TABLES 7 and 9 are examples of data obtained. With these factors in mind we have recently suggested and have used a model (Kartman 1957) for deriving categorical values that compare flea species in a hierarchy of relative importance in plague epizootiology. This scheme at

TABLE 9

PLAGUE INFECTION IN FLEAS FROM NESTS AND FROM HOSTS (JANUARY 1955)

Flea species	Flea species			
	<i>M. t. aureus</i> t. t. h. m.	<i>H. t. h. p. yll</i> l. d. l. o. y.	<i>A. t. phl</i> m. l. i. d. e. n. t. a. l.	<i>C. t. h. g.</i> s. y. m.
Mean fleas per nest	19.0	5.1	6.7	9.7
Mean fleas per host	4.7	0.6	0.4	0.6
Nest fleas per cent infected	4	33	15	0
Host fleas per cent infected	7	14	23	25

tempts to combine laboratory with field data in the following way: flea vector potential = experimental vector index \times field infection index \times field prevalence index \times 100

Although the numerical values thus derived represent a considerable amount of ecologic observation we must still agree with Girard (1955) that much obscurity still obtains in the problem of vector capacity. Hypothetically it can be postulated that in any set of ecologic conditions governing the epizootic wave or the enzootic trough the plague vector capacity of a flea population is ultimately a function of the hospitality of the individual flea to the plague bacillus. Thus the biochemical physiology of the flea-plague relationship is an absolutely necessary avenue of investigation that has been neglected almost completely. By the use of artificial flea feeding techniques (Kartman 1954) we have found for instance that certain strains of plague will grow normally and produce proventricular blockage in fleas when the bacteria are suspended in rat blood but will not multiply when the medium is rabbit blood. Preliminary studies have shown that blockage in the flea may depend upon the number of bacteria ingested during its infectious feeding; a definite threshold appears to exist. Temperatures of about 20 to 22 C favor the growth of *Yersinia pestis* in the flea, whereas temperatures of 25 C and above are not only unfavorable to the ingested bacilli but also shorten the life of the flea itself. The development of a proventricular block does not follow the same course in different flea species. In *Xenopsylla cheopis* blockage is fairly rapid and a high percentage of individuals show initial development of blockage in the proventriculus. On the other hand bacterial growth in most wild rodent fleas begins in the ventriculus and it may take as long as two months for proventricular blockage to occur under the same conditions.

Although plague transmission by the individual blocked flea is an established mechanism in the natural history of the disease, the phenomenon of mass transmission must be mentioned as another mechanism that has received insufficient attention. As an example it is known from the studies of Burroughs (1944, 1945) and of this laboratory that the wild rodent flea *Malareus telchinum* does not show proventricular blockage under conditions in which other species become blocked. Thus plague transmission by individuals of this species almost never occurs in laboratory tests of vector capacity.

If we compare this flea with another wild rodent species *Hystrichopsylla linsdalei* utilizing the concepts discussed above the following data (Kartman 1957) result

<i>Hystrichopsylla linsdalei</i>		<i>Malareus telchinum</i>	
Experimental vector index	= 0.07	Experimental vector index	= 0
Field infection index	= 0.11	Field infection index	= 0.04
Field prevalence index	= 0.08	Field prevalence index	= 0.43
Vector potential	= 0.02	Vector potential	= 0

Although these data appear significant they mean little, since we know that *M. telchinum* is quite prevalent in the field and has a comparatively high plague infection rate during epizootic conditions (TABLES 7 and 9). A further indication of its importance is obtained when groups of many individuals are allowed to feed *en masse* on clean mice very soon after having fed on a mouse with terminal plague bacteremia. In this case plague transmission is obtained quite consistently. *M. telchinum* is therefore considered an efficient mass transmitter presumably by mechanical transference of the plague bacillus on contaminated mouth parts.

This mechanism of mass transmission is undoubtedly more important in plague ecology than has heretofore been assumed. Our own studies suggest that mechanical transmission by a highly prevalent flea may be the means by which epizootics among wild rodents flare up once they have been initiated by primary vectors that transmit biologically via the blockage mechanism. Experimental studies with *Xenopsylla cheopis* and mice (Quan *et al.* 1953) showed that some of the fleas with at least partially blocked proventriculi were infective in the usual manner. However the work also suggested that during the first few feedings after the infectious blood meal infection of the clean mice took place through the mechanism of plague contaminated mouth parts being washed by saliva. Obviously the phenomenon of mass transmission must be taken into account in the ecology of wild rodent plague. Devignat (1946) working with fleas in the Belgian Congo was the first to attempt a quantitative estimate of mass transmission. Differences in vector capacity were determined by comparing the surface areas of the curves illustrating the mortality of flea infected mice with the surface area of a graph showing the mortality of mice infected with standard doses of *Pasteurella pestis*. Future work is faced with the problem of integrating the concept of mass transmission with current hypotheses regarding the mechanisms involved in sylvatic plague.

Transfer of Fleas Between Hosts

The literature contains numerous references to the so called exchange of fleas between rodent host species yet the evidence has been based primarily upon findings that certain flea species parasitize animals other than their normal or preferred hosts. The concept of preferred hosts is in itself beset with pitfalls for the unwary observer for a considerable number of factors must be known before the parasite host relationship can be evaluated ade-

quately. With respect to fleas such factors as seasonal fluctuations, nest populations, and availability of specific hosts are of critical importance. In personal correspondence H. B. Morlan has discussed the question of host preference in reference to his paper on mammal fleas in New Mexico (Morlan 1955). It is of interest here to quote from his remarks:

Where data from extensive collections of all possible hosts at all seasons of the year are available, the percentage of animals infested with one or more parasites and the mean number of parasites per examined animal are usually sufficient to indicate host preference. In cases of inadequate collections, host preference will remain unknown in spite of any additional manipulation of insufficient data.

My paper on Santa Fe fleas provides an actual example where the mean number of *Opisocrostitis hirsutus* from infested ground squirrels was 10.0 while the number from infested prairie dogs was 7.6. In this case only a few squirrels were collected, mainly in the vicinity of a dog town, and there was enough infestation of squirrels with the normal prairie dog flea (*O. hirsutus*) to give a false impression of host preference.

The records merely indicate a specific instance of an unusual potential means for the spread of disease from prairie dogs to ground squirrels. I am presently

inclined to classify the hosts of an ectoparasite as normal, secondary, and accidental. The normal host is that animal which has been demonstrated to be the most commonly encountered associate. A secondary host is one that is frequently and consistently associated with a particular ectoparasite. There may be more than one secondary host. An accidental host is one that is infrequently and inconsistently found associated with a parasite.

Morlan's remarks have been quoted at length because it is important to make clear-cut distinctions in host preferences when evaluating the problem of flea transfer between host species. In plague epizootiology it is of the utmost significance to determine the preferred host, especially in cases where certain rodent species are known to be the prime reservoirs of plague. Thus in our studies the native meadow vole *Microtus californicus* is thought to be the primary plague reservoir, and this host in most instances is parasitized most frequently and in greatest numbers with those flea species showing the highest plague infection rates. This relationship is true of the rodent nests as well as of the rodents, and any transfer of plague to other wild rodent species or to domestic rats would suggest that the disease had originated in *Microtus* and its fleas.

It should also be noted that, except in clear-cut instances, the finding of a wild rodent flea on a domestic rat, or the presence of a flea on a wild rodent other than its normal host, does not imply transfer of the fleas between hosts. Too little is known of the breeding capacities of wild rodent flea species in the nests of various hosts. In the laboratory we have been able to culture the *Microtus* flea *Malareus telchinum* on albino rats. This suggests that *Rattus norvegicus* in nature could possibly support the breeding of *Malareus telchinum* in its burrow. If this is true, the finding of *M. telchinum* on domestic rats would not necessarily imply transfer from *Microtus*, even if populations of the latter were prevalent in the vicinity of the rats. Recent studies in the city of Tacoma, Wash., showed that *Microtus* nests contained cultures

If we compare this flea with another wild rodent species *Hystrichopsylla linsdalei*, utilizing the concepts discussed above the following data (Kartman 1957) result

<i>Hystrichopsylla linsdalei</i>		<i>Malariaeus telchinum</i>	
Experimental vector index	= 0.02	Experimental vector index	= 0
Field infection index	= 0.11	Field infection index	= 0.04
Field prevalence index	= 0.03	Field prevalence index	= 0.43
Vector potential	= 0.02	Vector potential	= 0

Although these data appear significant they mean little since we know that *M. telchinum* is quite prevalent in the field and has a comparatively high plague infection rate during epizootic conditions (TABLES 7 and 9). A further indication of its importance is obtained when groups of many individuals are allowed to feed *en masse* on clean mice very soon after having fed on a mouse with terminal plague bacteremia. In this case plague transmission is obtained quite consistently. *M. telchinum* is therefore considered an efficient mass transmitter presumably by mechanical transference of the plague bacillus on contaminated mouth parts.

This mechanism of mass transmission is undoubtedly more important in plague ecology than has heretofore been assumed. Our own studies suggest that mechanical transmission by a highly prevalent flea may be the means by which epizootics among wild rodents flare up once they have been initiated by primary vectors that transmit biologically via the blockage mechanism. Experimental studies with *Xenopsylla cheopis* and mice (Quan *et al.* 1953) showed that some of the fleas with at least partially blocked proventriculi were infective in the usual manner. However the work also suggested that during the first few feedings after the infectious blood meal infection of the clean mice took place through the mechanism of plague contaminated mouth parts being washed by saliva. Obviously the phenomenon of mass transmission must be taken into account in the ecology of wild rodent plague. Devignat (1946), working with fleas in the Belgian Congo was the first to attempt a quantitative estimate of mass transmission. Differences in vector capacity were determined by comparing the surface areas of the curves illustrating the mortality of flea infected mice with the surface area of a graph showing the mortality of mice infected with standard doses of *Pasteurella pestis*. Future work is faced with the problem of integrating the concept of mass transmission with current hypotheses regarding the mechanisms involved in sylvatic plague.

Transfer of Fleas Between Hosts

The literature contains numerous references to the so called exchange of fleas between rodent host species yet the evidence has been based primarily upon findings that certain flea species parasitize animals other than their normal or preferred hosts. The concept of preferred hosts is in itself beset with pitfalls for the unwary observer for a considerable number of factors must be known before the parasite host relationship can be evaluated ade-

TABLE 10

TRANSFER OF Ce^{144} TAGGED WILD-RODENT FLEAS *MALAREUS TELCHINUM* FROM THE FIELD VOLE *MICROTUS CALIFORNICUS* TO THE DOMESTIC RAT *RATTUS NORVEGICUS*

No fleas placed on <i>Microtus</i>	60
No transfers to rats while <i>Microtus</i> were alive†	3
No fleas taken from rats after <i>Microtus</i> were killed‡	8
No fleas taken from rats allowed in same area with live <i>Microtus</i>	9
No fleas found in rat nests	27
Total number of fleas accounted for from rats	4
Percentage of recovery of fleas from rats	18.3

From data of W. V. Hartwell (unpublished)

† Rats in contiguous area but prevented from mingling with *Microtus*

‡ Rats allowed into area with dead *Microtus*

Radioactive fleas were also recovered from *Microtus* nests and radioactive feces were observed which showed that the rodents had ingested some of the fleas. In general a higher percentage of tagged female fleas was recovered.

Tests with domestic rats showed that tagged fleas would transfer most readily from *Microtus* to *R. norvegicus* when infested *Microtus* were killed and allowed to remain in the area or when rats were allowed to enter the *Microtus* area and compete with them. TABLE 10 summarizes the pertinent data. Some indication of the life span of Ce^{144} tagged fleas was given by the fact that 76.5 per cent of the fleas were recovered from rats and from their nests after 50 days (TABLE 11). In all the trials tagged fleas showed an initial

TABLE 11

TIMES OF RECOVERY OF Ce^{144} TAGGED FLEAS *MALAREUS TELCHINUM* FROM RATS *R. RATTUS NORVEGICUS* WHEN THE FLEAS WERE ORIGINALLY PLACED ON THE WILD RODENT *MICROTUS CALIFORNICUS*

Time (day)	Number fleas transferred	Test condition
Fleas combed from rats		
18	1	Rats allowed in area with dead <i>Microtus</i>
29	2	
33	1	
34	4	
38	1	Rats isolated in area contiguous with <i>Microtus</i>
42	2	
50	4	Rats allowed free access to area with live <i>Microtus</i>
51	2	
56	3	
Fleas from rat nests		
56	27	Nests removed after all above conditions terminated

From data of W. V. Hartwell (unpublished)

of the rat flea *Nosopsyllus fasciatus*. Thus the finding of this rat flea on trapped *Microtus* need not necessarily imply transfer of the fleas from rats.

In the introduction we have already noted the observations of McCoy (1910) Eskey (1938) Prince (1943) and Meyer and Holdenried (1949) on the subject of flea transfer between wild and domestic rodents. In addition Meyer (1942b) presented evidence of flea transfer and Pollitzer (1954) has summarized the pertinent literature on this subject. Recent studies at this laboratory (Miles *et al.* 1957) have revealed several instances of finding rat fleas on wild rodents and wild rodent fleas on commensal rats.

In plague epidemiology the significance of flea transfer is twofold. On the one hand enzootic plague foci and strictly wild rodent plague epizootics depend on the transmission of the organism by efficient vectors that transfer between individual hosts of the primary reservoir species. On the other hand the spread of plague either from one wild rodent species to another or from a wild to a domestic rodent species depends on transfer of infected fleas interspecifically between hosts. The existing evidence suggests that transfer to rats may be fairly rare since no cases of rat plague have been found as yet in the interior of the western states. That this is not due to a lack of commensal rats was shown during a plague epizootic of rabbits in New Mexico where plague free rats were not only present but also entirely free of flea infestation (Link 1950a).

On the basis of present knowledge two problems of prime importance must be solved. (1) direct observation of flea transfer between host species must be shown in order to substantiate the current hypothesis of plague spread based on circumstantial evidence and (2) quantitative data on the commingling of wild rodent and rat populations must be gathered to evaluate more adequately the potential for plague spread. Some data on the first of these problems is presented here. Recently this laboratory in cooperation with the University of California developed a simple method for the radioactive tagging of fleas and other arthropods with cerium¹⁴⁴ (Quan *et al.* 1951). Fleas were successfully tagged by bathing them for less than 5 minutes in aqueous solutions of Ce¹⁴⁴ Pr¹⁴⁴ containing 10 μ c/ml. Tagging was accomplished by a stable combination of cerium with the exoskeleton a reaction that is at present still obscure. The isotope Ce¹⁴⁴ has a half life of 282 days and the beta particles from its short lived daughter isotope Pr¹⁴⁴ (half life 1.5 minutes) can be detected readily with the usual Geiger Muller survey meter. Thus the means were developed to evaluate the problem of flea transfer between hosts in the field. W. V. Hartwell recently conducted at our laboratory a pilot experiment in simulated field plots; this was done on a small scale as a preliminary to actual field studies. The rodents *M. californicus* and *R. norvegicus* and tagged fleas *Xenopsylla conformis* were used in these tests. The experiments established the fact that under the more or less normal conditions of the rodents' existence the fleas transferred between individual *Microtus*. In a total of 5 trials 111 tagged fleas were placed on different individuals with observations lasting 15 days on the average and ranging from 6 to 21 days. A total of 99 actual flea transfers was observed or an average of 19 transfers per test and the percentage of tagged fleas accounted for ranged from 30 to 82.

TABLE 10

TRANSFER OF Ce^{144} TAGGED WILD-RODENT FLEAS *MALARAUS TELCHINUM* FROM THE FIELD VOLF *MICROTUS CALIFORNICUS* TO THE DOMESTIC RAT *RATTUS NORVEGICUS*

No fleas placed on <i>Microtus</i>	60
No transfers to rats while <i>Microtus</i> were alive†	3
No fleas taken from rats after <i>Microtus</i> were killed‡	8
No fleas taken from rats allowed in same area with live <i>Microtus</i>	9
No fleas found in rat nests	27
Total number of fleas accounted for from rats	47
Percentage of recovery of fleas from rats	78.3

From data of W. V. Hartwell (unpublished)

† Rats in contiguous area but prevented from mingling with *Microtus*‡ Rats allowed into area with dead *Microtus*

Radioactive fleas were also recovered from *Microtus* nests and radioactive feces were observed which showed that the rodents had ingested some of the fleas. In general a higher percentage of tagged female fleas was recovered.

Tests with domestic rats showed that tagged fleas would transfer most readily from *Microtus* to *R. norvegicus* when infested *Microtus* were killed and allowed to remain in the area or when rats were allowed to enter the *Microtus* area and compete with them. TABLE 10 summarizes the pertinent data. Some indication of the life span of Ce^{144} tagged fleas was given by the fact that 76.3 per cent of the fleas were recovered from rats and from their nests after 50 days (TABLE 11). In all the trial tagged fleas showed an initial

TABLE 11

TIMES OF RECOVERY OF Ce^{144} TAGGED FLEAS *MALARAUS TELCHINUM* FROM RATS *RATTUS NORVEGICUS* WHEN THE FLEAS WERE ORIGINALLY PLACED ON THE WILD RODENT *MICROTUS CALIFORNICUS*

Time (day)	Number of fleas	Test condition
Fleas collected from rats		
18	1	Rats allowed in area with dead <i>Microtus</i>
29	2	
33	1	
34	4	
38	1	Rats isolated in area contiguous with <i>Microtus</i>
42	2	
50	4	Rats allowed free access to area with live <i>Microtus</i>
51	2	
56	3	
Fleas found in rat nests		
56	27	Nests removed after all above conditions terminated

From data of W. V. Hartwell (unpublished)

of the rat flea *Nosopsyllus fasciatus*. Thus the finding of this rat flea on trapped *Microtus* need not necessarily imply transfer of the fleas from rats.

In the introduction we have already noted the observations of McCoy (1910), Eskey (1935), Prince (1943), and Meyer and Holdenried (1949) on the subject of flea transfer between wild and domestic rodents. In addition Meyer (1942b) presented evidence of flea transfer and Pollitzer (1954) has summarized the pertinent literature on this subject. Recent studies at this laboratory (Miles *et al.* 1957) have revealed several instances of finding rat fleas on wild rodents and wild rodent fleas on commensal rats.

In plague epidemiology the significance of flea transfer is twofold. On the one hand, enzootic plague foci and strictly wild rodent plague epizootics depend on the transmission of the organism by efficient vectors that transfer between individual hosts of the primary reservoir species. On the other hand, the spread of plague either from one wild rodent species to another or from a wild to a domestic rodent species depends on transfer of infected fleas interspecifically between hosts. The existing evidence suggests that transfer to rats may be fairly rare, since no cases of rat plague have been found as yet in the interior of the western states. That this is not due to a lack of commensal rats was shown during a plague epizootic of rabbits in New Mexico where plague free rats were not only present but also entirely free of flea infestation (Link 1950a).

On the basis of present knowledge two problems of prime importance must be solved. (1) direct observation of flea transfer between host species must be shown in order to substantiate the current hypothesis of plague spread based on circumstantial evidence, and (2) quantitative data on the commingling of wild rodent and rat populations must be gathered to evaluate more adequately the potential for plague spread. Some data on the first of these problems is presented here. Recently this laboratory in cooperation with the University of California developed a simple method for the radioactive tagging of fleas and other arthropods with cerium¹⁴⁴ (Quan *et al.* 1957). Fleas were successfully tagged by bathing them for less than 5 minutes in aqueous solutions of Ce¹⁴⁴ Pr¹⁴⁴ containing 10 μ c/ml. Tagging was accomplished by a stable combination of cerium with the exoskeleton, a reaction that is at present still obscure. The isotope Ce¹⁴⁴ has a half life of 282 days and the beta particles from its short lived daughter isotope Pr¹⁴⁴ (half life 17.5 minutes) can be detected readily with the usual Geiger Muller survey meter. Thus the means were developed to evaluate the problem of flea transfer between hosts in the field. W. V. Hartwell recently conducted at our laboratory a pilot experiment in simulated field plots; this was done on a small scale as a preliminary to actual field studies. The rodents *M. californicus* and *R. norvegicus* and tagged fleas *Malariaeus telchinum* were used in these tests. The experiments established the fact that under the more or less normal conditions of the rodents' existence the fleas transferred between individual *Microtus*. In a total of 5 trials, 111 tagged fleas were placed on different individuals with observations lasting 15 days on the average and ranging from 6 to 21 days. A total of 99 actual flea transfers was observed or an average of 19 transfers per test, and the percentage of tagged fleas accounted for ranged from 50 to 82.

TABLE 12

WILD RODENT FLEAS *XIPHOCHUS TYPICUS* *HYSTICHOPSYLLA LINDALEI*
 1) *PHILOPPUS MULTIDENTATUS* FOUND NATURALLY INFECTED WITH
PASTURELLA PESTIS BY BACTERIAL CULTIVATION OF INDIVIDUAL
 FLEAS AND BY ANIMAL INOCULATION OF FLEA POOLS

C llect	Fl it l l dually				Fl pool ocul t d					T tal fl p	P tag fl po t
	N t t l	N pos	P tag po	Eff y t (p t g)	N pool test l	N pos l	P tag po t	N fl po t	P tag fl po t	Eff y t (p t g)	
A	44	40	9 0	97	88	9	10 0	45	10 0	98	46
B	357	13	3 6	77	91	10	11 0	10	2 8	59	17
C	254	8	3 1	73	82	10	12 0	10	3 9	91	11
Total	1058	61	5 8	83	261	29	11 0	65	6 2	88	74

5 me positive pools contained more than one plague positive flea

† Individual culture checked on fleas used in pools—based on number of fleas shown in second column to the left

‡ Assuming that no positive fleas were missed by combination of both methods and assuming that only one positive flea was present in a positive pool from fleas found negative by culture

tion of the two methods revealed all plague positive fleas the culture method showed from 73 to 87 per cent of the positives and averaged 79 per cent the inoculation method showed from 59 to 98 per cent of the positives and averaged 83 per cent. Thus the efficiency of detecting plague in the fleas by a combination of the two methods was 96 per cent (on the average 3 naturally infected fleas were not detected when 74 plague positive fleas were collected)

If the fleas had been processed only by the individual culture method 61 of 74 positives or 83 per cent would have been detected. During these tests which will be reported in greater detail elsewhere it was felt that the efficiency of individual flea culture would be higher if trituration were accomplished by a tissue homogenizer rather than by manual crushing with a roughened glass rod in a test tube. If the fleas had been processed solely by the animal inoculation method 65 positives out of 74 or 88 per cent would have been found. In general a high level of efficiency can be expected with the animal inoculation method which is of value mainly in plague surveillance or when it is necessary to test for the presence of plague and to determine its extent on a rough scale. The individual culture of fleas not only reveals the presence of plague efficiently but also provides a natural plague infection rate in different flea species. Thus the latter method though tedious and time consuming is valuable in more detailed ecologic research on wild rodent plague and has provided types of data hitherto unknown.

The total absence of plague from the inhabitants of California for the past eight years could lead to the erroneous conclusion that control measures applied in this state for the past 50 years have completely eradicated the disease. It is wiser to see the present level as a plateau of unknown length and to resist the temptation to draw conclusions about the cause. —Karl F. Meyer (1955)

average count of $6.2 \text{ to } 8.5 \times 10^2$ counts/flea/min and after 56 days, $3.6 \text{ to } 5.1 \times 10^2$ counts/flea/min

These preliminary experiments are not conclusive but they suggest that a wild rodent flea *Malareus telchinum* will readily transfer between individuals of its normal host species whereas transfer to domestic rats may take place only under unusual circumstances. Nevertheless the transfer of wild rodent fleas to rats from dead wild rodent hosts is of special significance when applied to the problem of plague spread from sylvatic to domestic environments. The fact that *M. telchinum* will transfer to rats illustrates the most probable method by which domestic rats acquired plague during an epizootic among *Microtus* and other small field rodents (Kartman *et al.* 1958). Undoubtedly, other important wild rodent flea vectors of plague will also transfer to rats.

Detection of Plague in the Field

Winter and Moody (1957) have announced the adoption of Coons' fluorescent antibody technique for rapid identification of *Pasteurella pestis*. They observed that individual cells in a smear were brightly stained in bubo exudate and blood collected from a human case four days after onset of the clinical infection. The method certainly appears to hold great promise for the detection of plague in the field, especially of plague in flea specimens.

Currently detection of plague in nature still depends on general epizootiologic observations and on inoculation of guinea pigs or mice with pools of rodent tissues and fleas. In all probability die offs of larger colonial rodents still occur, and they may be due to plague. However in themselves die offs may be difficult to locate, especially those of the smaller native field mice. Furthermore when plague resistant rodents are present die offs may not occur at all. Thus in plague surveillance workers still rely chiefly on collecting fleas and inoculating susceptible laboratory animals with pools of them.

Eskey and Haas (1940) rediscovered and popularized inoculation with flea pools for the detection of wild rodent plague foci. Inoculation with flea pools for *Pasteurella pestis* has the following advantages: (1) fleas may be plentiful even when rodents are relatively scarce; (2) flea pools are less dangerous to handle than rodent tissues; (3) the specimens are less likely to contain gross contamination than are tissues; and (4) the fleas do not deteriorate as readily as tissues. Flea pool inoculation tests, however, are not suitable for determination of the extent or the intensity of plague infection in a given rodent population. It is rather surprising that in spite of its long and extensive usage the flea pool method has never been evaluated critically with regard to its efficiency as a method for detecting plague in naturally infected fleas.

In recent studies by Quan and his co-workers (unpublished) on wild rodent plague among native field rodents such as *Microtus* and *Peromyscus*, the flea pool inoculation method was compared with the bacteriological culture of individual fleas. The fleas were first individually crushed and triturated and then streaked on a blood agar plate. The remaining portions of individual flea triturates were then pooled and used to inoculate laboratory mice. The comparative results obtained by these two methods and an evaluation of their relative efficiencies are shown in TABLE 12. Assuming that the combina-

TABLE 12

WILD RODENT FLEA *MACALPUS TELCHINUM* *HYSTRICHOPSYLLA LIVINGSTONI*
1. PHLOPPAS MULTIPUNCTATUS FOUND NATURALLY INFECTED WITH
PASTEURILLA PESTIS BY BACTERIAL CULTIVATION OF INDIVIDUAL
 FLEAS AND BY ANIMAL INOCULATION OF FLEA POOLS

Cult	Individual Culture				Pool Inoculation						Total fleas pooled	Plague positive pooled
	Negative tested	Negative pooled	Plague positive pooled	Efficiency (% of total tested)	Negative tested	Negative pooled	Plague positive pooled	Negative pooled	Plague positive pooled	Efficiency (% of total pooled)		
A	447	40	90	87	89	9	100	45	100	98	46	103
B	357	13	36	77	91	10	110	10	28	51	17	48
C	254	8	31	73	87	10	120	10	39	91	11	43
Totals	108	61	58	83	261	29	110	65	62	88	74	70

Some positive pool contained more than one plague positive flea

† Individual culture checked on flea used in pools—based on number of fleas shown in second column to the left

‡ Assuming that no positive fleas were missed by combination of both methods and assuming that only one positive flea was present in a positive pool from fleas found negative by culture

tion of the two methods revealed all plague positive fleas the culture method showed from 73 to 87 per cent of the positives and averaged 79 per cent the inoculation method showed from 59 to 98 per cent of the positives and averaged 83 per cent. Thus the efficiency of detecting plague in the fleas by a combination of the two methods was 96 per cent (on the average 3 naturally infected fleas were not detected when 74 plague positive fleas were collected)

If the fleas had been processed only by the individual culture method 61 of 74 positives or 83 per cent would have been detected. During these tests which will be reported in greater detail elsewhere it was felt that the efficiency of individual flea culture would be higher if trituration were accomplished by a tissue homogenizer rather than by manual crushing with a roughened glass rod in a test tube. If the fleas had been processed solely by the animal inoculation method 65 positives out of 74 or 88 per cent would have been found. In general a high level of efficiency can be expected with the animal inoculation method which is of value mainly in plague surveillance or when it is necessary to test for the presence of plague and to determine its extent on a rough scale. The individual culture of fleas not only reveals the presence of plague efficiently but also provides a natural plague infection rate in different flea species. Thus the latter method though tedious and time consuming is valuable in more detailed ecologic research on wild rodent plague and has provided types of data hitherto unknown.

The total absence of plague from the inhabitants of California for the past eight years could lead to the erroneous conclusion that control measures applied in the state for the past 50 years have completely eradicated the disease. It is wiser to view the present level as a plateau of unknown length and to resist the temptation to draw conclusions about the causes.—Karl F. Meyer (1955)

average count of 62 to 85×10^4 counts/flea/min and after 56 days 36 to 51×10^4 counts/flea/min

These preliminary experiments are not conclusive but they suggest that a wild rodent flea, *Malaraeus telchinum* will readily transfer between individuals of its normal host species whereas transfer to domestic rats may take place only under unusual circumstances. Nevertheless the transfer of wild rodent fleas to rats from dead wild rodent hosts is of special significance when applied to the problem of plague spread from sylvatic to domestic environments. The fact that *M. telchinum* will transfer to rats illustrates the most probable method by which domestic rats acquired plague during an epizootic among *Microtus* and other small field rodents (Kartman *et al.* 1958). Undoubtedly other important wild rodent flea vectors of plague will also transfer to rats.

Detection of Plague in the Field

Winter and Moody (1957) have announced the adoption of Coons' fluorescent antibody technique for rapid identification of *Pasteurella pestis*. They observed that individual cells in a smear were brightly stained in bubo exudate and blood collected from a human case four days after onset of the clinical infection. The method certainly appears to hold great promise for the detection of plague in the field especially of plague in flea specimens.

Currently detection of plague in nature still depends on general epizootiologic observations and on inoculation of guinea pigs or mice with pools of rodent tissues and fleas. In all probability die offs of larger colonial rodents still occur, and they may be due to plague. However in themselves die offs may be difficult to locate especially those of the smaller native field mice. Furthermore when plague resistant rodents are present die offs may not occur at all. Thus in plague surveillance workers still rely chiefly on collecting fleas and inoculating susceptible laboratory animals with pools of them.

Eskey and Haas (1940) rediscovered and popularized inoculation with flea pools for the detection of wild rodent plague foci. Inoculation with flea pools for *Pasteurella pestis* has the following advantages: (1) fleas may be plentiful even when rodents are relatively scarce; (2) flea pools are less dangerous to handle than rodent tissues; (3) the specimens are less likely to contain gross contamination than are tissues; and (4) the fleas do not deteriorate as readily as tissues. Flea pool inoculation tests however are not suitable for determination of the extent or the intensity of plague infection in a given rodent population. It is rather surprising that in spite of its long and extensive usage the flea pool method has never been evaluated critically with regard to its efficiency as a method for detecting plague in naturally infected fleas.

In recent studies by Quan and his co-workers (unpublished) on wild rodent plague among native field rodents such as *Microtus* and *Peromyscus* the flea pool inoculation method was compared with the bacteriological culture of individual fleas. The fleas were first individually crushed and triturated and then streaked on a blood agar plate. The remaining portions of individual flea triturates were then pooled and used to inoculate laboratory mice. The comparative results obtained by these two methods and in evaluation of their relative efficiencies are shown in TABLE 12. Assuming that the combina-

through the mechanism of flea exchange discussed in a preceding section. Recent studies at this station have shown the importance of small native field mice (genera *Microtus*, *Peromyscus* and *Reithrodontomys*) not only as a source of plague but also as a reservoir of infected fleas that may transfer to domestic rats. The most important fact is that enzootic foci of plague are found near suburban developments, farms and in the environs of cities and that wild rodents commingle with rats in such areas as well as in city garbage dumps, small towns and camps. The increase in suburban populations and the extension of human habitations into rural areas is probably the most significant factor in the present wild rodent plague problem. Our studies in northern San Mateo County, just beyond the San Francisco limits, have shown that plague epizootics occur among the native field rodents and that plague actually transfers to domestic rats (Kartman *et al* 1958). Thus in the problem of wild rodent plague control, close attention must be given to situations in which plague foci in native field rodents are found in areas where commensal rats occur adjacent to human habitations and human activity.

It is undoubtedly true that under usual conditions in this country the sanitary practices that have evolved and that are continually being improved constitute effective barriers to the transfer of plague from wild to domestic rodents and thence to man. Furthermore, unlike the situation in other regions of the world, there is no evidence here that the domestic rat constitutes an independent reservoir of plague infection. In Madagascar, for instance, the periodic resurgence of localized murine plague foci has been noted without any evidence that rodents other than the rat are the primary source of the disease (Girard 1940). From the standpoint of public health, the critical factor in our western region is the maintenance of effective barriers between the wild rodent reservoirs of plague and the domestic rats. On the other hand, the normal sanitary barriers could conceivably be severed in times of natural disasters such as floods and earthquakes and also during national emergencies, especially in case of devastation due to war. The latter may seem a highly remote possibility, yet in the face of ever growing threats of international atomic conflict we must be prepared with the knowledge requisite for dealing with plague and other vector borne diseases. Thus resources for public health must include not only a cognizance of wild rodent plague ecology but also an awareness of the most effective means for the detection, diagnosis, treatment and control of plague.

Diagnosis and Treatment of Plague Cases

Today prognosis of plague is excellent provided the disease is diagnosed sufficiently early. Rapidity of diagnosis, however, still depends heavily on both epidemiologic intelligence and sensitive, rapid laboratory methods for detecting *Pasteurella pestis* and its antibodies. Fortunately, advances have been made along all these fronts. It has been shown recently that the plague organism in human blood, exudates and post mortem tissues is subject to rapid identification by use of fluorescent antibody techniques (Winter and Moody 1957). The detection of plague antibodies by the hemagglutination procedure (Chen 1952) has also proved useful (McCrumb *et al* 1953).

Public Health Aspects of Sylvatic Plague

In the introduction we have noted the large area in which evidence of wild rodent plague has been found in the western United States. In contrast to this it must be pointed out that the number of human infections derived directly from wild rodent foci has been of a low order. TABLE 13 which is modified after Pollitzer (1954) summarizes the incidence of human plague infection contracted from wild rodents or their fleas.

The reason for the low level of human plague directly associated with wild rodents is not difficult to assess. Foci of wild rodent plague are generally situated in sparsely populated and isolated rural districts where wild rodent fleas do not have the opportunity of biting man. Even when humans do enter areas harboring wild rodent plague various biological factors intervene to prevent infection. A person would have to be present during an epizootic that in most cases appears quite transitory. He would have to be bitten by an infective (or blocked) wild rodent flea or simultaneously by a number of wild rodent fleas that had very recently fed on a septicemic animal. The chances against this are great when the natural infection rate of fleas is taken into account and when the factor of host preference is considered. Further more in a large flea population the number of blocked individuals would probably be very small and of those infective fleas present that actually did bite very few would be likely to accomplish transmission. That plague transmission to humans by wild rodent fleas does occur is shown by TABLE 13 but according to the World Health Organization Expert Committee on Plague (1953). The unexpected occurrence of human plague in insufficiently explored areas draws attention to the need for an exact delimitation of wild rodent plague. We might add that the occurrence of sylvatic plague near human habitations makes the ecology of wild rodent plague a paramount subject for investigation.

The real danger from wild rodent plague occurs when wild and domestic rodents coexist in areas close to human habitation. It is here that the possibility of plague transfer from wild to commensal rodents may take place.

TABLE 13
INCIDENCE OF PLAGUE INFECTION ORIGINATING IN SYLVATIC FOCI

Y w h p l g w f i r t f m d			H m plagu 1908 t 1956	
Stat	W l d o d t	Huma	C	De th
California	1908	1908	81	53
Oregon	1935	1934	1	1
Utah	1936	1936	1	0
Nevada	1936	1937	1	0
Idaho	1936	1940	1	1
New Mexico	1938	1949	6	3
Arizona	1938	1950	1	0
Totals			92	58

* Plague suspected in ground squirrels in 1903

through the mechanism of flea exchange discussed in a preceding section. Recent studies at this station have shown the importance of small native field mice (genera *Microtus*, *Peromyscus* and *Reithrodontomys*) not only as a source of plague but also as a reservoir of infected fleas that may transfer to domestic rats. The most important fact is that enzootic foci of plague are found near suburban developments, farms, and in the environs of cities, and that wild rodents commingle with rats in such areas as well as in city garbage dumps, small towns, and camps. The increase in suburban populations and the extension of human habitations into rural areas is probably the most significant factor in the present wild rodent plague problem. Our studies in northern San Mateo County, just beyond the San Francisco limits, have shown that plague epizootics occur among the native field rodents and that plague actually transfers to domestic rats (Kartman *et al* 1958). Thus in the problem of wild rodent plague control, close attention must be given to situations in which plague foci in native field rodents are found in areas where commensal rats occur adjacent to human habitations and human activity.

It is undoubtedly true that under usual conditions in this country, the sanitary practices that have evolved and that are continually being improved constitute effective barriers to the transfer of plague from wild to domestic rodents and thence to man. Furthermore, unlike the situation in other regions of the world, there is no evidence here that the domestic rat constitutes an independent reservoir of plague infection. In Madagascar, for instance, the periodic resurgence of localized murine plague foci has been noted without any evidence that rodents other than the rat are the primary source of the disease (Girard 1940). From the standpoint of public health, the critical factor in our western region is the maintenance of effective barriers between the wild rodent reservoirs of plague and the domestic rats. On the other hand, the normal sanitary barriers could conceivably be severed in times of natural disasters such as floods and earthquakes, and also during national emergencies, especially in case of devastation due to war. The latter may seem a highly remote possibility, yet in the face of ever growing threats of international atomic conflict, we must be prepared with the knowledge requisite for dealing with plague and other vector-borne diseases. Thus, resources for public health must include not only a cognizance of wild rodent plague ecology, but also an awareness of the most effective means for the detection, diagnosis, treatment, and control of plague.

Diagnosis and Treatment of Plague Cases

Today, prognosis of plague is excellent provided the disease is diagnosed sufficiently early. Rapidity of diagnosis, however, still depends heavily on both epidemiologic intelligence and sensitive, rapid laboratory methods for detecting *Pasteurella pestis* and its antibodies. Fortunately, advances have been made along all these fronts. It has been shown recently that the plague organism in human blood, exudates, and post mortem tissues is subject to rapid identification by use of fluorescent antibody techniques (Winter and Moody 1957). The detection of plague antibodies by the hemagglutination procedure (Chen 1952) has also proved useful (McCrumb *et al* 1953).

In the effective treatment of plague the evaluations made by Meyer and his co workers (1952) still represent the most modern concepts of therapy. Smadel and his associates (1955) have added their experiences from the handling of pneumonic cases in Madagascar. Specific plague therapy calls for a judicious administration of antibiotics and, when necessary, the use of anti serum. The antibiotics of choice are streptomycin, chloramphenicol and the tetracyclines. 'Sulfa' drugs especially sulfadiazine and sulfamerazine are very useful both in treatment and for prophylaxis (Quan *et al.* 1955). Vaccination is a logical procedure under certain conditions.

Although further general discussion of plague therapy is beyond the scope of this paper, the fatal outcome in spite of effective treatment, of the 1956 bubonic plague case in California deserves comment. This case was of sylvatic origin, it was diagnosed early and treated so intensely, that no bacilli could be isolated at autopsy after but three days of therapy. The death of the patient indicates that therapy still presents problems. The patient (Sackacs) was a retired Chief Petty Officer in his middle forties. He had been in good health and there was no reason to suppose that he was unusually susceptible or sensitive to plague. Quan and Meyer (unpublished) have investigated the virulence and chemotherapy of the Sackacs strain of *P. pestis*, a succinct report of the findings follows.

From the bacteriological point of view, the plague bacilli isolated from the bubo exudate of Sackacs were undifferentiated from any other virulent *Pasteurella pestis* strains. Although the content of the fraction I protein (antigenic portion) of this California strain was not studied, that of its fraction II content (toxic portion) was extracted and compared with an Indian strain (195/P). Both strains were equally toxic. When compared with the Alexander and 195/P strains for growth in broth containing various concentrations of streptomycin, aureomycin or chloramphenicol, no differences were found in the sensitivity of the three strains of these antibiotics *in vitro* (TABLE 14). When mice were infected intracutaneously with about 500 bacilli of the above three strains and given therapy at 48 hours after infection, the success of the antibiotics against the Sackacs strain was similar to that against infection by the

TABLE 14
GROWTH OF *PASTEURELLA PESTIS* IN BROTH CONTAINING ANTIBIOTICS

P I II p t t	St pt my	A eomy	Chl mph l
	C	t t om (μg/ml) s h b t y t	t d y
Alexander	4 0	1 0	2 0
Sackacs	4 0	1 0	2 0
195/I	4 0	1 0	2 0
	C	t also (μg/ml) b t	d l d y
Alexan ler	8 0	32	4 0
Sackacs	8 0	32	4 0
195/P	8 0	32	4 0

* Broth contained 10⁶ microorganisms per ml

TABLE 15
SURVIVAL RATE OF MICE INOCULATED WITH *PASTEURELLA PESTIS* AND TREATED WITH ANTIBIOTICS

P _{teu} il p _{ti} tra nocolated	A t b i t sed th rapy			
	St ptomy	A eomy	Cbl ramph	N (t l)
		N mb	d mb t d	
Alexand r	20/20	6/20	14/20	0/11
Sackacs	16/20	5/21	12/20	0/11
195/1	13/ 0	3/20	15/20	0/11

Infection was produced intracutaneously by injection of 0.05 ml containing about 500 *Pasteurella pestis*. The course of therapy was 3 days beginning 48 hours after infection. The daily dose was 100 µg I.I. followed in 8 hours by 200 µg subcut.

other two strains (TABLE 15). Although in some tests the Sackacs strain appeared to kill mice slightly more rapidly than did the other two strains the difference was not significant. These investigations suggested that in the light of present knowledge the Sackacs strain was undifferentiated bacteriologically and chemotherapeutically from the New Mexico and the Indian strains.

The therapeutics may be a major factor in the fatal outcome of the California case. It was reported that in this particular case heroic dosages of all the effective plague medicinals were administered and that the patient was sterilized of *P. pestis*. Although this has not been heretofore observed in man its possibility was indicated when Quan and his co-workers (1941) sterilized mice of plague bacilli by the use of streptomycin (Meyer *et al* 1952). Nevertheless it was not known what effect if any such sterilization might have on the host animal.

We know that experiments in the chemotherapy of plague with laboratory mice and guinea pigs show that the use of increasing dosages of effective drugs or combinations of drugs up to toxic dosages have elevated survival rates (TABLE 16). However as may be seen from TABLE 16 even the intensive therapy was short of the sterilizing range of *P. pestis in vivo*. Thus to test for sterilizing effect a level higher than intensive therapy heroic therapy was formulated from the pattern of the clinical treatment used in the California case. In that case sulfisoxazole which is far less effective than sulfadiazine in experimental plague therapy (Quan *et al* 1955) and penicillin which is ineffective against *in vivo P. pestis* (Meyer *et al* 1952) were used with streptomycin chloramphenicol and oxytetracycline. In the heroic therapy sulfadiazine was substituted for sulfisoxazole but penicillin was included. All the drugs were adjusted for treating mice every six hours. The various dosages of the therapies are shown in TABLE 17. The moderate and heroic therapies were given to mice that had been infected intraperitoneally at a time when between 20 and 40 per cent of the animals were dead.

The results (TABLE 18) indicate a clear trend excessive dosage of drugs given in the heroic therapy was less effective and saved fewer mice than did the moderate dosage. It should be noted that these are the results of but

TABLE 16
CHEMOTHERAPY OF MICE INOCULATED INTRACUTANEOUSLY WITH SACKACS AND ALEXANDER
STRAINS OF *PASTEURELLA PESTIS*

Therapy	Intracutaneous					
	36-hour infection		51-hour infection		66-hour infection	
	Alexander	Sackacs	Alexander	Sackacs	Alexander	Sackacs
	(Percentage survival)					
Sulfadiazine	58	73	42	45	—	—
Sulfa and streptomycin	83	77	67	45	33	27
Intensive†	100	100	83	91	67	59

* Twenty two mice per group of Sackacs 12 mice per group of Alexander

† Intensive therapy consisted of a combination of sulfadiazine streptomycin terramycin and chloramphenicol

one experiment and are not conclusive. K. F. Meyer has commented that these results may have been due to a mechanism analogous to that in the Herxheimer reaction. Furthermore, it is possible that this strain of *P. pestis* is more readily lysed *in vitro* when under antibiotic treatment than are other strains. If these results are confirmed they suggest a new type of toxic effect of antibiotics in addition to those already known.

Civilization alone has destroyed the plague in Europe; it alone will destroy it in the East.—Aulert Roche (quoted by H. H. Scott 1942)

Sylvatic Plague Control

In any discussion of the control of an arthropod transmitted disease of man cognizance must be given to the effects of human social development upon its rise or fall. Hence the above quotation. We cannot enter here into a dissertation on philosophies of history but it should be mentioned that certain men prominent in their field—Ronald Ross, Angelo Celli, Hans Zinsser and others—have promulgated a monistic theory that assumes that disease makes history. Although we cannot deny that malaria, typhus, yellow fever and plague have often been strong influences against social progress we are inclined to accept the view of such scholars as Ackerknecht (1945) who discussed malaria in particular that the waves of these diseases can be more logically explained by historical events. In his famous Milroy Lectures on plague Liston (1924) cited the second agrarian revolution in England during the Eighteenth Century as the unrecognized beginning in the conquest of plague in that country. He discussed in detail the influences of social reforms, sanitary measures and the prosperity associated with the period of high farming. On the other hand Hirst (1953) explained this same decline in plague on the basis of a change in the rat and rat flea species without investigating the causal factors influencing reductions in black rat and *Xenopsylla cheopis* populations. Very recently it was observed by Fox (1956) that typhus in San Juan, Puerto Rico, showed a 100 per cent decrease

TABLE 17
DOSAGES OF VARIOUS THERAPIES

Therapy	Dose in (mg) (cc) (sets)	Mg/kg	Kilograms (lb) (kg)
Mild			
Sulfadiazine	2.00	100	40
Moderate			
Streptomycin	0.20 0.25	10 12.5	
Sulfadiazine	2.00 1.00	100 50	60
Intensive			
Streptomycin	0.20	10	
Chloramphenicol	0.40	20	80
Terramycin	0.40	20	
Sulfadiazine	2.00	100	
Heroic			
Streptomycin	1.00	50	
Chloramphenicol	0.50	25	
Terramycin	0.50	25	
Sulfadiazine	1.00	50	
Penicillin	1.00	50	
Serum			
Antitoxic (intraperitoneal 0.5 cc of 1 to 5 dilution of 40 units/cc)			0

Average survival rate of mice infected intracutaneously with about 100 plague bacilli for 51 hours and then treated

TABLE 18
SERO THERAPY AND CHEMOTHERAPY OF MICE INOCULATED INTRAPERITONEALLY WITH
SACKS STRAIN OF *PASTEURRELLA PESTIS*

Therapy	Mortality (death)	Survival (death)
Moderate With serum	9/25 (36)	5/12 (41) 7/12 (58)
Heroic With serum	5/25 (20) 10/25 (40)	4/11 (36) 5/11 (45)

Mice received about 5000 bacilli

at a time when absolutely no control operations were in effect as in the southern United States. Thus Fox concluded that a reduced typhus rate is not necessarily proof of the efficacy of control measures. Girard (1955) has evaluated the modern position on plague in the following nicely balanced terms. The modern pandemic is declining. This situation is legitimately attributed to the success of a prophylactic method, the result of fruitful research, but one must also include the contribution of natural circumstances in order to explain this regression in countries where the control has been too limited to be effective.

All evidence points to a spectacular decline in human plague throughout the world (Gear and Deutschman 1956). Plague has apparently been conquered

TABLE 16
CHEMOTHERAPY OF MICE INOCULATED INTRACUTANEOUSLY WITH SACKACS AND ALEXANDER
STRAINS OF *PASTEURILLA PESTIS*

Therapy	Fifty per cent					
	36 hours		31 hours		66 hours	
	Alexander	Sack	Alexander	Sack	Alexander	Sack
	(Percentage of survivors)					
Sulfadiazine	58	73	42	45	—	—
Sulfa and streptomycin	83	77	67	45	33	27
Intensive†	100	100	83	91	67	59

* Twenty two mice per group of Sackacs 12 mice per group of Alexander

† Intensive therapy consisted of a combination of sulfadiazine streptomycin terramycin and chloramphenicol

one experiment and are not conclusive. H. F. Meyer has commented that these results may have been due to a mechanism analogous to that in the Herxheimer reaction. Furthermore it is possible that this strain of *P. pestis* is more readily lysed *in vivo* when under antibiotic treatment than are other strains. If these results are confirmed they suggest a new type of toxic effect of antibiotics in addition to those already known.

Civilization alone has destroyed the plague in Europe it alone will destroy it in the East. —Aulert Roche (quoted by H. H. Scott 1942)

Sylvatic Plague Control

In any discussion of the control of an arthropod transmitted disease of man cognizance must be given to the effects of human social development upon its rise or fall. Hence the above quotation. We cannot enter here into a dissertation on philosophies of history but it should be mentioned that certain men prominent in their field—Ronald Ross, Angelo Celli, Hans Zinsser and others—have promulgated a monistic theory that assumes that disease makes history. Although we cannot deny that malaria, typhus, yellow fever and plague have often been strong influences against social progress we are inclined to accept the view of such scholars as Ackerknecht (1945) who discussed malaria in particular that the waves of these diseases can be more logically explained by historical events. In his famous Milroy Lectures on plague Liston (1924) cited the second agrarian revolution in England during the Eighteenth Century as the unrecognized beginning in the conquest of plague in that country. He discussed in detail the influences of social reforms, sanitary measures and the prosperity associated with the period of high farming. On the other hand Hirst (1953) explained this same decline in plague on the basis of a change in the rat and rat flea species without investigating the causal factors influencing reductions in black rat and *Xenopsylla cheopis* populations. Very recently it was observed by Fox (1956) that typhus in San Juan, Puerto Rico showed a 100 per cent decrease

small native field voles and meadow mice is difficult because their nests are hidden and do not occur in towns or community groupings as do those of the larger colonial rodents. Furthermore wholesale dispersal of insecticidal dusts may not be safe in many areas near human habitation or human activity where the smaller rodents are found and where they present the chief potential for the spread of plague. For these reasons flea control methods were developed in the plague endemic zone of Hawaii in which a DDT bait box was used (Kartman and Lonerhan 1955a). The apparatus consisted of a covered bait station with DDT filled sacks hung at opposite entrances so that the rodent would receive a shower of DDT powder on entering and leaving the box. The theory behind this method was that the rodent itself would become the vehicle by which the insecticide was dispersed to its nest and its mates. In practice this method not only controlled those fleas on the rats but provided evidence suggesting that fleas in the nests also were affected. Excavation of rat burrows showed a complete lack of fleas in the area of the DDT bait box whereas nests were 100 per cent infested in a check area and showed an average of about 25 fleas per nest. Analysis of nesting material in one instance only showed DDT residues amounting to 0.72 mg. In contrast to these results Wagle and Seal (1953) reported that DDT could be detected only in the first few inches of rat holes when the insecticide was applied by insufflation of the burrows. Five fleas were found in 14 of 44 nests examined.

Recent field trials have shown that a simplified insecticide bait box controls fleas on the heavily parasitized *Microtus californicus* as well as on other species of meadow mice both on the hosts and in their nests. FIGURE 6 shows an end view of the bait box which consists of the following: a floor board one half an inch thick, 12 inches long and 8 inches wide; a U shaped roof made by cutting in half lengthwise a lard tin 9¹/₂ inches in diameter and 12¹/₂ inches deep; and a bait container held by small nails in the center of the floor board. The bait boxes were placed at 50-foot intervals in a field grid 250 feet square. A similar check area in the same field was established 150 feet from the test area. After an initial prebaiting period with rolled oats about 80 gm of 5 per cent DDT powder was spread on the floor of each bait box. Data on fleas in rodent nests and on the rodents were taken periodically; examples of the results are summarized in TABLES 19 and 20. These preliminary observations suggest that the insecticide bait box method offers a cheap and efficient means of significantly reducing the plague vector flea populations affecting those native field rodents known to be implicated in sylvatic plague in the San Francisco region. Nevertheless as stated above actual field control of wild rodent plague remains to be demonstrated.

The control of fleas by contact insecticides having residual properties has correctly taken a place of primary importance in plague control. On the other hand workers in this field must be prepared for the development of flea resistance to some of the insecticides. Simmons (1954) has reviewed reports on such resistance and it is encouraging to note that no case of major resistance is known in important plague vectors in nature.

In the long view sanitary measures, construction of rat proof buildings, public education and legal pressures must be intensified to reduce the popu-

as an epidemic scourge but it remains entrenched as an animal infection external to man covering hundreds of square miles and perpetuating itself away from human habitation as a potential threat which may 'spill over' to humans under circumstances neither wholly understood nor predictable. The problem of plague control becomes more and more exclusively concerned with sylvatic plague control. To speak of eradication would be utopian. The so-called eradication campaigns against ground squirrels in California and in certain parts of the Union of Soviet Socialist Republics during the early decades of this century and the continuous program of rodent poisoning in the Hawaiian Islands have not yet resulted in any significant decline either in the rodent populations or in the incidence of wild rodent plague.

The ecologic picture of sylvatic plague developed in the foregoing sections of this paper suggests that control operations must be oriented to specific situations in which plague actually is transferred from wild to domestic sources or where a distinct threat of such transfer is obvious. For this reason the chief concern must be in areas where wild rodent reservoirs are located near harborages containing rats such as villages, suburban areas, camps, and garbage dumps in cities. Beyond that, control methods must be applied in specific situations where wild rodent epizootics in themselves threaten humans. An outbreak in a national park or in the vicinity of an army encampment would deserve immediate intensive control. It should be emphasized that control of wild rodent plague is primarily an emergency measure with clearly limited objectives. Plague foci that threaten humans must be located and kept under surveillance so that control may be instituted rapidly when the need arises. In certain cases prophylactic control of wild rodent plague foci may be justified.

The possible methods of control are based on specific situations. In cases where sylvatic foci impinge upon domestic rats, a systematic program of rat flea and rat control is the easiest and most effective point of attack. This can be accomplished by a routine program utilizing the anticoagulant poisons such as Warfarin and the newer insecticides including DDT, chlordane, and dieldrin. DDT powder (5 or 10 per cent) still remains the residual insecticide of choice in rodent flea control. Where small foci of plague occur near villages, camps, and other such areas, the control of wild rodents with fast acting poisons and the control of their fleas are measures complementary to rat and rat flea control.

Flea control in itself has become established as an important method of combating plague. The use of DDT in urban areas and during outbreaks of rat-borne plague has shown promising results (Kartman and Lonergan, 1955a). Unfortunately, there has been no adequate demonstration of plague control in wild rodent foci. Field trials have shown that flea populations on wild rodents can be reduced by means of insecticidal dusting of rodent burrows (Davis, 1951; Ryckman *et al.*, 1953) or by area dusting (Miles and Wilcomb, 1953) but no actual demonstration of plague control by these methods is available.

Whereas the insecticidal dusting of ground squirrel burrows is feasible because their entrances are easily located, the control of fleas affecting the

small native field voles and meadow mice is difficult because their nests are hidden and do not occur in towns or community groupings as do those of the larger colonial rodents. Furthermore wholesale dispersal of insecticidal dusts may not be safe in many areas near human habitation or human activity where the smaller rodents are found and where they present the chief potential for the spread of plague. For these reasons flea control methods were developed in the plague endemic zone of Hawaii in which a DDT bait box was used (Kartman and Loneragan 1955a). The apparatus consisted of a covered bait station with DDT filled sacks hung at opposite entrances so that the rodent would receive a shower of DDT powder on entering and leaving the box. The theory behind this method was that the rodent itself would become the vehicle by which the insecticide was dispersed to its nest and its mates. In practice this method not only controlled those fleas on the rats but provided evidence suggesting that fleas in the nests also were affected. Excavation of rat burrows showed a complete lack of fleas in the area of the DDT bait box whereas nests were 100 per cent infested in a check area and showed an average of about 25 fleas per nest. Analysis of nesting material in one instance only showed DDT residues amounting to 0.72 mg. In contrast to these results Wagle and Seal (1953) reported that DDT could be detected only in the first few inches of rat holes when the insecticide was applied by insufflation of the burrows. Five fleas were found in 14 of 44 nests examined.

Recent field trials have shown that a simplified insecticide bait box control fleas on the heavily parasitized *Microtus californicus* as well as on other species of meadow mice both on the hosts and in their nests. FIGURE 6 shows an end view of the bait box which consists of the following: a floor board one half an inch thick, 12 inches long and 8 inches wide; a U shaped roof made by cutting in half lengthwise a lard tin 9¹/₂ inches in diameter and 12¹/₂ inches deep; and a bait container held by small nails in the center of the floor board. The bait boxes were placed at 50-foot intervals in a field grid 250 feet square. A similar check area in the same field was established 150 feet from the test area. After an initial prebaiting period with rolled oats about 80 gm of 5 per cent DDT powder was spread on the floor of each bait box. Data on fleas in rodent nests and on the rodents were taken periodically; examples of the results are summarized in TABLES 19 and 20. These preliminary observations suggest that the insecticide bait box method offers a cheap and efficient means of significantly reducing the plague vector flea populations affecting those native field rodents known to be implicated in sylvatic plague in the San Francisco region. Nevertheless as stated above actual field control of wild rodent plague remains to be demonstrated.

The control of fleas by contact insecticides having residual properties has correctly taken a place of primary importance in plague control. On the other hand workers in this field must be prepared for the development of flea resistance to some of the insecticides. Simmons (1954) has reviewed reports on such resistance and it is encouraging to note that no case of major resistance is known in important plague vectors in nature.

In the long view sanitary measures, construction of rat proof buildings, public education and legal pressures must be intensified to reduce the popu-



FIGURE 6 End view of insecticide bait box in field operation. (Note bait container, DDT powder, and rodent trail in foreground.)

lations of domestic rats, especially those in the vicinity of wild rodent plague foci. If sylvatic plague can be isolated and returned to its former status of a purely wild disease, rising and falling in a state of nature in its delimited foci, the danger to man will have been reduced to the ultimate minimum. As the population of the country increases and our mechanical civilization encroaches

TABLE 19
EFFECT OF 5 PER CENT DDT POWDER IN BAIT BOXES ON THE INCIDENCE OF FLEAS ON
MICROTUS CALIFORNICUS IN SAN FRANCISCO

Period	Treated plot (DDT treated bait box Aug 27 to Sept 27)				Check plot (150 feet from DDT treated plot)			
	Date (1956- 1957)	No. of fleas	Mean no.	Percentage infected	Date (1956- 1957)	No. of fleas	Mean no.	Percentage infected
Pretreatment	8/15-8/23	64/343	5.4	92.2	8/8-8/9	11/49	4.4	54.5
Treatment	9/5-9/26	162/2	0.01	1.2	9/11-9/13	27/200	7.4	78.3
Post treatment	10/2-10/17	206/63	0.3	21.4	10/9-10/17	37/176	5.5	99.9
Post treatment	10/23-11/27	739/274	1.1	50.6	10/23-11/27	109/364	5.2	88.9
Post treatment	1/8-2/7	56/59	1.0	41.1	1/8-2/7	40/126	3.2	82.5
Post treatment	3/19-5/2	63/58	0.9	42.8	3/19-5/2	34/46	1.3	52.9

TABLE 20
EFFECT OF 5 PER CENT DDT POWDER IN BAIT BOXES ON FLEA INCIDENCE
IN NESTS OF *MICROTUS CALIFORNICUS* DURING THE POST TREATMENT PERIOD

Treated plot						Check plot					
DDT treated bait boxes Aug 27 to Sept 27						150 feet from DDT treated plot					
Date (1956-1957)	Day treated	No.	Mean no. fleas		Mean mt DDT (mg)	Date (1956-1957)	Day treated	No.	Mean no. fleas		Mean mt DDT (mg)
			Nest	Infected					Nest	Infected	
10-5	8	7	0.1	1.0	2.1	10-17	20	7	9.0	12.6	0
12-19	83	4	0.2	1.0	1.1	12-19	83	3	14.6	27.0	0
2-6	137	3	0	0	1.0	2-6	132	4	11.5	12.0	0
4-10	195	4	33.5	33.5	0	4-10	195	4	38.5	77.0	0

upon the suburban hinterland sylvatic plague will be increasingly subjected to the cleansing process that marks social progress

Summary and Conclusions

A general review of current problems and the present state of sylvatic plague in the western United States is given. Historically emphasis has shifted from the large colonial rodents such as ground squirrels to the small inconspicuous native field voles and mice such as *Microtus* and *Peromyscus*. The coexistence of these small rodents with domestic rats in the vicinity of human habitations and the prevalence of enzootic plague in these wild rodent populations poses a potential threat that has as yet not been fully elucidated. Some of the present concepts included in the International Sanitary Regulations are fast becoming obsolete since they place major emphasis on classic urban and ship borne plague. Cognizance must now be taken of the potential for transfer of wild rodent plague to domestic rats and thus to humans who could spread the infection abroad by modern rapid means of transportation.

Brief accounts are presented of the rodents and fleas thought to be of importance in plague ecology in the San Francisco Bay region. Of several field rodents, the vole *Microtus californicus* is thought to be the chief plague reservoir because of its high degree of resistance to *Pasteurella pestis* and the fact that it was found to be parasitized by the important vector species as well as by more than 90 per cent of all fleas of all species taken in the area. The fleas *Hystrichopsylla linsdalei* and *Malareus telchinum* the two most prevalent species, were both found to have relatively high natural plague infection rates during epizootic incidents. Of the two *H. linsdalei* is an efficient vector by blocked individuals and *M. telchinum* transmits primarily by mass mechanical means. The former is probably the primary plague vector in this region, whereas the latter may be a secondary vector of importance. The potential for transfer of wild rodent fleas to domestic rats was experimentally demonstrated with radioactively tagged *M. telchinum*, *M. californicus* and *Rattus norvegicus*. It was also shown that plague infection rates in individual fleas taken in the field could be determined and that this method compared favorably with the established method of plague detection by the inoculation of susceptible laboratory animals with flea pools.

Investigations on enzootic epizootic plague in this region revealed that foci are quite delimited, that the disease is actually transferred from wild rodents to rats, that an increase in plague incidence in fleas and in rodents is not necessarily accompanied by a decimation of the wild rodent population, that an apparent epizootic may occur simultaneously with a rise in the *Microtus* population, and that sylvatic plague may remain dormant and difficult, if not impossible, to demonstrate for periods of eighteen months or more and then suddenly become resurgent.

Although relatively few cases of human plague have been traced directly to sylvatic sources, today's increasing suburbanization is of primary concern since it implies a period of joint tenancy between wild rodents, domestic rats and humans. An account of modern methods of diagnosis and treatment of plague is given, and certain problems arising from a recent fatal human case are discussed.

Finally, sylvatic plague control is distinguished from urban control. The primary importance of the control of the domestic rat and the rat flea is emphasized. Sylvatic plague control is conceived of as an emergency measure or as a routine to be established in special instances as an adjunct to domestic rat and rat flea control. Recent field studies with an insecticide bait box method are summarized, and the current limitations of sylvatic plague control are discussed.

References

- ACKFRANCHEIT, F. H. 1945. Malaria in the upper Mississippi Valley, 160-1900. Bull. Hist. Med. Suppl. No. 4.
- BACOT, A. W. & C. J. MARTIN. 1914. Observations on the mechanism of the transmission of plague by fleas. J. Hyg. 13. Plague suppl. 3: 423-439.
- BALTAZAR, M., M. BAHMANYAR, C. MOJIDI & B. SEYDIAN. 1952. Le foyer de peste lu Kurdistan. Bull. World Health Organization 6: 441-472.
- ELAIR, W. I. 1940. Home ranges and population of the meadow vole in southern Michigan. J. Wildlife Mgt. 4: 149-161.

- BLANC C 1956 Une opinion non conformiste sur la mode de transmission de la peste
Rev hyg méd soc 4 535-567
- BURROUGHS A L 1944 The flea *Xenopsylla cheopis* a vector of *P. pestis* Proc Soc
Exptl Biol Med 55 10-11
- BURROUGHS A L 1947 Sylvatic plague studies. The vector efficiency of nine species of
fleas compared with *Xenopsylla cheopis* J Hyg 45 371-386
- CHEN T H 1952 Studies on immunization against plague. IV. The method of the hemag-
glutination test and some observations on the antigen J Immunol 69 587-596
- COSTA LIMA A DA & C R HATHAWAY 1946 Pulgas. Monogr Inst Oswaldo Cruz
No 4 Rio de Janeiro Brazil
- DAVIS D H S 1939 Some ecological methods in research on bubonic plague S Afr
can J Sci 36 438-444
- DAVIS D H S 1948 Sylvatic plague in South Africa. History of plague in man 1919-43
Ann Trop Med Parasitol 42 20-217
- DAVIS D H S 1951 Rodent control in gold fields area. Union of S Africa Dept Health
Plague Research Lab Johannesburg Spec Rept No 21
- DAVIS D H S 1953 Plague in Africa from 1935 to 1949: a survey of wild rodents in
African territories Bull World Health Organization 9 665-700
- DEIGNAT R 1946 Aspects de l'épidémiologie de la peste au Lac Albert Ann soc
belge méd trop 26 13-34
- FSKEY C R 1938 Flea infestation of domestic rats in San Francisco California Public
Health Repts U S 53 948-951
- FSKEY C R & V H HAAS 1940 Plague in the western part of the United States U
S Public Health Service Public Health Bull No 254
- FOX I 1956 Murine typhus fever and rat ectoparasites in Puerto Rico Am J Trop
Med Hyg 5 893-900
- GAISKI N A 1976 Plague in marmosets by periods of the year Rev Microbiol Sarato
5 3
- GARNHAM P C C 1949 Distribution of wild rodent plague Bull World Health Or-
ganization 2 271-278
- GEAR H S & Z DEUTSCHMAN 1956 Disease control and international travel: a review
of the international sanitary regulations Spec No Chronicle World Health Organi-
zation 10(9-10) 273-343
- GIRARD G 1940 Foyers persistants de peste murine à Tananarive Bull soc pathol
exotique 3 209-211
- GIRARD G 1955 Plague Ann Rev Microbiol 9 253-266
- GRUBBS S B 1977 Bubonic plague and maritime quarantine Public Health Repts
U S 42 2045-2055
- HAYNE D W 1950 Apparent home range of *Microtus* in relation to distance between
traps J Mammal 31 26-39
- HEISCH R B 1956 Zoonoses as a study in ecology with special reference to plague re-
lapsing fever and leishmaniasis Brit Med J 2 669
- HITCH R B W E GRAINGER & J ST A M D SOLZA 1953 Results of a plague in-
vestigation in Kenya Trans Roy Soc Trop Med Hyg 47 503-521
- HIRST I F 1976 Researches on the parasitology of plague Part I Ceylon J Sci D
1 155-271
- HIRST I F 1953 The Conquest of Plague Oxford Univ Press London England
- HOLDENRIED R 1951 Program of the Santa Fe Field Station New Mexico Health
Officer 19 17-20
- HOLDENRIED R & H B MORLAN 1955 Plague infected fleas from northern New Mexico
wild rodents J Infectious Diseases 96 133-137
- HOLDENRIED R & S F QUAN 1956 Susceptibility of New Mexico rodents to experi-
mental plague Public Health Repts U S 71 979-984
- HOLLAND G 1957 Notes on the genus *Hystriolopsylla* Roths in the New World with
description of one new species and two new sub-species (Siphonaptera: Hystri-
chopsyllidae) Can Entomologist 89 309-324
- HOOPER E T 1944 San Francisco Bay as a factor influencing prevalence in rodents
Univ Mich Museum Zool Misc Publ No 59
- HOOPER FOUNDATION 1946 University of California Medical Center San Francisco
Calif Collected Reprints 17 (1939-1946)
- HUBBARD C A 1947 Fleas of Western North America Iowa State College Press
Ames Ia
- JELLISON W I & N I GOOP 1942 Index to the literature of Siphonaptera of North
America Natl Inst Health Bull 178

- JELLISON W L B LOCKER & R F BACON 1953 Index to the literature of Siphonaptera of North America Rocky Mt Lab Natl Inst Health Suppl 1
- JONES R W & T M STREET 1956 A half century of plague in California. Bur Vector Control Calif State Dept Health Berkeley Calif Unpublished data
- JORDAN K 1948 Suctoria. In Smart A Handbook for the Identification of Insects of Medical Importance Brit Museum Natural Hist London 211
- KARTMAN L 1954 Studies on *Pasteurella pestis* in fleas I An apparatus for the experimental feeding of fleas Exptl Parasitol 3 525-537
- KARTMAN L 1957 The concept of vector efficiency in experimental studies of plague Exptl Parasitol 6 599-609
- KARTMAN L & R P LOVERGAN 1955a Wild rodent flea control in rural areas of an enzootic plague region in Hawaii a preliminary investigation of methods Bull World Health Organization 13 49-68
- KARTMAN L & R P LOVERGAN 1955b Observations on rats in an enzootic plague region of Hawaii Public Health Repts U S 70 585-593
- KARTMAN L & F M PRINCE 1956 Studies on *Pasteurella pestis* in fleas V The experimental plague vector efficiency of wild rodent fleas compared with *Xenopsylla cheopis* together with observations on the influence of temperature Am J Trop Med Hyg 6 1058-1070
- KARTMAN L F M PRINCE & S F QUAN 1956 Studies on *Pasteurella pestis* in fleas III Comparative plague vector efficiency of *Xenopsylla vexabilis hawaiiensis* and *Xenopsylla cheopis* Bull World Health Organization 14 681-704
- KARTMAN L V I MILES & F M PRINCE 1958 Ecological studies of wild rodent plague in the San Francisco Bay area of California I Introduction Am J Trop Med Hyg In press
- LINK V B 1950a Plague epizootic in cottontail rabbits Public Health Repts U S 65 696
- LINK V B 1950b Plague J Am Med Assoc 144 315-377
- LISTON W G 1924 The Milroy Lectures on the plague I The history of plague Brit Med J 1 900-903
- LOBANOV V W & V FEDEROV 1938 On the pathogenesis of experimental plague in southern gerbils (*Pallasiomys meridianus* Pall.) Izv Microbiol Saratov 17 57-71
- MCCOY G W 1908 Plague in ground squirrels Public Health Repts U S 23 1289-1293
- MCCOY G W 1910 Plague among ground squirrels in America J Hyg 10 589-601
- MCCRUMB F R S MERCIER J ROBIN M BOUILLAY J E SMADEL T F WOODWARD & K GOODNER 1953 Chloramphenicol and terramycin in the treatment of pneumonic plague Am J Med 14 284-293
- MACCHIAVELLO A 1948 Epidemiologia de la peste en las Américas Proc 4th Intern Congr Trop Med Malaria Washington D C 1 240-249
- MACCHIAVELLO A 1954 Reservoirs and vectors of plague J Trop Med Hyg 57
- MEYER K F 1942a The known and the unknown in plague Am J Trop Med 22 9-36
- MEYER K F 1942b The ecology of plague Medicine 21 143-174
- MEYER K F 1955 The modern outlook on plague in California Bur Vector Control Calif State Dept Public Health Berkeley Calif Vector Views 2 41 43
- MEYER K F & R HOLDENRIED 1949 Rodents and fleas in a plague epizootic in a rural area of California Puerto Rico J Public Health Trop Med 24 201-209
- MEYER K F S F QUAN F R MCCRUMB & A IARSON 1952 Effective treatment of plague Ann N Y Acad Sci 65(6) 1278-1274
- MEYER K F H SONMER L F FOSTER & S F QUAN 1950 Hooper Foundation Univ Calif Med Center San Francisco Calif Unpublished data
- MILES V I & M J WILCOB 1953 Control of native rodent fleas with DDT applied to simulate aerial dusting J Econ Entomol 46 255-257
- MILES V I A R KINNEY & H E STARR 1951 Flea host relationships of a so-called *Rattus* and native wild rodents in the San Francisco Bay area of California with special reference to plague Am J Trop Med Hyg 6 752-760
- MORLAN H B 1955 Mammal fleas of Santa Fe County New Mexico Texas Repts Biol and Med 13 93-125
- PAYLOVSKA E N 1955 The actual state of the doctrine of natural mortality of diseases of man (Abstr.) Pev Appl Entomol B44 145 146
- PETRIE G I 1929 In A System of Bacteriology C Brit Med Research Council
- PLAGUE RESEARCH COMMISSION IN INDIA 1906-1915 J Hyg 6-8 14 Mo Suppl 4
- POLLITZER R 1954 Plague World Health Organization Monogr Ser No 22

- PRINCE F M 1943 Species of fleas on rats collected in states west of the 103d meridian and their relation to the dissemination of plague. *Public Health Repts U S* 58: 400-408
- QUAN S F, L F FOSTER, A I ARON & K F MEYER 1947 Streptomycin in experimental plague. *Proc Soc Exptl Biol Med* 66: 578-532
- QUAN S F, A L BERROUGH, R HOLDENRIED & K F MEYER 1953 Studies on the prevention of experimental plague epizootics institute I among mice by infected fleas. *6th Congr Intern Microbiol* 5: 537-540
- QUAN S F, A G McMANUS & K F MEYER 1955 Comparisons of sulfisoxazole with sulfalazine and thiocymetin with chl ramphenicol in chemotherapy of experimental plague in mice. *Am J Trop Med Hyg* 4: 1078-1036
- QUAN S I, H VON FINTEL & A G McMANUS 1958 Ecological studies of wild rodent plague in the San Francisco Bay Area of California. II Efficiency of bacterial culture compared to animal inoculation as methods for detecting *Yersinia pestis* in wild rodent fleas. Unpublished data
- QUAN S F & L KARTMAN 1956 The resistance of *Microtus* and *Peromyscus* to infection by *Pasteurella pestis*. *Trans Roy Soc Trop Med Hyg* 50: 104-105
- QUAN S F, W A HARTWELL, K C SCOTT & C T PENG 1957 Cerium as a tag for arthropods of medical importance. *Trans Roy Soc Trop Med Hyg* 51: 8-88
- RYCKMAN R F, C T AMES & C C LINDT 1953 A comparison of aldrin, dieldrin, heptachlor and DDT for control of plague vectors on the California ground squirrel. *J Econ Entomol* 46: 598-601
- SCOTT H H 1942 *A History of Tropical Medicine*. 2: 767. Edward Arnold, London, England
- SIMMONS S W 1954 Public health significance of insect resistance to insecticides and future orientation of vector control programs. 1st Intern Symposium: Control of Insect Vectors of Disease. Kendall super sanit 97: 125
- SMUEL J E, K GOODNER & T F WOODWARD 1955 The control of plague. In *Recent Advances in Medicine and Surgery*. Army Med Service Grad School Med Sci Publ 1: No 4: 2: 234-244
- SNOODGRASS R E 1946 The skeletal anatomy of fleas (Siphonaptera). *Smithsonian Inst Misc Collections* 104: 1-89
- STARK H E, A I MILES, F M PRINCE, L KARTMAN & A R KINNEY 1958 Ecological studies of wild rodent plague in the San Francisco Bay Area of California. III Occurrence and interrelationships of fleas on coexisting wild and domestic rodents. Unpublished data
- STEWART M A & F C EVANS 1941 A comparative study of rodent and burrow flea populations. *Proc Soc Exptl Biol Med* 47: 140-142
- TINKER I S & A I KALABUKHOV 1934 Life cycle of small marmot in relation to development of epizootic modification in susceptibility of marmots as influenced by age and sex. *Rev Microbiol Saratov* 13: 299-307
- VARELA G & A VAZQUEZ 1954 Halla go de la peste selatica en la Republica Mexicana. *Infeccion del Cy-mys mexicanus* (perros llaneos) con *Pasteurella pestis*. *Rev inst salubridad y enfermedad trop Mex* 14: 219-223
- WAGLE P M & S C SEAL 1953 Application of DDT, BHC and cyan gas in the control of plague in India. *Bull World Health Organization* 9: 59-614
- WHEELER C M & J R DOUGLAS 1941 Transmission studies of sylvatic plague. *Proc Soc Exptl Biol Med* 47: 65-66
- WHEELER C M & J R DOUGLAS 1945 Sylvatic plague studies. V The determination of vector efficiency. *J Infectious Diseases* 77: 1-12
- WHERRY W B 1908 Plague among the ground squirrels of California. *J Infectious Diseases* 5: 483-506
- WINTER C C & M D MOODY 1957 Rapid identification of *Pasteurella pestis* with fluorescent antibody. *Bacteriol Proc* 146-147
- WORLD HEALTH ORGANIZATION 1951 International Sanitary Regulations. World Health Organization Regulations No 2. World Health Organization Tech Rept Ser No 41
- WORLD HEALTH ORGANIZATION 1953 Expert Committee on Plague. 2nd report. *Bull World Health Organization* 9: 707-729

ANIMAL DISEASE AND BIOLOGICAL WARFARE

By Jack H. Hempy

Office of the Surgeon General, United States Air Force, Washington, D. C.

Any discussion of biological warfare (BW) and defense against it must include a serious consideration of animal diseases.¹ The close and vital relationship of animal disease to human health and well being in the field of public health holds equally true in the field of BW.

By way of definition BW is the intentional use of living microorganisms or their toxic products for the purpose of destroying or reducing the military effectiveness of man. This includes damage to or destruction of food sources and supplies. The military objective, as in the case of all weapon systems, is to reduce the will or capability to wage war.

The statement has been made that BW is public health practice and procedure in reverse. This is a common misconception. The difference rests in the word deliberate. BW is the deliberate use of natural disease agencies the inherent potential of which has been exploited by scientific research and development resulting in the production of BW weapon systems.

Medical and military histories both provide us with the proper background for a review of modern day concepts of BW. They tell us that military campaigns and troop concentrations have always provided fertile fields for naturally occurring epidemic disease and furthermore that infectious disease has often been the critical factor in the outcome of a military campaign. Bubonic plague was credited with stopping the Crusaders at the very gates of Jerusalem. Dysentery probably caused more casualties in Napoleon's Grand Army than did enemy firearms. Typhoid fever and the dysenteries seriously hampered both sides in the Civil War and in the Spanish American War. Even as late as World War I infectious disease was a controlling factor in some campaigns. It is quite clear that typhus fever prevented the Germans from carrying through their Balkan campaign. In World War II infectious diseases continued to hamper military forces materially. Malaria ranked high as an enemy to United States forces in the Mediterranean and Southwest Pacific theaters. Scrub typhus alone accounted for about 7000 casualties in the latter area. Finally, despite our most modern sanitary and preventive medicine practices we know that enteric infectious Japanese B encephalitis and hemorrhagic fever posed serious problems to the armed forces of the United States in Korea.

What of the deliberate use of infectious disease for military purposes? History records many instances of it. In the Fourth Century B. C. for example Alexander the Great attempted such exploitation by catapulting the bodies of dead men and animals over the walls of besieged cities. It is recorded that smallpox was spread successfully among the American Indians during the French and Indian War by the distribution of blankets contaminated with purulent smallpox material. In more modern times during World War I the Germans infected with glanders horses that were consigned from the United States to the Romanian cavalry. In World War II German occupation forces

in Eastern Europe were said to have been the target of local sabotage efforts with bacteriological agents

Obviously these efforts were small local makeshift and unorganized. It was not until early in World War II that an officially planned program was devoted to research and development in biological warfare. This work has continued to the present time as a recognized activity of the United States Department of Defense. Responsibility for carrying out this program was delegated to the Department of the Army which in turn assigned the operational responsibility to the Army Chemical Corps. The major portion of the research and development is conducted at Fort Detrick, Frederick, Md. Close liaison is maintained with other federal agencies in the United States having defensive responsibilities including the Federal Civil Defense Agency, the Public Health Service, the Food and Drug Administration, and the Department of Agriculture. The fact that a program of research and development continues in permanent facilities constructed for the purpose can be considered as recognition of the potential effectiveness of this weapon and thus of the defense problems that we must be prepared to meet.

The nature of BW dictates its use primarily as a strategic weapon. The major reason for this is that it has no quick kill effect. The incubation period of infectious disease plus a variable period of illness even before a lethal effect render this weapon unsuitable for hand to hand encounter. A man can be an effective fighting machine throughout the incubation period of most infectious diseases. Thus it is logical to assume that an enemy would probably consider this weapon as primarily suited for attack on static population centers such as large cities. Consequently it would appear that a logical approach in BW for an enemy would be to create an aerosol or cloud of the agent over the target areas.

A number of unique medical problems will be created if and when man is exposed to an infectious agent through the respiratory route rather than by the natural portal of entry. Botulinal toxin, for example, is several thousandfold more toxic by this route than when given *per os*. In some instances a different clinical disease picture may result from this route of exposure making diagnosis difficult. In tularemia produced by aerosol exposure, for example, one would not expect to find the classic ulcer of rabbit fever on a finger.

What about agents? There are a number of them that an enemy might select from the several classes of microorganisms (bacteria, viruses, rickettsiae, and fungi). There are, however, certain general characteristics that would apply in making a selection.

An enemy obviously would choose an agent that is believed to be highly infectious. Agents that are known to cause frequent infections among laboratory workers such as those causing Q fever, tularemia, brucellosis, glanders, and coccidioidomycosis all belong in this category.

Any agent selected would probably possess sufficient viability and virulence stability to meet realistic minimal logistic requirements. In this connection it should be capable of being disseminated without undergoing excessive destruction. Moreover, it should not be so fastidious in its growth requirements as to make production on a militarily significant scale improbable.

ANIMAL DISEASE AND BIOLOGICAL WARFARE

By Jack H. Hempy

Office of the Surgeon General United States Air Force Washington D C

Any discussion of biological warfare (BW) and defense against it must include a serious consideration of animal diseases¹. The close and vital relationship of animal disease to human health and well being in the field of public health holds equally true in the field of BW.

By way of definition BW is the intentional use of living microorganisms or their toxic products for the purpose of destroying or reducing the military effectiveness of man. This includes damage to or destruction of food sources and supplies. The military objective as in the case of all weapon systems is to reduce the will or capability to wage war.

The statement has been made that BW is public health practice and procedure in reverse. This is a common misconception. The difference rests in the word deliberate. BW is the deliberate use of natural disease agencies the inherent potential of which has been exploited by scientific research and development resulting in the production of BW weapon systems.

Medical and military histories both provide us with the proper background for a review of modern day concepts of BW. They tell us that military campaigns and troop concentrations have always provided fertile fields for naturally occurring epidemic disease and furthermore that infectious disease has often been the critical factor in the outcome of a military campaign. Bubonic plague was credited with stopping the Crusaders at the very gates of Jerusalem. Dysentery probably caused more casualties in Napoleon's Grand Army than did enemy firearms. Typhoid fever and the dysenteries seriously hampered both sides in the Civil War and in the Spanish American War. Even as late as World War I infectious disease was a controlling factor in some campaigns. It is quite clear that typhus fever prevented the Germans from carrying through their Balkan campaign. In World War II infectious diseases continued to hamper military forces materially. Malaria ranked high as an enemy to United States forces in the Mediterranean and Southwest Pacific theaters. Scrub typhus alone accounted for about 7000 casualties in the latter area. Finally despite our most modern sanitary and preventive medicine practices we know that enteric infectious Japanese B encephalitis and hemorrhagic fever posed serious problems to the armed forces of the United States in Korea.

What of the deliberate use of infectious disease for military purposes? History records many instances of it. In the Fourth Century B. C. for example Alexander the Great attempted such exploitation by catapulting the bodies of dead men and animals over the walls of besieged cities. It is recorded that smallpox was spread successfully among the American Indians during the French and Indian War by the distribution of blankets contaminated with purulent smallpox material. In more modern times during World War I the Germans infected with glanders horses that were consigned from the United States to the Romanian cavalry. In World War II German occupation forces

in Eastern Europe were said to have been the target of local sabotage efforts with bacteriological agents

Obviously these efforts were small local makeshift and unorganized. It was not until early in World War II that an officially planned program was devoted to research and development in biological warfare. This work has continued to the present time as a recognized activity of the United States Department of Defense. Responsibility for carrying out this program was delegated to the Department of the Army which in turn assigned the operational responsibility to the Army Chemical Corps. The major portion of the research and development is conducted at Fort Detrick, Frederick, Md. Close liaison is maintained with other federal agencies in the United States having defensive responsibilities including the Federal Civil Defense Agency, the Public Health Service, the Food and Drug Administration, and the Department of Agriculture. The fact that a program of research and development continues in permanent facilities constructed for the purpose can be considered as recognition of the potential effectiveness of this weapon and thus of the defense problems that we must be prepared to meet.

The nature of BW dictates its use primarily as a strategic weapon. The major reason for this is that it has no quick kill effect. The incubation period of infectious disease plus a variable period of illness even before a lethal effect render this weapon unsuitable for hand-to-hand encounter. A man can be an effective fighting machine throughout the incubation period of most infectious diseases. Thus it is logical to assume that an enemy would probably consider this weapon as primarily suited for attack on static population centers such as large cities. Consequently, it would appear that a logical approach in BW for an enemy would be to create an aerosol or cloud of the agent over the target areas.

A number of unique medical problems will be created if and when man is exposed to an infectious agent through the respiratory route rather than by the natural portal of entry. Botulinal toxin, for example, is several thousandfold more toxic by this route than when given *per os*. In some instances a different clinical disease picture may result from this route of exposure, making diagnosis difficult. In tularemia produced by aerosol exposure, for example, one would not expect to find the classic ulcer of rabbit fever on a finger.

What about agents? There are a number of them that an enemy might select from the several classes of microorganisms (bacteria, viruses, rickettsiae, and fungi). There are, however, certain general characteristics that would apply in making a selection.

An enemy obviously would choose an agent that is believed to be highly infectious. Agents that are known to cause frequent infections among laboratory workers such as those causing Q fever, tularemia, brucellosis, glanders, and coccidioidomycosis, all belong in this category.

Any agent selected would probably possess sufficient viability and virulence stability to meet realistic minimal logistic requirements. In this connection it should be capable of being disseminated without undergoing excessive destruction. Moreover, it should not be so fastidious in its growth requirements as to make production on a militarily significant scale improbable.

An aggressor would seek minimal naturally acquired or artificially induced immunity in a target population. A solid immunity in the target population is the one effective circumstance whereby attack by a specific agent can be neutralized. It must be remembered, however, that there are many agents for which there is no solid immunity and that a partial or low grade immunity may be broken by an appropriate dose of the agent.

There is a broad spectrum of agents from which selection might be made for a specified military purpose. An enemy might choose an acutely debilitating agent, one that produces chronic disease, or one that is highly lethal.

It is possible that certain mutational forms, such as drug resistant strains, may be produced. Mutants may be developed with changes in biochemical properties that are of importance in identification. All these considerations are of critical importance in considering defensive measures. However, despite some of the unsubstantiated claims that have been made from time to time, the likelihood of creating an entirely new agent of unique virulence or new disease producing capacity is extremely remote. Even the remarkable genetic progress made in recent years in the production of bacterial transformation does not warrant deviation from this opinion at the present time.

There are other general considerations in connection with BW agents that merit some consideration.

Biological agents are, of course, highly host specific. They do not destroy physical structures, as is true of high explosives. This may be of overriding importance in considering some military objectives.

One must be unrelenting in emphasizing that there is no secrecy concerning agents that might be included in an overall BW arsenal. Only certain agents will meet the general and specific BW requirements, and we and any potential enemy both know them. The production of these agents is not like the invention or, rather, the synthesis of a new chemical poison. One frequently hears it said that, "If we only knew on what agents our potential enemies were working, we should know against what to defend ourselves." This is the kind of statement that might be made by an ostrich before burying its head in the sand. It would be more appropriate to ask ourselves, "What are we doing about it? Are we doing enough?"

In defense against such attacks, we must also consider that biological agents are suitable for delivery through enemy sabotage. We could let our imaginations run wild in this regard. I might mention a few obvious areas of significant importance. Food processing, including the preparation of soft drinks and the processing of milk and milk products, constitutes an enormous industry that is subject to sabotage. Huge industries are also involved in the production of biological products, drugs, and cosmetics that are vulnerable to this type of attack. I mention these areas because sabotage in them would be far reaching in its consequences.

The possibility of overt military delivery of biological agents from appropriate disseminating devices also must be considered. It should be no more difficult to deliver such devices than other weapons. The same means of delivery, whether airplanes, submarines, or guided missiles, should be usable.

If it is possible for an enemy to drop an atomic bomb on a city it should be equally possible to put a cloud of some biological agent over that same city.

Another defense consideration is the possible use of biological agents for the reduction or destruction of agricultural crops and domestic animals—in other words antifeed biological warfare. The importance of food particularly during war needs no emphasis. Actually food production is of major concern to most countries even during peacetime.

BW may find its greatest effectiveness when used against animals. We must also consider that the epizootic potential of antianimal agents would be exploited by an enemy. Antifeed including antianimal BW could play a decisive role in any war not decided at its onset.

Thus far we have touched on certain inherent capabilities of BW and the general requirements its agents should possess. In so doing we find that BW might be employed overtly or covertly against man, livestock, and crops. With this as a framework for defensive thinking let us now consider a few of the general features of the problem of defense.

It is worth while to emphasize that there is a vast amount of medical knowledge in existence that can be useful. We have had long medical and epidemiological experience with infectious diseases in both man and animal. We have a vast public health effort in existence at the federal, state, and local levels. Our preventive medicine practices and methods for disease control are at a high level of efficiency. All of these are positive values that can be drawn upon now and in an emergency.

This leaves no room for complacency, however, and we must avoid being lulled into thinking that these aids would render BW ineffective. Unfortunately this is not the case. All of these techniques have been developed over the years for dealing with naturally occurring infectious disease. The military exploitation of massive amounts of highly infectious agents through unusual portals of entry creates many new problems for which these procedures were not designed and against which we have no experience. For example, public health practices have been developed for delivering wholesome food to all inhabitants in a community, and today we take this for granted. On the other hand, there is no known public health procedure that will deliver sterile air to all inhabitants of a city. Defense against a massive biological aerosol is a new and critically serious problem.

In the light of such a possibility, prophylaxis by active biological immunization may well be considered the most important defensive procedure of all. A number of effective immunizing materials are already available for some infectious diseases. On the other hand, there are a number of potential BW agents against which there is no known method of immunization. In other cases the value of the immunizing material continues to be questionable.

Consequently we must encourage all possible research that is devoted to the development of new methods of active immunization or the improvement of existing ones. Ancillary research dealing with host-parasite relationships will also have positive defense value. As a further incentive, research of this nature has great peacetime value.

An aggressor would seek minimal naturally acquired or artificially induced immunity in a target population. A solid immunity in the target population is the one effective circumstance whereby attack by a specific agent can be neutralized. It must be remembered however that there are many agents for which there is no solid immunity and that a partial or low grade immunity may be broken by an appropriate dose of the agent.

There is a broad spectrum of agents from which selection might be made for a specified military purpose. An enemy might choose an acutely debilitating agent, one that produces chronic disease, or one that is highly lethal.

It is possible that certain mutational forms, such as drug resistant strains, may be produced. Mutants may be developed with changes in biochemical properties that are of importance in identification. All these considerations are of critical importance in considering defensive measures. However, despite some of the unsubstantiated claims that have been made from time to time, the likelihood of creating an entirely new agent of unique virulence or new disease producing capacity is extremely remote. Even the remarkable genetic progress made in recent years in the production of bacterial transformation does not warrant deviation from this opinion at the present time.

There are other general considerations in connection with BW agents that merit some consideration.

Biological agents are, of course, highly host specific. They do not destroy physical structures, as is true of high explosives. This may be of overriding importance in considering some military objectives.

One must be unremitting in emphasizing that there is no secrecy concerning agents that might be included in an over-all BW arsenal. Only certain agents will meet the general and specific BW requirements, and we and any potential enemy both know them. The production of these agents is not like the invention or rather the synthesis of a new chemical poison. One frequently hears it said that, "If we only knew on what agents our potential enemies were working, we should know against what to defend ourselves." This is the kind of statement that might be made by an ostrich before burying its head in the sand. It would be more appropriate to ask ourselves, "What are we doing about it? Are we doing enough?"

In defense against such attacks we must also consider that biological agents are suitable for delivery through enemy sabotage. We could let our imaginations run wild in this regard. I might mention a few obvious areas of significant importance. Food processing, including the preparation of soft drinks and the processing of milk and milk products, constitutes an enormous industry that is subject to sabotage. Huge industries are also involved in the production of biological products, drugs, and cosmetics that are vulnerable to this type of attack. I mention these areas because sabotage in them would be far reaching in its consequences.

The possibility of overt military delivery of biological agents from appropriate disseminating devices also must be considered. It should be no more difficult to deliver such devices than other weapons. The same means of delivery, whether airplanes, submarines, or guided missiles, should be usable.

If it is possible for an enemy to drop an atomic bomb on a city it should be equally possible to put a cloud of some biological agent over that same city.

Another defense consideration is the possible use of biological agents for the reduction or destruction of agricultural crops and domestic animals—in other words, antifeed biological warfare. The importance of food, particularly during war, needs no emphasis. Actually, food production is of major concern to most countries even during peacetime.

BW may find its greatest effectiveness when used against animals. We must also consider that the epizootic potential of antianimal agents would be exploited by an enemy. Antifeed, including antianimal BW, could play a decisive role in any war not decided at its onset.

Thus far we have touched on certain inherent capabilities of BW and the general requirements its agents should possess. In so doing we find that BW might be employed overtly or covertly against man, livestock, and crops. With this as a framework for defensive thinking, let us now consider a few of the general features of the problem of defense.

It is worth while to emphasize that there is a vast amount of medical knowledge in existence that can be useful. We have had long medical and epidemiological experience with infectious diseases in both man and animal. We have a vast public health effort in existence at the federal, state, and local levels. Our preventive medicine practices and methods for disease control are at a high level of efficiency. All of these are positive values that can be drawn upon now and in an emergency.

This leaves no room for complacency, however, and we must avoid being lulled into thinking that these aids would render BW ineffective. Unfortunately, this is not the case. All of these techniques have been developed over the years for dealing with naturally occurring infectious disease. The military exploitation of massive amounts of highly infectious agents through unusual portals of entry creates many new problems for which these procedures were not designed and against which we have no experience. For example, public health practices have been developed for delivering wholesome food to all inhabitants in a community and today we take this for granted. On the other hand, there is no known public health procedure that will deliver sterile air to all inhabitants of a city. Defense against a massive biological aerosol is a new and critically serious problem.

In the light of such a possibility, prophylaxis by active biological immunization may well be considered the most important defensive procedure of all. A number of effective immunizing materials are already available for some infectious diseases. On the other hand, there are a number of potential BW agents against which there is no known method of immunization. In other cases the value of the immunizing material continues to be questionable.

Consequently, we must encourage all possible research that is devoted to the development of new methods of active immunization or the improvement of existing ones. Ancillary research dealing with host-parasite relationships will also have positive defense value. As a further incentive, research of this nature has great peacetime value.

An aggressor would seek minimal naturally acquired or artificially induced immunity in a target population. A solid immunity in the target population is the one effective circumstance whereby attack by a specific agent can be neutralized. It must be remembered however that there are many agents for which there is no solid immunity and that a partial or low grade immunity may be broken by an appropriate dose of the agent.

There is a broad spectrum of agents from which selection might be made for a specified military purpose. An enemy might choose an acutely debilitating agent, one that produces chronic disease, or one that is highly lethal.

It is possible that certain mutational forms, such as drug resistant strains, may be produced. Mutants may be developed with changes in biochemical properties that are of importance in identification. All these considerations are of critical importance in considering defensive measures. However, despite some of the unsubstantiated claims that have been made from time to time, the likelihood of creating an entirely new agent of unique virulence or new disease-producing capacity is extremely remote. Even the remarkable genetic progress made in recent years in the production of bacterial transformation does not warrant deviation from this opinion at the present time.

There are other general considerations in connection with BW agents that merit some consideration.

Biological agents are of course highly host specific. They do not destroy physical structures, as is true of high explosives. This may be of overriding importance in considering some military objectives.

One must be unrelenting in emphasizing that there is no secrecy concerning agents that might be included in an overall BW arsenal. Only certain agents will meet the general and specific BW requirements, and we and any potential enemy both know them. The production of these agents is not like the invention or rather the synthesis of a new chemical poison. One frequently hears it said that 'If we only knew on what agents our potential enemies were working, we should know against what to defend ourselves.' This is the kind of statement that might be made by an ostrich before burying its head in the sand. It would be more appropriate to ask ourselves, 'What are we doing about it? Are we doing enough?'

In defense against such attacks we must also consider that biological agents are suitable for delivery through enemy sabotage. We could let our imaginations run wild in this regard. I might mention a few obvious areas of significant importance. Food processing, including the preparation of soft drinks and the processing of milk and milk products, constitutes an enormous industry that is subject to sabotage. Huge industries are also involved in the production of biological products, drugs, and cosmetics that are vulnerable to this type of attack. I mention these areas because sabotage in them would be far reaching in its consequences.

The possibility of overt military delivery of biological agents from appropriate disseminating devices also must be considered. It should be no more difficult to deliver such devices than other weapons. The same means of delivery, whether airplanes, submarines, or guided missiles, should be usable.

Part VII Selected Chronic Diseases

A COMPARISON OF RHEUMATOID LIKE ARTHRITIS IN SWINE AND RHEUMATOID ARTHRITIS IN MAN

By Dennis Sikes

Department of Pathology and Parasitology, University of Georgia, Athens, Ga

The problem of arthritis in man, his domestic animals and other members of the vertebrate subkingdom dates to antiquity. The earliest known case of multiple arthritis is in a fossil vertebrate of a large swimming reptile, the plate carpus, which is calculated to have lived approximately 100 000 000 years ago, a skeleton of which is in the Museum of Natural History at the University of Kansas¹ Lawrence, Kan. According to Hollander² chronic arthritis of the spine was present in the ape man of 2 000 000 years ago, as well as in the Java and Lansing men of 500 000 years ago, and in Egyptian mummies dating to 8000 B.C. The Romans built baths throughout their empire because of this disease.

Hollander² estimates that more than 10 000 000 people in the United States have some form of rheumatic disease, and that more than 4 500 000 of these have some form of arthritis. Of those with arthritis, 190 000 are completely disabled and another 1 143 000 are partially disabled.

Swine erysipelas is an infectious and contagious disease of swine that causes a relatively high morbidity with a low mortality rate in most affected herds. The causative agent *Erysipelothrix rhusiopathiae* derives its name from the Greek *erysipelas* = red skin, *thrix* = a hair or thread, *rhusis* = reddish and *pathos* = a disease. According to Bergey³ the swine erysipelas organism has been assigned to the family Corynebacteriaceae and genus *Erysipelothrix*. The organism was described by Pasteur and Dumas⁴ and Loefler⁵ is credited with the first observation of the organism in swine. He observed the organism in the blood vessels of a pig that apparently had died from swine erysipelas and thought that it was similar to Koch's *Erysipelothrix muriseptica*. Later Rosenbach⁶ reported his classic studies of the organism and suggested the names *E. muriseptica*, *E. porci* and *E. erysipeloides* for the mouse, pig and human organisms, respectively.

Konst⁶ indicates the possibility that two variants of *E. rhusiopathiae* exist, the highly virulent one prevalent in Europe and a less virulent one in North America. The observations of Breed⁷ do not support this concept for the disease is spreading in North America and is of great economic importance. It may be that the more virulent strains have been imported in swine originating in Europe. The relationships of the microorganisms isolated from man, sheep, chickens, turkeys and swine have been studied by Beaudette and Hudson⁸, Klauder and Harkens⁹ and Van Roekel, Bullis and Clarke¹⁰ who were unable to show any significant cultural differences between strains.

The work reported in this paper was supported in part by Research Grant A 1120 (C2) from the National Institute of Arthritis and Metabolic Diseases, Public Health Service, Bethesda, Md.

Another defense procedure of potential value lies in the possible prophylactic use of antibiotics and chemotherapeutic drugs. However here again we must avoid being lulled into a false sense of security. For example published articles have expounded the thesis that our enormous antibiotic industry has made BW obsolete. This idea verges on the ridiculous for several reasons. To permit such reasoning to guide our defense planning would be dangerous indeed. There are many potential BW agents for which there is no known effective antibiotic or drug. Among these may be mentioned *Coccidioides immitis*, *Histoplasma capsulatum* and more importantly from a BW standpoint most of the filtrable viruses. Moreover what of the not unlikely possibility of drug resistant agents?

Much more space than is available here could be devoted to still other and equally significant defensive problems including detection, disease reporting, decontamination and protective equipment. As stated above my purpose here is merely to suggest some areas of investigation that are pertinent to the problems of defense against BW. Unfortunately in the world as it is today we must give serious attention to the negative aspect of the ever growing body of evidence that man's health and well being progress in direct relationship to his successes in the prevention, control and treatment of animal diseases.

Reference

1. FOTHERGILL, L. D. 1957. Biological warfare and its defense. Public Health Repts U S 72(11) 865-871.



FIGURE 1 Extensive erythema of the skin of a white hog with acute swine erysipelas following intensive exposure to *Erysipel thure rhust palliae*

glomerulosa was infiltrated with many leukocytes and complete or almost complete disappearance of this zone occurred in some cases. Numerous foci of round cells were often scattered throughout the fascicular and reticular zones. The medulla usually appeared normal but some showed atrophic or degenerative changes. In advanced cases of chronic arthritis a disruption of the normal architecture of the glomerulosa and a reduction in thickness occurred.

Collins and Goldie¹⁸ and Sikes, Neher and Doyle¹⁹ observed in the acute stages of *E. rhusiopathiae* arthritis that the principal changes in the joints were characterized by vascular engorgement of the joint capsule and synovial tissues and more or less edema. The effusions were turbid or serosanguineous or mucinous. The synovial villi showed beginning proliferation, extensive vasodilation, engorgement and incipient lymphocytic infiltration.

In arthritis of two months' duration greater proliferation and less edema were observed. Much proliferation of the cells covering the hypertrophied synovial villi was present. The villi contained young highly vascular connective tissue infiltrated with plasma cells and lymphocytes. A striking feature was the dense collection of lymphocytes (FIGURE 3). The villi of arthritic joints of six months' or longer duration frequently resembled granulomatous polyps (FIGURE 4). Very few polymorphonuclear leukocytes were observed and the infiltrating cells were principally lymphocytes, plasma cells and eosinophils. Less exten-

Natural outbreaks of *E. rhusiopathiae* infections have been recorded in sheep cattle horses turkeys ducks peacocks pheasants and farm raised mink (Beaudette and Hudson⁸ Graham Levine and Hester¹¹ Marsh¹ Paterson and Heatley¹²) Also according to Klauder Righter and Harkens¹⁴ human infections have occurred from handling fish

For more than a century in Europe the economic importance of swine erysipelas has been emphasized, but it has reached serious proportions only in the last two decades in the United States According to a study made by the Organization for European Economic Cooperation (OEEC)¹⁵ all of the European nations reported that the disease was affecting many herds of swine

In the United States arthritis in swine has rated second or third in responsibility for the number of swine carcasses condemned at federally inspected meat packing plants for the last ten years according to the annual reports of the Chief of the Bureau of Animal Industry United States Department of Agriculture¹⁶ The number of carcasses condemned in 1952 was 22 196 This represents only a small fraction of the true incidence of the disease complex because the animals sent to slaughter for human consumption are thought to be healthy It is generally known that lame animals are condemned and swine producers do not ordinarily send arthritic animals to market Obviously, the great majority of affected animals remain at the farm until they recover or die

Pathological Changes in Acute and Chronic Swine Erysipelas

The lesions of swine erysipelas are varied Sikes *et al*¹⁷ have shown that in animals that die of the acute septicemic form of the disease pathological changes observed at necropsy may be no different from those observed in any other septicemic disease The skin lesions perhaps are more nearly pathognomonic than any other finding (FIGURE 1) Damage to the blood vessels of the skin resulting in the peculiar type of erythema presumably was due to the action of the organism and its by products The histopathological changes occurred principally in the papillae and in the subpapillary and upper layers of the dermis The blood vessels were distended and the papillae were enlarged and showed marked hyperemia Frequently small hemorrhages were observed in the connective tissue elements of the papillae (FIGURE 2) Numerous foci of round cells and necrotic areas were present In the urticarial lesions marked edema of the skin occurred in the wheals Dilation of the vessels and numerous thrombi were present Capillary hemorrhages appeared especially in the papillary layer The chief difference in the skin lesions of swine with the acute septicemic form and those showing urticaria was in the degree of damage done

They also observed pronounced white blood cell dyscrasia A significant and marked leukopenia lymphopenia neutropenia and a monocytosis occurred in the acute phase of the disease This appears to be the most critical phase of the disease as related to survival or death of the animal In acute swine erysipelas the adrenal glands were enlarged and frequently subcapsular hemorrhages were observed In chronic cases lasting two months or more the adrenal glands were often swollen to twice the normal size and were very soft the zona



FIGURE 3 Proliferation of the synovial villi with accumulation of round cells $\times 110$

changes first then lesions of muscles nerves tendons bursae integument and serous membranes of the heart kidneys reticuloendothelial structures and intestines Collins⁹ also places great importance on the joint changes supplemented by other tissue change Both of these authors agree that the microscopic pathology of synovial membranes in rheumatoid arthritis is quite characteristic and that the basic pattern may be considered pathognomonic of rheumatoid arthritis as it is not exactly duplicated in any other articular disease The basic pattern consists of hypertrophy vascularization cellular infiltration accumulations of inflammatory cells in follicle like collections and pannus formation

Several components of the synovial membrane undergo hypertrophy The endothelial lining may reach several cells in thickness The stroma is increased in amount being fibrillar infiltrated with fat and obscured by cellular filtrate in some cases The villous folds of the lining become hypertrophied causing the villi to be enlarged

Vascularity of the synovium may be profuse or modest The walls may be thickened and whorls of fibrous tissue surround the medial coats The veins frequently are dilated

The cellularity is variable from case to case Lymphocytes plasma cells reticuloendothelial cells polymorphonuclear leukocytes fibroblasts and a few giant cells appear in the subendothelial layer



FIGURE 2 Hemorrhages in the skin of an experimentally produced case of acute swine erysipelas $\times 200$

sive destruction of cartilage and bone than that which characterizes suppurative arthritis in other species was observed. The synovial effusions were nonpurulent and contained mostly monocytes, lymphocytes, and a few synovial cells.

In advanced chronic arthritis with pannus formation of subchondral origin, marked fibrosis of the adjacent marrow spaces, increased vascularity, and collections of lymphocytes occurred. In those cases in which the pannus apparently originated from the synovial membrane, subchondral fibrosis was less likely to occur (FIGURE 5). Pannus attachment to the articular cartilage was accompanied by erosions and destruction of the cartilage. In advanced cases with little or no synovial fluid, intra-articular fibrous adhesions formed apparently in an attempt to immobilize the joint.

Radiograms revealed osteophyte formation with apparent ankylosis of the carpal and tarsal joints (FIGURE 6). Also, the articular processes of the vertebrae frequently appeared to be fused. Irregular ossification and rarefaction of bones adjacent to the joints were frequently observed. In hogs that spontaneously recovered, no gross abnormalities were observed.

Pathological Changes in Rheumatoid Arthritis in Man

The pathological alterations of the tissues in rheumatoid arthritis are widespread throughout the body, according to Hollander¹ who emphasizes joint



FIGURE 6 Photograph of an X ray showing closure of carpal spaces

Follicle like collections of chronic inflammatory cells are present in the synovial membranes of rheumatoid arthritis. These follicles consist mainly of lymphocytes and some reticuloendothelial cells. In some cases they are sparse in others they are collected densely throughout subendothelial tissues.

The formation of pannus is a most important phenomenon in the pathological process. It may arise from the lateral margins of the synovium or occasionally subchondrally. In any case it may appear reddened and rough with irregular margins resulting in the destruction of cartilage. In these cases fibrous tissue covers the epiphyseal bone thus leading to fibrous intra articular adhesions.

The pathological changes in epiphyseal bone near the affected joint are osteoporosis, follicle like accumulations of lymphocytes in marrow spaces and granulations, absorption of bone, eburnation of subchondral bone and production of osteoid tissue in the marginal zone.

In cases of spondylitis the discs show progressive atrophy with narrowing of intervertebral spaces. Osteoid tissue grows through the degenerated discs and later forms bone resulting in bony adherence of the vertebrae. In late cases intervertebral ligaments may become calcified. Other bones such as ischium and pubis may be affected.

Discussion

Erysipelothrix rhusiopathiae infection in swine may be acute, subacute or chronic. Frequently arthritis is a sequel to these forms of infection (Sikes, Neher and Doyle)¹⁷. A distinctive feature of the pathological process is its chronic proliferative nature without the formation of pus and destruction of



FIGURE 4 Granulomatous polyp formation in advanced arthritis $\times 140$



FIGURE 5 Section of carpal bone showing lateral pannus formation extending between articular surfaces

- 8 BEAUDOITTE F R & C B HUDON 1936 An outbreak of acute swine erysipelas in section in turkeys. J Am Vet Med Assoc 88 475-488
- 9 KLAUDER J V & M J HARKINS 1937 Erysipeloid in the United States. J Am Med Assoc 96 1205
- 10 VAN ROEKEL H K L BULLIS & M R CLARK 1938 Erysipelas outbreak in turkey flocks. J Am Vet Med Assoc 92 403-419
- 11 GRAHAM H K D LEVINE & H R HESTER 1939 Erysipell septic pathiae associated with a fatal disease in ducks. J Am Vet Med Assoc 95 211-216
- 12 MARSH H 1931 The bacillus of swine erysipelas associated with arthritis in lambs. J Am Vet Med Assoc 78 57-63
- 13 PATTERSON J S & T G HEATLEY 1939 A case of infection of the horse with *Erysipell septic pathiae*. Vet J 94 33-34
- 14 KLAUDER J V L RIGTER & M J HARKINS 1936 A distinctive and severe form of erysipeloid among fish handlers. Arch Dermatol Syphil 14 667-678
- 15 ORGANIZATION FOR EUROPEAN ECONOMIC COOPERATION (O.E.C.) 1952 The control of livestock diseases in European countries. Tech Assistance Mission No 63 Paris France
- 16 REPORT OF THE CHIEF OF THE BUREAU OF ANIMAL INDUSTRY 1952 Vet Research Adm. U.S. Dept Agr
- 17 SIKES D C M NEHER & I I DOYLE 1955 Studies on arthritis in swine. I. Experimental erysipelas and chronic arthritis in swine. Am J Vet Research 16(6) 342-366
- 18 COLLINS D H & W GOLDIE 1940 Observations on polyarthritis and on experimental Erysipell septic infection of swine. J Pathol Bacteriol 50(2) 323-353
- 19 SIKES D C M NEHER & I I DOYLE 1955 Studies on arthritis in swine. II The effects of hormonal therapy on advanced chronic polyarthritis experimentally induced by *Erysipell septic pathiae* infections. J Am Vet Research 16(6) 367-373
- 20 COLLINS D H 1949 The Pathology of Articular and Spinal Diseases. 11 ed. Wilkins & Wilkin Baltimore Md

bone that is characteristic of other bacterial infections of joints. Few or no polymorphonuclear leukocytes are present. The synovial effusions are sero-sanguineous and mucinous, similar to those observed in rheumatoid arthritis in man.

The formation of hypertrophied synovial fringes whose villi become bulbous with focal collections of lymphocytes resembling germinal centers is common to both diseases. The villi become greatly enlarged and granulomatous later in the disease and are very active even in the absence of an infectious agent. Sikes, Neher, and Doyle¹ were not able to isolate an infectious agent after 226 days after exposure, yet the disease persisted in an active form with periods of remission and exacerbation. Collins and Goldie¹⁸ believe that one of three things may happen in chronic arthritis due to *E. rhusiopathiae* infections: (1) the lesions may resolve and the joint may recover; (2) healing of the damaged tissue may take place by fibrosis, and this may include the formation of intra-articular adhesions; or (3) the synovial granulation tissue may assume certain properties of a benign neoplasm and persist and extend in the absence of the exciting microorganisms, thus maintaining the arthritis in a state of clinical activity. Just what factors are responsible for this excitement of the arthritic condition in the absence of an infectious agent remains to be discovered. Sikes, Neher, and Doyle¹⁹ have suggested that perhaps adrenal and kidney pathology play an important role in this syndrome.

Pannus formation, destruction of cartilage at the site of pannus attachment and subchondral cellular reaction are common to *E. rhusiopathiae* arthritis of swine and rheumatoid arthritis of man. Intra-articular fibrous adhesions and rarefaction of the bone as revealed by X-ray are also common to both diseases. Gross destruction of bone does not occur in either disease. Perhaps no further analogy should be drawn between the two maladies until the cause of rheumatoid arthritis in man is known.

Summary

A description of the pathology of *Erysipelothrix rhusiopathiae* infections in swine is given; the major pathological joint changes in rheumatoid arthritis in man are discussed, and a comparison of the pathology of *E. rhusiopathiae* arthritis in swine and rheumatoid arthritis in man is made.

References

1. HOLLANDER, J. L. 1953. *Comroe's Arthritis and Allied Conditions*. 5th ed. Lea and Febiger, Philadelphia, Pa.
2. BERGEY, D. H. 1948. *Manual of Determinative Bacteriology*. 6th ed. Williams & Wilkins, Baltimore, Md.
3. PASTEUR, L. & M. DUMAS. 1887. Sur le rouget ou mal rouge des pores. *Compt. rend.* 95: 1120-1121.
4. IOEFFLER, D. 1886. Experimentelle Untersuchungen über Schweinerotlauf. *Archiv für klin. u. exp. Med.* 1: 46-56.
5. KOFENBACH, F. J. 1909. Experimentell-morphologische und klinische Studien der Erreger der Schweinerotlaufs-Erysipelose und Mauseseptikämie. *Z. Hyg. Infektionskrankh.* 63: 343-371.
6. KONST, H. 1940. Chemotherapy of swine erysipelas. *Can. J. Comp. Med. Vet. Sci.* 9: 135-139.
7. BREED, F. 1938. Swine erysipelas: its distribution, increasing importance and control. *J. Am. Vet. Med. Assoc.* 92: 344-355.

workers on a variety of experimental animal neoplasms there has emerged a tentative explanation for some at least of the apparent contradictions between the observed epidemiology of neoplasm in natural populations and the concept of viral transmission. A group of transmissible neoplastic diseases of sheep has been chosen to serve as a possible model for the study of neoplasms occurring in natural populations.

These sheep diseases occur in a variety of forms ranging from infiltrative pneumonia to metastasizing carcinoma. In this paper they are considered to constitute together a complex of related diseases; for convenience they are referred to here as sheep pulmonary adenomatosis or SPA even though a few forms do not show adenomatous features. The reason for considering these various diseases as members of a single group will be discussed below.

Considered from either viewpoint the SPA complex is of particular interest since (1) at least some and probably all forms are infectious, (2) members of the SPA complex range in pathology from inflammatory or infiltrative to carcinomatous and (3) various forms of SPA while almost certainly of viral etiology can occur in both epizootic and enzootic form thus resembling both the picture of classic viral transmission (epizootic) and that of natural tumor epidemiology (occurring at low rates or enzootic) in which viruses are not ordinarily thought to be implicated. Examples of the last case are certain tumor conditions such as mouse leukemia,⁷ fowl lymphomatosis,⁸ and the mouse mammary gland carcinoma,⁹ which occur at enzootic levels in wild populations and yet were shown after the development of inbred strains of the host species involved to be propagable by means of viruses just as are many of the traditional epizootic inflammatory diseases. The study of SPA therefore may be useful in formulating unifying theories concerning the pathology and the etiology of inflammatory and neoplastic diseases and also in understanding the etiology of natural tumors.

The History of SPA

SPA was first recognized by J. M. Fadyean in 1888 as a rather rare adenomatous condition occurring in certain flocks of sheep in England.¹⁰ A report from Germany followed in 1899.¹¹ A commercially significant infiltrative pneumonia was observed in South Africa in 1904 by W. Robertson¹² and reported there again in 1915 by D. Mitchell.¹³ A chronic progressive pneumonia with some adenomatous overtones was reported from North America by H. Marsh¹⁴ in 1922 and a more adenomatous form was reported from France by M. Lesbouyries and A. Bonnac in 1940.¹⁵ Perhaps the best studied forms of SPA from the viewpoint of epidemiology and mode of transmission are two diseases of Icelandic sheep. One pulmonary adenomatosis is largely adenomatous in character; the other *Mædi* (an Icelandic word meaning dyspnea) is of the infiltrative type. Both forms which occur as separate but presumably related diseases were introduced by two karakul rams in a single shipment from Halle (Germany) in 1933. The diseases were not widely disseminated in Germany at that time and did not appear in Iceland until 1936; nevertheless it was possible to trace the infections back to the original carriers that had arrived three years earlier.¹⁶ The two diseases subsequently became grave commercial problems

THE PULMONARY ADENOMATOSIS COMPLEX IN SHEEP*

By Francisco Duran Reynals†

Department of Microbiology School of Medicine Yale University New Haven Conn

Erwin Jungherr

University of Connecticut Storrs Conn

Alberto Cuba Caparo

*University of San Marcos Lima Peru and the Department of Microbiology
School of Medicine Yale University*

Keen A Rafferty Jr

Department of Microbiology School of Medicine Yale University

and Charles Helmboldt

University of Connecticut

Introduction

With the discovery that certain tumors can be transmitted by means of cell free extracts interest naturally arose in possible comparisons between neoplastic viruses on the one hand and viruses of the traditional sort on the other. As a consequence of this interest attention has been drawn to situations in which viruses of classic inflammatory disease can give rise to disease that is histologically neoplastic. For example interest has revived in A. Borrel's studies of sheep pox and of fowl pox.¹ This investigator showed that the viruses of these diseases induced both necrotic and proliferative lesions and were in fact once designated as contagious epithelioma.¹ E. Roux observed pulmonary adenomatosis in sheep infected with sheep pox virus² and papillomas have been observed at the sites of the lesions of contagious pustular dermatitis in sheep.³ Smallpox cowpox rabbit pox mouse pox and other members of the pox group of viruses also can show a proliferative phase as well as the better known necrotizing phase.

Conversely viruses that ordinarily cause tumor formation can produce inflammatory and necrotizing disease under certain conditions according to dosage inoculation route host age and host resistance. F. Duran Reynals has demonstrated the inflammatory character of the Shope fibroma virus in newborn rabbits⁴ and of the Rous sarcoma virus in young chickens.⁵ In some recent work Duran Reynals⁶ has shown that mice pretreated with methyl cholanthrene and cortisone and then inoculated intradermally with vaccinia virus develop tumors rapidly in the sites of the healed vaccinia lesions. Although at present it is not possible to say whether vaccinia participates directly in the formation of these neoplasms it can be said at least that the vaccinia lesion appears to be an optimal ground for the development of malignancy.

A second important question for students of neoplasia is the role of viruses in the production of tumors in general. From observations made by many

The work reported in this paper was supported in part by the American Cancer Society, Inc., New York, N. Y.

† Deceased

workers on a variety of experimental animal neoplasms there has emerged a tentative explanation for some at least of the apparent contradictions between the observed epidemiology of neoplasm in natural populations and the concept of viral transmission. A group of transmissible neoplastic diseases of sheep has been chosen to serve as a possible model for the study of neoplasms occurring in natural populations.

These sheep diseases occur in a variety of forms ranging from infiltrative pneumonia to metastasizing carcinoma. In this paper they are considered to constitute together a complex of related diseases; for convenience they are referred to here as sheep pulmonary adenomatosis or SPA even though a few forms do not show adenomatous features. The reason for considering these various diseases as members of a single group will be discussed below.

Considered from either viewpoint the SPA complex is of particular interest since (1) at least some and probably all forms are infectious, (2) members of the SPA complex range in pathology from inflammatory or infiltrative to carcinomatous, and (3) various forms of SPA while almost certainly of viral etiology can occur in both epizootic and enzootic form thus resembling both the picture of classic viral transmission (epizootic) and that of natural tumor epidemiology (occurring at low rates or enzootic) in which viruses are not ordinarily thought to be implicated. Examples of the last case are certain tumor conditions such as mouse leukemia,⁷ fowl lymphomatosis,⁸ and the mouse mammary gland carcinoma,⁹ which occur at enzootic levels in wild populations and yet were shown after the development of inbred strains of the host species involved to be propagable by means of viruses just as are many of the traditional epizootic inflammatory diseases. The study of SPA therefore may be useful in formulating unifying theories concerning the pathology and the etiology of inflammatory and neoplastic diseases and also in understanding the etiology of natural tumors.

The History of SPA

SPA was first recognized by J. M. Fadyean in 1888 as a rather rare adenomatous condition occurring in certain flocks of sheep in England.¹⁰ A report from Germany followed in 1899.¹¹ A commercially significant infiltrative pneumonia was observed in South Africa in 1904 by W. Robertson¹² and reported there again in 1915 by D. Mitchell.¹³ A chronic progressive pneumonia with some adenomatous overtones was reported from North America by H. Marsh¹⁴ in 1922 and a more adenomatous form was reported from France by M. Lesbouyries and A. Bonnac in 1940.¹⁵ Perhaps the best studied forms of SPA from the viewpoint of epidemiology and mode of transmission are two diseases of Icelandic sheep. One pulmonary adenomatosis is largely adenomatous in character, the other *Maedi* (an Icelandic word meaning dyspnea) is of the infiltrative type. Both forms which occur as separate but presumably related diseases were introduced by two karakul rams in a single shipment from Halle, Germany in 1933. The diseases were not widely disseminated in Germany at that time and did not appear in Iceland until 1936; nevertheless it was possible to trace the infections back to the original carriers that had arrived three years earlier.¹⁶ The two diseases subsequently became grave commercial problems

THE PULMONARY ADENOMATOSIS COMPLEX IN SHEEP*

By FRANCISCO DURAN REYNALS†

Department of Microbiology School of Medicine Yale University New Haven Conn

ERWIN JUNGHERR

University of Connecticut Storrs Conn

ALBERTO CUBA CAPARO

*University of San Marcos Lima Peru and the Department of Microbiology
School of Medicine Yale University*

KEEN A. RAFFERTY JR

Department of Microbiology School of Medicine Yale University

and CHARLES HELMBOLDT

University of Connecticut

Introduction

With the discovery that certain tumors can be transmitted by means of cell free extracts interest naturally arose in possible comparisons between neoplastic viruses on the one hand and viruses of the traditional sort on the other. As a consequence of this interest attention has been drawn to situations in which viruses of classic inflammatory disease can give rise to disease that is histologically neoplastic. For example interest has revived in A. Borrel's studies of sheep pox and of fowl pox.¹ This investigator showed that the viruses of these diseases induced both necrotic and proliferative lesions and were in fact once designated as contagious epithelioma.¹ E. Roux observed pulmonary adenomatosis in sheep infected with sheep pox virus² and papillomas have been observed at the sites of the lesions of contagious pustular dermatitis in sheep.³ Smallpox, cowpox, rabbit pox, mouse pox, and other members of the pox group of viruses also can show a proliferative phase as well as the better known necrotizing phase.

Conversely viruses that ordinarily cause tumor formation can produce inflammatory and necrotizing disease under certain conditions according to dosage, inoculation route, host age and host resistance. F. Duran Reynals has demonstrated the inflammatory character of the Shope fibroma virus in newborn rabbits⁴ and of the Rous sarcoma virus in young chickens.⁵ In some recent work Duran Reynals⁶ has shown that mice pretreated with methyl cholanthrene and cortisone and then inoculated intradermally with vaccinia virus develop tumors rapidly in the sites of the healed vaccinia lesions. Although at present it is not possible to say whether vaccinia participates directly in the formation of those neoplasms, it can be said at least that the vaccinia lesion appears to be an optimal ground for the development of malignancy.

A second important question for students of neoplasia is the role of viruses in the production of tumors in general. From observations made by many

The work reported in this paper was supported in part by the American Cancer Society, Inc., New York, N. Y.

† Deceased

another form of SPA (FIGURE 1b) that has been named epizootic pulmonary adenomatosis or Icelandic IA. Unlike *Maedi* which seems to have a somewhat longer development period Icelandic IA shows considerable proliferative change with slight to pronounced infiltration¹⁹ much of which does not appear to be of an irritative nature. Metastasis has not been observed. Spread of the disease is thought to have been facilitated by the Icelandic practice of huddling or closely confining the sheep in barns during the winter. This disease together with *Maedi* had rendered commercial sheep raising all but impossible until both diseases were eradicated within a few years by a radical slaughter program following their recognition as contagious diseases. At this writing neither disease has been observed in Iceland for more than 4 years although it was not uncommon in the mid 1940s for 60 per cent of infected flocks to be lost within the span of a few years.¹⁶

A situation similar to that in Iceland exists in South Africa where a form of SPA very similar to *Maedi* Graaff Reinet disease occurs side by side with *jaagsiekte* a form more similar to or identical with Icelandic PA.²⁰⁻²¹ Graaff Reinet disease (FIGURE 2a) seems to involve infiltration of rather more uniform character than in the case of *Maedi* there being less tendency toward the formation of clusters of lymphocytic cells and a somewhat shorter development period. Histologically Graaff Reinet disease so named in recognition of the high incidence of that form near the Graaff Reinet experimental station in the Union of South Africa resembles progressive pneumonia and also lacks obvious features of irritation. On the other hand *jaagsiekte* (FIGURE 2b) or




FIGURE 2 (a) Graaff Reinet disease South Africa. Infiltrate with little sign of endothelial proliferation. $\times 200$. (b) *Jaagsiekte* South Africa. Adenoma with or without marked infiltrative change. $\times 200$. (Specimens furnished by G. deKock, Liesbeck Clinic, Cape Town, and G. leRou, University of Pretoria, Onderstepoort, Union of South Africa.)

in Iceland Finally a metastasizing adenocarcinoma somewhat resembling *jaagsiekte* (see below) in section was reported from Peru in 1945 by A. Cuba Caparo¹⁷

The Pathology of SP 1

Beginning at the inflammatory end of the SPA spectrum is an Icelandic disease *Maedi* (FIGURE 1a). A highly contagious illness *Maedi* requires about three years from the time of infection to manifest clinical signs; death usually follows a few months later. B. Sigurdsson¹⁶ and Sigurdsson *et al.*¹⁸ have shown, however, that development of the disease actually begins within a few months of exposure in the form of numerous infiltrative nodules that progressively increase in number and size until respiration is impaired sufficiently for the appearance of external signs. These consist of emaciation, rapid and shallow breathing, moist rales on auscultation, and general weakness. The signs invariably progress until death results. On necropsy the lungs are massively infiltrated, presenting large grayish areas that seem to consist of small confluent nodules. The cut surface is firm or friable, often dry, while the weight of the lungs is threefold to fourfold that of normal lungs. The tissue is often sterile. On section (FIGURE 1a) the lungs are seen to be engorged with cells of lymphocytic type, with a tendency toward the formation of clusters. There is no sign of adenoma or of irritation. Necrosis is not prominent. A series of carefully performed experiments has shown clearly that *Maedi* is highly contagious by means of air and fecal contamination of drinking water.¹⁶⁻¹⁸ Also in Iceland, in territory largely different from that of *Maedi* but overlapping it, there occurs




FIGURE 1 (a) *Maedi* of Icelandic. Infiltrative necrosis lacking. $\times 200$. (b) Contagious pulmonary adenomatosis of Icelandic (Icelandic I.A.). Infiltration plus adenoma. $\times 200$. (Specimens furnished by Björn Sigurdsson, University of Iceland, Reykjavik, Iceland.)

monia showing a marked tendency toward the early formation of peribronchiolar clusters of cells resembling lymphocytes. As the disease progresses invasion becomes more generalized resulting in greatly thickened interalveolar walls and cuboidal transition of the alveolar lining. In the later stages of the disease a striking adenomatous change often occurs as an accompaniment of the characteristic pneumonia (FIGURE 3b). Marsh²¹ believes that Montana SPA like *Maedi* is transmissible on contact and although this author does not consider the adenomatous condition to be neoplastic it has not been possible in our experience to distinguish Marsh's material from lesions that have long been classed as neoplasms in human pathology. Furthermore Montana SPA whether the pneumonia or adenoma form prevails is invariably progressive as in Iceland and in South Africa which does not seem to suit the hypothesis that the changes are hyperplastic reactions to nonspecific stimulation of normal lung tissue. Although it is well known that in man at least many irritants such as poisons can cause pulmonary hyperplasia that somewhat resembles adenoma^{24, 25} hyperplasia does not in our opinion closely resemble legitimate cases of SPA. The crux of the nomenclature problem seems to depend in the case of pulmonary adenoma upon whether metastasis is observed. The fact that neither Icelandic South African nor Montana SPA has been observed to form metastasis is of somewhat diminished consequence however since closely related forms of SPA from France and from Peru do show metastasis.

The French form of SPA *bouhite* derives its name from a patois term meaning panting. Histologically it is close to Montana SPA with the reservation that adenoma generally accompanies infiltration and is often in fact the dominant feature of the disease. Metastasis to the mediastinal lymph nodes or to subcutaneous lumbar lymphatic tissue has been shown in two cases^{21, 22}. The infiltrative aspect of *bouhite* closely resembles lymphomatosis and the term malignant pulmonary lymphomatosis has been suggested by Lucam for this disease²³ and for others of the complex that show parenchymal infiltration by cells of the lymphocytic series in the absence of massive necrosis or obvious irritation. Actually all forms of the complex show this leukotic aspect to some degree. Predisposition toward *bouhite* may be associated with the ingestion of resinous matter in pine forests judging from the distribution of the disease in France although there is no positive evidence to support this theory²⁴. One case in fact has been reported from the Department of Seine et Marne where contact with pine resin is less likely²⁵.

As it appears in Peru (FIGURE 4) SPA cannot escape classification as carcinoma since lymph node metastasis has been observed⁷ (FIGURE 4b). Infiltrative aspects are subdued. The disease is recognized in the field by respiratory distress following exercise and by discharge from the nostrils of large amounts (often a liter or more a day) of mucous exudate presumably produced by the adenoma or adenocarcinoma¹⁷. Except for the sparseness of the inflammatory change Peruvian SPA closely resembles *jaagstekte* and other adenomatous forms of SPA. Like them Peruvian SPA is invariably progressive. The disease occurs preferentially at altitudes of several thousand feet.

driving sickness so named following the observation that the first clinical signs of the disease commonly appear during exercise seems to be variable in the comparative amounts of adenomatous versus infiltrative changes that occur *Jaagsiekte* the prototype of SPA is referred to here as representing the more adenomatous form of the disease Both forms however, occur in the same territory and it does not seem possible to separate them into distinct entities since there is considerable gradation of one form into the other Adenomatous cases however can often be distinguished clinically from infiltrative cases because the former tend toward the production of more or less large amounts of mucous exudate which is discharged through the nostrils Metastasis has not been observed in either form

The North American form of SPA called progressive pneumonia or chronic catarrhal pneumonia is confined to Montana and a few other western states The disease appears in range sheep and is commonly first observed in the spring of the year infection possibly occurring during the winter The subsequent development of the disease is typical of that of the SPA complex afflicted sheep (lungers) dying of asphyxiation or secondary bacterial pneumonia 3 to 12 months after the first appearance of clinical signs¹¹

As in Iceland and in South Africa SIA of Montana presents both the infiltrative and adenomatous aspects of this disease but both forms are more likely to occur in the same animal the adenomatous form being less conspicuous After a study of both forms E Cowdry and H Marsh¹² concluded that *jaagsiekte* and Montana SIA were probably essentially the same disease Histologically the inflammatory aspect (FIGURE 3a) is an infiltrating pneu




FIGURE 3 (a) Chronic catarrhal pneumonia United States Infiltrative form $\times 200$
 (b) Chronic catarrhal pneumonia United States Adenomatous form accompanying the infiltrative form (Specimens furnished by Haleigh Marsh Livestock Station Boz Helena Mont) $\times 100$

by specimens taken from heavily diseased hosts⁴³. A virus came to be suspected as the etiological agent when SI A was observed to occur in epizootics in South Africa and Iceland. At the present time there remains no doubt that at least some forms are virus induced. Transmission by extracts of diseased lung has been effected with the Montana⁴⁴ Icelandic^{45, 46} and Central African⁴⁷ forms of SI A. The Icelandic form *Maedi* apparently can be transmitted by air and by means of fecal contamination of drinking water⁴⁸ and Shirlaw⁴⁹ claims to have recovered a specifically antigenic virus after two serial passages of extracts of lung from cases of Laikipia lung disease (of Central Africa) in the yolk sacs of hen's eggs. Many attempts to transmit the disease experimentally, however, have been complicated by the fact that adequate isolation of control animals is extremely difficult. The difficulty of accomplishing effective isolation implies that contagiousness may be of a high order in spite of the fact that the attack rate is usually well below 30 per cent. Another factor making for uncertainty in the results has been the length of the latent period which has been shown in some forms to be from 1 to 3 years in duration. Some of the negative experiments were undoubtedly terminated before the development of clinical disease. A few attempts have been made to transmit the disease by means of filtrates but it is felt that the results are not entirely conclusive because of the small numbers of animals used in the experiments or because of difficulties in maintaining strict isolation of inoculated sheep.

In one experiment Dungal observed SPA in 3 of 8 Icelandic sheep receiving a Gradacol filtrate (0.9 μ pore diameter) of glycerine solution through which sick sheep had been required to breathe. In another experiment 6 of 38 sheep developed SPA 4 to 9 months after receiving injections of Berkefeld Chamberland or Seitz filtrates of infected lung. In both experiments the uninoculated control animals remained free of disease⁴³. In an experiment performed by Creech and W. Gochenour⁴¹ 23 sick sheep were shipped from Montana to Maryland where SPA is unknown and 96 Maryland sheep were exposed to the disease either by inoculation (81 sheep) or by contact (15 sheep). None of the sheep exposed by contact developed the disease although 4 of the 81 sheep inoculated with sterile extracts of diseased lung developed SPA in 3 to 12 months and 4 other cases of slight pneumonic lesions were noted on histological section at autopsy.

The Serology of SI A

The attempt to establish several geographically separated and histologically varying diseases as members of a single complex certainly should include serologic studies of the agents involved and the distribution of antibodies in infected and uninfected flocks. Unfortunately the agents involved have been and remain extremely elusive with the exception of the previously mentioned case reported by Shirlaw⁴. The partially purified agent obtained by inoculating lung extract from sheep ill with Laikipia lung disease into the yolk sacs of embryonated hen's eggs fixed complement with high dilutions of serum from sick sheep but not with that from healthy sheep of the same area. Formalin treatment of the preparation is reported to result in a vaccine that reduces the




FIGURE 4 (a) Pulmonary adenomatosis of Peru. Adenomatous infiltration slight to moderate $\times 250$. (b) Lymph node metastasis pulmonary adenomatosis of Peru. The same sheep as in (a). $\times 250$.

The Epidemiology of SPA

Beginning in the 1920s SIA seemed to increase sharply in world wide incidence and distribution although part of the apparent increase was undoubtedly due to its recognition as an important disease in commercial flocks. SPA has also been reported at one time or another from The Netherlands³⁸ Scotland^{37,39} Israel⁴⁰ and Central Africa⁴¹. In recent years SPA may have begun a natural decline in incidence at least in Peru⁴, the United States³ and Iceland (even before the artificial eradication of the disease there)⁴² possibly showing the declination of a long term epidemic curve. In addition many parts of the world have never reported the disease. The principal factor in distribution does not seem to be entirely one of host susceptibility since sheep from regions free from SIA have been shown to be susceptible when inoculated with extract of diseased lung⁴³ or when moved to infected areas⁴⁵. Certain breeds however such as the Adalbol were discovered to be partially refractory to the Icelandic disease⁴⁴ and others including the Merino to the South African disease⁴⁶.

The contagiousness of SPA was suspected by its first describer M Gadyern who believed it to be a form of verminous pneumonia caused by *Strongylus* infestation. Other workers however showed that worm infestation did not necessarily accompany the disease and studies in Iceland failed to show a correlation between occurrence of SIA and infection with any particular parasite or microorganism^{41,47}. At one time the sheep ked *Melophagus ovinus* was suspected as a vector in Iceland but *Maedi* was not induced in sheep bitten

ment were allowed to mingle with the inoculated group and a few with those receiving only methylcholanthrene. Sheep from these groups were sacrificed and examined at intervals.

Eight sheep sacrificed at 4 months showed no significant lung lesions. In 3 sheep sacrificed at 8 months, however, apparent incipient proliferative lesions of the pulmonary alveolar epithelium were found in both of the sheep that received infected lung extract (FIGURE 5 a and b). Here the alveolar sacs are




FIGURE 5 (a) Lung of an apparently healthy sheep sacrificed 8 months after the intravenous injection of extract from lungs of sheep with chronic catarrhal pneumonia (see FIGURE 3a) $\times 250$ (b) Lung of similar sheep that received the same injection also sacrificed 8 months later $\times 250$ (c) Lung of apparently healthy sheep sacrificed 11 months after the intravenous injection of the same extract $\times 250$

natural incidence of disease from 30 per cent to about 1 per cent when the vaccine is administered within the first month of life. Further evidence for the antigenicity of SPA agents is the observation made in Peru by one of us (A. Cuba Caparo) that formalin treated suspensions of diseased lung emulsion are effective as vaccines. No complement fixation tests have been done with the Peruvian form. In a careful search no complement fixing activity was found in the Icelandic form *Maedi*. All attempts to cultivate the agents of *Maedi* and of Icelandic SPA have failed and extracts of diseased lung prepared in various ways have been found to be devoid of the ability to fix complement with serum from sick sheep or to produce specific hemagglutinins.¹⁶

Unfortunately, no general method for the demonstration of specific antigen in the lungs of diseased animals is available; consequently, it has been impossible to determine whether serologic relationships exist between geographically separated forms of SPA.

The Host Range of SPA

All evidence available indicates that the host range of SPA is extremely narrow, the disease being transmissible only from sheep to sheep, although pulmonary adenomatosis or infiltrative pneumonia have been observed in man,^{49, 55} cats,⁵⁶ guinea pigs,⁵⁷ mice,⁵⁸ chinchillas (E. J.), goats,⁵ horses,⁵⁹ hogs,²⁴ cattle,⁵ a tiger,²⁴ a deer,⁶⁰ and many other animals. Our attempts to transmit Montana SPA to Swiss mice, hamsters, and chinchillas, as well as to induce replication of an agent in embryonated hen's eggs, have failed up to the present time. Early speculation that pneumonia occurring in sheepherders may be derived from SPA infected sheep is now believed to be unfounded.³ I. A. has not been reported in man in Iceland, although that country has suffered the most devastating attack rate of this disease in sheep.¹⁹ Over a period of a few weeks glycerol stored material from a case of PA in man did not give rise to lesions when inoculated into a monkey, into rabbits, and into guinea pigs.⁶¹ The possibility exists that cultivation of an agent of SPA in embryonated eggs or in tissue cultures might allow the demonstration of a wider host range than has been shown to exist.

Experimental

In our experiments attempts have been made to transmit the North American form, sheep progressive pneumonia, to other sheep using lung tissue from diseased sheep. The lung tissue was taken from diseased Montana sheep that were sacrificed for the purpose, frozen on dry ice, and shipped to our laboratory by Hadleigh Marsh of the Livestock Sanitary Board, Helena, Mont. Sterile saline extracts were inoculated intravenously in lambs 1 to 7 days old, some of which were simultaneously treated with methylcholanthrene by intraperitoneal injection in sesame oil to determine whether a carcinogenic agent might accelerate the development of the disease or shift its expression from infiltrative and inflammatory to a more neoplastic type. Other groups received methylcholanthrene with no extract of infected lung, and were separated from the inoculated lambs by means of a board fence. A few lambs receiving no treat

ment were allowed to mingle with the inoculated group and a few with those receiving only methylcholanthrene. Sheep from these groups were sacrificed and examined at intervals.

Eight sheep sacrificed at 4 months showed no significant lung lesions. In 5 sheep sacrificed at 8 months, however, apparent incipient proliferative lesions of the pulmonary alveolar epithelium were found in both of the sheep that received infected lung extract (FIGURE 5, a and b). Here the alveolar sacs are




FIGURE 5. (a) Lung of an apparently healthy sheep sacrificed 8 months after the intra-uterine injection of fetal cells from lungs infected with chronic catarrhal pneumonia (see FIGURE 3). $\times 250$. (b) Lung of similar sheep that received the same injection, also sacrificed 8 months later. $\times 250$. (c) Lung of apparently healthy sheep sacrificed 11 months after the intra-uterine injection of the same extract. $\times 250$.

lined in places by cuboidal epithelium. Of those animals receiving lung extract one showed the infiltrative condition characteristic of Montana progressive pneumonia (FIGURE 5a). The nature of the lesions was the same regardless of whether methylcholanthrene was administered. The 3 sheep that did not receive extract of infected lung showed no such proliferation.

It was suspected at first that this early proliferation might develop into adenoma in sheep receiving both lung extract and methylcholanthrene, later sacrifices however, did not bear out that assumption. Five sheep were sacrificed approximately 11 months after inoculation. Of the 2 that received extract of infected lung, the one that did not receive methylcholanthrene showed slight but probably significant proliferation of alveolar and bronchiolar epithelium (FIGURE 5c). In 8 sheep (4 controls and 4 receiving lung extract) sacrificed 13 months after inoculation, the inoculated group showed nodes and consolidation in gross aspect and circumscribed nodules of lymphocytic infil




FIGURE 6 Lungs of 4 apparently healthy sheep sacrificed 13 months after the intra-venous injection of the same extract used to induce the lesions shown in FIGURE 5. $\times 250$

tration on section (these 4 cases are pictured in FIGURE 6) The picture represented is that typically seen in pulmonary adenomatosis of Montana The 4 control sheep receiving methylcholanthrene alone showed only normal tissue Periodic sacrifices are continuing at the present time

It seems evident that although the disease has probably been successfully transmitted in these experiments more than one year is required before clinical signs appear after inoculation This period cannot properly be termed a latent one since proliferative changes were observed in infected lungs after eight months to be replaced by the thirteenth month with the nodular lymphocytic congestion characteristic of pulmonary adenomatosis of Montana It was pointed out by Sigurdsson¹⁶ that this is not a chronic disease but a progressive one Proliferative changes in the alveolar and bronchiolar cells actually may begin a few months after inoculation and the earliest stages of the Montana form of SPA closely resemble the earliest stages of the highly neoplastic Peruvian form Methylcholanthrene has shown no effect on the course of infection to date however the carcinogen was administered intraperitoneally and not at the site of virus action and in comparatively low dosage (100 mg per lamb)

Experiments are under way to effect transmission using various routes of administration intraperitoneal intratracheal intranasal and intrapulmonary inoculations have been made Material from apparently infected lungs of the first experiment has been used in an attempt at serial transmission and an additional group of lambs has been inoculated with new material from symptomatic Montana sheep At the present time no sheep from these groups have been sacrificed

Discussion

Of great interest is the relationship between infiltrative or inflammatory disease and adenomatous disease since these two aspects have been kept separate by custom yet each aspect in the case of the SPA complex can occur singly or in company with the other A possible parallel exists in fowl lymphomatosis in which experimental infection with one form may give rise to such conditions as erythroblastosis myeloblastosis or visceral or neural lymphomatosis all related serologically to the agents of the Rous sarcoma and the Fujimami myxosarcoma¹⁷ In another case the formation of sarcomas or of salivary gland tumors in mice occurs in mice inoculated with the leukemia agent¹⁸ The results imply that the agents involved are highly mutable maintaining a mixture of slightly differing viruses or that expression of disease depends to a large extent upon variations in cell or organ susceptibility in different host individuals

Either explanation serves to account for the mixed occurrence of infiltrative pneumonia and adenoma in sheep although both have been transmitted in relatively unmixed form *Maedi* (an infiltrative form) has been transmitted experimentally as has Icelandic PA (largely adenomatous) Only *Maedi* seems to occur as an unmixed condition and *Maedi* itself is found in areas overlapping that of the adenomatous Icelandic PA Thus it has not been

lined in places by cuboidal epithelium. Of those animals receiving lung extract one showed the infiltrative condition characteristic of Montana progressive pneumonia (FIGURE 3a). The nature of the lesions was the same regardless of whether methylcholanthrene was administered. The 3 sheep that did not receive extract of infected lung showed no such proliferation.

It was suspected at first that this early proliferation might develop into adenoma in sheep receiving both lung extract and methylcholanthrene; later sacrifices, however, did not bear out that assumption. Five sheep were sacrificed approximately 11 months after inoculation. Of the 2 that received extract of infected lung the one that did not receive methylcholanthrene showed slight but probably significant proliferation of alveolar and bronchiolar epithelium (FIGURE 5c). In 8 sheep (4 controls and 4 receiving lung extract) sacrificed 13 months after inoculation, the inoculated group showed nodes and consolidation in gross aspect and circumscribed nodules of lymphocytic infil-




FIGURE 6. Lungs of 4 apparently healthy sheep sacrificed 13 months after the intravenous injection of the same extract used to induce the lesions shown in FIGURE 5. $\times 250$.

tration on section (these 4 cases are pictured in FIGURE 6) The picture represented is that typically seen in pulmonary adenomatosis of Montana The 4 control sheep receiving methylcholanthrene alone showed only normal tissue Periodic sacrifices are continuing at the present time

It seems evident that although the disease has probably been successfully transmitted in these experiments more than one year is required before clinical signs appear after inoculation This period cannot properly be termed a latent one since proliferative changes were observed in infected lungs after eight months to be replaced by the thirteenth month with the nodular lymphocytic congestion characteristic of pulmonary adenomatosis of Montana It was pointed out by Sigurdsson¹⁶ that this is not a chronic disease but a progressive one Proliferative changes in the alveolar and bronchiolar cells actually may begin a few months after inoculation and the earliest stages of the Montana form of SPA closely resemble the earliest stages of the highly neoplastic Peruvian form Methylcholanthrene has shown no effect on the course of infection to date however the carcinogen was administered intraperitoneally and not at the site of virus action and in comparatively low dosage (100 mg per lamb)

Experiments are under way to effect transmission using various routes of administration intraperitoneal intratracheal intranasal and intrapulmonary inoculations have been made Material from apparently infected lungs of the first experiment has been used in an attempt at serial transmission and an additional group of lambs has been inoculated with new material from symptomatic Montana sheep At the present time no sheep from these groups have been sacrificed

Discussion

Of great interest is the relationship between infiltrative or inflammatory disease and adenomatous disease since these two aspects have been kept separate by custom yet each aspect in the case of the SPA complex can occur singly or in company with the other A possible parallel exists in fowl lymphomatosis in which experimental infection with one form may give rise to such conditions as erythroblastosis myeloblastosis or visceral or neural lymphomatosis all related serologically to the agents of the Rous sarcoma and the Fujinami myxosarcoma⁴¹ In another case the formation of sarcomas or of salivary gland tumors in mice occurs in mice inoculated with the leukemia agent⁴² The results imply that the agents involved are highly mutable maintaining a mixture of slightly differing viruses or that expression of disease depends to a large extent upon variations in cell or organ susceptibility in different host individuals

Either explanation serves to account for the mixed occurrence of infiltrative pneumonia and adenoma in sheep although both have been transmitted in relatively unmixed form *Maedi* (an infiltrative form) has been transmitted experimentally as has Icelandic PA (largely adenomatous) Only *Maedi* seems to occur as an unmixed condition and *Maedi* itself is found in areas overlapping that of the adenomatous Icelandic IA Thus it has not been



FIGURE 7 Human pulmonary adenomatosis (FIGURE 1b) and to *jaagstekte* (FIGURE 2b) of sheep (Specimen loaned by V. A. Liebow, Yale University School of Medicine, New Haven, Conn.) $\times 250$. Note the similarity to Icelandic PA.

demonstrated that either adenoma or infiltrative pneumonia is completely separable from the other and the entire question as to whether these are different diseases may be meaningless for the present. In the end it may be necessary to resolve such differences by some arbitrary definition.

Many animals including man show conditions essentially similar to SPA both in histology and in clinical course. In man PA was first described in 1876 by Malassez⁶² but judging from reports in the literature this disease has increased greatly in incidence during roughly the same period of time in which SPA has increased. It has been estimated that about 1 per cent of all human lung neoplasm is IA.⁶ The disease in man differs from that in sheep almost solely in the incidence of metastasis which is much higher in the former. Other features of PA in man are very similar to those encountered in SIA: dyspnea, rapid and shallow breathing, moist rales, emaciation, exhaustion, little or no fever in uncomplicated cases, and eventual termination in death over about the same period as in SIA unless supervened by early surgery in cases lacking bilateral dissemination. Histologically the diseases in animals and man are indistinguishable (FIGURE 7) with the qualification that in man no evidence has been brought forward to indicate that sterile progressive pneumonias are related to adenoma. The similarities between IA of man and sheep, however, have been observed by many workers.^{4, 55}

Epidemiologically there is no reason to believe that IA in man is infectious. Irregularities in distribution are clearly accountable for by occupational environment. Thus silicosis may be followed by PA.⁵

Many bacterial and other conditions in man give rise to lung lesions that bear some resemblance to PA.⁵³ Proliferation in bovine lungs is said to follow nutritional irregularity or the ingestion of large amounts of certain minerals.⁵ Although the role of nonspecific stimulation in the genesis of tumors is not well understood, it is difficult to account for the inexorable course of neoplastic diseases such as IA solely in terms of irritation, particularly when metastasis to distant organs occurs. Consequently it appears that IA can

be considered an entity separate and distinguishable from pneumonia and metaplasia resulting from irritation

On the basis of its clinical progression and its general similarity to SPA C. Bonne⁴⁴ and J. Sims⁴⁵ have suggested the possibility of a viral origin for human IA. Since those suggestions were made further study of SIA by workers in South Africa, Iceland and the United States plus many recent experiments with animal neoplasia in general have revealed mechanisms of viral epidemiology that explain how a viral disease might be very difficult of detection as such as indeed is often the case. Since many of these mechanisms tend to obscure the viral etiology of SIA they are of particular interest in consideration of the origins of neoplasms generally. Some significant features of SPA may be described as follows:

(1) The latency or slow progressive development of SPA has been well established experimentally by several groups of workers. An asymptomatic period of several years between the time of infection and the appearance of clinical signs obviously tends to obscure infectiousness as a factor in development of the disease. There is some reason to believe that natural SPA infection may occur early in life followed by a lapse of years before the disease is recognizable. In studies on formalin killed vaccine against SIA Shirlaw⁴¹ reports that vaccination after the first month of life is not considered effective in preventing the disease which subsequently develops in sheep several years old. Infection of the young followed by long periods of latency is known to occur naturally in the case of the mouse mammary carcinoma⁴⁶ of fowl lymphomatosis⁴⁷ and of mouse leukemia.⁴⁸

(2) In these cases active virus can be recovered from the tissues of animals not showing the disease. There is good reason to suspect that infection by the agents of SPA may occur at a higher rate than development of recognizable disease a factor also tending to obscure infectiousness. Sigurdsson¹⁶ has observed that lung lesions which seem to be abortive *Maedi* are found in healthy exposed sheep at incidences higher than those expected from the observed incidences of the diseases. Marsh³ has suggested that some early lesions regress without producing disease whereas Sigurdsson in experiments on *Maedi* believes that development of the disease is very slowly progressive. If so many individuals should die from natural causes or be slaughtered before the disease appears.

(3) The isolation of viruses giving rise to neoplastic disease is classically difficult to accomplish especially when inbred lines of the animal in question are not available for experiment. In the case of SPA numerous methods for isolation of the causative agent have been attempted by several groups without success until the claim of Shirlaw was made⁴¹. It is evident that failure to isolate the causative agent of any disease in no way excludes an infectious origin. Initial failure in the isolation of neoplastic agents is in fact more the rule than the exception.

In the case of SPA infectiousness would doubtless not have been considered had the disease not occurred in epizootic form in certain areas. In the case of PA in animals and in man although the incidence in the latter has increased somewhat in the last few decades the disease has not been observed in cycles

or epidemics, hence any attempt to extend the known infectious etiology of SPA to its counterpart in other hosts is highly speculative. On the other hand information gained from the study of SPA should prompt closer examination of animal tumors generally, with the infectious viewpoint in mind as an etiological possibility.

Acknowledgment

The authors are indebted to Hadleigh Marsh Livestock Sanitary Board Helena Mont for providing infected lung material for experiment and to Bjorn Sigurdsson University of Iceland, Reykjavik Iceland to Gilles de Kock Liesbeck Clinic, Cape Town Union of South Africa to G LeRoux University of Pretoria Onderstepoort Union of South Africa and to A A Liebow Yale University School of Medicine who furnished slides and specimens of diseased lungs for study and reproduction.

References

- 1 BORREL A. Epithélioses infectieuses et épithéliomas 1903 Ann inst Pasteur 17 81-137
- 2 ROUX E. 1903 Sur les microbes dits invisibles Bull inst Pasteur 1 49-56
- 3 GLOVER R. E. 1928 Contagious pustular dermatitis of sheep J Comp Pathol Therap 41 318-340
- 4 DURAN REYNALS F. 1940 Production of degenerative inflammatory or neoplastic effects in the newborn rabbit by the Shope fibroma virus Yale J Biol Med 13 99-110
- 5 DURAN REYNALS F. 1940 A hemorrhagic disease occurring in chicks inoculated with the Rous and Fujinami viruses Yale J Biol Med 13 11-98
- 6 DURAN REYNALS F. 1937 Preliminary studies on the development of neoplasia in the skin of mice painted with methylcholanthrene and injected with cortisone and vaccine virus Ann N Y Acad Sci 68(2) 430-440
- 7 GROSS L. 1937 Mouse leukemia Ann N Y Acad Sci 64(6) 1184-1196
- 8 BURMESTER B. R. 1937 Studies on fowl lymphomatosis Ann N Y Acad Sci 64(6) 992-1003
- 9 ANDERSON H. B. 1952 Biological studies on the mammary tumor inciter in mice Ann N Y Acad Sci 64(6) 1004-1011
- 10 MACDYAN J. 1888 Lung disease in sheep caused by the *Strongylus rufescens* J Comp Pathol Therap 1 139-146
- 11 FIEBIGER A. 1899 Über multiple Adenombildung in den Lungen der Schafe Z Tiermed 3 161-172
- 12 ROBERTSON W. 1904 Jaagsiekte or chronic catarrhal pneumonia (sheep) J Comp Pathol Therap 17 221-224
- 13 MITCHELL D. T. 1915 Investigations into jaagsiekte or chronic catarrhal pneumonia of sheep 3rd & 4th Rept Director Vet Lduc Research Union of South Africa 585-614
- 14 MARSH H. 1923 Progressive pneumonia in sheep J Am Vet Med Assoc 62(15) 458-473
- 15 LESBOUYRIES M. & A BONNAC. 1940 La bouhite du mouton Recueil méd vét 116 2 7 290
- 16 SIGURDSSON B. 1934 Maedi a slow progressive pneumonia of sheep an etiological and pathological study Brit Vet J 110 275-270
- 17 CUBA CAPARÓ A. 1945 La poliadenomatosis pulmonar del carnero Bol escuela natl ciencias vet 1 77-87
- 18 SIGURDSSON B. I. A. PALLSON & A. TRYGGVADOTTIR. 1933 Transmission experiments with Maedi J Infectious Diseases 93 166-175
- 19 DUNN N. 1938 Epizootic adenomatosis of the lungs of sheep its relation to chronic pneumonia and jaagsiekte Proc Roy Soc Med 31 49-50
- 20 DEKOCK G. 1939 Are the lesions of jaagsiekte in sheep of the nature of a neoplasia 15th Ann Rept Director Vet Services Union of South Africa 611-641
- 21 DEKOCK G. 1939 Further observations on the etiology of jaagsiekte in sheep 15th Ann Rept Director Vet Service Union of South Africa 1163-1193

- 22 COWDRY E V & H MARSH 1937 Comparative pathology of South African jaagsiekte and Montana progressive pneumonia of sheep J Exptl Med 45 571-585
- 23 MARSH H 1955 Personal communication
- 24 DEKOCK G 1939 Pathology of phenol poisoning in sheep induced by certain dips 15th Ann Rept Director Vet Services Union of South Africa 643-656
- 25 MONTELL W J FITTE G KENDRICK & H DEBLISSON 1933 Progressive pulmonary adenomatosis in cattle Southwestern Vet 6 267-269
- 26 ROBSON W K D A IRWIN & E J KING 1934 Experimental silicosis quartz sericite and irritating gases Can Med Assoc J 31 237-245
- 27 SEATON V A 1937 Pulmonary adenomatosis in cattle produced by nitrogen dioxide poisoning North Am Veterinarian 38 109-111
- 28 GEEVER L F K T NEUBUERGER & C L DAVIS 1943 The pulmonary alveolar lining under various pathologic conditions in man and animals Am J Pathol 19 913-933
- 29 NEUBUERGER K T & L F GEEVER 1949 Alveolar cell tumor of the human lung A M A Arch Pathol 33 551-569
- 30 ROSS S 1933 Pulmonary epithelium and proliferative reactions in the lungs A M A Arch Pathol 27 478-496
- 31 AYNARD M 1936 Origine vermineuse du cancer pulmonaire de la brebis Compt rend soc biol 95 1540-1542
- 32 AYNARD M PYRON & FALCHETTI 1932 Sur le cancer du poulmon chez le mouton et ses liens étologiques avec les lésions parasitaires et infectieuses Compt rend soc biol 195 342-345
- 33 LUCAS F 1942 La bouhite ou lymphomatose pulmonaire maligne du mouton Rec méd vét 118 273-284
- 34 BERTHELON M H DRIEUX & P PALLAS 1944 La bouhite expérimentale Bull acad vét France 17 144-146
- 35 GUILHON J & C CHAUVIER 1952 La bouhite du mouton existe t elle en France en dehors du foyer landais? Bull acad vét France 25 205-209
- 36 BOS S E 1931 Over Alveolaire Cellumoren Thesis Amsterdam Holland
- 37 BLAKEMORE F & T J BOSWORTH 1941 The occurrence of jaagsiekte in England Vet Record 53 35-37
- 38 BLAKEMORE F 1955 Personal communication
- 39 HARBOUR H E & S JAMIESON 1946 Jaagsiekte in Scotland Vet Record 58 6
- 40 PATTISON I H 1946 Observations of a possible case of jaagsiekte in a sheep in Palestine J Comp Pathol 56(1) 63-67
- 41 SHIRLAW J D 1956 A preliminary note on successful vaccination of sheep in Kenya against progressive enzootic pneumonia (Laiapia lung disease) Bull Epizootic Diseases Africa 4 57-59
- 42 CUBA CAPARÓ A & T PAREDES 1955 Personal communication
- 43 SIGURDSSON B 1957 Adenomatosis of sheep's lungs Experimental transmission In press
- 44 DUNGAL N G GISLASON & E L TAYLOR 1938 Epizootic adenomatosis in the lungs of sheep Comparisons with jaagsiekte verminous pneumonia and progressive pneumonia J Comp Pathol & Therap 51 46-68
- 45 DUNGAL N 1946 Experiments with jaagsiekte Am J Pathol 22 737-750
- 46 DEKOCK G 1937 Private communication
- 47 TAYLOR F L 1937-8 The lungworm theory for the origin of epizootic adenomatosis and the question of the existence of the adenomatosis in Great Britain Proc Roy Soc Med 31 505-508
- 48 CREECH G T & W S GOCHENOUR 1936 Chronic progressive pneumonia of sheep with particular reference to its etiology and transmission J Agr Research 52 667-679
- 49 FISHER J H & W J HALLEY 1953 Primary alveolar cell carcinoma of the lung A M A Arch Pathol 55 162-170
- 50 LAIPPLY T C & C I FISHER 1949 Primary alveolar cell tumors of the lung A M A Arch Pathol 48 107-118
- 51 SIM J L 1943 Multiple bilateral pulmonary adenomatosis in man A M A Arch Intern Med 71 403-409
- 52 SWANN L 1949 Pulmonary adenomatosis of man a review of the literature and a report of nine cases A M A Arch Pathol 47 517-544
- 53 TILLET R S & D F HIRSCH 1952 So called pulmonary adenomatosis of the lungs Illnot Med J 101 232-239
- 54 WATSON W L & R R SMITH 1951 Terminal bronchiolar or alveolar cell cancer of the lung J Am Med Assoc 147, 7-13

or epidemics hence, any attempt to extend the known infectious etiology of SPA to its counterpart in other hosts is highly speculative. On the other hand information gained from the study of SPA should prompt closer examination of animal tumors generally with the infectious viewpoint in mind as an etiological possibility.

Acknowledgment

The authors are indebted to Hadleigh Marsh Livestock Sanitary Board Helena Mont for providing infected lung material for experiment and to Bjorn Sigurdsson University of Iceland Reykjavik Iceland, to Gilles de Kock, Liesbeck Clinic Cape Town Union of South Africa to G LeRoux University of Pretoria, Onderstepoort Union of South Africa and to A A Liebow, Yale University School of Medicine who furnished slides and specimens of diseased lungs for study and reproduction.

References

- 1 BORREL, A. Epithélioses infectieuses et épithéliomas. 1903. Ann inst Pasteur 17 81-137
- 2 ROUX E. 1903. Sur les microbes dits invisibles. Bull inst Pasteur 1 49-56
- 3 GLOVER R E. 1928. Contagious pustular dermatitis of sheep. J Comp Pathol Therap 41 318-340
- 4 DURAN REYNALS F. 1940. Production of degenerative inflammatory or neoplastic effects in the newborn rabbit by the Shope fibroma virus. Yale J Biol Med 13 99-110
- 5 DURAN REYNALS F. 1940. A hemorrhagic disease occurring in chicks inoculated with the Rous and Fujinami viruses. Yale J Biol Med 13 77-98
- 6 DURAN REYNALS F. 1957. Preliminary studies on the development of neoplasia in the skin of mice painted with methylcholanthrene and injected with cortisone and vaccine virus. Ann N Y Acad Sci 68(2) 430-440
- 7 GROSS I. 1952. Mouse leukemia. Ann N Y Acad Sci 54(6) 1184-1196
- 8 BURMESTER B R. 1952. Studies on fowl lymphomatosis. Ann N Y Acad Sci 54(6) 992-1003
- 9 ANDERVONT H B. 1952. Biological studies on the mammary tumor inciter in mice. Ann N Y Acad Sci 54(6) 1004-1011
- 10 M FADVEAN J. 1888. Lung disease in sheep caused by the *Strongylus rufescens*. J Comp Pathol Therap 1 139-146
- 11 EBER A. 1899. Über multiple Adenombildung in den Lungen der Schafe. Z Tiermed 3 161-172
- 12 ROBERTSON W. 1904. Jagziekte or chronic catarrhal pneumonia (sheep). J Comp Pathol Therap 17 221-224
- 13 MITCHELL D T. 1915. Investigations into jagziekte or chronic catarrhal pneumonia of sheep. 3rd & 4th Rept Director Vet Educ Research Union of South Africa 585-614
- 14 MARSH H. 1923. Progressive pneumonia in sheep. J Am Vet Med Assoc 62(15) 458-463
- 15 LESBOUVRIES M & A BONVAC. 1940. La touxite du mouton. Recueil méd vet 116 277-280
- 16 SIGURDSSON B. 1954. Maedi a slow progressive pneumonia of sheep an etiological and pathological study. Brit Vet J 110 225-270
- 17 CUBA CAPABO A. 1945. La poladenomatosis pulmonar del carnero. Bol escuela natl ciencias vet 1 27-57
- 18 SIGURDSSON B P A PALLSON & A TRYGGVADOTTIR. 1953. Transmission experiments with Maedi. J Infectious Diseases 93 166-175
- 19 DUNGAL N. 1938. Epizootic adenomatosis of the lungs of sheep its relation to verminous pneumonia and jaagsiekte. Proc Roy Soc M 1 31 497-505
- 20 DEKOCK G. 1939. Are the lesions of jaagsiekte in sheep of the nature of a neoplasia? 15th Ann Rept Director Vet Services Union of South Africa 611-641
- 21 DEKOCK C. 1939. Further observations on the etiology of jaagsiekte in sheep. 15th Ann Rept Director Vet Services Union of South Africa 1169-1183

- 22 COWDRY F V & H MARSH 1977 Comparative pathology of South African jaagsiekte and Montana progressive pneumonia of sheep J Exptl Med 45 571-585
- 23 MARSH H 1955 Personal communication
- 24 DEKOCK C 1929 Pathology of phenol poisoning in sheep induced by certain dips 15th Ann Rept Director Vet Services Union of South Africa 643-656
- 25 MONLUX W J FITTE G KENDRICK & H DELBUISSON 1953 Progressive pulmonary adenomatosis in cattle Southwestern Vet 6 261-269
- 26 ROBSON W K D A IRWIN & I J KING 1934 Experimental silicosis quartz senite and irritating gases Can Med Assoc J 31 237-245
- 27 SEATON V A 1957 Pulmonary adenomatosis in cattle produced by nitrogen dioxide poisoning North Am Veterinarian 38 109-111
- 28 GEEVER I F K T NEUBURGER & C L DAVIS 1943 The pulmonary alveolar lining under various pathologic conditions in man and animals Am J Pathol 19 913-933
- 29 NEUBURGER K T & F F GEEVER 1949 Alveolar cell tumor of the human lung A M A Arch Pathol 33 551-569
- 30 ROSS S 1939 Pulmonary epithelium and proliferative reactions in the lungs A M A Arch Pathol 27 478-496
- 31 VYNAUD M 1936 Origine vermineuse du cancer pulmonaire de la brebis Compt rend soc biol 95 1540-1547
- 32 VYNAUD M PEYRON & FALCHETTI 1932 Sur le cancer du poulmon chez le mouton et ses lésions étiologiques avec les lésions parasitaires et infectieuses Compt rend soc biol 195 342-345
- 33 LUCAM I 1942 La boubite ou lymphomatose pulmonaire maligne du mouton Rec méd vét 118 213-284
- 34 BERTHELON M H DRIEUX & P LALLAS 1944 La boubite expérimentale Bull acad vét France 17 144-146
- 35 GLILION J & G CHAUVIER 1952 La boubite du mouton existe t elle en France en dehors du fover landais? Bull acad vét France 25 203-209
- 36 BOS S F 1951 Oer Alveolair Celumoren Thesis Amsterdam Holland
- 37 BLAKEMORE F & T J BOSWORTH 1941 The occurrence of jaagsiekte in England Vet Peco d 53 33-37
- 38 BLAKEMORE F 1955 Personal communication
- 39 HARBOUR H E & S JAMIESON 1946 Jaagsiekte in Scotland Vet Record 58 6
- 40 IATHON I H 1946 Observations of a possible case of jaagsiekte in a sheep in Palestine J C mp Pathol 56(1) 63-67
- 41 SHIRLAW J D 1956 A preliminary note on successful vaccination of sheep in Kenya against progressive enzootic pneumonia (Lauki lung disease) Bull Epizootic Diseases Africa 4 57-59
- 42 CUBA CAPARÓ A & T PAREDES 1955 Personal communication
- 43 SIGURDSSON B 1957 Adenomatosis of sheep's lungs Experimental transmission In press
- 44 DUNGAL N G CIELASON & F L TAYLOR 1938 Epizootic adenomatosis in the lungs of sheep Comparisons with jaagsiekte verminous pneumonia and progressive pneumonia J C mp Pathol & Therap 51 46-68
- 45 DUNGAL N 1946 Experiments with jaagsiekte Am J Pathol 22 737-750
- 46 DEKOCK G 1957 Private communication
- 47 TAYLOR E L 1937-8 The lungworm theory for the origin of epizootic adenomatosis and the question of the existence of the adenomatosis in Great Britain Proc Roy Soc Med 31 505-508
- 48 CREECH G T & W S GOCHENOLD 1936 Chronic progressive pneumonia of sheep with particular reference to its etiology and transmission J Agr Research 62 667-69
- 49 FISHER J H & W J HALLEY 1953 Primary alveolar cell carcinoma of the lung A M A Arch Pathol 55 167-170
- 50 LAIPPLY T C & C I FISHER 1949 Primary alveolar cell tumors of the lung A M A Arch Pathol 48 107-118
- 51 SIM J L 1943 Multiple bilateral pulmonary adenomatosis in man A M A Arch Internal Med 71 403-409
- 52 SWANN I 1949 Pulmonary adenomatosis of man a review of the literature and a report of nine cases A M A Arch Pathol 47 517-544
- 53 TILLET R S & D F HIRSCH 1952 So-called pulmonary adenomatosis of the lungs Illinois Med J 101 232-239
- 54 WATSON W L & R R SMITH 1951 Terminal bronchiolar or alveolar cell cancer of the lung J Am Med Assoc 147, 7 13

- 55 WOOD D A & P H PIERSON 1945 Pulmonary alveolar adenomatosis in man Is this the same disease as jaggsiekte in sheep? *Am Rev Tuberc* **51**(3) 203-224
- 56 OLAFSEN A & W MONLUN 1942 *Cornell Vet* **32** 176-178
- 57 NORRIS R F 1948 Pulmonary adenomatosis resembling jaggsiekte in the guinea pig A M A *Arch Pathol* **43** 553-558
- 58 JOBLING J W 1910 Spontaneous tumors of the mouse *Monographs Rockefeller Inst Med Research* **1** 81-119
- 59 THEILER A 1918 7th & 8th Rept Director Vet Educ Union of South Africa 59
- 60 GRUBER G B 1939 Über Veränderungen der Lungen nach Wurmbefall sog Pneumonia verminosa *Arch pathol Anat u Physiol Virchows* **304** 591-607
- 61 BEARD J W 1957 Etiology of avian leukosis *Ann N Y Acad Sci* **68**(2) 473-486
- 62 LAW L W 1954 Recent advances in experimental leukemia research *Cancer Research* **14** 695-709
- 63 MALASSEZ L 1876 *Arch Physiol* **3** 353
- 64 BONNF C 1939 Morphological resemblance of pulmonary adenomatosis (*jaggsiekte*) in sheep and certain cases of cancer of the lung in man *Am J Cancer* **35** 491-501

Part VIII International Programs

GLOBAL PRIORITIES IN ANIMAL-DISEASE CONTROL

By Martin M. Kaplan

World Health Organization Geneva Switzerland

In this paper I propose to discuss aspects and possibilities of certain human-animal disease relationships. These relationships constitute far more of a challenge than do the essentially technological solutions to the well-defined diseases.

The permanent agencies concerned with animal disease control on an international scale are the Food and Agriculture Organization (FAO) and the World Health Organization (WHO) with the Pan American Sanitary Bureau (IASB) which serves as the Regional Office for the Americas of WHO. Their function in field operations, coordination of research and international collaboration are described in official publications, and I refer anyone interested to those documents. The Office International des Epizooties (OIE) established in 1924 is another intergovernmental agency. It meets annually, has working commissions on special subjects with FAO and WHO participation and publishes and circulates official statistics on epizootic diseases. The only other group that can perhaps squeeze into the category of permanency is the Point Four program of the United States government which is devoted to bilateral assistance agreements with various governments.

Another area of WHO-FAO effort is in food hygiene, particularly with respect to meat and milk. I pass over this aspect of our work, even though considerable attention is given to this field, because its interest lies in the application of primarily technological procedures (including the magic word, radioactivity) to populations economically and socially unprepared as yet to receive them.

The Major Diseases

Now a glance at the animal-disease problems having the greatest visible impact on public health and the economics of food production. Here we shall use capsule summaries of the major diseases as they affect the less developed countries and as they may affect the more technically advanced ones.

Foot and mouth disease The difficulties in the control of this malady may be summarized thus: extreme communicability, with such bizarre features as human and avian migrants as vectors; at least seven distinct major types of virus with intratype variation; and costly and cumbersome methods of control such as slaughter and ring vaccination with inactivated vaccines of short and often doubtful efficacy. A light on the horizon appears to be the successful development of living attenuated strains for vaccination purposes, the only practical answer for most countries apart from North America, northern Europe and Oceania.

Rinderpest In distinction to foot and mouth disease, rinderpest is a rather

- 55 WOOD D A & P H PIERSON 1945 Pulmonary alveolar adenomatosis in man Is this the same disease as jaagsiekte in sheep? *Am Rev Tuberc* **51**(3) 203-224
- 56 OLAFSEN A & W MONLUCY 1942 *Cornell Vet* **32** 176-178
- 57 NORRIS R F 1948 Pulmonary adenomatosis resembling jaagsiekte in the guinea pig A M A Arch Pathol **43** 553-558
- 58 JOBLING J W 1910 Spontaneous tumors of the mouse Monographs Rockefeller Inst Med Research **1** 81-119
- 59 THEILER A 1918 7th & 8th Rept Director Vet Educ Union of South Africa 59
- 60 CRUBER G B 1939 Über Veränderungen der Lungen nach Wurmbefall sog Pneumonia verminosa Arch pathol Anat u Physiol Virchows **304** 597-607
- 61 BEARD J W 1957 Etiology of avian leukosis Ann N Y Acad Sci **68**(2) 473-486
- 62 LAW L W 1954 Recent advances in experimental leukemia research Cancer Research **14** 695-709
- 63 MALASSEZ L 1876 Arch Physiol **3** 353
- 64 BONNF C 1939 Morphological resemblance of pulmonary adenomatosis (*jaagsiekte*) in sheep and certain cases of cancer of the lung in man Am J Cancer **35** 491-501

problem but we are not optimistic about finding an early answer because relatively few laboratories are working in this field

Diseases of Less Importance

Now let us take a quick jump from these obvious diseases to more shadowy areas where future developments in human animal disease relationships are likely to be most interesting and fruitful. Here we find that actual and potential zoonoses occupy only a part of these relationships while essentially noncommunicable entities begin to call for increased attention. In keeping to my assignment however I shall deal only with the communicable diseases having an animal component known or suspected in their epidemiology. To remain within the limits of the space at my disposal I shall use the device of question and no answer.

Do the findings of *apparently* specific antibodies to the polio, Coxsackie and adenoviruses in domestic animals and to canine distemper in humans have any significance in the epidemiological cycles of these diseases?

Do animal reservoirs play any part in the epidemiology of human influenza? Of toxoplasmosis?

What is the evolutionary history of the arthropod borne (arbor) viruses (or of any virus)?

To what extent are the orphan viruses being isolated from man and animals species specific? The psittacosis like organisms? What is their significance as disease producing entities?

Is rabies in nonvampire bats a new phenomenon or have these animals long been involved in the natural history of this disease? Are they of any importance in its epidemiology? Apart from vampire bats are there other asymptomatic carriers of the rabies virus in wildlife?

The common denominator of these questions is that we are dealing with viruses and their natural history a rich field for speculation with commensurate rewards in pragmatic achievement and intellectual satisfaction to those providing the keys to their solution.

WHO is approaching these problems by taking advantage of its unusual opportunity to coordinate laboratory research efforts and field investigations. This is illustrated by animal serologic studies on human influenza antibodies now under way in twenty five countries to determine whether animals are involved in the epidemiology of human influenza and by a world wide survey for Q fever prevalence recently completed. In addition networks of laboratories are collaborating with WHO with respect to poliomyelitis, treponematoses, tuberculosis, malaria and schistosomiasis apart from influenza, rabies, brucellosis, leptospirosis and hydatidosis all mentioned previously.

A Project of Interest

One further project as yet unimplemented may be worth mentioning. It is clear to everyone that host-parasite relationships are continuously changing and it is probable that new disease entities or unsuspected human animal disease relationships will evolve. Until the present time one of the failures

easy disease to control. Effective and inexpensive living attenuated vaccines have been developed and are available and even rudimentary sanitary measures can limit its spread. The problem is how to use these tools under the difficult conditions prevailing in Africa and Asia but even on these continents the progress has been considerable for which we thank FAO for its energetic efforts. In general, the same thing may be said about contagious pleuropneumonia, hemorrhagic septicemia, and anthrax diseases of great destructiveness in these areas.

Parasitic diseases. The many ailments in this category unquestionably take the greatest economic toll. Helminths, flukes, and blood parasites lead the roster. Textbook recommendations for the control of these diseases often fail in the face of the realities of field conditions in underdeveloped countries. The magnitude of the problems posed by these diseases, involving as they do the social organization and economic structure of these countries, is such as to make impossible any early solution.

Brucellosis and leptospirosis. These two diseases illustrate the close working relationship between WHO and FAO as they deal with zoonoses problems of economic and public health importance.

The major problem at this time with respect to brucellosis lies in the development and application of a vaccine effective in sheep and goats. During the past five years WHO and FAO have coordinated and supported research in this field and results are encouraging. In October 1957 the joint FAO-WHO Expert Committee on Brucellosis will have met in Lima, Peru, to evaluate the experimental work carried out on brucellosis in the fifteen FAO-WHO brucellosis centers located in different parts of the world and the report of this committee should be useful as a guide to our member countries.

In leptospirosis WHO and FAO have concentrated their efforts on typing procedures with reference to *Leptospira* and in developing standard diagnostic tests that can be used without too much difficulty in hospitals and laboratories in the routine diagnosis of this disease in man and animals. This work is being carried out through six WHO-FAO Reference Leptospirosis Laboratories, one each in Australia, Holland, Italy, Japan, the United Kingdom, and the United States.

Rabies. Most of you are aware of the considerable attention WHO has given to rabies with respect to the improvement of vaccines, investigations of hyperimmune serum in prophylaxis, and in wildlife studies. Perhaps one of the greatest obstacles in the way of advances in laboratory research connected with rabies has been the inability to cultivate the virus on tissue culture to any satisfactory degree.

Hydatidosis. This disease deserves brief mention because of the extent of human illness it produces as well as the very great economic losses it causes in livestock products in South America, in the Mediterranean area, and in Oceania. Effective control of this disease cannot be achieved under conditions found in most affected areas until a better drug than arecoline is found for the treatment of dogs infested with the *Echinococcus* tapeworm and until more practical ovacides than creolin are developed. We are encouraging research on this

problem but we are not optimistic about finding an early answer because relatively few laboratories are working in this field

Diseases of Less Importance

Now let us take a quick jump from these obvious diseases to more shadowy areas where future developments in human animal disease relationships are likely to be most interesting and fruitful. Here we find that actual and potential zoonoses occupy only a part of these relationships while essentially noncommunicable entities begin to call for increased attention. In keeping to my assignment however I shall deal only with the communicable diseases having an animal component known or suspected in their epidemiology. To remain within the limits of the space at my disposal I shall use the device of question and no answer.

Do the findings of *apparently* specific antibodies to the polio Coxsackie and adenoviruses in domestic animals and to canine distemper in humans have any significance in the epidemiological cycles of these diseases?

Do animal reservoirs play any part in the epidemiology of human influenza? Of toxoplasmosis?

What is the evolutionary history of the arthropod borne (arbor) viruses (or of any virus)?

To what extent are the 'orphan' viruses being isolated from man and animals species specific? The psittacosis like organisms? What is their significance as disease producing entities?

Is rabies in nonvampire bats a new phenomenon or have these animals long been involved in the natural history of this disease? Are they of any importance in its epidemiology? Apart from vampire bats are there other asymptomatic carriers of the rabies virus in wildlife?

The common denominator of these questions is that we are dealing with viruses and their natural history a rich field for speculation with commensurate rewards in pragmatic achievement and intellectual satisfaction to those providing the keys to their solution.

WHO is approaching these problems by taking advantage of its unusual opportunity to coordinate laboratory research efforts and field investigations. This is illustrated by animal serologic studies on human influenza antibodies, now under way in twenty five countries to determine whether animals are involved in the epidemiology of human influenza and by a world wide survey for Q fever prevalence recently completed. In addition networks of laboratories are collaborating with WHO with respect to poliomyelitis, treponematoses, tuberculosis, malaria and schistosomiasis apart from influenza, rabies, brucellosis, leptospirosis and hydatidosis all mentioned previously.

A Project of Interest

One further project as yet unimplemented may be worth mentioning. It is clear to everyone that host-parasite relationships are continuously changing and it is probable that new disease entities or unsuspected human animal disease relationships will evolve. Until the present time one of the failures

in studies on communicable diseases has been our inability to follow the progression of disease in geographical areas at different points of time. This failure cannot be attributed to human oversight but rather to the limited opportunities and tools available for instituting appropriate studies. While conditions are better in this connection it will be many years before there will be sufficient facilities to cope with the many problems associated with progression parameters in the epidemiology of communicable diseases. Nevertheless in the meantime certain steps are possible that would no doubt be of invaluable assistance later when more detailed studies will be feasible. This thought underlines the reasoning behind projected work in epidemiological studies based on serology. Briefly it is planned to arrange for serum specimens to be taken from different age groups in human beings and animals in various parts of the world within a given span of time to catalogue these specimens and to store them in a refrigerated or freeze dried state for examination perhaps many years later. Thus where a particular study will be undertaken on one or more diseases serum specimens from the human and animal population of a geographical area of interest would be available for examination by qualified laboratories.

Such work can be done uniquely by an organization such as WHO with the collaboration of FAO and should be of immense benefit in improving our knowledge in control efforts of communicable diseases both in the near and more distant future. WHO's role would be principally that of a coordinator for it should be stressed that the real contribution and effort would come from the collaborating laboratories and scientists throughout the world who have so generously contributed their services and facilities in the past and who we do not doubt will continue to do so.

THE PAN AMERICAN ZOONOSIS CENTER

By Benjamin D. Blood

Pan American Zoonoses Center, Azul, Argentina

The Pan American Zoonoses Center located in Azul province of Buenos Aires, Argentina, is an international institution with the basic objective of promoting and strengthening activities against the zoonoses in the countries of the Americas. The center serves these countries in the education and training of professional and lay technical personnel in techniques and methods to be used in combating diseases transmitted between animals and man. It stimulates zoonoses control and eradication campaigns in the different countries and coordinates such campaigns on an international basis. Its staff is available for consultation on problems and activities concerned with those diseases. It conducts research with the aim of improving diagnosis and control of the zoonoses and it promotes, aids and coordinates such research in governmental and private institutions throughout the Americas. It works toward standardization of diagnostic methods and procedures for making and testing vaccines, sera and other biological products used for the zoonotic diseases. It prepares and disseminates information about those diseases. All of these services are aimed at helping the people of the Americas with the understanding and support of their governments to reduce the social and economic burden caused by uncontrolled diseases common to people and animals.

The Pan American Zoonoses Center was formally established on August 10, 1956, with the signing of the agreement between the government of Argentina as host country and the Pan American Sanitary Bureau (PASB), Regional Office of the World Health Organization (WHO), Washington, D. C., as the responsible international agency. The first staff members were assigned late in that year and the work of organizing and installing the center was begun. The staff has since been enlarged according to definite schedule until at the present time there are four international officials and sixteen local employees at their posts. The number of local employees will have increased to at least twenty by the end of 1957.

Location and Facilities

A number of countries expressed interest in becoming host to the center. Upon careful study of each of the different proposals it was judged that the facilities offered in Argentina were the most adequate.

Azul, a city of about 25,000 population located 185 miles south of Buenos Aires, the Argentine capital, is situated in an important livestock raising region. The cooperation of the medical and veterinary professions, livestock breeders, hospitals, municipal authorities and other local groups is an important asset of this international institution.

The building provided by the Argentine government is very well suited for housing the Pan American Zoonoses Center. Although built for another

purpose it was never occupied until transferred for its current use. It is a two story structure with space adequate for present needs in laboratories, offices, classrooms, and library, all of which are now complete or in final stages of remodeling. There is land remaining on either side for additional construction as needed. The host government provides the funds required for public utility services and for the maintenance and servicing of the building, and has agreed to finance new construction as necessary.

As an adjunct to the central facilities, the Argentine government has also purchased a 370-acre farm annex for the exclusive use of the center. This annex is less than three miles from the central site and provides the location for isolation units, holding pens, the laboratory animal colony, and supporting services.

Administration

The Pan American Zoonoses Center was created in response to desires of the American governments, as expressed both individually and collectively in several international meetings, political as well as technical. Its financial and material support are presently derived from three sources: the host government, the Pan American Sanitary Bureau, and the United Nations Technical Assistance Program. Negotiations are now in progress to broaden the financial structure and to assure its operation on a permanent basis.

The center is sponsored and administered by the PASB and is subject to the rules, regulations, and procedures of that organization. Other agencies interested in the objectives and activities of the center may participate in it by accepting the terms of the agreement by which it was established.

The permanent technical staff members, including veterinarians, physicians, and members of allied professions, are international employees; that is, their posts are supported with international funds. In addition, specialists in various aspects of zoonosology are under contract for limited periods and are paid with international or special funds. The permanent supporting staff (that is, the lay and nonskilled personnel) is recruited locally in Argentina and at this time is supported by funds contributed by the host government.

Functions and Operations

The diseases classified as zoonoses are so vast in number that the center will not be able, at least initially, to give due attention to all of them. Priorities are being established in accordance with the relative impact on human welfare and the geographical extension of the different zoonotic diseases in the Americas. Accordingly, brucellosis, hydatidosis, and rabies are being given major attention, followed by virus encephalitis, anthrax, leptospirosis, tuberculosis, psittacosis, trichinosis, and salmonellosis. This does not mean that the remaining diseases are to be ignored completely, since the center is intended to be a clearinghouse for information on all the zoonoses. Furthermore, the priorities may be revised from time to time to coincide with epidemiological/epizootiological changes and recognized needs of the countries.

The center is not intended to replace government zoonoses activities, nor even to compete with them. Its role is to complement those activities by

providing services that the individual countries have not found it possible to develop as well as to coordinate those activities internationally. The functions and services of the center are essentially education and training, consultation and coordination, laboratory services, demonstrations, research and information.

Education and training. The educational and training activities are probably the most important part of the work of the center. The lack of adequate numbers of well trained personnel, both professional workers and lay assistants, is a problem recognized throughout the Americas, although the need is somewhat more acute in some countries than in others. It is obviously impossible to carry out successful diagnostic, field control, eradication, treatment or research work without a qualified staff. In order to help provide the personnel required, the center offers a variety of training courses and opportunity for specialized study.

Training methods will make maximum use of demonstration and of trainee participation. Lectures are supplemented with audio-visual aids. Laboratory work is arranged so that each trainee actually practices the specific techniques under study, and field demonstrations make it possible for participants to become thoroughly familiar with methods for applying recognized measures of disease control.

Fellowships for training at the center are provided by individual governments and by PASB and WHO; they will undoubtedly also be provided by other agencies. Training is offered on both short term and long term programs. Special courses, seminars and workshops of from one to twelve weeks duration are being scheduled, with attention given to special topics or fields of work. The individuals accepted for such training are those who already have the necessary professional or technical preparation to continue.

One course that will be repeated on an annual basis is for a three to four week period devoted to zoonoses control for recent graduates from schools of public health. The course will be for veterinarians as well as for physicians preparing for epidemiological or rural health assignments. Other courses to be held are, for example, on the laboratory procedures in brucellosis, including diagnosis and vaccine production; on the diagnosis of rabies; and on control methods for hydatidosis. An annual two-month course will be offered for training nonprofessional field assistants for antizoonoses work. Other courses and seminars will be held as needs and possibilities arise.

Long term training is offered for individuals to undertake special study in one or more of the many phases of work with zoonoses. These individuals are expected to remain at the center for at least 6 to 12 months for postgraduate specialization and research. It is anticipated that this type of training will be developed in many cases in cooperation with universities.

It is highly desirable that the Pan American Zoonoses Center operate in close cooperation with colleges and universities. Although the exact patterns for such cooperation remain to be developed, there are excellent possibilities for the center to work with schools of public health, medicine and veterinary medicine at both the undergraduate and postgraduate levels of study. One of the several forms that collaborations might take would be for students to

perform their field (thesis) work at the Zoonoses Center after having completed the required formal courses at a university

The educational activities of the center also include the collection and distribution of audiovisual materials for use in programs dealing with the zoonoses. Such materials will in addition be developed at the center and they also will be produced there on a limited scale when not already available.

Consultation and coordination Upon request from the government concerned the center is prepared to send members of its staff to visit countries in a consultative capacity on matters related to the zoonoses and their control. In addition consultation services are provided through correspondence. Special attention is being given to the development in each country of the mechanism necessary for planning and operating control programs for the zoonoses. Such mechanisms based on the concept of united effort should provide for coordination and cooperation between public health and animal sanitation agencies with maximum participation of the community. Visits by consultants also provide the opportunity for international coordination of control and eradication programs.

Laboratory services These services include reference diagnosis, standardization of antigens, vaccines, the production of sera and other biological products, confirmatory testing of such products for potency and safety, and the distribution of strains of viruses and bacteria for production or challenge purposes and of standard antigens, vaccines, and sera for use in check testing.

The center will operate a model colony of laboratory animals that will serve for training, demonstration and study purposes, as well as for supplying foundation stock for other institutions.

Demonstrations The center will utilize field campaigns against various zoonoses in pursuance of its functions. Trainees are taken to the site of the work and actually participate in zoonoses control programs. Existing anti-zoonoses programs of the city of Azul and of the host country (at both provincial and national levels) serve for such demonstration purposes. Demonstration work will also include the development of special control programs for various zoonoses in different parts of the Americas, the number, type and extent of such programs being dependent upon the needs, capabilities and desires of the country concerned.

Research Research at the center is of an applied nature, that is, the evaluation of a vaccine under certain field conditions, the practical application of a diagnostic procedure, the assessment of drugs for their antiparasitic effect and similar projects. Surveys and epidemiological/epizootiological studies will be undertaken with respect to certain of the zoonoses. Also included under research is the collection and processing of statistical data on the occurrence and overall importance of the diseases in man and animals.

The needs and opportunities for research in connection with the zoonoses are vast and the center itself can be expected to cover only a very small part of them. It is thus quite evident that public and private institutions in all countries must continue and intensify their research work in this field. It is a responsibility of the center to stimulate and coordinate such work.

Information Facilities are being developed so that the center may serve

as the source of both technical and popular information on the zoonoses. Library service will be an important activity with microfilms and photocopies of technical references made available to investigators and control officials. Special priority is being given to the collection and cataloguing of books, periodicals, bulletins, and reports on diseases common to man and animals.

The film library will be made as complete as possible with a collection of all available films, film strips, and slide sets concerning the zoonoses and their control. When possible, new films will be produced on subjects on which no suitable material already exists.

A periodic information bulletin is to be published; this will contain abstracts of the world literature on zoonotic diseases, as well as brief references to the latest developments in this field as they may occur in the Americas and elsewhere.

A collection of bulletins, pamphlets, illustrated booklets, posters, exhibits, and other items used with success in the public education aspects of zoonoses control is maintained for demonstration and loan. New material of this type will be prepared at the center as needs and resources permit.

Summary

The recently established Pan American Zoonoses Center is an international institution sponsored by the Pan American Sanitary Organization (Regional Office of the World Health Organization) and is dedicated to promoting and strengthening activities against the zoonoses in the countries of the Americas. Its services are available to health departments, agriculture departments, educational institutions, and other agencies having an interest in zoonoses, the diseases naturally transmitted between animals and man.

The center, which has its central site in Azul, Argentina, is designed to serve all of the Americas in the education and training of professional and lay technical personnel in techniques and methods to be used in attacking the zoonoses. It will also (1) conduct research with the aim of improving diagnosis, epizootiological-epidemiological knowledge, and control procedures; (2) promote and coordinate such research in governmental and private institutions throughout the Americas; (3) work toward the standardization of diagnostic methods and procedures for making and testing vaccines, sera, antigens, and other biological products; (4) prepare and disseminate information on the zoonotic diseases; and (5) develop field demonstration activities in accordance with the needs and desires of participating countries.

The principal aim of the center is to promote and strengthen governmental activities against the zoonoses in the countries of the Americas. All of its activities, whether in the field of training, demonstration, research, information, or coordination, have been planned for that purpose. It is fully recognized that this objective can be reached only insofar as there are sound technical activities and services in each country dedicated to research, control, and eradication of diseases common to man and animals.

perform their field (thesis) work at the Zoonoses Center after having completed the required formal courses at a university.

The educational activities of the center also include the collection and distribution of audiovisual materials for use in programs dealing with the zoonoses. Such materials will in addition be developed at the center and they also will be produced there on a limited scale when not already available.

Consultation and coordination Upon request from the government concerned the center is prepared to send members of its staff to visit countries in a consultative capacity on matters related to the zoonoses and their control. In addition consultation services are provided through correspondence. Special attention is being given to the development in each country of the mechanism necessary for planning and operating control programs for the zoonoses. Such mechanisms based on the concept of united effort should provide for coordination and cooperation between public health and animal sanitation agencies with maximum participation of the community. Visits by consultants also provide the opportunity for international coordination of control and eradication programs.

Laboratory services These services include reference diagnosis, standardization of antigens, vaccines, the production of sera and other biological products, confirmatory testing of such products for potency and safety and the distribution of strains of viruses and bacteria for production or challenge purposes and of standard antigens, vaccines and sera for use in check testing.

The center will operate a model colony of laboratory animals that will serve for training, demonstration and study purposes as well as for supplying foundation stock for other institutions.

Demonstrations The center will utilize field campaigns against various zoonoses in pursuance of its functions. Trainees are taken to the site of the work and actually participate in zoonoses control programs. Existing anti-zoonoses programs of the city of Azul and of the host country (at both provincial and national levels) serve for such demonstration purposes. Demonstration work will also include the development of special control programs for various zoonoses in different parts of the Americas, the number, type and extent of such programs being dependent upon the needs, capabilities and desires of the country concerned.

Research Research at the center is of an applied nature, that is, the evaluation of a vaccine under certain field conditions, the practical application of a diagnostic procedure, the assessment of drugs for their antiparasitic effect and similar projects. Surveys and epidemiological, epizootiological studies will be undertaken with respect to certain of the zoonoses. Also included under research is the collection and processing of statistical data on the occurrence and overall importance of the diseases in man and animal.

The needs and opportunities for research in connection with the zoonoses are vast and the center itself can be expected to cover only a very small part of them. It is thus quite evident that public and private institutions in all countries must continue and intensify their research work in this field. It is a responsibility of the center to stimulate and coordinate such work.

Information Facilities are being developed so that the center may serve

Among the basic principles of epidemiology is the concept of reservoirs of infection. It is always desirable to have our problems appear in simple form so that understanding them easily we may achieve simple solutions. In the area of the zoonoses however the relationship of source and/or reservoir of infection is seldom direct and rarely simple. To paraphrase Cameron's words the complicated life cycles of parasites their facilitation of interchange between alternative hosts by close association their varying pathogenicity dependent upon recency of host acquisition the favorability of the host environment for multiplication the forms of the invading parasite their essential or alternate modes of transmission which in turn are dependent upon their ability to survive in transit or in contiguous species and the phylogenetic or physiologic kinship of the several hosts—all complicate animal human disease ecology and make our search for reservoirs of infection both frustrating and revealing. It is obviously this continuous challenge coupled with the necessities of human health and economy that leads to our explorations in the complex areas of the viral encephalitides rabies leptospiroses salmonellosis plague and typhus to name but a few.

If I were asked to select the most ecologically complicated entity from among these my task would not be simple. Whether it be the phenomenon of wintering over of encephalitis viruses the reasons for geographical differences in endemic and epidemic vectorial spread of these viruses the significance of bat rabies to human health the multiplicity of sources of salmonella or the reservoirs of plague the complexity of these problems is exceeded only by our ignorance of them. Deep inroads however have been made. Hess and Holden in their paper on The Natural History of the Arthropod Borne Encephalitides in the United States and Chamberlain in his contribution on Vector Relationships of the Arthropod Borne Encephalitides in North America have presented certain areas of agreement that help substantially to clarify some aspects of this complex ecologic problem. They are apparently agreed on birds as the most common vertebrate hosts on *Culex tarsalis* as both the endemic and epidemic vector of western equine encephalitis (WEE) and St. Louis encephalitis (SLE) in the western United States on separate endemic and epidemic vectors for WEE SLE and eastern equine encephalitis (EEE) in the eastern part and on *Culex pipiens* as the vector of SLE in the east and of urban SLE in the west. In addition Hess and Holden have presented evidence that both birds and hibernating mosquitoes are potential overwintering mechanisms for WEE and SLE viruses. Speculating on the relative infrequency of encephalitis epidemics in the east these investigators have postulated that natural environmental factors must concurrently provide optimum conditions both for vector breeding and longevity and that irrigation practices in the west provide these optimum conditions. Kissing in his paper on Host Relationship of the Arthropod Borne Encephalitides has helped us to visualize the determinants of geographical spread among reservoir hosts by categorizing the encephalitis viruses of group A according to their abilities to propagate in certain species with WEE propagating in birds Venezuelan equine encephalitis (VEE) in mammals and EEE intermediate with birds however affording the best viremia levels for transmission.

Summary

By Leonard M. Schuman

School of Public Health University of Minnesota Minneapolis Minn

The role of summarizer of a monograph as diversified in subject matter as this is a difficult one especially when the entire tone and orientation of the work have been so ably keynoted by the introductory presentation of J. D. Martin. A summarizer, furthermore, is constantly subject to the risk of routine repetition and the pronouncement of impressions already obvious to the reader. Despite the hazards of such a role, it offers the humble privilege of pointing out the highlights of the monograph, attempting to assess its impact and interjecting bits of philosophy and speculation while distilling the essential truths from the matrix of contributions.

Although the admirable presentations of each of the contributors guarantee the value of this publication, a significant share of the credit must go to the conference chairman and the members of his organizing committee for adroitly bringing together the talents of the several disciplines—laboratory, epidemiological, clinical, and administrative—of human and veterinary medicine. From this observer's point of view, the subject matter is happily balanced for topics of practical and continuing importance on the one hand and provocative newer knowledge at the frontiers of the life sciences on the other.

The very title of this publication conjures up an image of human ecology, serving to efface the customary categorical but nonetheless artificial distinctions between human and animal disease. It immediately denies the treatment of human and animal diseases as isolates and serves to set the pace for an approach to the complex interdependence of all living matter in a common environment. From the papers by Hsiung and Melnick on Orphan Viruses of Man and Animals and by M. Klein on The Significance of Human Antiviral Neutralizing Substances in Animal Sera to the philosophically stimulating presentation by Cameron on Parasites of Animals and Human Disease, we are even more fully impressed with the underlying interrelationships of basic living matter in the form of etiological agents of disease despite the emerging host specificity of the parasites. The contribution by Hsiung and Melnick not only reiterates these relationships but perforce broadens our concern with respect to the potentialities of nonhuman sources of orphan viruses. Now that these original orphans have found a haven of security in the form of human disease, their importance takes on increasing significance. Klein's contribution is of a similar character philosophically, for the provoking presence of human antiviral neutralizing substances in the sera of animals in which the related disease states apparently do not exist and from which the specific antigens have not been isolated leads to wholesome speculation. Whether ultimately a common antigen is discovered to explain these antibodies or antibody-like substances or a mechanism of formation is elucidated to revolutionize our concepts of antigen-antibody relationships, the pursuit of this investigation bears great promise in furthering our as yet inadequate understanding of animal-human disease ecology.

Among the basic principles of epidemiology is the concept of reservoirs of infection. It is always desirable to have our problems appear in simple form so that understanding them easily we may achieve simple solutions. In the area of the zoonoses however the relationship of source and/or reservoir of infection is seldom direct and rarely simple. To paraphrase Cameron's words the complicated life-cycles of parasites their facilitation of interchange between alternative hosts by close association their varying pathogenicity dependent upon recency of host acquisition the favorability of the host environment for multiplication the forms of the invading parasite their essential or alternate modes of transmission which in turn are dependent upon their ability to survive in transit or in contiguous species and the phylogenetic or physiologic kinship of the several hosts—all complicate animal human disease ecology and make our search for reservoirs of infection both frustrating and revealing. It is obviously this continuous challenge coupled with the necessities of human health and economy that leads to our explorations in the complex areas of the viral encephalitides rabies leptospiroses salmonellosis plague and typhus to name but a few.

If I were asked to select the most ecologically complicated entity from among these my task would not be simple. Whether it be the phenomenon of wintering over of encephalitis viruses the reasons for geographical differences in endemic and epidemic vectorial spread of these viruses the significance of bat rabies to human health the multiplicity of sources of salmonella or the reservoirs of plague the complexity of these problems is exceeded only by our ignorance of them. Deep inroads however have been made. Hess and Holden in their paper on The Natural History of the Arthropod Borne Encephalitides in the United States and Chamberlain in his contribution on Vector Relationships of the Arthropod Borne Encephalitides in North America have presented certain areas of agreement that help substantially to clarify some aspects of this complex ecologic problem. They are apparently agreed on birds as the most common vertebrate hosts on *Culex tarsalis* as both the endemic and epidemic vector of western equine encephalitis (WEE) and St. Louis encephalitis (SLE) in the western United States on separate endemic and epidemic vectors for WEE SLE and eastern equine encephalitis (EEE) in the eastern part and on *Culex pipiens* as the vector of SLE in the east and of urban SLE in the west. In addition Hess and Holden have presented evidence that both birds and hibernating mosquitoes are potential overwintering mechanisms for WEE and SLE viruses. Speculating on the relative infrequency of encephalitis epidemics in the east these investigators have postulated that natural environmental factors must concurrently provide optimum conditions both for vector breeding and longevity and that irrigation practices in the west provide these optimum conditions. Kissling in his paper on Host Relationship of the Arthropod Borne Encephalitides has helped us to visualize the determinants of geographical spread among reservoir hosts by categorizing the encephalitis viruses of group A according to their abilities to propagate in certain species with WEE propagating in birds Venezuelan equine encephalitis (VEE) in mammals and EEE intermediate with birds however affording the best viremia levels for transmission.

Summary

By Leonard M. Schuman

School of Public Health University of Minnesota Minneapolis Minn

The role of summarizer of a monograph as diversified in subject matter as this is a difficult one especially when the entire tone and orientation of the work have been so ably keynoted by the introductory presentation of J. D. Martin. A summarizer furthermore is constantly subject to the risk of routine repetition and the pronouncement of impressions already obvious to the reader. Despite the hazards of such a role it offers the humble privilege of pointing out the highlights of the monograph attempting to assess its impact and interjecting bits of philosophy and speculation while distilling the essential truths from the matrix of contributions.

Although the admirable presentations of each of the contributors guarantee the value of this publication a significant share of the credit must go to the conference chairman and the members of his organizing committee for adroitly bringing together the talents of the several disciplines—laboratory, epidemiological, clinical and administrative—of human and veterinary medicine. From this observer's point of view, the subject matter is happily balanced for topics of practical and continuing importance on the one hand and provocative newer knowledge at the frontiers of the life sciences on the other.

The very title of this publication conjures up an image of human ecology serving to efface the customary categorical but nonetheless artificial distinctions between human and animal disease. It immediately denies the treatment of human and animal diseases as isolates and serves to set the pace for an approach to the complex interdependence of all living matter in a common environment. From the papers by Hsiung and Melnick on Orphan Viruses of Man and Animals and by M. Klein on The Significance of Human Antiviral Neutralizing Substances in Animal Sera to the philosophically stimulating presentation by Cameron on Parasites of Animals and Human Disease we are even more fully impressed with the underlying interrelationships of basic living matter in the form of etiological agents of disease despite the emerging host specificity of the parasites. The contribution by Hsiung and Melnick not only reiterates these relationships but perforce broadens our concern with respect to the potentialities of nonhuman sources of orphan viruses. Now that these original orphans have found a haven of security in the form of human disease their importance takes on increasing significance. Klein's contribution is of a similar character philosophically for the provoking presence of human antiviral neutralizing substances in the sera of animals in which the related disease states apparently do not exist and from which the specific antigens have not been isolated leads to wholesome speculation. Whether ultimately a common antigen is discovered to explain these antibodies or antibody like substances or a mechanism of formation is elucidated to revolutionize our concepts of antigen antibody relationships the pursuit of this investigation bears great promise in furthering our as yet inadequate understanding of animal human disease ecology.

Among the basic principles of epidemiology is the concept of reservoirs of infection. It is always desirable to have our problems appear in simple form so that understanding them easily we may achieve simple solutions. In the area of the zoonoses however the relationship of source and/or reservoir of infection is seldom direct and rarely simple. To paraphrase Cameron's words the complicated life-cycles of parasites their facilitation of interchange between alternative hosts by close association their varying pathogenicity dependent upon recency of host acquisition the favorability of the host environment for multiplication the forms of the invading parasite their essential or alternate modes of transmission which in turn are dependent upon their ability to survive in transit or in contiguous species and the phylogenetic or physiologic kinship of the several hosts—all complicate animal human disease ecology and make our search for reservoirs of infection both frustrating and revealing. It is obviously this continuous challenge coupled with the necessities of human health and economy that leads to our explorations in the complex areas of the viral encephalitides rabies leptospiroses salmonelloses plague and typhus to name but a few.

If I were asked to select the most ecologically complicated entity from among these my task would not be simple. Whether it be the phenomenon of wintering over of encephalitis viruses the reasons for geographical differences in endemic and epidemic vectorial spread of these viruses the significance of bat rabies to human health the multiplicity of sources of salmonella or the reservoirs of plague the complexity of these problems is exceeded only by our ignorance of them. Deep inroads however have been made. Hess and Holden in their paper on The Natural History of the Arthropod Borne Encephalitides in the United States and Chamberlain in his contribution on Vector Relationships of the Arthropod Borne Encephalitides in North America have presented certain areas of agreement that help substantially to clarify some aspects of this complex ecologic problem. They are apparently agreed on birds as the most common vertebrate hosts on *Culex tarsalis* as both the endemic and epidemic vector of western equine encephalitis (WEE) and St. Louis encephalitis (SLE) in the western United States on separate endemic and epidemic vectors for WEE SLE and eastern equine encephalitis (EEE) in the eastern part and on *Culex pipiens* as the vector of SLE in the east and of urban SLE in the west. In addition Hess and Holden have presented evidence that both birds and hibernating mosquitoes are potential overwintering mechanisms for WEE and SLE viruses. Speculating on the relative infrequency of encephalitis epidemics in the east these investigators have postulated that natural environmental factors must concurrently provide optimum conditions both for vector breeding and longevity and that irrigation practices in the west provide these optimum conditions. Kissling in his paper on Host Relationship of the Arthropod Borne Encephalitides has helped us to visualize the determinants of geographical spread among reservoir hosts by categorizing the encephalitis viruses of group A according to their abilities to propagate in certain species with WEE propagating in birds Venezuelan equine encephalitis (VEE) in mammals and EEF intermediate with birds however affording the best viremia levels for transmission.

Although neither as complex in its host parasite relationship nor as productive of human epidemics as the viral encephalitides the leptospiroses represent a group of etiologically related diseases with a multiplicity not only of reservoirs and sources but of clinical syndromes in animals and man. Stockard and Woodward's contributions to our clinical understanding in their paper 'Leptospirosis Infections in Man' make it obvious that despite their relatively infrequent occurrence in man in outbreak form the leptospiroses have so broad a spectrum of clinical features (ranging from short term pyrexias in influenza like illnesses through central nervous system infections simulating nonparalytic polio to fulminating disease resembling fatal cases of yellow fever) that coupled with the ubiquity of animal reservoirs these etiologically definitive entities must constantly be considered in the differential diagnosis of human disease.

The dissertation on Animal Reservoirs of Leptospirae by Babudieri reminds us that efficient carrier states are produced only when a satisfactory equilibrium between the leptospire and the host has been established. He notes that while domestic animals and rodents are particularly important reservoirs of infection for man, pathogenic leptospires recently have been isolated from migratory birds. Babudieri re-emphasizes the need for concurrent conjoint and integrated epizootiological and epidemiological investigations. In their paper on Epidemiological Patterns of Leptospirosis Galton Menges and Steele reiterated the universality of these infections among several animal species their pathogenicity for man and the role of the animal carrier. Their presentation of epidemiological patterns in the United States and the human epidemic potential inherent in existing serotypes as well as the discovery of new sylvatic hosts serve to augment this observer's appeal for a concerted multidisciplinary approach to the problem.

Among the earliest recognized zoonoses rabies has continued to constitute a serious threat to human populations. As such and because of recent developments both in the epidemiology and prophylaxis of the disease the prominence given it in this monograph is justified. In reviewing Recent Developments in the Epidemiology of Rabies Tierkel has pointed to the decline in canine rabies as evidence of the success of control measures applied but he views with concern the increase in the disease among sylvatic vectors. Although it is difficult to separate reports based on field investigations and research or stimulated by animal morbidity reporting systems from real increases in incidence the growing awareness of ever unfolding wildlife reservoirs is salutary to the definition of the problem despite its growing complexity. Tierkel thus correctly presented the need for expanding investigational activities in the epidemiology of the disease in wildlife.

Of provocative interest was the paper Bat Rabies Experimental Host Transmission Studies by Burns. This investigator's experiments on animal transmission of bat rabies virus by induced association with resultant inability to demonstrate transmission from experimentally and naturally infected bats to monkeys guinea pigs and white mice as well as the lack of evidence of clinical recovery or of the development of efficient carrier states among inoculated insectivorous bats will provoke I am sure a flurry of laboratory and field

investigations of this potential reservoir of infection. Burns admonition to treat bat bites as potential exposures to rabies until more extensive studies are available is well made.

Salmonellosis is rarely discussed in works on animal diseases related to human health. Consequently interest in this disease appears to be confined to the general public which often fails to avail itself of the care of the practicing physician who in turn rarely undertakes the etiological definition of diarrhea when patients do come for treatment and of the health officers who all too often make ineffectual gestures toward control measures. It is therefore noteworthy that no less an expert than Edwards has presented his paper *Salmonellosis: Observations on Incidence and Control* in this volume. Presenting available information on cases and outbreaks in man, this investigator very properly calls attention to the gross inadequacy of these data as indicators of true incidence and points to the similarity of serotypes in human and animal populations. He reiterates the frequently indigenous character of salmonellae in animal foods and appeals to medical and paramedical personnel involved in human and animal health to institute effective measures for the control of infections in animals.

Despite the rarity of even sporadic cases of plague in the United States today the existence of heavily infected reservoirs among sylvatic rodents denotes our failure thus far to control sylvatic plague and to remove this potential threat to human populations. As worthy a challenge to ecologic investigation as any disease sylvatic plague embodies complex relationships among the reservoir host, the ectoparasitic vector and the bacterial parasite relationships that have been studied very inadequately. Kartman and his co-workers in their paper on *New Knowledge on the Ecology of Sylvatic Plague* indicate studies in progress that will remedy these inadequacies in large measure. Of highly significant import is their finding that a replacement of emphasis on specific reservoirs must be made. Whereas prior to 1950 ground squirrels and prairie dogs received prime emphasis, recent years have revealed that field mice, principally of the species *Microtus*, are the important reservoir. The high infection rates of *Microtus* fleas, their prevalence in the field and their high efficiency as vectors make the species *Microtus* a primary source of plague transmission. Kartman emphasized the significance of this threat to human populations by pointing to the suburban habitat of *Microtus* and its coexistence with rats in dumps and parks and around homes where flea transfer to domestic rodents can occur.

Murine typhus has also presented problems in ecology although they are not nearly as complex as those of plague. Pratt in *The Changing Picture of Murine Typhus in the United States* reviewed the reported incidence of the disease in two four year periods, noting an increase from 1941 to 1944 and a marked decrease from 1953 to 1956. With all due respect for the sanitation programs that undoubtedly contributed to the control of rodents and their ectoparasites, we cannot overlook the possibility of natural downward trends in rat and flea populations or the acceleration of the decline in the disease by more exact diagnosis with laboratory confirmation stimulated by the murine typhus surveillance program of the Communicable Disease Center, Atlanta, Ga.

Although neither as complex in its host parasite relationship nor as productive of human epidemics as the viral encephalitides the leptospiroses represent a group of etiologically related diseases with a multiplicity not only of reservoirs and sources but of clinical syndromes in animals and man. Stockard and Woodward's contributions to our clinical understanding in their paper *Leptospirosis Infections in Man* make it obvious that despite their relatively infrequent occurrence in man in outbreak form the leptospiroses have so broad a spectrum of clinical features (ranging from short term pyrexias in influenza like illnesses through central nervous system infections simulating nonparalytic polio to fulminating disease resembling fatal cases of yellow fever) that coupled with the ubiquity of animal reservoirs these etiologically definitive entities must constantly be considered in the differential diagnosis of human disease.

The dissertation on 'Animal Reservoirs of Leptospire' by Babudieri reminds us that efficient carrier states are produced only when a satisfactory equilibrium between the leptospire and the host has been established. He notes that while domestic animals and rodents are particularly important reservoirs of infection for man pathogenic leptospires recently have been isolated from migratory birds. Babudieri re-emphasizes the need for concurrent conjoint and integrated epizootiological and epidemiological investigations. In their paper on 'Epidemiological Patterns of Leptospirosis' Calton Menges and Steele reiterated the universality of these infections among several animal species their pathogenicity for man and the role of the animal carrier. Their presentation of epidemiological patterns in the United States and the human epidemic potential inherent in existing serotypes as well as the discovery of new sylvatic hosts serve to augment this observer's appeal for a concerted multidisciplinary approach to the problem.

Among the earliest recognized zoonoses rabies has continued to constitute a serious threat to human populations. As such and because of recent developments both in the epidemiology and prophylaxis of the disease the prominence given it in this monograph is justified. In reviewing 'Recent Developments in the Epidemiology of Rabies' Tierkel has pointed to the decline in canine rabies as evidence of the success of control measures applied but he views with concern the increase in the disease among sylvatic vectors. Although it is difficult to separate reports based on field investigations and research or stimulated by animal morbidity reporting systems from real increases in incidence the growing awareness of ever unfolding wildlife reservoirs is salutary to the definition of the problem despite its growing complexity. Tierkel thus correctly presented the need for expanding investigational activities in the epidemiology of the disease in wildlife.

Of provocative interest was the paper 'Bat Rabies: Experimental Host Transmission Studies' by Burns. This investigator's experiments on animal transmission of bat rabies virus by induced association with resultant inability to demonstrate transmission from experimentally and naturally infected bats to monkeys guinea pigs and white mice as well as the lack of evidence of clinical recovery or of the development of efficient carrier states among inoculated insectivorous bats will provoke I am sure a flurry of laboratory and field

Traum present their paper on *The Exotic Zoonoses* cautiously indicating that diseases of man presently *not known to infect animals and diseases of animals presently not known to infect man* may be shown as explorations continue to do otherwise. A point made by these investigators that different species may demonstrate different manifestations of the same infection is well taken.

In this same respect Koprowski who was given a most difficult task in finding Counterparts of Human Viral Disease in Animals stresses the need for expanded research in the area of comparative pathology and epidemiology warning us however to avoid equating the unknown with the unknown and overlooking species differences in clinical response. Certainly his curious finding of poliovirus Type I in calf feces should be explored thoroughly.

Thus far I have dealt with the epidemiological considerations of animal hosts, reservoirs and agents of disease discussed or implied in the several presentations. However animal diseases that do not necessarily occur in man may have equal if subtle importance in human ecology. I refer now to a group of papers dealing with the study of purely animal diseases for their value as research tools in eliciting pathogenetic truths common to human disease. Pollard's *Comparative Studies on Viral Hepatitis in Animals and in Man* deals with the use of mouse hepatitis virus as a possible prototype of the human disease in lieu of work with the human virus which is not transmissible to other animals and which thus far has not been grown in tissue culture. This work is illustrative of the methodology necessary to circumvent obstacles in the study of human disease.

It is gratifying that this monograph does not confine itself entirely to *acute* communicable disease but seeks to explore the comparative pathology of certain chronic diseases or chronic sequelae to infectious disease in animals that show histological configurations similar to those of important diseases in man. The contributions of Sikes on *A Comparison of Rheumatoid Like Arthritis in Swine and Rheumatoid Arthritis in Man* and of Duran Reynals and Rafferty on *The Pulmonary Adenomatosis Complex in Sheep* are cases in point. An approach via comparative pathology might not reveal identical underlying etiologies for the counterparts of human disease but it should yield information of great value in the study of neoplastic and other chronic diseases of man.

I should not overlook the area of diagnostic tests in the zoonoses. Gochenour and his co-workers in reviewing the *Laboratory Diagnosis of Leptospirosis* presented the merits and disadvantages of the several tests employed. In their hands the killed antigen agglutination test gave results identical to those with live antigen. In discussing more recently developed tests these investigators noted excellent correlation of the capillary tube test of Stoenner with live antigen. More nearly genus-specific antigens such as the sonic vibrated complement fixation (CF) tests of Randall the erythrocytic sensitizing test of Chang and the hemolytic test of Cox requiring but a limited number of antigens yielded only fair correlation with agglutination tests. Gochenour and his collaborators expressed the need for a more rapid means of recovering leptospire and for a simple inexpensive sensitive genus-specific test for local laboratories. It is gratifying however that work in this direction is proceeding.

The paper on 'Recent Developments in Animal Ringworm and Their Public Health Implications' by William Kaplan and his co workers was in this reviewer's opinion very welcome. The epidemiology of mycotic infections is an all too neglected field and there is great need for inquiry and important contributions in this area. Kaplan's proposal for a functional classification of dermatophytes based on epidemiological relationships and comprising three groups—anthropophilic, zoophilic, and geophilic—appears to be a practical one.

If the public health importance of a disease were dependent upon the completeness of our understanding of its epidemiology, then a number of diseases presently engaging our intense attention would be relegated to the unimportant. The lack of attention to a number of diseases that are common to animals and man is frustrating at times. All too frequently the lack of awareness of a disease determines that it shall fail of recognition and intensive study. In 'Listeriosis: A Potential Public Health Problem' Winn and his co workers assert this for a disease whose ubiquitous agent is well known, its pathology in animals well documented in veterinary medical science, but whose epidemiology is most inadequately understood. To me this represents primarily, a failure to apply an interdisciplinary approach. It is my hope that Winn's presentation will stimulate our thinking in this direction.

It is a rare disease indeed for which the epidemiology is relatively completely known. In man the several modes of disease transmission have not been satisfactorily worked out and their natural histories are poorly understood. The value of the contribution of Welsh and his co workers in their extremely fine paper on the 'Air Borne Transmission of Q Fever: The Role of Parturition in the Generation of Infective Aerosols' lies not only in the excellent review material presented but in the original work described which adds to our understanding of the modes of transmission of the disease in the area of air borne infection. The experimental work of these investigators on the production of aerosols during the parturition of inoculated sheep and the persistence of these aerosols in the environment is classic in form and once again it emphasizes the contribution that animal disease study can make to human epidemiology.

In dealing with animal reservoirs of human disease we customarily expect disease counterparts in the animal host or at least serologic evidence of infection to denote a host-parasite state of equilibrium. Cat scratch fever according to Prier is geographically widespread, quite common, presents a typical clinical picture and is expected to increase in incidence because of man's firm association with these domestic pets. Of special epidemiological interest is the fact that no definite clinical disease in the animal host has been recognized.

Although Hanson and Brandy in their discussion of Newcastle Disease minimize its role as an occupational hazard among poultry producers and handlers and speak of its indirect impact on human economy principally as an avian disease, to the epidemiologist the rare occurrence of a disease is equal in importance to a wide scale outbreak in the attempt to arrive at a more complete understanding of the agent-host-environment relationship. Not long ago Newcastle disease could have been included in the list of animal diseases that are known not to infect man. It is in this philosophical vein that Shahan and

Traum present their paper on *The Exotic Zoonoses* cautiously indicating that diseases of man presently not known to infect animals and diseases of animals presently not known to infect man may be shown as explorations continue to do otherwise. A point made by these investigators that different species may demonstrate different manifestations of the same infection is well taken.

In this same respect Koprowski who was given a most difficult task in finding Counterparts of Human Viral Disease in Animals stresses the need for expanded research in the area of comparative pathology and epidemiology warning us however to avoid equating the unknown with the unknown and overlooking species differences in clinical response. Certainly his curious finding of poliovirus Type I in calf feces should be explored thoroughly.

Thus far I have dealt with the epidemiological considerations of animal hosts reservoirs and agents of disease discussed or implied in the several presentations. However animal diseases that do not necessarily occur in man may have equal if subtle importance in human ecology. I refer now to a group of papers dealing with the study of purely animal diseases for their value as research tools in eliciting pathogenetic truths common to human disease. Pollard's *Comparative Studies on Viral Hepatitis in Animals and in Man* deals with the use of mouse hepatitis virus as a possible prototype of the human disease in lieu of work with the human virus which is not transmissible to other animals and which thus far has not been grown in tissue culture. This work is illustrative of the methodology necessary to circumvent obstacles in the study of human disease.

It is gratifying that this monograph does not confine itself entirely to *acute* communicable disease but seeks to explore the comparative pathology of certain chronic diseases or chronic sequelae to infectious disease in animals that show histological configurations similar to those of important diseases in man. The contributions of Sikcs on *A Comparison of Rheumatoid Like Arthritis in Swine and Rheumatoid Arthritis in Man* and of Duran Reynals and Rafferty on *The Pulmonary Adenomatosis Complex in Sheep* are cases in point. An approach via comparative pathology might not reveal identical underlying etiologies for the counterparts of human disease but it should yield information of great value in the study of neoplastic and other chronic diseases of man.

I should not overlook the area of diagnostic tests in the zoonoses. Gochenour and his co-workers in reviewing the *Laboratory Diagnosis of Leptospirosis* presented the merits and disadvantages of the several tests employed. In their hands the killed antigen agglutination test gave results identical to those with live antigen. In discussing more recently developed tests these investigators noted excellent correlation of the capillary tube test of Stoenner with live antigen. More nearly genus specific antigens such as the sonic vibrated complement fixation (CF) tests of Randall the erythrocytic sensitizing test of Chang and the hemolytic test of Cox requiring but a limited number of antigens yielded only fair correlation with agglutination tests. Gochenour and his collaborators expressed the need for a more rapid means of recovering leptospire and for a simple inexpensive sensitive genus-specific test for local laboratories. It is gratifying however that work in this direction is proceeding.

The paper on 'Recent Developments in Animal Ringworm and Their Public Health Implications' by William Kaplan and his co-workers was in this reviewer's opinion very welcome. The epidemiology of mycotic infections is an all too neglected field, and there is great need for inquiry and important contributions in this area. Kaplan's proposal for a functional classification of dermatophytes based on epidemiological relationships and comprising three groups—anthropophilic, zoophilic, and geophilic—appears to be a practical one.

If the public health importance of a disease were dependent upon the completeness of our understanding of its epidemiology, then a number of diseases presently engaging our intense attention would be relegated to the unimportant. The lack of attention to a number of diseases that are common to animals and man is frustrating at times. All too frequently the lack of awareness of a disease determines that it shall fail of recognition and intensive study. In 'Listeriosis: A Potential Public Health Problem' Winn and his co-workers assert this for a disease whose ubiquitous agent is well known, its pathology in animals well documented in veterinary medical science, but whose epidemiology is most inadequately understood. To me this represents primarily a failure to apply an interdisciplinary approach. It is my hope that Winn's presentation will stimulate our thinking in this direction.

It is a rare disease indeed for which the epidemiology is relatively completely known. In man the several modes of disease transmission have not been satisfactorily worked out and their natural histories are poorly understood. The value of the contribution of Welsh and his co-workers in their extremely fine paper on the 'Air Borne Transmission of Q Fever: The Role of Parturition in the Generation of Infective Aerosols' lies not only in the excellent review material presented but in the original work described which adds to our understanding of the modes of transmission of the disease in the area of air borne infection. The experimental work of these investigators on the production of aerosols during the parturition of inoculated sheep and the persistence of these aerosols in the environment is classic in form and once again it emphasizes the contribution that animal disease study can make to human epidemiology.

In dealing with animal reservoirs of human disease we customarily expect disease counterparts in the animal host or at least serologic evidence of infection to denote a host-parasite state of equilibrium. Cat scratch fever according to Prier is geographically widespread, quite common, presents a typical clinical picture, and is expected to increase in incidence because of man's firm association with these domestic pets. Of special epidemiological interest is the fact that no definite clinical disease in the animal host has been recognized.

Although Hanson and Brandly in their discussion of Newcastle Disease minimize its role as an occupational hazard among poultry producers and handlers and speak of its indirect impact on human economy principally as an avian disease, to the epidemiologist the rare occurrence of a disease is equal in importance to a wide scale outbreak in the attempt to arrive at a more complete understanding of the agent-host-environment relationship. Not long ago Newcastle disease could have been included in the list of animal diseases that are known not to infect man. It is in this philosophical vein that Shahan and

symptom of our failure to achieve a safe and efficient vaccine for humans. Serving not as a substitute but in a sense as an adjuvant to vaccine, hyperimmune serum augments our treatment but fails to obviate the hazard of reactions. It was inevitable then that chick embryo adapted living virus vaccines should be applied to volunteer human groups.

Although at this moment field experiments on seroimmunity reveal phenolized vaccines to be more efficient, the demonstrations by Fox in collaboration with the World Health Organization (WHO) of the practicality of pre bite long term immunization by primary and booster vaccinations with high egg passage Flury vaccines and the rapid antibody response in Pasteur treatment conditioned individuals to boosters of the avianized vaccine hold great promise for human prophylaxis. Certainly their implications for persons at high and relatively constant risk is quite obvious when we consider the reaction potential in repeated immunizations effected by Pasteur techniques.

Brucellosis which has suffered an unwarranted loss of emphasis in many conferences on the zoonoses is not neglected in this monograph. Despite the optimism expressed by Mingle in his paper on Brucellosis in Livestock Control and Eradication with respect to bovine disease and despite his prediction that there will be less than 1 per cent infected cattle in the United States by 1960, a note of pessimism was injected by McCullough who although he noted the great strides made in control of bovine brucellosis stressed the lack of attention paid to swine brucellosis and the virtual lack of efforts toward controlling it. When one considers the fact that today this disease is found predominantly among packing house workers and others who handle animals, one must be concerned over the possibility that a significant level of human brucellosis will be maintained far short of the irreducible minimum (not to mention complete eradication) even after bovine brucellosis is virtually eradicated.

The complexity that animal-human disease relationships may assume and the gross alterations in epidemiology and control applications that may be induced by economic factors is well illustrated by anthrax. In their paper, Industrial Anthrax, Brachman and Fekety have quite competently demonstrated the shifting sources of infection through improvement of animal health and have described the impact of war which changes the source of contaminated raw materials and technological advances such as the development of plastics and synthetic fibers that completely remove certain animal materials from the list of hazards. The bulk of anthrax today is industrial rather than agricultural and in industry goat hair processing presently constitutes the principal source. Here is an admirable example of the need for constant epidemiological appraisal of a problem so that we may remain poised for attack from any quarter. Of additional significance in Brachman and Fekety's contribution is their interesting approach to the question of achieving human immunity by prolonged exposure. The data of these investigators suggest that such immunity does not develop.

Bovine tuberculosis has received little attention in more recent years chiefly because of the tremendous success of its control program. In his contribution on The Public Health Importance of Animal Tuberculosis, Anderson has cited the gains made by the now historic program of tuberculin testing of herds.

The detection of ornithosis in domestic fowl is grossly inadequate at best. Benedict, exploring the indirect CF test and modifications of the direct CI test along with an intradermal test, found 10 to 85 per cent agreement between the allergic and serologic tests and proposed the application of the intradermal test as the most practicable for ornithosis surveys. It is hoped that field tests of the method will provide us with the necessary tool for augmenting control of this economically devastating and at times human epidemic disease.

The ultimate goal of laboratory workers, clinicians, and epidemiologists is the control of disease. This monograph includes reports of satisfying achievements in several disease areas. Sussman, Cohen, and their co-workers applied EEE vaccine to flocks of pheasants in New Jersey during the 1956 epizootic with surprisingly good results. Of significance is the fact that appreciably less than 100 per cent vaccination of a flock during an outbreak will provide reasonably high protection for the flock as a whole, with evidence that nonvaccinates in a partially treated flock also enjoy relative protection. Applied prior to an outbreak, partial treatment of a flock gave even higher protection rates. To an epidemiologist this work represents more than the achievement of a practical tool to control an animal disease—that is both of economic importance and a potential hazard to man. In it I can see a striking example of experimental epidemiology in the field, a sadly neglected area of inquiry since the pioneering work of Greenwood. Furthermore, the study on partial flock immunization represents a direct counterpart of human population immunization phenomena, with the significant exception that the latter have involved conjecture more often than analytic observation. As such, the pheasant study represents an epidemiological model with aspects that may throw light upon human epidemic dynamics.

In turkey ornithosis, Delaplane succeeded in establishing prophylactic levels of chlortetracycline with apparent total suppression of virus when dosage of more than 200 gm. per ton of feed for two weeks or 200 gm. per ton for three weeks was employed. Furazolidone was found to be ineffective. Of collateral epidemiological interest was Delaplane's finding that virus was apparently not transmitted via the egg even during the viremic stage when infection potential is highest. The application of these observations may well augment our efforts at control despite our lack of an active immunizing agent. Its indirect significance in human disease prevention is quite obvious when we realize the epidemic potential of this disease reservoir.

In charting the development of prophylactic measures for the control of rabies in domestic animals, Kaerberle very properly stressed the fact that the continuous attempts to develop attenuated living virus vaccines were indicative of the inadequacy and disadvantages of vaccines of nervous tissue origin. The effectiveness of avianized low egg passage Flury strain vaccine has now been well demonstrated for dogs, and the application of high egg passage vaccines to cattle is a recent bright spot in our progress.

What has been said of rabies vaccines for domestic animals applies equally, if not more so, to human prophylaxis. Here a sword of Damocles has hung on a finer thread, and the formidable power of treatment reactions has plagued us since the days of Pasteur. The advent of hyperimmune serum was another

symptom of our failure to achieve a safe and efficient vaccine for humans. Serving not as a substitute but in a sense as an adjuvant to vaccine, hyperimmune serum augments our treatment but fails to obviate the hazard of reactions. It was inevitable then that chick embryo-adapted living virus vaccines should be applied to volunteer human groups.

Although at this moment field experiments on seroimmunity reveal phenolized vaccines to be most efficient, the demonstrations by Fox in collaboration with the World Health Organization (WHO) of the practicality of pre bite long term immunization by primary and booster vaccinations with high-egg passage Flury vaccines and the rapid antibody response in Pasteur treatment-conditioned individuals to boosters of the avianized vaccine hold great promise for human prophylaxis. Certainly their implications for persons at high and relatively constant risk is quite obvious when we consider the reaction potential in repeated immunizations effected by Pasteur techniques.

Brucellosis, which has suffered an unwarranted loss of emphasis in many conferences on the zoonoses, is not neglected in this monograph. Despite the optimism expressed by Mingle in his paper on "Brucellosis in Livestock: Control and Eradication" with respect to bovine disease and despite his prediction that there will be less than 1 per cent infected cattle in the United States by 1960, a note of pessimism was injected by McCullough who, although he noted the great strides made in control of bovine brucellosis, stressed the lack of attention paid to swine brucellosis and the virtual lack of efforts toward controlling it. When one considers the fact that today this disease is found predominantly among packing house workers and others who handle animals, one must be concerned over the possibility that a significant level of human brucellosis will be maintained far short of the irreducible minimum (not to mention complete eradication) even after bovine brucellosis is virtually eradicated.

The complexity that animal-human disease relationships may assume and the gross alterations in epidemiology and control applications that may be induced by economic factors is well illustrated by anthrax. In their paper "Industrial Anthrax," Brachman and Fekety have quite competently demonstrated the shifting sources of infection through improvement of animal health and have described the impact of war which changes the source of contaminated raw materials and technological advances such as the development of plastics and synthetic fibers that completely remove certain animal materials from the list of hazards. The bulk of anthrax today is industrial rather than agricultural and in industry goat hair processing presently constitutes the principal source. Here is an admirable example of the need for constant epidemiological appraisal of a problem so that we may remain poised for attack from any quarter. Of additional significance in Brachman and Fekety's contribution is their interesting approach to the question of achieving human immunity by prolonged exposure. The data of these investigators suggest that such immunity does not develop.

Bovine tuberculosis has received little attention in more recent years, chiefly because of the tremendous success of its control program. In his contribution on "The Public Health Importance of Animal Tuberculosis," Anderson has cited the gains made by the now historic program of tuberculin testing of herds.

slaughter of reactors and tracing of sources of infection. The relatively rare cases of bovine tuberculosis in humans speak well not only for the success of the control program but unfortunately also connote our failure to eradicate this disease completely. It must be remembered that Modified Tuberculosis Free Area Certification still implies a maximum of one half of 1 per cent individual animal infection. It is the history of human populations that complacency leads to apathy and apathy to disaster. Our attitude is unfortunately tied to our definition of the word 'problem'. To some this means only clinical disease either in man or in animal reservoirs; to me (and I hope to many others) it means a great deal more for it includes the potential of individual and epidemic hazard even after mass control has been achieved. It matters not whether it be diphtheria or bovine tuberculosis for it is the very essence of life that new generations without immunity, without protection and without knowledge shall be born. Thus eternal vigilance directed toward unrelenting maintenance of control becomes the second phase of every problem of disease control; bovine tuberculosis is no exception.

The interdependence of animals and man and their ecological relationships in disease can lead literally to rather inhuman applications. Hempy in his fine summary of 'Animal Disease and Biological Warfare' presented the general philosophy of the use of disease agents in biological warfare as a method of reducing a nation's military effectiveness. He stressed the fact that aerosols are readily deliverable by any country and that there are no secrets among nations relative to potential infectious agents of disease. Because of the attributes that such microorganisms must possess to be effective agents in warfare, the number available is actually limited. These attributes include high infectivity, simple dissemination, ease of production, inefficiency in the production of solid immunity, high lethality (or at least pathogenicity) and mutability toward resistance to chemotherapy. In addition to the direct aerosol application of suitable zoonotic microorganisms to human populations, the application of aerosols to crops and food animals could result in fatal reduction of food supplies. Hempy emphasized that our preventive techniques are an advantage in this area but they certainly provide no cause for complacency. More agents of effective active immunization, both for man and for animals, are needed. Hempy's precise delineation of the problem should serve as a stimulus in this vital field.

This monograph includes problems not only of national but of global importance. For the vast number of diseases of man, whether of human or animal origin, it has frequently been stated that as long as a disease exists anywhere in the world, it remains a constant threat to populations in more favorable positions. The underprivileged countries of the world, with their more numerous problems of animal disease of direct or indirect importance to human health and food availability, must be treated altruistically and with humanitarianism. These were the implications of Martin Kaplan's discussion of 'Global Priorities in Animal Disease Control'. The aggressive and well-conceived programs of animal-disease control of the World Health Organization (WHO) in collaboration with the Food and Agriculture Organization (FAO) and the Office International des Epizooties (OIE) should achieve not

only the objectives of animal and human health but should provide valuable field data contributory to even newer epidemiological knowledge and control techniques. In this regard it is quite admirable that the integration of the several disciplines involved in animal and human health has progressed to the point that animal sera will be collected and stored for future titrations to determine possible relationships between swine and human influenza as human epidemics rise and fall. In the same spirit of international cooperation and joint solution of problems of animal human disease is Blood's paper on The Pan American Zoonoses Center.

In conclusion at the risk of being redundant may I say that far and above the valuable technical contributions to our understanding of animal disease and human health there must have come to the reader as it has to me an appreciation of the vastness of the interrelationships between animal and human disease and of a compelling need for a concerted integrated multidisciplinary team approach to their problems which can no longer due to our growing awareness be separated into distinct categories of animal as opposed to human disease. No one discipline can provide the technical prowess necessary for the solution of problems of such complex ecology this has been made abundantly clear throughout this publication. Furthermore no one discipline can afford the luxury of neglecting to seek the aid of others. Animal and human health problems reside within the same continuum they cannot be removed and studied as isolates. Their mutually inclusive interdependence thrusts itself upon us daily. It is my firm conviction that if we took the energies that we expend in duplicating inquiries and studies and all too often in exalting one discipline while disparaging another and converted a portion of these energies to the organization of a horizontal partnership we should much more rapidly fill in the missing gaps in our knowledge of disease and more quickly solve the problems that face us. This I am sure is what J. D. Martin meant by the sharing of problems as well as knowledge. This then I hope to leave with you in summary.

slaughter of reactors and tracing of sources of infection. The relatively rare cases of bovine tuberculosis in humans speak well not only for the success of the control program but, unfortunately, also connote our failure to eradicate this disease completely. It must be remembered that Modified Tuberculosis Free Area Certification still implies a maximum of one half of 1 per cent individual animal infection. It is the history of human populations that complacency leads to apathy, and apathy to disaster. Our attitude is unfortunately tied to our definition of the word "problem." To some this means only clinical disease either in man or in animal reservoirs, to me (and I hope to many others) it means a great deal more, for it includes the potential of individual and epidemic hazard even after mass control has been achieved. It matters not whether it be diphtheria or bovine tuberculosis, for it is the very essence of life that new generations without immunity, without protection, and without knowledge shall be born. Thus eternal vigilance directed toward unrelenting maintenance of control becomes the second phase of every problem of disease control; bovine tuberculosis is no exception.

The interdependence of animals and man and their ecological relationships in disease can lead literally to rather inhuman applications. Hempy in his fine summary of "Animal Disease and Biological Warfare" presented the general philosophy of the use of disease agents in biological warfare as a method of reducing a nation's military effectiveness. He stressed the fact that aerosols are readily deliverable by any country and that there are no secrets among nations relative to potential infectious agents of disease. Because of the attributes that such microorganisms must possess to be effective agents in warfare, the number available is actually limited. These attributes include high infectivity, simple dissemination, ease of production, inefficiency in the production of solid immunity, high lethality (or at least pathogenicity), and mutability toward resistance to chemotherapy. In addition to the direct aerosol application of suitable zoonotic microorganisms to human populations, the application of aerosols to crops and food animals could result in fatal reduction of food supplies. Hempy emphasized that our preventive techniques are an advantage in this area, but they certainly provide no cause for complacency. More agents of effective active immunization, both for man and for animals, are needed. Hempy's precise delineation of the problem should serve as a stimulus in this vital field.

This monograph includes problems not only of national but of global importance. For the vast number of diseases of man, whether of human or animal origin, it has frequently been stated that as long as a disease exists anywhere in the world, it remains a constant threat to populations in more favorable positions. The underprivileged countries of the world, with their more numerous problems of animal disease of direct or indirect importance to human health and food availability, must be treated altruistically and with humanitarianism. These were the implications of Martin Kaplan's discussion of "Global Priorities in Animal Disease Control." The aggressive and well-conceived programs of animal-disease control of the World Health Organization (WHO) in collaboration with the Food and Agriculture Organization (FAO) and the Office International des Epizooties (OIE) should achieve not

